

# Molecular data for *Crenavolva* species (Gastropoda, Ovulidae) reveals the synonymy of *C. chiapponii*

Bastian T. Reijnen<sup>1</sup>

<sup>1</sup> Department of Marine Zoology, Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands

Corresponding author: Bastian T. Reijnen ([Bastian.Reijnen@naturalis.nl](mailto:Bastian.Reijnen@naturalis.nl))

---

Academic editor: T. Backeljau | Received 16 December 2014 | Accepted 13 April 2015 | Published 30 April 2015

<http://zoobank.org/4C35416C-73F6-4E97-8394-44D5115FFFDD>

---

**Citation:** Reijnen BT (2015) Molecular data for *Crenavolva* species (Gastropoda, Ovulidae) reveals the synonymy of *C. chiapponii*. ZooKeys 501: 15–26. doi: 10.3897/zookeys.501.9144

---

## Abstract

During fieldwork in Indonesia and Malaysia, eight lots containing 33 specimens belonging to the genus *Crenavolva* (Ovulidae) were collected. Species were initially identified as *C. aureola*, *C. chiapponii*, *C. striatula* and *C. trailli*, respectively. For *C. chiapponii* this is the second record. In contrast to the ecological data available from the original description of this species, it was found in shallow water on a gorgonian host coral, i.e. *Acanthogorgia* sp. A molecular analysis based on COI and 16S mtDNA markers, including sequence data obtained from GenBank, showed that *C. chiapponii* should be considered a junior synonym of *C. aureola* and that previously identified ovulid specimens are probably misidentified.

## Keywords

*Acanthogorgia*, host association, molecular phylogeny, Octocorallia, 16S, COI

## Introduction

The nominal taxon *Crenavolva* was introduced as a subgenus by Cate (1973), together with the subgenera *Crenavolva*, *Cuspivolva* and *Serratovolva*. In the most recent overview regarding Ovulidae these three taxa are considered genera (Lorenz and Fehse 2009). At present 18 nominal species are recognized within *Crenavolva* (Rosenberg 2014), most of which are considered rare (Lorenz and Fehse 2009). These species are

considered rare because few specimens have been collected, probably because they occur at depths greater than standard recreational diving depth of c. 30 m and/or are only known from a limited geographical area, usually just the type locality. This also accounts for *C. chiapponii* Lorenz & Fehse, 2009, which is only known from Balicasag Isl., Bohol, Philippines, where specimens were trawled from 70–120 m depth and, therefore, were considered rare and confined to deeper water (Lorenz and Fehse 2009). Like almost all other ovulids, species of *Crenavolva* are associated with octocoral hosts (Schiaparelli et al. 2005; Reijnen 2010) belonging to several families (e.g. Melithaeidae, Ellisellidae, Subergorgiidae and Plexauridae). However, the host species are usually not collected or are disregarded and therefore unknown, which is also the case for *C. chiapponii*.

Molecular data (16S and COI) obtained from *Crenavolva* was used by Meyer (2003) to root the phylogeny of the Cypraeidae. Later, the 16S sequence data were used by Schiaparelli et al. (2005) to produce the first molecular phylogenetic reconstruction of the Ovulidae, which included two *Crenavolva* species: *C. cf. rosewateri* (Cate, 1973) and *C. tokuoi* Azuma, 1989. In the present study, material of four additional nominal *Crenavolva* species, amongst other ovulids, have been used to reconstruct a phylogeny. The newly acquired molecular data are for *C. aureola* (Fehse, 2002), *C. chiapponii* Lorenz & Fehse, 2009, *C. striatula* (Sowerby I, 1828) (type species), and *C. trailli* (Adams, 1855). In addition to this phylogenetic reconstruction, data on host species and distributional records are given for this group of rarely recorded ovulid snails.


## Materials and methods

### Collection and identification

During fieldwork in Indonesia (Halmahera, Ternate; Sulawesi, Lembeh Strait) and Malaysia (Borneo, Semporna and Kudat) specimens of *Crenavolva* species were collected by SCUBA diving (Table 1). The snails and their octocoral hosts were photographed in situ (Fig. 1) whenever possible and subsequently fixed in 80% ethanol. The holotype of *C. chiapponii* was studied at the Muséum national d'Histoire naturelle (MNHN) in Paris. For the identification of the other ovulid species, Cate (1973), Fehse (2002b) and Lorenz and Fehse (2009) were used. For the identification of the host species, microscopy slides of their calcareous skeletal parts (sclerites) were made by dissolving the samples in a 4% solution of household bleach. The residual sclerites were rinsed with tap water followed by demineralised water before mounting on a slide or on a stub for Scanning Electron Microscopy (SEM). Stubs with sclerites were coated with Au/Pd before SEM images were made with a JEOL 6480 LV. Identification of the octocorals to genus level was based on Stiasny (1947) and Fabricius and Alderslade (2001).

**Table 1.** Specimens used in the analyses, including locality, host, and GenBank accession data.

| Collection number | Species  | Locality (Locality code)   | Coordinates                   | Date collected | Host species             | GenBank Accession number (16S; COI) | Reference                |
|-------------------|--|--|-------------------------------|----------------|--------------------------|-------------------------------------|--------------------------|
| RMNH.MOL.164072   | <i>Crenavolva aureola</i> (Fehse, 2002)              | Malaysia, Semporna, Si. Amil Island (SEM.16)                         | 4°19'02.1"N;<br>118°52'30.7"E | 4-12-2010      | <i>Acanthogorgia</i> sp. | KP033143; KP033151                  | This publication         |
| RMNH.MOL.164085   | <i>Crenavolva aureola</i> (Fehse, 2002)              | Indonesia, Halmahera, Tidore, N of Desa Rum (TER.18)                 | 0°44'35.8"N;<br>127°23'06.3"E | 4-11-2009      | <i>Acanthogorgia</i> sp. | KP033144; KP033152                  | This publication         |
| RMNH.MOL.164209   | <i>Crenavolva aureola</i> (Fehse, 2002)              | Indonesia, Halmahera, Tanjung Ratermu (S of river)(TER.21)           | 0°54'24.7"N;<br>127°29'17.7"E | 5-11-2009      | <i>Acanthogorgia</i> sp. | KP033148; KP033156                  | This publication         |
| RMNH.MOL.164211   | <i>Crenavolva chiapponii</i> Lorenz & Fehse, 2009    | Indonesia, Halmahera, Tanjung Ratermu (S of river)(TER.27)           | 0°54'44.5"N;<br>127°29'09.9"E | 8-11-2009      | <i>Acanthogorgia</i> sp. | KP033157                            | This publication         |
| RMNH.MOL.164217   | <i>Crenavolva chiapponii</i> Lorenz & Fehse, 2009    | Indonesia, Lembeh, Tanjung Kusukusu (LEM.31)                         | 1°27'13.8"N;<br>125°14'13.0"E | 16-2-2012      | <i>Acanthogorgia</i> sp. | KP033149; KP033158                  | This publication         |
| RMNH.MOL.164062   | <i>Primonula rosewateri</i> (Cate, 1973)             | Malaysia, Semporna, Kulapuan Island 2, N side (SEM.31)               | 4°32'07.4"N;<br>118°50'18.2"E | 9-12-2010      | <i>Panutelesto</i> sp.   | KP033142; KP033150                  | This publication         |
| RMNH.MOL.164186   | <i>Crenavolva striatula</i> (Sowerby I, 1828)        | Malaysia, Sabah, S Pulau Banggi, E Molleangan Besar Island, (TMP.37) | 7°05'07.2"N;<br>117°03'33.8"E | 19-9-2012      | <i>Echinogorgia</i> sp.  | KP033146; KP033154                  | This publication         |
| RMNH.MOL.164144   | <i>Crenavolva trailii</i> (Adams, 1855)              | Malaysia, Sabah, Kalang, (TMP.41)                                    | 6°59'48.1"N;<br>117°03'13.4"E | 18-9-2012      | <i>Subergorgia</i> sp.   | KP033145; KP033153                  | This publication         |
| RMNH.MOL.164189   | <i>Crenavolva trailii</i> (Adams, 1855)              | Malaysia, Sabah, Kalang, (TMP.41)                                    | 6°59'48.1"N;<br>117°03'13.4"E | 18-9-2012      | <i>Panaplexaura</i> sp.  | KP033147; KP033155                  | This publication         |
| -                 | <i>Crenavolva</i> cf. <i>rosewateri</i> (Cate, 1973) | Philippines, Bohol, Balicasag Island                                 | -                             | -              | -                        | AY161394; AY161627                  | Meyer 2003               |
| -                 | <i>Crenavolva tokuoi</i> Azuma, 1989                 | Philippines, Bohol, Balicasag Island                                 | -                             | -              | -                        | AY161390; AY161623                  | Meyer 2003               |
| -                 | <i>Primonula beckeri</i> (Sowerby III, 1900)         | Indonesia, Sulawesi  | -                             | -              | -                        | AJ868555; -                         | Schiaparelli et al. 2005 |
| -                 | <i>Ovula ovum</i> (Linnaeus, 1758)                   | Indonesia, Sulawesi, Spermonde Archipelago                           | -                             | -              | -                        | AY161399; AY161632                  | Meyer 2003               |



**Figure 1.** **A** *In situ* image of *Crenavolva aureola* (Fehse, 2002) (RMNH.MOL.164209) and **B** *C. chiapponii* Lorenz & Fehse, 2009 (RMNH.MOL.164211) on *Acanthogorgia* sp. at Halmahera, Indonesia at 21 m and 17 m depth respectively.


### Barcoding Ovulidae

Specimens were barcoded for the COI barcoding region and for additional phylogenetic research also for the 16S marker. Tissue samples obtained from the foot and/or mantle were extracted with the Machery-Nagel DNA extraction kit on a KingFisher Flex. The standard COI barcoding primers by Folmer et al. (1994) and the Palumbi (1996) 16S primers were used. PCR amplification was performed on a C1000 Touch Thermal Cycler (Bio-RAD). Sequencing of the PCR products was performed at Macrogen Europe on an ABI 3730xl Automated Sequencer. Sequences were edited in Sequencher 4.10.1 and aligned with GUIDANCE (Penn et al. 2010) using the MAFFT algorithm (Katoh et al. 2005). Selecting an evolutionary model was done with jModeltest based on the Akaike Information Criterion score. MEGA 6.0.6 (Tamura et al. 2013) was used to perform Maximum Likelihood (ML) and Maximum Parsimony (MP) analyses and to calculate p-distances. Bayesian analyses were performed in MrBayes 3.2.0 (Ronquist and Huelsenbeck 2003). MrBayes was run for 4,000,000 generations with six chains. Data were sampled every 100 generations. Sequence data for *Ovula ovum* (Linnaeus, 1758) from GenBank was used as an outgroup. GenBank data for *Crenavolva* cf. *rosewateri* (Cate, 1973), *C. tokuoi* Azuma, 1989 and *Primovula beckeri* (Sowerby III, 1900) was also included in the phylogenetic analyses.

## Results

### Collecting and morphology

Eight lots, containing 33 specimens representing four nominal *Crenavolva* species (*C. aureola*, *C. chiapponii*, *C. striatula* and *C. trailli*) were collected in Indonesia and Malaysia (Table 1; Fig. 2). For *C. chiapponii* this is the first record from shallow water. The specimens were assigned to these nominal species based on shell shape (rhomboid, inflated or slender) and the colour bands on the dorsum, which in case of *C. striatula* were



**Figure 2.** Dorsal and ventral views of shells. **A** Holotype of *Crenavolva chiapponii* Lorenz & Fehse, 2009 (MNH 21244) **B** *C. chiapponii* Lorenz & Fehse, 2009 (RMNH.MOL.164211) **C** *C. chiapponii* Lorenz & Fehse, 2009 (RMNH.MOL.164217) **D** *C. aureola* (Fehse, 2002) (RMNH.MOL.164085) **E** *C. aureola* (Fehse, 2002) (RMNH.MOL.164072) **F** *C. aureola* (Fehse, 2002) (RMNH.MOL.164209) **G** *C. trailli* (Adams, 1855) (RMNH.MOL.164144) **H** *C. striatula* (Sowerby I, 1828) (RMNH.MOL.164186) **I** *Primovula rosewateri* (Cate, 1973) (RMNH.MOL.164062). Scale bars: 5 mm.


also present on the labrum. For *C. aureola* and *C. chiapponii* the absence or presence of a white dorsal band on the shell is allegedly the most obvious character to distinguish the species. After examination of the illustrations presented by Lorenz and Fehse (2009) and the newly collected material, minor morphological differences (strongly or weakly pronounced dentation, keeling angle, strongly or weakly produced funiculum, position of the widest part of the shell) do not clearly separate between *C. aureola* and *C. chiapponii* and can be considered morphological variation in a single species. The soft tissue

colouration of both *C. aureola* and *C. chiapponii* is very similar (e.g. Fig 1; Lorenz and Fehse 2009: A106, A107 p. 527). Both have a semi-transparent mantle which is entirely covered with small, irregularly placed, white dots, and both have a completely black or white foot, black tentacles with white tips, and a black siphon.

## Molecular data

Nine specimens representing five species were sequenced for COI and 16S. For one sample of *C. chiapponii* (RMNH.MOL.164211) the 16S marker could not be amplified. Sequences were concatenated and aligned (GUIDANCE alignment score: 0.965034) which resulted in an alignment length of 1080 base pairs per specimen including indels. Sequences obtained from GenBank are slightly shorter (~40 base pairs), these missing base pairs were coded as ‘missing data’. The program jModeltest yielded in HKY+G as most optimal evolutionary model. This evolutionary model was implemented in the Bayesian and ML analysis. The results from the different phylogenetic reconstructions were congruent, therefore only the ML tree is shown (Fig. 3).

In the phylogenetic reconstructions, specimens of *Crenavolva striatula* and *C. tokuoi* form an unresolved trichotomy with the other *Crenavolva* specimens. The two *Primovula* species cluster together and are well-supported sister species to all the *Crenavolva* species (with *C. striatula* as type species for the genus). This implies that the *Crenavolva* species used herein form a monophyletic group. The clustering of two *C. trailli* specimens is highly supported. Another well-supported clade holds three nominal species: *Crenavolva aureola*, *C. chiapponii* and *C. cf. rosewateri*. The pairwise p-distances between these three species are very low (16S: 0.2%; COI: 0.7%; concatenated: 0.9%).



**Figure 3.** Maximum Likelihood cladogram with support values for the ML/MP/BP analyses. Numbers preceding the species names represent RMNH.MOL. collection numbers of Naturalis Biodiversity Center; species names without numbers are obtained from GenBank for which additional data can be found in Table 1.

In contrast, the sequence divergence between *C. trailli* and the *C. chiapponii* / *C. aureola* clade is almost ten times larger (16S: 5.2%; COI: 8.7%; concatenated: 8.2%). The sequence divergence between the two *C. trailli* specimens (16S: 0.6%; COI: 0.8%; concatenated: 0.8%) is almost equal to that between *C. aureola* and *C. chiapponii*. With the help of the Automatic Barcode Gap Discovery tool (ABGD) (Puillandre et al. 2011), the data were analysed to identify the MOTU's within the dataset. The results of this analysis showed that the barcode gap to identify the different species is 5–6% sequence divergence. This resulted in five groups containing the following species: 1, *C. aureola*, *C. chiapponii*, *C. cf. rosewateri*; 2, *C. trailli*; 3, *C. tokuoi*; 4, *C. striatula*; 5, *P. rosewateri*. One of the samples obtained from GenBank, viz. *Crenavolva cf. rosewateri* (= *Primovula cf. rosewateri*), clusters surprisingly within the clade containing *C. aureola* and *C. chiapponii* and not with the other *Primovula rosewateri* specimen. Instead, *Primovula beckeri* proves to be identical to the newly sequenced specimen of *Primovula rosewateri* from Malaysia.

## Octocoral hosts

Almost all Ovulidae species are associated with Octocorallia hosts. By examining the sclerites and the habitus of the host corals, several new host species for ovulids of the genus *Crenavolva* could be identified. An overview of previously identified host species and new records is provided in Table 2. Some of the former host identifications were published with obsolete generic names, and therefore their names in the current literature are also provided. Before *C. chiapponii* was synonymised, *Acanthogorgia* would have been a new host record. Yet, Reijnen (2010) already recorded *Acanthogorgia* sp. as a host for *C. aureola* and therefore it is not a new host record. Morphologically at least two different species of *Acanthogorgia* could be distinguished but these could not be identified since a revision of the family Acanthogorgiidae is lacking.

**Table 2.** Literature overview of the octocoral hosts of selected *Crenavolva* species including new records. Updated names of the octocoral hosts are provided between parentheses.

| Ovulid species   | Host genera   | Reference  |
|--|---|--|
| <i>Crenavolva aureola</i>                              | <i>Euplexaura</i> ; <i>Astromuricea</i> (= <i>Echinogorgia</i> );<br><i>Acanthogorgia</i>   | Lorenz and Fehse 2009; Reijnen 2010                            |
| <i>Crenavolva chiapponii</i><br>(= <i>C. aureola</i> ) | <i>Acanthogorgia</i>  | this publication; Reijnen 2010                                 |
| <i>Crenavolva striatula</i>                            | <i>Ellisella</i> ; <i>Euplexaura</i> ; <i>Echinogorgia</i>  | Lorenz and Fehse 2009; Yamamoto 1973; Cumming 1997; Mase 1989; |
| <i>Crenavolva trailli</i>                              | <i>Echinogorgia</i> ; <i>Anthoplexaura</i> (= <i>Astrogorgia</i> );<br><i>Plexauroides</i> (= <i>Echinogorgia</i> ); <i>Euplexaura</i> ; <i>Subergorgia</i> | Goh et al. 1999; Mase 1989                                     |
| <i>Primovula rosewateri</i>                            | <i>Subergorgia</i> ; <i>Dendronephthya</i> ; <i>Stereonephthya</i> ;<br><i>Paratelesto</i>  | Goh et al. 1999; Lorenz and Fehse 2009; this publication       |
| <i>Primovula beckeri</i>                               | <i>Acanthogorgia</i> ; <i>Acabaria</i> (= <i>Melithaea</i> ); <i>Unicella</i> [sic]<br>(= <i>Eunicella</i> ); <i>Lophogorgia</i> (= <i>Leptogorgia</i> )    | Schiaparelli et al. 2005; Lorenz and Fehse 2009                |

Furthermore, examination of the ovulid species and their octocoral hosts revealed that in two instances individuals formerly identified as *C. chiapponii* and *C. aureola* would have co-occurred on the same host coral, in both cases *Acanthogorgia* sp.

## Discussion

Based on the molecular data and morphological observations listed above, *C. chiapponii* is considered a junior synonym of *C. aureola*. The systematic account is therefore as follows:

## Systematic part

### Family Ovulidae Fleming, 1822

### Genus *Crenavolva* Cate, 1973

### *Crenavolva aureola* (Fehse, 2002)

*Primovula aureola* Fehse 2002: 37, pl. 1, fig. 1

*Delonovolva formosa*. — Gosliner et al. 1996: 136, fig. 469. Not *Delonovolva formosa* (Sowerby II in Adams and Reeve 1848) [= *Cuspivolva formosa* (Sowerby II in Adams and Reeve 1848)]

*Primovula* sp. — Coleman 2003: 51, fig. (Ovul: 121).

*Crenavolva chiapponii* Lorenz and Fehse 2009: 69, pl. 74, fig. 7–11.

The occurrence of *C. chiapponii* (= *C. aureola*) on Indonesian shallow water coral reefs would have represented new distribution records, both geographically and bathymetrically, before it was synonymised. However *C. chiapponii* proved to be a junior synonym of *C. aureola* and the new distribution records fall within the distribution range already known for *C. aureola*. Apparently, the dorsal white band and the minor morphological differences in shell shape are not indicative of species-level differences between *C. aureola* and *C. chiapponii*.

## Molecular data

The species *Primovula rosewateri* was previously placed in the genus *Crenavolva* by Cate (1973) but Fehse (2002a) moved it to *Primovula*, primarily based on the triangular shape of the funiculum. The results of the molecular analyses (Fig. 3) support this decision. There is great genetic similarity between *C. cf. rosewateri* (= *Primovula cf. rosewateri*) obtained from GenBank, and *C. aureola*. However, the specimen from GenBank was collected from Balicasag Island, near Bohol, Philippines, which is the

type locality of *C. chiapponii*. This location is approximately 85 km from Mactan Island of Cebu, Philippines which is the type locality of *C. aureola*. It is not unlikely that the so-called *C. cf. rosewateri* from GenBank (AY161394 (16S), AY161627 (COI)) was misidentified and actually represents *C. aureola*. Moreover, the newly sequenced specimen of *P. rosewateri* from Malaysia convincingly clusters with *Primovula beckeri*. According to Lorenz and Fehse 2009, *P. beckeri* has an E African distribution and was originally described from South Africa. The specimen obtained from GenBank is from Sulawesi, Indonesia (Schiaparelli et al. 2005). It is therefore unlikely that this sequence represents *P. beckeri* but instead is the quite similar species from the central Indo-Pacific, *P. rosewateri*.

### Host species and distribution records

The ranges of the presently discussed species all fit within the Coral Triangle (see Hoeksema 2007) and depend on the ranges of their host species. Species of the genus *Acanthogorgia* are not unique hosts for just *Crenavolva* spp. Reijnen (2010) already mentioned *Acanthogorgia* spp. as a host for *Dentiovula eizoi* Cate & Azuma, 1973 (in Cate 1973) and *D. colobica* (Azuma & Cate, 1971). *Acanthogorgia* species and their ovulid associates are both known to occur from shallow to deep water in the Coral Triangle. In an overview of the Acanthogorgiidae by Stiasny (1947) the deepest record for an *Acanthogorgia* species is 4239 m, collected SE of Seram, Indonesia (*Acalycigorgia densiflora* = *Acanthogorgia densiflora* (Kükenthal & Gorzawsky, 1908). Nevertheless, Stiasny (1947) doubts the identification and compared it to congeneric species which are found in waters not exceeding 400 m depth. As a result Stiasny (1947) doubts the entire record. Therefore the deepest reliable record for an *Acanthogorgia* species in the Malayan Archipelago is 1254 m for *Acanthogorgia multispina* (Kükenthal & Gorzawsky, 1908). The deepest record for *Crenavolva* species is from approximately 1000 m, which is the deepest record for any ovulid species found to date (Lorenz and Fehse 2009).

### Acknowledgements

Many thanks to Dr Bert W. Hoeksema who organized the Ternate and Lembah Strait expedition together with Ir M.I. Yosephine Tuti Hermanlimianto under the umbrella of E-Win (Ekspedisi Widya Nusantara). LIPI Ternate and LIPI Bitung are acknowledged for accommodating the research at their field stations. LIPI and RISTEK granted research permits. The Semporna Marine Ecological Expedition (SMEE2010) was jointly organized by WWF-Malaysia, Universiti Malaysia Sabah's Borneo Marine Research Institute, Naturalis Biodiversity Center and Universiti Malaya's Institute of Biological Sciences, while research permission was granted by the Economic Planning Unit, Prime Minister's Department, Economic Planning Unit Sabah, Sabah Parks

and Department of Fisheries Sabah. The MV Celebes Explorer accommodated the research in Semporna. The 2012 Tun Mustapha Park expedition (TMP) was jointly organized by WWF-Malaysia, Universiti Malaysia Sabah (UMS), Sabah Parks and Naturalis Biodiversity Center, the Netherlands. The research permits were granted by the Economic Planning Unit, Prime Minister's Department Malaysia and Sabah Biodiversity Centre. Both expeditions to Malaysia were co-organised by Ms Zarinah Waheed and WWF Malaysia, which was greatly appreciated. Dr Philippe Bouchet and Ms Virginie Heros were so kind to accommodate a visit to Muséum national d'Histoire naturelle (MNHN) in Paris to investigate the Ovulidae (type) collections. Dr Leendert P. van Ofwegen kindly provided help by identifying the octocoral hosts and Sancia van der Meij was a perfect dive buddy and ovulid hunter. Sequencing of the barcoding region of the ovulids was part of one of the barcoding initiatives within the Naturalis Biodiversity Center. Funding for fieldwork was provided by the Jan-Joost ter Pelkwijkfonds and A.M. Buitendijkfonds (Naturalis Biodiversity Center), Malacological Society of Australasia and the Percy Sladen Fund. I also would like to thank Prof Dr Catherine S. McFadden, an anonymous reviewer and Dr Thierry Backeljau for their constructive comments and remarks which improved the manuscript greatly.

## References

- Adams A (1855) Descriptions of twenty-five new species of shells from the collection of Hugh Cumming. *Proceedings of the Zoological Society London* 23: 221–226.
- Adams A, Reeve LA (1848–1850) Mollusca. In: Adams A (Ed.) *The zoology of the voyage of H.M.S. Samarang under the command of Sir Edward Belcher, during the years 1843–1846*. Reeve and Benham, London, 1–87.
- Azuma M, Cate CN (1971) Sixteen new species and one new genus of Japanese Ovulidae. *The Veliger* 13: 261–268.
- Azuma M (1989) Systematic studies on the recent Japanese family Ovulidae (Gastropoda) – XIV. Supplement (3) Description of a new species of the genus *Crenavolva* Cate, 1973. *Venus* 48: 161–166.
- Cate CN (1973) A systematic revision of the recent cypraeid family Ovulidae. *The Veliger* suppl. 15: 1–116.
- Coleman N (2003) 2002 Seashells: Catalogue of Indo Pacific Mollusca. Neville Coleman's Underwater Geographic Pty Ltd., Australia, 144 pp.
- Cumming RL (1997) Ovulids (Gastropoda) associated with gorgonians (Anthozoa: Gorgonoidea) at Cape D'Aguilar, Hong Kong: species, hosts, distributions and feeding ecology. In: Morton B (Ed.) *The Marine Flora and Fauna of Hong Kong and Southern China (IV)*. Proceedings of the eight International Marine Biological Workshop, Hong Kong, April 1995. Hong Kong University Press, Hong Kong, 285–301.
- Fabricius KE, Alderslade P (2001) Soft corals and sea fans: A comprehensive guide to the tropical shallow water genera of the central-west Pacific, the Indian Ocean and the Red Sea. Australian Institute of Marine Science, Townsville, 264 pp.

- Fehse D (2002a) Contributions to the knowledge of Ovulidae (Mollusca: Gastropoda). XI. A new species of the genus *Primovula* Thiele, 1925. *La Conchiglia* 34: 36–40.
- Fehse D (2002b) Contributions to the knowledge of Ovulidae (Mollusca: Gastropoda). X. The genus *Crenavolva* Cate, 1973. *La Conchiglia* 34: 23–55.
- Fleming J (1822) The philosophy of zoology, or a general view of the structure, functions, and classification of animals. Hurst, Robinson & Co, Edinburgh, 432 pp. doi: 10.1037/11823-000
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3: 294–299.
- Goh NGC, Ng PKL, Chou LM (1999) Notes on the shallow water gorgonian-associated fauna on coral reefs in Singapore. *Bulletin of Marine Science* 65: 259–282.
- Gosliner TM, Behrens DW, Williams GC (1996) Coral reef animals of the Indo-Pacific. *Sea Challengers*, Monterey, 314 pp.
- Hoeksema BW (2007) Delineation of the Indo-Malayan Centre of Maximum Marine Biodiversity: The Coral Triangle. In: Renema W (Ed.) *Biogeography, Time and Place: Distributions, Barriers and Islands*. Springer, Dordrecht, 117–178. doi: 10.1007/978-1-4020-6374-9\_5
- Katoh K, Kazutaka K, Kuma K, Toh H, Miyata T (2005) MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research* 33: 511–518. doi: 10.1093/nar/gki198
- Kükenthal W, Gorzawsky H (1908) Japanische Gorgoniden. 1. Teil: Die Familien der Primnoiden, Muriceiden und Acanthogorgiiden. In: Doflein F (Ed.) *Beiträge zur Naturgeschichte Ostasiens. Abhandlungen Mathematisch-Physikalischen Klasse der Königlich Bayerischen Akademie der Wissenschaften Supplement Bd. 1*, München, 1–71.
- Lorenz F, Fehse D (2009) The living Ovulidae, a manual of the families of allied cowries: Ovulidae, Pediculariidae and Eocypraeidae. *Conchbooks*, Hackenheim, 651 pp.
- Mase K (1989) Taxonomic significance of color patterning of the soft body in the family Ovulidae – Descriptions of soft body of 26 species. *Venus suppl.* 1: 75–120.
- Meyer CP (2003) Molecular systematics of cowries (Gastropoda: Cypraeidae) and diversification patterns in the tropics. *Biological Journal of the Linnean Society* 79: 401–459. doi: 10.1046/j.1095-8312.2003.00197.x
- Palumbi SR (1996) PCR and molecular systematics. In: Hillis D, Moritz C, Mable B (Eds) *Molecular Systematics*. Sinauer Press, Sunderland, 205–247.
- Penn O, Privman E, Ashkenazy H, Landan G, Graur D, Pupko T (2010) GUIDANCE: a web server for assessing alignment confidence scores. *Nucleic Acids Research* 38 (suppl 2): W23–W28. doi: 10.1093/nar/gkq443
- Puillandre N, Lambert A, Brouillet S, Achaz G (2012) ABGD, Automatic Barcode Gap Discovery for primary species delimitation. *Molecular Ecology* 21(8): 1864–1877. doi: 10.1111/j.1365-294X.2011.05239.x
- Reijnen BT (2010) Camouflage hampering the taxonomy of Ovulidae (Mollusca: Gastropoda) in the centre of marine biodiversity (Halmahera, Indonesia). *Australian Shell News* 137: 5–7.
- Ronquist F, Huelsenbeck JP (2003) MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574. doi: 10.1093/bioinformatics/btg180

- Rosenberg G (2014) *Crenavolva* Cate, 1973. Available from World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=204661> [accessed 26 November 2014]
- Schiaparelli S, Barucca M, Olmo E, Boyer M, Canapa A (2005) Phylogenetic relationships within Ovulidae (Gastropoda: Cypraeoidea) based on molecular data from the 16S rRNA gene. *Marine Biology* 147: 411–420. doi: 10.1007/s00227-005-1566-0
- Sowerby I GB (1828) On the recent species of the genus *Ovulum*. *Zoological Journal* 4: 145–162.
- Sowerby III GB (1900) On some marine shells from Pondoland and the Kowie, with descriptions of seventeen new species. *Proceedings of the Malacological Society of London* 4: 1–7.
- Stiasny G (1947) De gorgonarien-familie Acanthogorgiidae Kükenthal & Gorzawsky, 1908 met bijzondere inachtneming van het materiaal der Siboga-expeditie. *Verhandelingen der Koninklijke Nederlandsche Akademie der Wetenschap, Afd. Natuurkunde. Tweede Sectie* 43: 1–93.
- Tamura K, Stecher G, Peterson D, Filipiski A, Kumar S (2013) MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution* 30: 2725–2729. doi: 10.1093/molbev/mst197
- Yamamoto T (1973) Molluscs symbiotic with coelenterates in Japan, with special reference to Ovulidae and allied forms. *Publications from the Seto Marine Biological Laboratory* 20: 567–581.