

# Two new species of *Oobius* Trjapitzin (Hymenoptera, Encyrtidae) egg parasitoids of *Agrilus* spp. (Coleoptera, Buprestidae) from the USA, including a key and taxonomic notes on other congeneric Nearctic taxa

Serguei V. Triapitsyn<sup>1</sup>, Toby R. Petrice<sup>2</sup>, Michael W. Gates<sup>3</sup>, Leah S. Bauer<sup>2</sup>

**1** Entomology Research Museum, Department of Entomology, University of California, Riverside, California, 92521, USA **2** United States Department of Agriculture Forest Service, Northern Research Station, 3101 Technology Blvd., Suite F, Lansing, Michigan, 48910, USA **3** Systematic Entomology Laboratory c/o National Museum of Natural History, P.O. Box 37012, Washington, DC, 20013-7012, USA

Corresponding author: *Serguei V. Triapitsyn* (serguei@ucr.edu)

Academic editor: *M. Engel* | Received 7 February 2015 | Accepted 3 March 2015 | Published 21 April 2015

<http://zoobank.org/480DEF98-A22C-4253-8479-6222FD3E9E2F>

**Citation:** Triapitsyn SV, Petrice TR, Gates MW, Bauer LS (2015) Two new species of *Oobius* Trjapitzin (Hymenoptera, Encyrtidae) egg parasitoids of *Agrilus* spp. (Coleoptera, Buprestidae) from the USA, including a key and taxonomic notes on other congeneric Nearctic taxa. ZooKeys 498: 29–50. doi: 10.3897/zookeys.498.9357

## Abstract

*Oobius* Trjapitzin (Hymenoptera, Encyrtidae) species are egg parasitoids that are important for the biological control of some Buprestidae and Cerambycidae (Coleoptera). Two species, *O. agrili* Zhang & Huang and *O. longoi* (Siscaro), were introduced into North America for classical biocontrol and have successfully established. Two new native North American species that parasitize eggs of *Agrilus* spp. (Buprestidae) are described and illustrated from the USA: *O. minusculus* Triapitsyn & Petrice, **sp. n.** (Michigan), an egg parasitoid of both *A. subcinctus* Gory on ash (*Fraxinus* spp.) and *A. egenus* Gory on black locust (*Robinia pseudoacacia* L.) trees, and *O. whiteorum* Triapitsyn, **sp. n.** (Pennsylvania), an egg parasitoid of *A. anxius* Gory on European white birch (*Betula pendula* Roth). A taxonomic key and notes on the Nearctic native and introduced *Oobius* species are also included.

## Keywords

Emerald ash borer, new species, congener identification key, *Oobius agrili*, Nearctic, egg parasitoid, biological control

## Introduction

The rather poorly known encyrtid genus *Oobius* Trjapitzin (Hymenoptera: Encyrtidae) currently includes 41 species worldwide, and seven are known from North America (Noyes 2014). Noyes (2010) recently described 20 of these species from Costa Rica and compared some of these new species to similar Nearctic taxa. Also, in Noyes (2010) the genera *Avetianella* Trjapitzin, *Szelenyiola* Trjapitzin, and *Oophagus* Liao were synonymized under *Oobius*.

As egg parasitoids of Buprestidae, Cerambycidae (Coleoptera; Noyes 2014) and Asilidae (Diptera; Annecke 1967), species of *Oobius* are important for the natural and classical biological control of some coleopteran species. Two species are being used as biological control agents in the USA where they are successfully established: *O. agrili* Zhang & Huang and *O. longoi* (Siscaro). The former was collected from China and was first released into the USA in 2007 as a biocontrol agent of the emerald ash borer, *Agrilus planipennis* Fairmaire (Buprestidae) (Bauer et al. in press). *Agrilus planipennis* is an invasive pest from Asia that attacks ash trees (*Fraxinus* spp.) (Haack et al. 2002; Bray et al. 2011). Releases of *O. agrili* are ongoing throughout infested regions of the USA. As of March 2015, establishment of *O. agrili* has been confirmed in Indiana, Maryland, Michigan, New York, Ohio, and Pennsylvania (Abell et al. 2014; Bauer et al. in press). *Oobius longoi* was introduced from Australia to California, USA, as a biocontrol agent for management of *Phoracantha recurva* Newman and *P. semipunctata* (Fabricius) (Cerambycidae), which are invasive pests of *Eucalyptus* trees in the USA (Hanks et al. 1995; Luhring et al. 2000).

Here two new species of *Oobius* are reported and a taxonomic key to the known native and introduced species of *Oobius* in North America is provided. One of the newly described species was reared initially from eggs of the native buprestid *Agrilus subcinctus* Gory in Michigan, whose larvae feed on the dead twigs of ash trees. This parasitoid was previously reported by Petrice et al. (2009) as *Avetianella* sp. Later, the second author of the current paper also reared this same species from eggs of *A. egenus* Gory on black locust trees (*Robinia pseudoacacia* L.) in Michigan. *Agrilus egenus* is a native species that oviposits on dead or dying branches of black locust (Nelson et al. 2008; MacRae 1991). The second newly described *Oobius* species was reared from *A. anxius* Gory eggs on European white birch (*Betula pendula* Roth) in Pennsylvania, based on the voucher specimens from the study by Loerch and Cameron (1983). *Agrilus anxius* is a native species that attacks both native and introduced birch trees (*Betula* spp.) in North America (Nelson et al. 2008).

## Material and methods

**Collecting and rearing new species of *Oobius*.** Ash tree twigs with *A. subcinctus* eggs and black locust twigs with *A. egenus* eggs were collected in the field in Ingham and Clinton counties, Michigan in 2013 and 2014. Eggs were monitored in the labora-

tory for parasitoid emergence. Voucher specimens of the parasitoids were preserved in 95% ethanol and sent to the senior author for identification. See Loerch and Cameron (1983) for collection of parasitoids from *A. anxius*.

**Taxonomic studies.** Parasitoid specimens used in the taxonomic studies were critical point dried from ethanol and point-mounted. Selected specimens were then dissected and slide-mounted in Canada balsam, examined under a Zeiss<sup>®</sup> Axioskop 2 plus compound microscope using Nomarski differential interference contrast optics. Stereomicroscopic images were compiled with Auto-Montage 4.02 (Synchroscope<sup>®</sup>) to illustrate select specimens. Images of specimens were produced by scanning electron microscopy (SEM) and an EntoVision Imaging Suite. A Nikon<sup>®</sup> SMZ1500 and Leica<sup>®</sup> MZ 9.5 stereomicroscope with 10X oculars (Nikon C-W10X/22) and Chiu Technical Corp.<sup>®</sup> Lumina 1 FO-150 and fiber optic light source was used for pinned specimen observation. Mylar film was placed over the ends of the light source to reduce glare. Scanning electron microscope (SEM) images were taken with a Hitachi<sup>®</sup> TM3000 desktop unit (Tungsten source). Some specimens were manually cleaned of external debris with forceps or brushes and affixed to 12.7X 3.2 mm Leica/Cambridge aluminum SEM stubs with carbon adhesive tabs (Electron Microscopy Sciences, #77825-12). Stub-mounted specimens were imaged uncoated or sputter coated using a Cressington Scientific 108 Auto with a gold-palladium mixture from at least three different angles to ensure complete coverage (~20–30nm coating). Color images were obtained using an EntoVision Imaging Suite, which includes a firewire JVC KY-75 3CCD digital camera mounted to a Leica M16 zoom lens via a Leica z-step microscope stand. Slides of *O. buprestidis* and *O. dahlsteni* were imaged with a Leica DMRB compound microscope fitted with Leica HCX PL “Fluotar” 5× and 10× metallurgical grade lenses. Both systems fed image data to a desktop computer where Cartograph 5.6.0 (Microvision Instruments<sup>®</sup>, France) was used to capture a fixed number of focal planes (based on magnification); the resulting focal planes (manually captured via Archimed 5.5.0 on the DMRB) were merged into a single, in-focus composite image. Uniform lighting was achieved using a LED illumination dome with all four quadrants set to 99.6% intensity. The images were then retouched where necessary using Adobe Photoshop<sup>®</sup> CS4/CS6 with plates assembled using InDesign CS4/CS6.

Terms used for morphological features are those of Gibson (1997). Abbreviations used are: F = antennal funicle segment; mps = multiporous plate sensillum or sensilla on the antennal flagellar segments (= longitudinal sensillum or sensilla or sensory ridge(s) of authors). Body length was measured without the exerted part of the ovipositor.

Acronyms for depositories of specimens are as follows: BMNH, The Natural History Museum, London, England, UK; EMEC, Essig Museum of Entomology, University of California, Berkeley, California, USA; IZCAS, Institute of Zoology, Chinese Academy of Sciences, Beijing, China; MSUC, Albert J. Cook Arthropod Research Collection, Department of Entomology, Michigan State University, East Lansing, Michigan, USA; PSUC, Frost Entomological Museum, Pennsylvania State University, University Park, State College, Pennsylvania, USA; UANL, Universidad Autónoma de

Nuevo León, San Nicolás de los Garza, Monterrey, Mexico; UCRC, Entomology Research Museum, University of California, Riverside, California, USA; UNCA, Institute of Agricultural Entomology, University of Catania, Catania, Sicily, Italy; USNM, National Museum of Natural History, Washington, District of Columbia, USA.

## Taxonomy

### *Oobius* Trjapitzin, 1963

*Oobius* Trjapitzin 1963: 544–545. Type species: *Tyndarichus rudnevi* Nowicki, by original designation.

*Avetianella* Trjapitzin 1968: 97–99. Type species: *Avetianella capnodiobia* Trjapitzin, by monotypy. Synonymized under *Oobius* by Noyes 2010: 660–662.

*Szelenyiola* Trjapitzin 1977: 160. Type species: *Szelenyiola nearctica* Trjapitzin, by original designation and monotypy. Synonymized under *Oobius* by Noyes 2010: 660–662.

*Oobius*: Trjapitzin 1977: 161 (key to genera of the subtribe Oobiina of the tribe Discodini of the subfamily Encyrtinae); Noyes 2010: 660–662 (synonymy, diagnosis, host associations, comments); Trjapitzin and Volkovitsh 2011: 670–672 (diagnosis of *Oobius* s. str., taxonomic position, key to world species).

*Avetianella*: Gordh and Trjapitzin 1981: 6 (comments); Trjapitzin 2001: 734–735 (comments, key to world species).

*Oophagus* Liao in Liao et al. 1987: 184. Type species: *Oophagus batocerae* Liao, by original designation and monotypy. Synonymized under *Avetianella* by Zhang and Huang 2004: 34–35, and under *Oobius* by Noyes 2010: 660.

*Szelepyiola*: Trjapitzin and Volkovitsh 2011: 671 (misspelled).

**Comments.** *Oobius* is a cosmopolitan genus as defined by Noyes (2010) who provided its detailed diagnosis, which is omitted here for brevity. One extralimital species, *O. striatus* Annecke, is also known from eggs of Asilidae (Diptera) in Montenegro and Zimbabwe (Annecke 1967; Noyes 2010, 2014).

### Key to the Nearctic species of *Oobius*, females (both native and introduced)

(*Oobius depressus* (Girault) not included)

- |      |   |   |
|------|---|---|
| 1    | Tarsi 4-segmented (Fig. 1) .....  | <b><i>O. agrili</i> Zhang &amp; Huang</b> |
| –    | Tarsi 5-segmented (Figs 7, 13) .....  | <b>2</b>                                  |
| 2(1) | Clava entire (Figs 9, 10) .....   | <b><i>O. nearcticus</i> (Trjapitzin)</b>  |
| –    | Clava 3-segmented (Figs 2, 6, 8, 12, 22) .....                                      | <b>3</b>                                  |
| 3(2) | Body length (dry-mounted specimens) at most 0.53 mm; mps only on F6 (Fig. 12) ..... | <b><i>O. minusculus</i> sp. n.</b>        |

- Body length (dry-mounted specimens) at least 0.66 mm; mps on F6 and other funicle segments (Figs 6, 20, 22) ..... 4
- 4(3) Mps on F5 and F6 (Fig. 6) ..... *O. buprestidis* (Gordh & Trjapitzin)
- Mps on F4–F6 (Figs 8, 22) ..... 5
- 5(4) Linea calva “open” posteriorly (Fig. 23), uninterrupted by row of setae .....  
..... *O. longoi* (Siscaro)
- Linea calva interrupted posteriorly by a line (or lines) of setae (Figs 19, 27)... 6
- 6(5) F5 and F6 each notably longer than F4 (Fig. 26), F4 0.8× length of F5 .....  
..... *O. whiteorum* sp. n.
- F5 and F6 each subequal in length to F4 (Fig. 8), F4 more than 0.9× length of F5 ..... *O. dahlsteni* (Trjapitzin)

### Alphabetical synopsis of the Nearctic species

#### *Oobius agrili* Zhang & Huang, 2005

Figures 1–4

*Oobius agrili* Zhang & Huang in Zhang et al. 2005: 254–258. Holotype female [IZ-CAS], not examined. Type locality: Changchun, Jilin, China.

*Oobius agrili* Zhang & Huang; Trjapitzin and Volkovitsh 2011: 671 (key), 672–673 (taxonomic history, host associations, use in biological control for *A. planipennis*).

**Material examined.** USA, Michigan, Ingham Co., East Lansing, United States Department of Agriculture (USDA) Forest Service Northern Research Station, laboratory culture of *O. agrili* reared in *Agrilus planipennis* eggs: 37<sup>th</sup>-generation progeny, emerged 10.viii.2014, D.L. Miller, originally from CHINA, Jilin (Jingyuetan Forest Park, Changchun), 2006, T. Zhao (Zhao Tonghai), from eggs of *A. planipennis* [10 ♀, UCRC]; 6–7<sup>th</sup>-generation progeny, emerged 31.vii.2014, D.L. Miller, originally from CHINA, Jilin (Jingyuetan Forest Park, Changchun), 2008, T. Zhao, from eggs of *A. planipennis* [11 ♀, UCRC]; 4–8<sup>th</sup>-generation progeny, emerged 10.viii.2014, D.L. Miller, originally from CHINA, Jilin (Jingyuetan Forest Park, Changchun), 2009, T. Zhao, from eggs of *A. planipennis* [16 ♀, UCRC]; 9<sup>th</sup>-generation progeny, emerged 18.vii.2014, D.L. Miller, originally from CHINA, Jilin (Jingyuetan Forest Park, Changchun), 2008, T. Zhao, from eggs of *A. planipennis* [11 ♀, UCRC].

**Distribution.** China (Zhang et al. 2005; Liu et al. 2007); USA (introduced): Indiana, Maryland, Michigan, New York, Ohio, and Pennsylvania, as of March 2015 (Abell et al. 2014; Bauer et al. in press).

**Host.** *Agrilus planipennis* Fairmaire.

**Comments.** *Oobius agrili* is a solitary thelytokous egg parasitoid of *A. planipennis*, discovered in 2004 during foreign exploration for natural enemies in northeast China (Zhang et al. 2005; Liu et al. 2007; Trjapitzin and Volkovitsh 2011). Adults *O. agrili* were reared from eggs at the USDA Forest Service Northern Research Station laboratory



**Figures 1–8.** 1–4 *Oobius agrili* female (from USDA Forest Service laboratory colony, East Lansing, Michigan, USA; of China origin), 1 hind leg 2 antenna 3 lateral habitus 4 forewing base 5–6 *Oobius buprestidis* female (holotype), 5 lateral habitus 6 antenna 7–8 *Oobius dahlsteni* female (holotype) 7 dorsal habitus 8 antenna.

in East Lansing, Michigan, USA. Rearing stock for this colony originated from parasitized *A. planipennis* eggs collected from *Fraxinus pennsylvanica* trees in Jingyuetan Forest Park, Changchun, Jilin Province, China in 2004–2009. In 2007, *O. agrili* introductions began in Michigan, USA, for classical biological control of *A. planipennis*. As of fall 2014, releases of *O. agrili* had expanded to 19 states (Bauer et al. in press). Abell et al. (2014) reported parasitism of *A. planipennis* eggs averaged approximately 20% in 2012–2013 at some sites where *O. agrili* was established, however, more studies are needed to assess the impact of *O. agrili* and other *A. planipennis* biocontrol agents on ash recovery in the USA. Since 2010, stock cultures of *O. agrili* have been provided to the USDA Animal and Plant Health Inspection Service Emerald Ash Borer Biocontrol Facility, Brighton, Michigan, USA, for mass-rearing and releasing as a biocontrol agent of *A. planipennis* in infested regions of the USA (Mapbiocontrol 2014). To distinguish *O. agrili* from the known native and the other introduced *Oobius* species, we provide illustrations of its metatarsus (Fig. 1), female antenna (Fig. 2), lateral habitus of the female (Fig. 3), and base of the forewing (Fig. 4).

### ***Oobius buprestidis* (Gordh & Trjapitzin, 1981)**

Figures 5–6

*Avetianella buprestidis* Gordh and Trjapitzin 1981: 7–8, 9 (key), 59 (illustrations). Type locality: Portland, Multnomah Co., Oregon, USA.

*Avetianella buprestidis*: Trjapitzin 2001: 735 (key), 736 (list).

*Oobius buprestidis* (Gordh & Trjapitzin): Noyes 2010: 662.

**Type material examined.** Holotype female [USNM] on point mount labeled with following seven labels: “Ex egg of *Buprestus* [sic] *aurulentus*”, “Portland, Ore., F.D. Keen Colr.”, “Hopk. US No. 33150-D”, “Lot No. 41-14524”, “*Habrolepoidea* n. sp. det. Gahan”, “*Avetianella* sp.n. Det Trjapitzin et Gordh”, [red] “Holotypus *Avetianella buprestidis* G. & T.”. The head and antenna are slide mounted separately: [left label] “♀ Holotype, Head & antenna, *Avetianella buprestidis* Gordh & Trjapitzin”, [right label] “Portland, Oregon, Hopkins #33150-D, Lot #41-14524, Ex eggs *Buprestus aurulentus*”. The forewing is mounted on an additional slide with the forewing of a male paratype: [left label] “♂ Forewing, top, *Avetianella buprestidis* G.&T., Portland, Ore., Lot # 41-14524, Hopkins # 33150-D, ♂ paratype”, [right label] “♀ Forewing, bottom, (Holotype) Ex. eggs *Buprestus aurulentus*, F.P. Keen, col. Head & antenna, *Avetianella buprestidis* Gordh & Trjapitzin, [right label] “Portland, Oregon, Hopkins #33150-D, Lot #41-14524, Ex eggs *Buprestus aurulentus*”.

**Distribution.** USA (Oregon) (Gordh and Trjapitzin 1981).

**Host.** *Buprestis aurulenta* L. (Gordh and Trjapitzin 1981 [as *Buprestus aurulentus*]; Trjapitzin 2001 [as *Cypriacus aurilentus* L.]).

**Comments.** The point-mounted portion of the type (Fig. 5) is positioned at the apex of the point. Co-mounted proximally is a complete male paratype. A sliver of wood is pinned in the main collection (USNM) on which are eight eggs of *B. aurulenta* (7 of which have parasitoids emergence holes) bearing the 33150-D Hopkins number designation.

### ***Oobius dahlsteni* (Trjapitzin, 1971)**

Figures 7–8

*Avetianella dahlsteni* Trjapitzin 1971: 890–892. Type locality: McCloud [Flat], Siskiyou Co., California, USA.

*Avetianella dahlsteni*: Gordh and Trjapitzin 1981: 9 (key); Trjapitzin and Gordh 1984: 1275; Trjapitzin 2001: 735 (key), 736 (list).

*Oobius dahlsteni* (Trjapitzin): Noyes 2010: 662.

**Type material examined.** Holotype female [EMEC] on slide labeled: [left label] “*Avetianella dahlsteni* Trjapitzin ♀, Trjapitzin 1970, CFL III-69, Ch. phenol gum damar, Div. Biol. Conn. Univ. Calif [“holotype” handwritten at top, middle, and bottom of label in red ink]”, [right label] “McCloud Flat, Siskiyou Co. Calif., July, 1968, *D. brevicornis* rearing carton, A900, MF2-5 SR., D. L. Dahlsten”.

**Distribution.** USA (California) (Trjapitzin 1971).

**Hosts.** Unknown.

**Comments.** The holotype (Fig. 7) is complete and whole mounted; its antenna (Fig. 8) is also illustrated to facilitate recognition of this species.

### ***Oobius depressus* (Girault, 1916)**

Figures 17–18

*Habrolepoidea depressa* Girault 1916: 343–344. Type locality: Morrystown, Henry Co., Illinois, USA.

*Avetianella depressa* (Girault): Gordh and Trjapitzin 1981: 8; Trjapitzin and Gordh 1984: 1275 (lectotype designation, comments); Trjapitzin 2001: 735 (mentioned), 736 (list).

*Oobius depressus* (Girault): Noyes 2010: 662, 690.

**Type material examined.** Lectotype female [USNM], designated by Trjapitzin & Gordh (1984), on point with following six labels: “Morrystown XII-8-14 III”, “Ex-Eggs Cylene robinae”, “JRMalloch Coll.”, [red] “Paratype No. 20328 U. S. N. M.”, “*Avetianella* Det. Trjapitzin et Gordh”, [red] “*Lectotypus* ♀ *Habrolepoidea depressa* Grlt Des. Trjapitzin et Gordh”. Paralectotypes, 2 males, 2 females: 1 female [USNM] on point with following six labels: “Morrystown XII-8-14 III”, “ExEggs Cylene robinae”,

“JRMalloch Coll.”, [red] “Paratype No. 20328 U. S. N. M.”, [red] “*Paralectotypus* ♀ *Habrolepoidea depressa* Grlt Des. Trjapitzin et Gordh”, “*Avetianella depressa* (Girault) ♀ Det. V. Trjapitzin May 1997”; 1 female [USNM] on point with following six labels: “Morristown XII-8-14 III”, “ExEggs *Cylene robiniae*”, “JRMalloch Coll.”, [red] “Paratype No. 20328 U. S. N. M.”, 5. “*Habrolepoidea depressa* Gir Type”, “LECTOTYPE *Habrolepoidea depressa* Girault By B.D. Burks”; 2 males [USNM] on points, each with following four labels: “Morristown XII-8-14 III”, “ExEggs *Cylene robiniae*”, “JRMalloch Coll.”, [red] “Paratype No. 20328 U. S. N. M.”. All specimens of the type series lack the heads and antennae (Trjapitzin and Gordh 1984; Trjapitzin 2001).

**Distribution.** USA (Illinois) (Girault 1916).

**Host.** *Megacyllene robiniae* (Forster) (Cerambycidae) (Girault 1916 [as *Cyllene robiniae*]).

**Comments.** The identity of this species remains unclear because the original description is poor and without any illustrations; unfortunately, the slide with a head and a forewing of each sex (Girault 1916) could not be found in the USNM and is presumed lost. The lectotype label affixed by B. D. Burks was not validly designated and is merely a paralectotype. To facilitate identification of this species, we provide illustrations of its scutellum (Fig. 17) and habitus of the female in dorsal view (Fig. 18).

### *Oobius longoi* (Siscaro, 1992)

Figures 22–23

*Avetianella longoi* Siscaro 1992: 206–211. Holotype female [UNCA], not examined.

Type locality: Grammichele, Catania Prov., Sicily, Italy.

*Avetianella longoi*: Trjapitzin 2001: 735 (key), 737–738 (taxonomic history, host associations); Wang et al. 2008: 1772–1777 (host associations, morphological and molecular data).

*Oobius longoi* (Siscaro): Noyes 2010: 662, 692.

**Material examined.** Australia, New South Wales, Corowa, 22.i.2006, Q. Wang, from eggs of *Phoracantha recurva* [1 ♀, 1 ♂, UCRC]. Portugal: Lisboa, Montijo, Pegões, viii.1992, P. Albino, M. R. Paiva, from eggs of *Phoracantha semipunctata* [9 ♀, 11 ♂, UCRC]. Viseu, Villa Cova à Coelheira, viii.1992, P. Albino, M. R. Paiva, from eggs of *P. semipunctata* [11 ♀, UCRC]. USA, California, Riverside Co., Riverside, University of California campus, Department of Entomology Insectary, laboratory culture on eggs of *P. semipunctata* on *Eucalyptus* sp.: 29.ix.1994, L. Hanks (originally from Australia) [4 ♀, UCRC]; 1998, S. McElfresh, J. Gould (originally from: Australia, Victoria, Melbourne, Bundoora, La Trobe Wildlife Sanctuary, i.1992, Q. Wang, from eggs of *P. semipunctata* on fallen *Eucalyptus* sp.) [25 ♀, 22 ♂, UCRC].

**Distribution.** Australia (indigenous); introduced (in some cases possibly unintentionally) into Hungary, Italy, Portugal, South Africa, Spain, USA (California), and Zambia (Trjapitzin 2001; Noyes 2014).

**Hosts.** *Phoracantha recurva* Newman and *P. semipunctata* (Fabricius) (Cerambycidae) in California, USA (Wang et al. 2008); its other longhorned beetle hosts in Australia are listed by Trjapitzin (2001) and Noyes (2014).

**Comments.** *Oobius longoi* is well known as an effective biological control agent and a successfully established parasitoid of *P. recurva* and *P. semipunctata* in California and elsewhere in the world (Hanks et al. 1995; Lühring et al. 2000; Trjapitzin 2001).

***Oobius minusculus* Triapitsyn & Petrice, sp. n.**

<http://zoobank.org/A7698FE3-D6BF-4AB1-B796-9D006B040D45>

Figures 11a, 12–16, 25, 31, 33–34

*Avetianella* sp.: Petrice et al. 2009: 179–180 (egg parasitoid of *A. subcinctus* in Livingston Co., Michigan, USA).

**Type material.** Holotype female [UCRC] on slide (Fig. 11a) with following four labels: “USA: Michigan, Clinton Co., near Bath, 42.812°N, 84.410°W, 255 m, parasitized *Agrilus subcinctus* Gory eggs collected 12.vii.2013, T.R. Petrice, emerged 16-22.vii.2013 in laboratory (Lansing, MI)”, “Mounted by V. V. Berezovskiy 2014 in Canada balsam”, [magenta] “*Oobius minusculus* Triapitsyn & Petrice HOLOTYPE ♀”, [database label] “Univ. Calif. Riverside Ent. Res. Museum UCRC ENT 142420”. The holotype is in good condition, complete, dissected under 3 coverslips.


Paratypes: USA, Michigan: Clinton Co. (same data as the holotype), 2 ♀ on points [MSUC, UCRC] and 1 ♀, 1 ♂ on slides [UCRC]. Ingham Co., Michigan State University Tree Research Center, 42°40'12"N, 84°28'12"W, 267 m, 14.viii.2014, T. R. Petrice, emerged in laboratory (East Lansing) from parasitized *Agrilus egenus* Gory eggs on black locust, *Robinia pseudoacacia*, twigs: emerged 22.viii.2014 [3 ♀ on points, MSUC, UCRC, USNM]; emerged 29.viii.2014 [3 ♀ on points, MSUC, UCRC, USNM, and 1 ♂ on slide, UCRC]; emerged 6.ix.2014 [1 ♀ on point, UCRC]; emerged 17.ix.2014 [1 ♂ on slide, UCRC].

**Description.** FEMALE (holotype). Body dark brown to black except scutellum and propodeum brown; scape and pedicel brown, flagellum light brown; legs whitish or pale yellowish with wide brown bands on coxae, femora, and tibiae.

Frontovertex and mesonotum with faint mesh-like or lineolate sculpture [very difficult to see in dry-mounted specimens, best observed in slide-mounted ones (as in Fig. 25)]. Pronotum, mesoscutum, axillae, and scutellum with short, dusky setae; scutellum also with a pair of long, fine setae near posterior margin.

Head (as in Fig. 15, collapsed when air-dried) with ocelli in an obtuse triangle, posterior ocellus a little less than its diameter away from eye margin. Transfacial and inner orbital sutures present. Mandible 3-dentate, the inner tooth with two denticles; maxillary palpus 4-segmented, labial palpus 1-segmented (i.e., palpal formula 4–1).

Antenna (Fig. 12) inserted below lower eye margin. Radicle about 0.3× total scape length, rest of scape slender, 4.5× as long as wide, a little wider in the middle, with



**Figures 9–16.** 9–10 *Oobius nearcticus* female (holotype), 9 antenna 10 clava 11a–11b: 11a holotype slide of *O. minusculus* 11b holotype slide of *O. whiteorum* 12–16 *Oobius minusculus* 12 antenna (holotype female) 13 mesosoma and metasoma (holotype female) 14 metasoma (paratype male) 15 head (paratype female) 16 pedicel and flagellum (paratype male).

faint longitudinal sculpture. Pedicel longer than any funicle segment; F1–F5 slightly transverse, F1–F4 subequal in length, F5 a little longer and slightly wider than long; F6 the longest funicle segment, longer than wide; F1–F5 without mps, and F6 with 2 mps. Clava 3-segmented, about 2.3× as long as wide and almost as long as funicle; first claval segment with 1 mps, second and third segments each with 3 mps.

Mesosoma a little shorter than gaster (Fig. 13). Mesoscutum about 1.7× as wide as long. Scutellum a little wider than long, a little shorter than mesoscutum; scutellar placoid sensilla closer to the posterior margin of scutellum and close to each other.

Wings (Fig. 31) not abbreviated, forewing extending far beyond apex of gaster. Forewing 2.1× as long as wide, hyaline; marginal setae very short; disc densely setose, linea calva interrupted posteriorly by an irregular row of setae, filum spinosum present. Hindwing 4.2× as long as wide, hyaline; longest marginal seta 0.3× maximum wing width.

Mesotibial spur a little longer than mesobasitarsus.

Ovipositor occupying a little more than 0.5× length of gaster, exerted markedly beyond gastral apex (by 0.2× own length) (Fig. 13); ovipositor length:metatibia length ratio 1.2:1. Outer plate of ovipositor with two subapical setae.


Measurements of the holotype (mm, as length or length:width). Body (of the dry-mounted specimen prior to slide-mounting): 0.462; mesosoma: 0.233; gaster: 0.245; ovipositor: 0.173. Antenna: radicle: 0.03; rest of scape: 0.103; pedicel: 0.045; F1: 0.012; F2: 0.012; F3: 0.011 (0.012); F4: 0.012; F5: 0.015; F6: 0.03; clava: 0.103. Forewing: 0.495:0.234; longest marginal seta: 0.021; hindwing: 0.357:0.085; longest marginal seta: 0.025.

Variation (paratypes). Body length 0.43–0.46 mm (dry-mounted specimens from *A. subcinctus*, Fig. 33) or 0.46–0.53 mm (critical-point dried specimens from *A. egenus*, Fig. 34). In the latter specimens, legs (except tarsi) are somewhat darker (mostly brown), scape (minus radicle) of the female antenna is about 5.0× as long as wide, and clava is about 2.5× as long as wide. Mandibles are identical for specimens reared from both host species, and there is no doubt that they are conspecific. In all specimens, F6 is sometimes slightly paler than other flagellomeres but not contrastingly, still almost concolorous or often concolorous.

MALE (paratype from *A. subcinctus*). Head dark brown, mesosoma and gaster dark brown to black except mesoscutum with a brownish tinge, base of gaster whitish; antenna with scape and pedicel brown to dark brown, flagellum light brown. Antenna (Fig. 16) with scape minus radicle 2.9× as long as wide; F2–F4 more or less subequal in length, F1 and F5 slightly longer, F6 the longest funicle segment; F2–F4 without mps, F1, F5, F6 and clava with mps; flagellar segments with very long setae (slightly longer than each funicle segment's width); clava entire, 2.6× as long as wide, a little wider than funicle segments. Mesosoma (Fig. 25) about as long as gaster. Forewing 2.0× as long as wide, hyaline. Genitalia (Fig. 14) typical for the genus.

Variation (paratypes from *A. egenus*). Body length 0.4–0.5 mm (critical-point dried specimens).

**Diagnosis.** This species is similar to the European *O. zahaikovitshi* Trjapitzin (Figs 19, 30), whose type locality is Zhuravlivka, Vinnytsia Oblast, Ukraine, where it was



**Figures 17–24.** 17–18 *Oobius depressus* female 17 dorsal scutellum (lectotype) 18 dorsal habitus (paralectotype) 19 *Oobius zahaikevitchi* female (environs of Volgograd, Krasnoarmeyskiy District, Volgograd Province, Russia), forewing 20–21 *Oobius hasmik* female (paratype) 20 antenna 21 forewing 22–23 *Oobius longoi* female (from University of California laboratory colony, Riverside, California, USA; of Australia origin), 22 antenna 23 forewing 24 *Oobius nearcticus* female (holotype), lateral habitus.

reared from eggs of *Agrilus viridis* (Linnaeus) on *Carpinus betulus* (Trjapitzin 1963). *Oobius zahaikovitshi* was recently well illustrated by Gumovsky et al. (2013). It was recorded from several European countries and *Agrilus* spp. hosts, listed by Trjapitzin and Volkovitsh (2011) and Noyes (2014). However, we are not absolutely confident that all these records are correct: it is quite possible that they might represent a complex of more than one cryptic species that are difficult to distinguish without supporting molecular data and thorough morphological studies based on good quality slide-mounted specimens. Proportions of funicle segments of the female antenna seem to be somewhat different between the specimens of *O. zahaikovitshi* from Ukraine illustrated by Trjapitzin (1963) and Gumovsky et al. (2013), in which F5 is about as long as wide, and the examined specimens from Volgograd Province of Russia, in which F5 is a little wider than long (Fig. 30).

*Oobius minusculus* differs from *O. zahaikovitshi* in having the palpal formula 4–1, a relatively smaller F5 of the female antenna and also by F6 being longer than wide and almost concolorous or often concolorous with other flagellomeres (Fig. 12). In contrast, the palpal formula for *O. zahaikovitshi* is 3–1, F5 is relatively larger, and F6 is about as long as wide and contrastingly lighter than other flagellomeres (Fig. 30), as also described and illustrated in Trjapitzin (1963) and Gumovsky et al. (2013).


*Oobius minusculus* is the only described native Nearctic species of *Oobius* s. str., as characterised by Noyes (2010) in having the outer plate of the ovipositor being relatively short and apically rounded with paired subapical setae (one long and one short), in which this new taxon fits well. In the key by Trjapitzin and Volkovitsh (2011) to the world species of *Oobius* (s. str.), it keys to *O. zahaikovitshi*. In Noyes (2010), *O. minusculus* tentatively keys (although it really does not key to any of the included Neotropical species) to the same couplet with *O. xochipili* Noyes and *O. zagan* Noyes from Costa Rica, from both of which it differs by F5 of the female antenna being much less transverse, just slightly wider than long (Fig. 12) whereas in *O. xochipili* and *O. zagan* F5 is anelliform, much wider than long (Noyes 2010).

**Etymology.** The name of this new taxon is an adjective referring to its small size.

**Hosts.** *Agrilus subcinctus* on ash (*Fraxinus* spp.) and *A. egenus* on black locust (*Robinia pseudoacacia*).

**Notes on biology.** Originally reported by Petrice et al. (2009) as *Avetianella* sp. that parasitized *A. subcinctus* eggs. The second author has never found this parasitoid to overwinter in *A. subcinctus* eggs. However, collections of *A. egenus* eggs found overwintering *O. minusculus* larvae in eggs. This species likely attacks other *Agrilus* spp. in North America, and has multiple generations per year.

**Comments.** The following specimens of *O. zahaikovitshi* were examined: Bulgaria, Plovdiv Prov., Klisura, 27.vi–8.vii.1975, A. Atanasov, from eggs of *Agrilus cuprescens* (Ménétriés) on *Rosa* sp. [1 ♂, BMNH] (det. V. A. Trjapitzin 1977). Russia, Volgograd Prov. (oblast'), Krasnoarmeyskiy District (rayon), environs of Volgograd, vi.1971, A. M. Makhmadziyoev (Makhmadziev), from eggs of *A. viridis* on *Acer tataricum* [1 ♀, BMNH; 5 ♀, UCRC] (det. V. A. Trjapitzin 1977 and 1975, respectively).



**Figures 25–32.** **25** *Oobius minusculus* male (paratype), mesosoma **26–29** *Oobius whiteorum* **26** antenna (holotype female) **27** a pair of wings (holotype female) **28** dorsal habitus (holotype female) **29** genitalia (paratype male) **30** *Oobius zahaikovitshi* female (environs of Volgograd, Krasnoarmeyskiy District, Volgograd Province, Russia), antenna **31** *Oobius minusculus* female (holotype), a pair of wings **32** *Oobius whiteorum* (paratype male), antenna.

***Oobius nearcticus* (Trjapitzin, 1977)**

Figures 9–10, 24

*Szelenyiola nearctica* Trjapitzin 1971: 160–161. Type locality: Blodgett Forest (8 mi. E. of Georgetown), University of California Blodgett Forest Research Station, El Dorado Co., California, USA.

*Oobius nearcticus* (Trjapitzin): Noyes 2010: 662, 668, 671.

**Type material examined.** Holotype female [EMEC] on point mount with following four labels: “UC Blodgett Forest 8 mi E. Georgetown, El Dorado Co., California Coll. F. M. Stephen 1970”, “Traps A-1094”, [red] “Holotypus *Szelenyiola nearctica* Trjapitzin”, “U.C. Berkeley EMEC 82,322”. Paratype female [EMEC] on point with following five labels: “UC Blodgett Forest 8 mi E. Georgetown, El Dorado Co., California Coll. F. M. Stephen 1970 A-1094 Traps”, “Head with appendages on slide No. 1955”, “Also forewing”, “Paratypus”, [red] “*Szelenyiola nearctica* Trjapitzin ♀”.

**Distribution.** USA (California) (Trjapitzin 1977).

**Hosts.** Unknown.

**Comments.** The holotype (Fig. 24) is missing its left hindwing and the apical 2/3 of the left forewing.

***Oobius whiteorum* Triapitsyn, sp. n.**

<http://zoobank.org/00F395AF-FD46-4102-A70E-BBD69B5176C1>

Figures 11b, 26–29, 32, 35

*Avetianella* sp.: Loerch and Cameron 1983: 1798–1799 (egg parasitoid, host information); Trjapitzin 2001: 738 (list).

**Type material.** Holotype female [UCRC] on slide (Fig. 11b) with following five labels: “USA, Pennsylvania, Venango Co., Bullion, 8.VII.1982, C. R. Loerch, Ex. *Agrilus anxius* Gory eggs”, “Mounted by V. V. Berezovskiy 2014 in Canada balsam”, “*Avetianella* sp. (Encyrtidae) Det. J. LaSalle”, [magenta] “*Oobius whiteorum* Triapitsyn HOLOTYPE ♀”, [database label] “Univ. Calif. Riverside Ent. Res. Museum UCRC ENT 401252”. The holotype is in good condition, complete, dissected under 3 coverslips.

Paratypes: same data as the holotype, 4 ♀ on points and 1 ♂ on slide [UCRC].

**Description.** FEMALE (holotype). Body somewhat flattened, dark brown to black; appendages brown except tarsi light brown; scape and pedicel a little darker than flagellum, and F6 just slightly lighter than other flagellar segments but still brown.

Frontoververtex and mesonotum with faint mesh-like sculpture [very difficult to see in dry-mounted specimens]. Pronotum, mesoscutum, axillae, and scutellum with short, dusky setae; scutellum also with a pair of long, fine setae near posterior margin.



**Figures 33–35.** 33–34 *Oobius minusculus* (paratype females) 33 dorsal habitus (from *Agrilus subcinctus*) 34 lateral habitus (from *Agrilus egenus*) 35 *Oobius whiteorum* (paratype female), dorsal habitus.

Head (Fig. 28) with ocelli in an obtuse triangle, posterior ocellus about its diameter away from eye margin. Transfacial and inner orbital sutures absent. Mandible 3-dentate; palpal formula 4–3.

Antenna (Fig. 26) inserted below lower eye margin. Radicle about 0.2× total scape length, rest of scape slender, 4.1–4.2× as long as wide, a little wider in the middle, with faint longitudinal sculpture. Pedicel longer than any funicle segment. F1–F3 about as long as wide, F4–F6 longer than wide; F1–F3 subequal, F4–F6 each progressively a little longer than the preceding funicle segment; F1–F3 without mps, F4 with 1 mps, F5 with 2 mps, and F6 with 3 or 4 mps. Clava 2.8× as long as wide, and slightly shorter than combined length of F2–F6; each claval segment with 3 mps; apical claval segment obliquely truncate ventrally.

Mesosoma (Fig. 28) shorter than gaster. Mesoscutum about 1.6× as wide as long. Scutellum wider than long, almost as long as mesoscutum.

Wings (Fig. 27) not abbreviated, forewing extending far beyond apex of gaster. Forewing 2.1× as long as wide, hyaline; marginal setae very short; disc densely setose, lineae calva interrupted posteriorly by rows of setae, filum spinosum present. Hindwing 3.7–3.8× as long as wide, hyaline; longest marginal seta 0.18× maximum wing width.

Mesotibial spur as long as mesobasitarsus.

Ovipositor occupying about 0.5× length of gaster, exerted markedly beyond gastral apex (by 0.36× total ovipositor length); ovipositor length:metatibia length ratio 1.3:1. Outer plate of ovipositor with 1 subapical seta.

Measurements of the holotype (mm, as length or length:width). Body (of the dry-mounted specimen prior to slide-mounting): 0.66; head: 0.19; mesosoma: 0.313; gaster: 0.35; ovipositor: 0.283. Antenna: radicle: 0.039; rest of scape: 0.151; pedicel: 0.06; F1: 0.021; F2: 0.021; F3: 0.021; F4: 0.028; F5: 0.035; F6: 0.044; clava: 0.155. Forewing: 0.677:0.314; longest marginal seta: 0.021; hindwing: 0.5:0.133; longest marginal seta: 0.024.

Variation (paratypes). Body (Fig. 35) length 0.66–0.75 mm (dry-mounted specimens).

MALE (paratype). Body length (of the dry-mounted specimen prior to slide-mounting) 0.66 mm. Head and mesosoma dark brown, gaster brown; scape and pedicel brown, flagellum light brown; legs light brown to brown. Antenna (Fig. 32) with scape minus radicle 3.2× as long as wide; funicle segments longer than wide, more or less subequal in length (F5 and particularly F6 slightly longer), F1 and F2 without mps, F3 with or without mps, F4–F6 and clava with mps; flagellar segments with very long setae (slightly longer than each funicle segment's width and about as long as width of clava); clava entire, 2.6–2.7× as long as wide, a little wider than funicle segments. Mesosoma about as long as gaster. Forewing 1.9× as long as wide, hyaline. Hindwing 3.5× as long as wide, hyaline. Genitalia (Fig. 29) typical for the genus.

**Diagnosis.** Among the Nearctic species of *Oobius*, *O. whiteorum* is most similar to *O. dahlsteni*, from which it differs by the proportions of the funicle segments of the female antenna, as indicated in the key. In Trjapitzin's (2001) key to the world species of the former genus *Avetianella* (s. str.) in which this new species mostly fits, as characterised by Noyes (2010) in having the outer plate of the ovipositor being conspicuously distally

elongate and ribbon-like and always with only a single subapical seta, it keys to *O. dahlsteni*. *Oobius whiteorum* differs from *O. depressus*, to which it is also somewhat similar, by a relatively less flattened body and by the much smaller body size in females; according to Girault (1916), the body length of the latter species is 1.15 mm. *Oobius whiteorum* differs from the North American species, but Neotropical species *O. hasmik* (Trjapitzin), known from Mexico (Trjapitzin 2001) and also Costa Rica (Noyes 2010), by the “closed” linea calva (Fig. 27) on the forewing (“open”, not interrupted, in *O. hasmik*, Fig. 21) and also by the different proportions of the scape of the female antenna (Figs 26 and 20, respectively). In Noyes (2010), *O. whiteorum* keys to the same couplet with *O. lutron* Noyes from Costa Rica and Brazil, from which it differs by each of F4–F6 of the female antenna being of different length and longer than wide (Fig. 26) whereas in *O. lutron* F4–F6 are subequal and each quadrate or hardly longer than broad (Noyes 2010).

**Host.** *Agrilus anxius* Gory on European white birch (*Betula pendula*).

**Etymology.** This species is named in honor of Lisa and Michael White of Chicago, Illinois, USA, good friends of the author’s family.

**Comments.** According to Loerch and Cameron (1983), additional voucher specimens of the egg parasitoids of *A. anxius* were deposited by them in PSUC; any of them belonging to this species are non-type specimens. Unfortunately, due to a renovation of the museum, point-mounted specimens in that collection are now inaccessible (A. Deans, personal communication).

The following paratypes [UCRC] of *O. hasmik* were examined, all collected at Las Barracas (~30 km E of Santiago, 23°28'02"N, 109°27'01"W, 50 m), Baja California Sur, Mexico: 1 ♀ on point with following five labels: “Mex. Baja Cal. Sur Las Barracas 17 - V - 1985”, “Coll. P. DeBach Pan trap”, “*Avetianella* ♀ Det. V. Trjapitzin May 1997”, [red] “Paratypus ♀ *Avetianella hasmik* Trjapitzin”, “Praep. micr. 22M” (an antenna, head, and a forewing were detached from this specimen; they are mounted on a slide with following two labels: “*Avetianella hasmik* ♀ Trjapitzin México: Baja California Sur, Las Barracas. Pan trap 17.V.1985 (Coll. P. DeBach) 22M Antena, cabeza, ala anterior”, [red] “Paratypus *Avetianella* ♀ *hasmik* Trjapitzin”); also 16 ♀, 1 ♂ on points, all collected by P. DeBach during 1985 and 1986, as indicated by Trjapitzin (2001).

### ***Oobius* sp.**

(Not included in the key)

*Oobius* sp. n.: Trjapitzin et al. 2008: 186 (record from Mexico).

*Oobius* sp. n. aff. *rudnevi* (S. Nowicki, 1928): Trjapitzin and Volkovitsh 2011: 674 (list, Mexico).

**Comments.** One female [UANL] of this undescribed species from Mexico, which has no host information, was mentioned by Trjapitzin and Volkovitsh (2011); however, they did not indicate the collecting locality so it is unknown from which part of that country it was found (Nearctic or Neotropical).

## Acknowledgments

We thank Natalie Dale-Skey Papillod (BMNH) and Robert L. Zuparko (EMEC) for the loans of specimens, Vladimir V. Berezovskiy (UCRC) for making excellent point- and slide-mounts, and Deborah L. Miller (USDA Forest Service, NRS, Lansing, Michigan, USA) for providing laboratory-reared specimens of *O. agrili*. Thanks to Taina Litwak (USDA-SEL) for adjusting the slide images and Dylan Johnston-Jordan for imaging *O. agrili*. Valuable comments by Robert Haack, Therese Poland, and two anonymous reviewers have contributed to the improvement of the manuscript. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the USDA.

## References

- Abell KJ, Bauer LS, Duan JJ, Van Driesche R (2014) Long-term monitoring of the introduced emerald ash borer (Coleoptera: Buprestidae) egg parasitoid, *Oobius agrili* (Hymenoptera: Encyrtidae), in Michigan, USA and evaluation of a newly developed monitoring technique. *Biological Control* 79 (1): 36–42. doi: 10.1016/j.biocontrol.2014.08.002
- Annecke DP (1967) Three new southern African species of *Oobius* Trjapitzin, 1963 (Hymenoptera: Encyrtidae). *Journal of Natural History* 1(3): 319–325. doi: 10.1080/00222936700770291
- Bauer LS, Duan JJ, Gould J (2014) XVII Emerald ash borer (*Agrilus planipennis* Fairmaire) (Coleoptera: Buprestidae). In: Van Driesche R, Reardon R (Eds) The use of classical biological control to preserve forests in North America. United States Department of Agriculture, Forest Service, FHTET-2013-02, Morgantown, West Virginia, 189–209. <http://www.fs.fed.us/foresthealth/technology/pdfs/FHTET-2013-2.pdf>
- Bauer LS, Duan JJ, Gould JG, Van Driesche RG (in press) Progress in the classical biological control of *Agrilus planipennis* Fairmaire (Coleoptera: Buprestidae) in North America. *The Canadian Entomologist*.
- Bray AM, Bauer LS, Poland TM, Haack RA, Cognato AI, Smith JJ (2011) Genetic analysis of emerald ash borer (*Agrilus planipennis* Fairmaire) populations in Asia and North America. *Biological Invasions* 13: 2869–2887. doi: 10.1007/s10530-011-9970-5
- Gibson GAP (1997) Chapter 2. Morphology and terminology. In: Gibson GAP, Huber JT, Woolley JB (Eds) Annotated keys to the genera of Nearctic Chalcidoidea (Hymenoptera). NRC Research Press, Ottawa, Ontario, Canada, 16–44.
- Girault AA (1916) Descriptions of and observations on some chalcidoid Hymenoptera – II. *The Canadian Entomologist* 48(10): 337–344. doi: 10.4039/Ent48337-10
- Gordh G, Trjapitzin VA (1981) Taxonomic studies of the Encyrtidae with descriptions of new species and a new genus (Hymenoptera, Chalcidoidea). *University of California Publications in Entomology* 93: 1–64.
- Gumovsky AV, Simutnik SA, Prokhorov AV (2013) Life-history review of *Oobius zabaikewitshi* Trjapitzin, 1963 (Hymenoptera: Encyrtidae), an egg parasitoid of jewel beetles (Coleoptera: Buprestidae). *Russian Entomological Journal* 22(3): 181–188.

- Haack RA, Jendek E, Liu HP, Marchant K, Petrice T, Poland TM, Ye H (2002) The emerald ash borer: a new exotic pest in North America. Newsletter of the Michigan Entomological Society 47(3–4): 1–5.
- Hanks LM, Gould JR, Paine TD, Millar JG, Wang Q (1995) Biology and host relations of *Avetianella longoi* (Hymenoptera: Encyrtidae), an egg parasitoid of the eucalyptus longhorned borer (Coleoptera: Cerambycidae). Annals of the Entomological Society of America 88(5): 666–671. doi: 10.1093/aesa/88.5.666
- Liao DX, Li XL, Pang XF, Chen TL (1987) Hymenoptera: Chalcidoidea (1). Economic Insect Fauna of China No 34: x+241 pp.
- Liu HP, Bauer LS, Miller DL, Zhao TH, Gao RT, Song L, Luan Q, Jin R, Gao C (2007) Seasonal abundance of *Agrilus planipennis* (Coleoptera: Buprestidae) and its natural enemies *Oobius agrili* (Hymenoptera: Encyrtidae) and *Tetrastichus planipennis* (Hymenoptera: Eulophidae) in China. Biological Control 42: 61–71. doi: 10.1016/j.biocontrol.2007.03.011
- Loerch CR, Cameron EA (1983) Natural enemies of immature stages of the bronze birch borer, *Agrilus anxius* (Coleoptera: Buprestidae), in Pennsylvania. Environmental Entomology 12: 1798–1801. doi: 10.1093/ee/12.6.1798
- Luhring KA, Paine TD, Millar JG, Hanks LM (2000) Suitability of the eggs of two species of eucalyptus longhorned borers (*Phoracantha recurva* and *P. semipunctata*) as hosts for the parasitoid *Avetianella longoi* Siscaro. Biological Control 19: 95–104. doi: 10.1006/bcon.2000.0853
- MacRae TC (1991) The Buprestidae (Coleoptera) of Missouri. Insecta Mundi 5: 101–126.
- Mapbiocontrol (2014) Emerald ash borer biocontrol release and recovery. <http://www.mapbiocontrol.org/>
- Nelson GH, Walters GC, Haines RD, Bellamy CL (2008) A catalog and bibliography of the Buprestoidea of America North of Mexico. The Coleopterists Society Special Publ. No. 4: 1–274.
- Noyes JS (2010) Encyrtidae of Costa Rica (Hymenoptera: Chalcidoidea), 3. Subfamily Encyrtinae: Encyrtini, Echthroplexiellini, Discodini, Oobiini and Ixodiphagini, parasitoids associated with bugs (Hemiptera), insect eggs (Hemiptera, Lepidoptera, Coleoptera, Neuroptera) and ticks (Acari). Memoirs of the American Entomological Institute 84: 1–848.
- Noyes JS (2014) Universal Chalcidoidea database. WWW publication. The Natural History Museum, London. <http://www.nhm.ac.uk/research-curation/projects/chalcidoids/index.html> [accessed 12.i.2015]
- Petrice TR, Haack RA, Strazanac JA, Lelito JP (2009) Biology and larval morphology of *Agrilus subcinctus* (Coleoptera: Buprestidae), with comparisons to the emerald ash borer, *Agrilus planipennis*. The Great Lakes Entomologist 42: 173–184.
- Siscaro G (1992) *Avetianella longoi* sp. n. (Hymenoptera Encyrtidae) egg parasitoid of *Phoracantha semipunctata* F. (Coleoptera Cerambycidae). Bollettino di Zoologia Agraria e Bachicoltura (2) 24(2): 205–212.
- Trjapitzin VA (1963) Species of the genus *Oobius* gen. n. (Hymenoptera, Encyrtidae) in the USSR. Acta Entomologica Musei Nationalis Pragae [Sborník Entomologického Oddelení Národního Musea v Praze] 35: 543–547.
- Trjapitzin VA (1968) [A survey of the encyrtid fauna (Hym. Encyrtidae) of the Caucasus]. Trudy Vsesoyuznogo Entomologicheskogo Obshchestva 52: 43–125. [In Russian]

- Trjapitzin VA (1971) [A Nearctic representative of the genus *Avetianella* Trjapitzin, 1968 (Hymenoptera, Encyrtidae)]. Entomologicheskoe Obozrenie 50 (4): 890–892. [In Russian]
- Trjapitzin VA (1977) New genera and species of parasitic Hymenoptera of the family Encyrtidae (Hymenoptera: Chalcidoidea). Folia Entomologica Hungarica 30 (1): 153–166.
- Trjapitzin VA (2001) [A review of encyrtids of the genus *Avetianella* Trjapitzin, 1968 (Hymenoptera, Encyrtidae) of the world fauna with description of a new species from Mexico]. Entomologicheskoe Obozrenie 80 (3): 734–739. [In Russian]
- Trjapitzin VA, Gordh G (1984) [Taxonomic notes on the Neotropic genus *Habrolepoidea* (Hymenoptera, Encyrtidae) and on species erroneously referred to it]. Zoologicheskii Zhurnal 63(8): 1273–1277. [In Russian]
- Trjapitzin VA, Myartseva SN, Ruíz-Cancino E, Coronado-Blanco JM (2008) Clave de géneros de Encyrtidae (Hymenoptera: Chalcidoidea) de México y un catálogo de las especies. Serie Avispas parasíticas de plagas y otros insectos. Universidad Autónoma de Tamaulipas, Ciudad Victoria, Tamaulipas, México, 265 pp.
- Trjapitzin VA, Volkovitsh MG (2011) A review of species of the genus *Oobius* Trjapitzin, 1963 (Hymenoptera, Encyrtidae) egg parasitoids of jewel beetles, longicorn beetles (Coleoptera, Buprestidae, Cerambycidae), and robber flies (Diptera, Asilidae). Entomological Review 91(5): 670–676. [English translation, originally published in Russian in Entomologicheskoe Obozrenie 90 (1): 226–234]
- Zhang Y, Huang D (2004) A review and an illustrated key to genera of Encyrtidae (Hymenoptera: Chalcidoidea) from China. Science Press, Beijing, China, 166 pp.
- Zhang Y-Z, Huang D-W, Zhao T-H, Liu H-P, Bauer LS (2005) Two new species of egg parasitoids (Hymenoptera: Encyrtidae) of wood-boring beetle pests from China. Phytoparasitica 33(3): 253–260. doi: 10.1007/BF02979863