

# Phytoseiid mites from tropical fruit trees in Bahia State, Brazil (Acari, Phytoseiidae)

Izabel Vieira de Souza<sup>1</sup>, Poliane Sá Argolo<sup>1</sup>, Manoel Guedes Correa Gondim Júnior<sup>2</sup>,  
Gilberto José de Moraes<sup>3</sup>, Maria Aparecida Leão Bittencourt<sup>1</sup>,  
Anibal Ramadan Oliveira<sup>1</sup>

**1** Universidade Estadual de Santa Cruz, Rodovia Ilhéus – Itabuna, km 16, 45.662-900, Ilhéus, BA, Brazil  
**2** Universidade Federal Rural de Pernambuco, Área de Fitossanidade, 52171-900, Recife, PE, Brazil  
**3** CNPq Researcher; Departamento de Entomologia e Acarologia, Escola Superior de Agricultura “Luiz de Queiroz”, Universidade de São Paulo, 13418-900, Piracicaba, SP, Brazil

Corresponding author: Anibal Ramadan Oliveira ([aroliveir@gmail.com](mailto:aroliveir@gmail.com))

---

Academic editor: F. Faraji | Received 27 February 2015 | Accepted 17 September 2015 | Published 9 October 2015

---

<http://zoobank.org/BA561653-53E9-448B-A21B-B72417B4E246>

---

**Citation:** Souza IV, Argolo PS, Gondim Jr MGC, Moraes GJ, Bittencourt MAL, Oliveira AR (2015) Phytoseiid mites from tropical fruit trees in Bahia State, Brazil (Acari, Phytoseiidae). ZooKeys 533: 99–131. doi: 10.3897/zookeys.533.5981

---

## Abstract

The cultivation of tropical fruit trees has grown considerably in the state of Bahia, northeastern Brazil. Some of these have been severely attacked by phytophagous mites, which are usually controlled by the use of chemical pesticides. However, there is today a growing interest for the adoption of less aggressive measures of pest control, as for example the use of predatory mites. Most of the plant-inhabiting predatory mites belong to the family Phytoseiidae. The objective of this paper is to report the phytoseiid species found in an intensive survey conducted on cultivated tropical fruit trees in fifteen localities of the southern coast of Bahia. Measurements of relevant morphological characters are provided for each species, to complement the understanding of the morphological variation of these species. Twenty-nine species of sixteen genera were identified. A key was elaborated to assist in the separation of these species. Fifteen species are reported for the first time in the state, raising to sixty-six the number of species of this family now known from Bahia. Seventy-two percent of the species collected belong to Amblyseinae, followed by Typhlodrominae (21%) and Phytoseiinae (7%). The most diverse genus was *Amblyseius*. *Amblyseius operculatus* De Leon was the most frequent and abundant species. Studies should be conducted to evaluate the possible role of the most common predators as control agents of the phytophagous mites co-occurring with them.

**Keywords**

Acarology, Predatory mites, Taxonomy, Biological Control

**Introduction**

Cultivation of tropical fruit trees has grown considerably in the state of Bahia, north-eastern Brazil, in the last years (Santos-Serejo et al. 2009). Several mite species have been reported on those plants, some causing economic losses (Moraes and Flechtmann 2008). These are usually controlled by the use of chemical pesticides.

However, there is today a growing interest on the use of less aggressive and less toxic strategies to control those organisms. Predatory mites of the family Phytoseiidae are considered important biological control agents of pest mites, and some phytoseiids are commercially available for the control of pest mites in several countries (Hoy 2011). There is an interest to implement the use of phytoseiids for the biological control of pest mites in orchards of tropical fruit trees in coastal Bahia, and the determination of the naturally occurring phytoseiids in that area is considered the first step in the implementation of a biological control program.


The objective of this paper is to report the phytoseiid species found in an intensive survey conducted on cultivated tropical fruit trees in the southern coastal region of Bahia, providing a key to help the separation of the species collected.

**Materials and methods**

Samples were collected from March 2007 to January 2010 in fifteen localities of eight municipalities (Figure 1, Table 1). These consisted mainly of leaves and, when present, flowers and fruits of 21 species of tropical fruit trees (Table 2).

Phytoseiid mites were mounted in Hoyer's medium, identified and measured under a phase-contrast microscope (Motic® B3 Professional Series). Under each species mentioned in the Results section, information concerning the specimens examined is given in the following order: sampling locality, plant species, month and year of the collection, number and sex of specimens. Measurements are given in micrometers, corresponding to the average for the structures measured followed in parentheses by the respective ranges. Numbers of teeth on the fixed and movable cheliceral digits do not include the respective apical teeth. Setae not referred to in the Results section should be considered absent.

Idiosomal setal notation adopted is that of Lindquist and Evans (1965), as applied to phytoseiids by Rowell et al. (1978) and Chant and Yoshida-Shaul (1989) for the dorsal surface, and by Chant and Yoshida-Shaul (1991) for the ventral surface. Macrosetal notation is that of Muma et al. (1970). The system of classification follows that of Chant and McMurtry (2007). The name adopted for each species is that mentioned in the Phytoseiidae Database (Demite et al. 2014), followed by the names attributed



**Figure 1.** Map locating the municipalities in the State of Bahia, Brazil, where collections were conducted.

to the species in the original description, in the catalog of Moraes et al. (2004) and in the comprehensive work of Chant and McMurtry (2007).

Voucher specimens were deposited in the mite reference collection of Universidade Estadual de Santa Cruz (UESC), Ilhéus, Bahia.

**Table 1.** Localities in the State of Bahia, Brazil, from which phytoseiid mites were collected between March 2007 and January 2010.

Municipality	Sampling locality	Coordinates
Camamu	Fazenda Bela Vista	13°58'35"S, 39°09'23"W
Ilhéus	UESC	14°47'53"S, 39°10'20"W
	Fazenda Terra Nova	14°43'52"S, 39°09'16"W
	Sítio Agrotropical	14°47'00"S, 39°14'42"W
	Fazenda Monte Alegre	14°43'29"S, 39°20'42"W
Ituberá	Colônia de Japoneses	13°46'29"S, 39°11'04"W
	Fazenda Frupical	13°45'23"S, 39°10'58"W
	Fazenda Kamuí	13°45'11"S, 39°10'50"W
Taperoá	Fazenda Nossa Senhora Auxiliadora	13°33'52"S, 39°12'06"W
Una	Estação Lemos Maia-CEPLAC	15°21'18"S, 38°59'56"W
Uruçuca	Fazenda Liberdade	14°35'52"S, 39°19'55"W
Valença	Fazenda Barra	13°21'05"S, 39°19'57"W
	Fazenda Formiga	13°18'01"S, 39°15'02"W
	Sítio Sabino	13°18'59"S, 39°15'46"W
	Sítio São Jorge	13°17'23"S, 39°18'05"W

**Table 2.** Species of tropical fruit trees from which phytoseiid mites were collected in the State of Bahia, Brazil, between March 2007 and January 2010.

Family	Plant species
Anacardiaceae	<i>Anacardium occidentale</i> L.
	<i>Mangifera indica</i> L.
	<i>Spondias mombin</i> L.
Annonaceae	<i>Annona muricata</i> L.
	<i>Annona squamosa</i> L.
Arecaceae	<i>Cocos nucifera</i> L.
	<i>Elaeis guineensis</i> Jacq.
	<i>Euterpe oleracea</i> Mart.
Caricaceae	<i>Carica papaya</i> L.
Ebenaceae	<i>Diospyros kaki</i> L.
Lauraceae	<i>Persea americana</i> L.
Malpighiaceae	<i>Malpighia emarginata</i> D.C.
Moraceae	<i>Artocarpus integrifolia</i> Lam.
Musaceae	<i>Musa sapientum</i> L.
Myrtaceae	<i>Psidium guajava</i> L.
	<i>Syzygium malaccense</i> (L.) Merr. & L.M. Perry
Passifloraceae	<i>Passiflora edulis</i> Sims.
Rubiaceae	<i>Genipa americana</i> L.
Sapotaceae	<i>Pouteria caimito</i> (Ruiz & Pav.) Radlk.
Sterculiaceae	<i>Theobroma cacao</i> L.
	<i>Theobroma grandiflorum</i> Schum.

## Results

In total, 564 phytoseiid specimens were collected, belonging to twenty-nine species of sixteen genera, as subsequently reported. Most specimens were collected from *C. nucifera* (41%), followed by *T. cacao* (24%), and *P. guajava* (16%) (Tables 3 and 4).

### Amblyseiinae Muma

#### *Amblydromalus manihoti* (Moraes)

*Amblyseius manihoti* Moraes et al., 1994: 211.

*Typhlodromalus manihoti*: Moraes et al. 2004: 200.

*Amblydromalus manihoti*: Chant and McMurtry 2007: 117.

**Specimens examined.** UESC, *C. papaya*, I-2008 (2♀); Fazenda Bela Vista, *T. cacao*, XI-2007 (2♀); Fazenda Frupical, *T. grandiflorum*, VII-2007 (1♀); Fazenda Liberdade, *C. papaya*, V-2007 (2♀).

**Table 3.** Numbers of specimens and of species of phytoseiid mites collected on each sampled plant species in the State of Bahia, Brazil, between March 2007 and January 2010.

Family	Plant species	Nr. *	Specimens	Species
Anacardiaceae	<i>Anacardium occidentale</i>	1	3	3
	<i>Mangifera indica</i>	2	15	6
	<i>Spondias mombin</i>	3	1	1
Annonaceae	<i>Annona muricata</i>	4	12	4
	<i>Annona squamosa</i>	5	2	1
Arecaceae	<i>Cocos nucifera</i>	6	234	17
	<i>Elaeis guineensis</i>	7	7	5
	<i>Euterpe oleracea</i>	8	6	6
Caricaceae	<i>Carica papaya</i>	9	9	4
Ebenaceae	<i>Diospyros kaki</i>	10	3	3
Lauraceae	<i>Persea americana</i>	11	2	1
Malpighiaceae	<i>Malpighia emarginata</i>	12	1	1
Moraceae	<i>Artocarpus integrifolia</i>	13	6	3
Musaceae	<i>Musa sapientum</i>	14	9	8
Myrtaceae	<i>Psidium guajava</i>	15	92	13
	<i>Syzygium malaccense</i>	16	1	1
Passifloraceae	<i>Passiflora edulis</i>	17	5	2
Rubiaceae	<i>Genipa americana</i>	18	3	1
Sapotaceae	<i>Pouteria caimito</i>	19	2	1
Sterculiaceae	<i>Theobroma cacao</i>	20	137	15
	<i>Theobroma grandiflorum</i>	21	14	6

\*Corresponding to plant species numbers given in Table 4.

**Table 4.** Numbers of phytoseiid mites and diversity of host plants on which each species was collected in the State of Bahia, Brazil, between March 2007 and January 2010.

Phytoseiid species	Nr. of specimens	Plant species nr.*
<b>Amblyseinae</b>		
<i>Amblyseius operculatus</i>	133	1, 2, 4, 6, 7, 10, 11, 12, 13, 14, 15, 20, 21
<i>Amblyseius perditus</i>	57	2, 6, 7, 13, 14, 15, 20, 21
<i>Iphiseiodes metapodalis</i>	53	2, 4, 5, 6, 7, 8, 14, 15, 20, 21
<i>Iphiseiodes zuluagai</i>	47	1, 2, 4, 6, 7, 8, 9, 19, 20, 21
<i>Amblyseius aeralis</i>	32	6, 9, 15, 20, 21
<i>Amblyseius tamatavensis</i>	32	6, 8, 14, 17
<i>Amblyseius igarassuensis</i>	28	6, 15, 20
<i>Typhlodromips theobromae</i>	15	2, 20
<i>Amblyseius impeltatus</i>	10	15
<i>Iphiseiodes setillus</i>	8	8, 20
<i>Paraamblyseius multicircularis</i>	8	8, 20
<i>Amblydromalus manihoti</i>	7	9, 20, 21
<i>Typhlodromalus peregrinus</i>	7	6, 9, 15, 20
<i>Proprioseiopsis neotropicus</i>	6	6, 10, 14, 20
<i>Proprioseiopsis dominigos</i>	4	6, 14, 15
<i>Arrenoseius urquharti</i>	3	6, 14
<i>Proprioseiopsis ovatus</i>	3	8, 14, 17
<i>Paraphytoseius orientalis</i>	2	15
<i>Phytoscutus sexpilis</i>	1	15
<i>Proprioseiopsis pentagonalis</i>	1	20
<i>Typhlodromips mangleae</i>	1	6
<b>Phytoseiinae</b>		
<i>Phytoseius latinus</i>	21	15
<i>Phytoseius woodburyi</i>	12	6, 15
<b>Typhlodrominae</b>		
<i>Leonseius regularis</i>	56	2, 3, 4, 6, 7, 13, 15, 16, 18, 20
<i>Cocoseius palmarum</i>	9	6
<i>Cocoseius elsalvador</i>	5	6, 20
<i>Metaseiulus ferlai</i>	1	1
<i>Typhlodromina subtropica</i>	1	10
<i>Typhlodromus transvaalensis</i>	1	6

\*Corresponding to plant species numbers given in Table 3.

**Female.** Four specimens measured. Dorsal shield 328 (320–338) long, 214 (205–223) wide, *j1* 29 (28–31), *j3* 38 (35–41), *j4* 10 (8–11), *j5* 9 (8–10), *j6* 12 (11–12), *J2* 14 (12–15), *J5* 9 (7–10), *z2* 11 (10–13), *z4* 12 (11–14), *z5* 9 (8–10), *Z1* 14 (13–15), *Z4* 14 (13–17), *Z5* 50 (47–51), *s4* 39 (35–44), *S2* 16 (15–17), *S4* 15 (14–16), *S5* 13 (11–14), *r3* 13 (12–13), *RI* 13 (11–14); distances between *St1–St3* 58 (55–60), *St2–St2* 67 (64–73) and *St5–St5* 69 (66–71); ventrianal shield 96 (92–105) long, 59 (56–

62) wide at level of ZV2 and 67 (64–70) wide at level of anus; movable cheliceral digit 32 (31–32) long, with 4 teeth; fixed cheliceral digit 26 (25–26) long, with 11 teeth; calyx of spermatheca 20 (16–23) long; *Sge I* 36 (34–38), *Sge II* 34 (32–37), *Sge III* 38 (36–41), *Sti III* 34 (32–36), *Sge IV* 61 (56–66), *Sti IV* 46 (40–51), *St IV* 80 (75–82).

**Remarks.** Measurements of the specimens collected are similar to those of the original description. They also agree with those of Moraes et al. (2013), except for the shorter Z5 [64 (52–70) in the latter].

### *Amblyseius aerialis* (Muma)

*Amblyseiopsis aerialis* Muma, 1955: 264.

*Amblyseius aerialis*: Moraes et al. 2004: 13.

*Amblyseius aerialis*: Chant and McMurtry 2007: 75.

**Specimens examined.** UESC, *T. grandiflorum*, IV-2008 (1♀); Fazenda Bela Vista, *C. nucifera*, IV-2007 (12♀), *P. guajava*, VII-2007 (1♀), *T. cacao*, XI-2007 (1♀); Fazenda Frupical, *T. grandiflorum*, VII-2007 (2♀); Fazenda Liberdade, *C. papaya*, V-2007 (1♀); Fazenda Nossa Senhora Auxiliadora, *C. papaya*, VII-2007 (1♀), *C. nucifera*, XI-2007 (2♀); Fazenda Terra Nova, *C. nucifera*, V-2007 (5♀); Sítio Agrotropical, *C. nucifera*, III-2007 (1♂); Sítio São Jorge, *C. nucifera*, XI-2007 (1♀, 4♂).

**Female.** Ten specimens measured. Dorsal shield 386 (364–422) long, 284 (266–309) wide, *j1* 34 (30–37), *j3* 55 (52–59), *j4* 5 (4–8), *j5* 5 (4–6), *j6* 7 (6–8), *J2* 9 (7–11), *J5* 8 (7–8), *z2* 14 (13–16), *z4* 9 (8–13), *z5* 6 (5–7), *Z1* 10 (10–12), *Z4* 133 (120–143), *Z5* 296 (271–315), *s4* 108 (105–111), *S2* 12 (11–14), *S4* 12 (11–14), *S5* 13 (12–13), *r3* 16 (14–17), *RI* 12 (11–13); distances between *St1–St3* 65 (63–71), *St2–St2* 83 (79–88) and *St5–St5* 81 (74–87); ventrianal shield 131 (120–142) long, 85 (78–91) wide at level of ZV2 and 73 (69–77) wide at level of anus; movable cheliceral digit 42 (39–44) long, with 4 teeth; fixed cheliceral digit 34 (33–35) long; calyx of spermatheca 18 (16–20) long; *Sge I* 44 (42–48), *Sge II* 42 (39–45), *Sge III* 61 (55–67), *Sti III* 40 (37–43), *Sge IV* 131 (118–138), *Sti IV* 90 (83–94), *St IV* 80 (75–84).

**Male.** Five specimens measured. Dorsal shield 309 (294–320) long, 217 (189–248) wide, *j1* 31 (28–32), *j3* 44 (43–45), *j4* 5 (5–6), *j5* 5 (4–6), *j6* 8 (6–9), *J2* 9 (8–10), *J5* 8 (7–8), *z2* 14 (13–15), *z4* 13 (12–13), *z5* 6 (5–7), *Z1* 10 (9–11), *Z4* 103 (102–105), *Z5* 229 (215–241), *s4* 85 (80–90), *S2* 12 (11–14), *S4* 9 (7–11), *S5* 10 (6–12), *r3* 17 (14–20), *RI* 12 (11–13); ventrianal shield 131 (122–137) long and 160 (156–162) wide at anterior corners; shaft of spermatodactyl 19 (16–20) long; *Sge I* 37 (32–39), *Sge II* 35 (30–39), *Sge III* 40 (37–41), *Sti III* 34 (32–35), *Sge IV* 82 (75–90), *Sti IV* 63 (56–70), *St IV* 61 (55–66).

**Remarks.** The calyx of spermatheca of the specimens collected is about 1.5 times longer than reported by Denmark and Muma (1989) (11–12), but it is comparable to measurements of Brazilian specimens given by Gondim Jr. and Moraes (2001).

***Amblyseius igarassuensis* Gondim Jr. & Moraes**

*Amblyseius igarassuensis* Gondim Jr. & Moraes, 2001: 71.

*Amblyseius igarassuensis*: Moraes et al. 2004: 30.

*Amblyseius igarassuensis*: Chant and McMurtry 2007: 78.

**Specimens examined.** UESC, *P. guajava*, I-2008 (3♂); Fazenda Bela Vista, *C. nucifera*, IX-2007 (2♀, 3♂); Fazenda Monte Alegre, *T. cacao*, V-2007 (3♀), I-2008 (3♀, 1♂), fruit, IX-2007 (1♀); Fazenda Terra Nova, *T. cacao*, IX-2007 (1♀), I-2008 (6♀, 5♂).

**Female.** Eight specimens measured. Dorsal shield 331 (307–346) long, 216 (197–241) wide, *j1* 25 (23–26), *j3* 37 (35–39), *j4* 8 (6–9), *j5* 7 (6–8), *j6* 10 (10–11), *J2* 10 (9–11), *J5* 9 (8–10), *z2* 11 (9–12), *z4* 11 (11–12), *z5* 7 (6–7), *Z1* 10 (9–11), *Z4* 64 (60–66), *Z5* 141 (130–155), *s4* 59 (56–61), *S2* 11 (10–13), *S4* 10 (10–11), *S5* 10 (9–11), *r3* 12 (11–14), *R1* 10 (9–11); distances between *St1–St3* 62 (60–63), *St2–St2* 66 (60–68) and *St5–St5* 62 (59–64); ventrianal shield 110 (102–118) long, 75 (68–79) wide at level of *ZV2* and 65 (62–70) wide at level of anus; movable cheliceral digit 34 (32–35) long, with 3 teeth; fixed cheliceral digit 29 (25–30) long; calyx of spermatheca 10 (9–13) long; *Sge I* 35 (34–38), *Sge II* 34 (33–37), *Sge III* 33 (30–35), *Sti III* 28 (26–29), *Sge IV* 57 (55–60), *Sti IV* 44 (41–46), *St IV* 57 (51–61).

**Male.** Seven specimens measured. Dorsal shield 252 (243–261) long and 158 (141–174) wide, *j1* 23 (17–28), *j3* 32 (31–34), *j4* 8 (7–9), *j5* 8 (7–8), *j6* 9 (9–10), *J2* 9 (8–10), *J5* 7 (6–8), *z2* 10 (9–12), *z4* 10 (8–11), *z5* 8 (7–8), *Z1* 10 (9–11), *Z4* 48 (41–52), *Z5* 109 (90–138), *s4* 41 (38–44), *S2* 11 (9–12), *S4* 10 (8–11), *S5* 8 (7–10), *r3* 12 (11–13), *R1* 10 (8–11); ventrianal shield 103 (98–110) long and 133 (122–150) wide at anterior corners; shaft of spermatodactyl 18 (17–20) long; *Sge I* 30 (29–34), *Sge II* 28 (26–31), *Sge III* 27 (23–32), *Sti III* 23 (21–27), *Sge IV* 39 (35–47), *Sti IV* 32 (27–38), *St IV* 47 (39–60).

**Remarks.** Measurements of females of this species are similar to those of the original description, except for the longer *Z5* (91–118 in the original description). This is the first record of this species in Bahia, and the first report of the measurements of males.

***Amblyseius impeltatus* Denmark & Muma**

*Amblyseius impeltatus* Denmark & Muma, 1973: 241.

*Amblyseius impeltatus*: Moraes et al. 2004: 30.

*Amblyseius impeltatus*: Chant and McMurtry 2007: 78.

**Specimens examined.** UESC, *P. guajava*, VII-2007 (10♀).

**Female.** Six specimens measured. Dorsal shield 367 (335–410) long, 228 (207–256) wide, *j1* 18 (16–19), *j3* 25 (22–28), *j4* 8 (8–9), *j5* 7 (4–8), *j6* 8 (7–11), *J2* 10 (8–13), *J5* 7 (5–8), *z2* 13 (12–15), *z4* 11 (10–12), *z5* 8 (7–9), *Z1* 12 (10–13), *Z4* 31



(30–33), *Z5* 125 (113–133), *s4* 42 (41–44), *S2* 13 (12–14), *S4* 12 (10–14), *S5* 10 (9–11), *r3* 12 (11–12), *RI* 12 (11–12); distances between *St1–St3* 53 (51–54), *St2–St2* 68 (66–69) and *St5–St5* 78 (77–82); ventrianal shield 90 (85–100) long, 80 (74–86) wide at level of *ZV2* and 68 (65–72) wide at level of anus; movable cheliceral digit 30 long, with 3 teeth; fixed cheliceral digit 27 (26–29) long; calyx of spermatheca 6 (5–6) long; *Sge I* 30 (29–30), *Sge II* 32 (31–33), *Sge III* 38 (36–40), *Sti III* 26 (25–27), *Sge IV* 63 (61–65), *Sti IV* 47 (45–48), *St IV* 49 (47–52).

**Remarks.** Measurements of the specimens collected fit those of the original description. This is the first record of this species in Bahia.

### *Amblyseius operculatus* De Leon

*Amblyseius operculatus* De Leon, 1967: 26.

*Amblyseius operculatus*: Moraes et al. 2004: 45.

*Amblyseius operculatus*: Chant and McMurtry 2007: 80.

**Specimens examined.** UESC, *M. sapientum* VII-2007 (1♀), *C. nucifera*, VII-2007 (4♀), I-2008 (3♂), IV-2008 (1♂), *A. muricata*, VII-2007 (1♀, 1♂), *E. guineensis*, VII-2007 (1♀), *P. guajava*, VII-2007 (3♀, 2♂), *A. integrifolia*, VII-2007 (1♀), *T. grandiflorum*, VIII-2007 (1♀), *P. americana*, I-2008 (2♀); CEPLAC, *C. nucifera*, IV-2008 (1♀), fruits, VI-2008 (1♀), IV-2009 (1♀), V-2009 (1♂), VII-2009 (2♀, 4♂), VIII-2009 (7♀), IX-2009 (2♀, 1♂), X-2009 (5♀, 1♂), XI-2009 (5♀, 1♂), XII-2009 (10♀, 4♂), I-2010 (3♀, 1♂); Colônia de Japoneses, *C. nucifera*, XI-2007 (2♀, 4♂); Fazenda Barra, *A. occidentale*, VIII-2007 (1♀), *T. cacao*, XI-2007 (2♀); Fazenda Bela Vista, *C. nucifera*, IV-2007 (1♀), IX-2007 (2♀); Fazenda Formiga, *C. nucifera*, VIII-2007 (5♀), fruit, VIII-2007 (1♀); Fazenda Frupical, *M. sapientum*, VII-2007 (1♀), *T. cacao*, VII-2007 (3♀), fruit, VII-2007 (1♀); Fazenda Liberdade, *T. cacao*, V-2007 (1♀); Fazenda Monte Alegre, *P. guajava*, V-2007 (1♀), IX-2007 (1♀), V-2008 (1♀), *M. indica*, IX-2007 (1♀), V-2008 (1♀), *M. emarginata*, IX-2007 (1♀), *C. nucifera*, I-2008 (1♀, 1♂), V-2008 (1♀, 2♂), *T. cacao*, V-2008 (1♀); Fazenda Nossa Senhora Auxiliadora, *T. cacao*, VII-2007 (1♀), *C. nucifera*, VII-2007 (1♀), XI-2007 (1♂), *P. guajava*, VII-2007 (1♀), XI-2007 (1♀), *T. grandiflorum*, XI-2007 (5♀); Fazenda Terra Nova, *T. cacao*, V-2007 (2♀), *D. kaki*, V-2007 (1♀), *C. nucifera*, IX-2007 (6♀, 1♂); Sítio São Jorge, *C. nucifera*, XI-2007 (5♀), XI-2007 (1♀).

**Female.** Thirteen specimens measured. Dorsal shield 389 (340–428) long, 259 (205–289) wide, *j1* 37 (33–42), *j3* 48 (42–55), *j4* 7 (5–8), *j5* 6 (5–7), *j6* 8 (6–10), *J2* 8 (7–11), *J5* 8 (7–9), *z2* 14 (12–16), *z4* 12 (10–13), *z5* 7 (6–9), *Z1* 10 (8–13), *Z4* 131 (115–148), *Z5* 283 (223–307), *s4* 109 (91–120), *S2* 13 (11–15), *S4* 11 (11–12), *S5* 12 (11–13), *r3* 17 (15–22), *RI* 12 (10–13); distances between *St1–St3* 69 (64–71), *St2–St2* 77 (72–80) and *St5–St5* 72 (67–77); ventrianal shield 125 (117–135) long, 77 (70–82) wide at level of *ZV2* and 78 (66–85) wide at level of anus; movable cheli-

ceral digit 45 (42–48) long, with 4 teeth; fixed cheliceral digit 33 (30–36) long; calyx of spermatheca 10 (8–12) long; *Sge I* 48 (45–51), *Sge II* 44 (42–47), *Sge III* 57 (50–63), *Sti III* 41 (37–44), *Sge IV* 121 (105–136), *Sti IV* 81 (65–91), *St IV* 82 (75–89).

**Male.** Eight specimens measured. Dorsal shield 294 (269–323) long, 217 (197–251) wide, *j1* 29 (27–31), *j3* 44 (42–47), *j4* 6 (5–7), *j5* 5 (5–6), *j6* 8 (7–9), *J2* 9 (7–10), *J5* 7 (6–9), *z2* 13 (10–15), *z4* 11 (10–13), *z5* 7 (6–7), *Z1* 10 (8–11), *Z4* 101 (90–115), *Z5* 218 (208–236), *s4* 82 (77–88), *S2* 12 (11–13), *S4* 11 (10–13), *S5* 12 (10–13), *r3* 15 (14–17), *R1* 11 (10–12); ventrianal shield 129 (115–140) long and 165 (147–181) wide at anterior corners; shaft of spermatodactyl 23 (21–25) long; *Sge I* 38 (35–40), *Sge II* 35 (31–37), *Sge III* 39 (35–43), *Sti III* 31 (27–33), *Sge IV* 81 (72–94), *Sti IV* 60 (55–70), *St IV* 65 (61–70).

**Remarks.** Measurements of the specimens collected are similar to those of the original description, as well as to those of Gondim Jr. and Moraes (2001).

### *Amblyseius perditus* Chant & Baker

*Amblyseius perditus* Chant & Baker, 1965: 16.

*Amblyseius perditus*: Moraes et al. 2004: 47.

*Amblyseius perditus*: Chant and McMurtry 2007: 80.

**Specimens examined.** UESC, *C. nucifera*, VII-2007 (1♀), IV-2008 (1♀), *E. guineensis*, VII-2007 (1♀), *P. guajava*, VII-2007 (1♀), I-2008 (3♀), *T. grandiflorum*, VII-2007 (1♀), *A. integrifolia*, I-2008 (2♀); Fazenda Barra, *T. cacao*, XI-2007 (1♀); Fazenda Frutical, *M. sapientum*, VII-2007 (1♀), *T. cacao*, IV-2007 (1♀), VII-2007 (1♀), IX-2007 (2♀); Fazenda Liberdade, *C. nucifera*, V-2007 (5♀), *M. indica*, V-2007 (1♀), *T. cacao*, V-2007 (3♀); Fazenda Monte Alegre, *P. guajava*, V-2007 (2♀), IX-2007 (2♀, 1♂), I-2008 (5♀); Fazenda Nossa Senhora Auxiliadora, *T. cacao*, VII-2007 (14♀); Fazenda Terra Nova, *C. nucifera*, V-2007 (1♀), IX-2007 (2♀), *P. guajava*, I-2008 (5♀).

**Female.** Fifteen specimens measured. Dorsal shield 345 (333–358) long, 217 (197–238) wide, *j1* 34 (29–36), *j3* 38 (35–43), *j4* 7 (6–8), *j5* 6 (5–7), *j6* 9 (8–10), *J2* 9 (8–11), *J5* 8 (7–9), *z2* 7 (7–8), *z4* 9 (8–10), *z5* 5 (4–7), *Z1* 10 (8–12), *Z4* 59 (54–69), *Z5* 155 (145–174), *s4* 57 (54–67), *S2* 12 (11–13), *S4* 10 (9–11), *S5* 8 (8–10), *r3* 16 (13–17), *R1* 11 (9–12); distances between *St1–St3* 60 (57–63), *St2–St2* 68 (65–71) and *St5–St5* 69 (62–74); ventrianal shield 44 (40–49) long, 65 (61–70) wide; anal shield 50 (46–55) long, 67 (64–70) wide; movable cheliceral digit 33 (31–34) long, with 3 teeth; fixed cheliceral digit 26 (25–28) long; calyx of spermatheca 21 (18–28) long; *Sge I* 43 (41–46), *Sge II* 41 (38–45), *Sge III* 45 (40–48), *Sti III* 34 (27–38), *Sge IV* 74 (71–85), *Sti IV* 57 (52–61), *St IV* 65 (62–71).

**Male.** One specimen measured. Dorsal shield 253 long and 182 wide, *j1* 22, *j3* 31, *j4* 7, *j5* 7, *j6* 8, *J2* 10, *J5* 7, *z2* 8, *z4* 8, *z5* 7, *Z1* 11, *Z5* 50, *s4* 33, *S2* 10, *S4* 9, *S5* 8, *r3*

12, *RI* 8; ventrianal shield 102 long and 166 wide at anterior corners; shaft of spermatodactyl 22 long; *Sge I* 30, *Sge II* 27, *Sge III* 27, *Sti III* 25, *Sge IV* 43, *Sti IV* 32, *St IV* 65.

**Remarks.** Measurements of the specimens collected fit the redescription given by Denmark and Muma (1989) and those of Gondim Jr. and Moraes (2001). As also reported in those publications, the specimens collected have longer *RI* (5 in the holotype). This is the first record of this species in Bahia.

### *Amblyseius tamatavensis* Blommers

*Amblyseius tamatavensis* Blommers, 1974: 144.

*Amblyseius tamatavensis*: Moraes et al. 2004: 52.

*Amblyseius tamatavensis*: Chant and McMurtry 2007: 81.

**Specimens examined.** CEPLAC, *C. nucifera*, II-2008 (2♀), fruits, VI-2008 (2♀, 1♂), I-2009 (1♂), II-2009 (2♀), IV-2009 (2♀), VIII-2009 (1♀), X-2009 (1♀), XII-2009 (13♀); Fazenda Frupical, *M. sapientum*, VII-2007 (1♀); Fazenda Liberdade, *E. oleracea*, V-2007 (1♀); Sítio Agrotropical, *C. nucifera*, III-2007 (1♀); Sítio Sabino, *P. edulis*, flowers, XI-2007 (2♀, 2♂).

**Female.** Seven specimens measured. Dorsal shield 352 (323–379) long and 216 (182–238) wide; *j1* 33 (31–36), *j3* 53 (50–57), *j4* 5 (4–5), *j5* 4 (3–4), *j6* 5 (5–6), *J2* 6 (5–6), *J5* 7 (6–7), *z2* 7 (6–8), *z4* 8 (7–8), *z5* 4 (3–4), *Z1* 6 (6–7), *Z4* 108 (100–115), *Z5* 235 (227–246), *s4* 91 (90–92), *S2* 7 (6–7), *S4* 6 (6–7), *S5* 6 (5–6), *r3* 14 (13–16), *RI* 8 (7–8); distances between *St1–St3* 60 (58–63), *St2–St2* 69 (67–72) and *St5–St5* 73 (70–77); ventrianal shield 113 (108–118) long, 95 (89–99) wide at level of *ZV2* and 85 (80–87) wide at level of anus; movable cheliceral digit 37 (36–38) long; fixed cheliceral digit 30 (27–32) long; calyx of spermatheca 17 (16–18) long; *Sge I* 40 (37–42), *Sge II* 39 (35–41), *Sge III* 57 (55–61), *Sti III* 47 (46–47), *Sge IV* 103 (100–105), *Sti IV* 77 (68–80), *St IV* 71 (70–72).

**Male.** Two specimens measured. Dorsal shield 259, 292 long, 177, 179 wide, *j1* 27, 29, *j3* 43, 45, *j4* 4, 5, *j5* 3, 4, *j6* 5, *J2* 5, *J5* 7, *z2* 6, *z4* 7, *z5* 3, 4, *Z1* 5, 6, *Z4* 82, 90, *Z5* 172, *s4* 67, 69, *S2* 6, 7, *S4* 6 (5–6), *S5* 5, *r3* 12 (11–12), *RI* 7; ventrianal shield 113 (108–118) long and 144 (137–150) wide at anterior corners; shaft of spermatodactyl 17 long; *Sge I* 32 (31–32), *Sge II* 31 (30–31), *Sge III* 38, *Sti III* 34 (33–34), *Sge IV* 67, *Sti IV* 49 (47–50), *St IV* 56 (55–56).

**Remarks.** Measurements of the specimens collected fit the original description. Moraes et al. (2013) reported shorter *j1*, *j3*, *Sti III*, *Sti IV*, *St IV*, fixed and movable cheliceral digits and the width of ventrianal shield at level of *ZV2* [respectively, 26 (22–30), 42 (40–43), 28 (27–30), 48 (45–50), 51 (50–52), 27, 30 and 75] and longer *J2* and *z2* [respectively, 7 (7–8) and 10 (9–10)] in specimens collected in São Paulo state, southeastern Brazil.

***Arrenoseius urquharti* Yoshida-Shaul & Chant**

*Amblyseius urquharti* Yoshida-Shaul & Chant, 1988: 2055.

*Fundiseius urquharti*: Moraes et al. 2004: 89.

*Arrenoseius urquharti*: Chant and McMurtry 2007: 98.

**Specimens examined.** Fazenda Monte Alegre, *M. sapientum*, V-2007 (1♀); Fazenda Nossa Senhora Auxiliadora, *C. nucifera*, XI-2007 (1♀, 1♂).

**Female.** Two specimens measured. Dorsal shield 384 (361–407) long, 378 (366–390) wide, *j1* 17, *j3* 27 (21–32), *j4* 5 (4–5), *j5* 4 (3–5), *j6* 6, *J2* 5 (4–5), *J5* 8 (7–9), *z2* 8 (6–9), *z4* 12 (10–13), *z5* 4 (3–4), *Z1* 4, *Z4* 111 (105–116), *Z5* 104 (95–113), *s4* 86 (79–93), *S2* 12 (11–12), *S4* 8 (7–8), *S5* 7, *r3* 12, *R1* 12; distances between *St1*–*St3* 49 (48–49), *St2*–*St2* 79 (77–80) and *St5*–*St5* 128 (124–132); ventrianal shield 143 (140–146) long, 220 (217–223) wide at level of *ZV2* and 142 (133–150) wide at level of anus; movable cheliceral digit 40 long, with 2 teeth; fixed cheliceral digit 32 long; calyx of spermatheca 17 (15–18) long; *St IV* 37 (35–39).

**Male.** One specimen measured. Dorsal shield 279 long and 212 wide, *j1* 19, *j3* 25 (20–30), *j4* 4, *j5* 4, *j6* 5, *J2* 3, *J5* 9, *z2* 9, *z4* 11, *z5* 3, *Z1* 6, *Z4* 74, *Z5* 59, *s4* 57 (56–58), *S2* 11, *S4* 9, *S5* 10, *r3* 12, *R1* 10; ventrianal shield 136 long and 189 wide at anterior corners; shaft of spermatodactyl 17 long; *St IV* 33.

**Remarks.** Measurements of the females collected are generally similar to those of the original description, except for the longer *Z4*, *Z5* and *s4* in the females measured in the present work [respectively, 77, 86 and 72 in the original description]. Measurements of the females collected differ from those of Moraes et al. (2013) for the shorter *j3* and the longer width of dorsal shield, *z4*, *Z4*, *S2*, *R1* and calyx of spermatheca [respectively, 33–35, 300–305, 8, 88–92, 98–103, 7–8, 7, 8, 22–23 in the latter]. This is the first record of this genus in Bahia.

***Iphiseiodes metapodalis* (El-Banhawy)**

*Amblyseius metapodalis* El-Banhawy, 1984: 132.

*Iphiseiodes metapodalis*: Moraes et al. 2004: 90.

*Iphiseiodes metapodalis*: Chant and McMurtry 2007: 98.

**Specimens examined.** UESC, *E. oleracea*, VII-2007 (1♀), *C. nucifera*, VII-2007 (5♀, 1♂), I-2008 (2♀, 1♂), IV-2008 (1♀), *E. guineensis*, VII-2007 (2♀), *P. guajava*, VII-2007 (8♀, 1♂); CEPLAC, *C. nucifera*, fruit, IV-2009 (1♀); Fazenda Barra, *A. muricata*, XI-2007 (1♀), *T. cacao*, XI-2007 (1♀); Fazenda Bela Vista, *C. nucifera*, IV-2007 (2♀), IX-2007 (1♀, 1♂), XI-2007 (1♀); Fazenda Frupical, *A. muricata*, XI-2007 (1♀); Fazenda Liberdade, *T. cacao*, V-2007 (2♀), *M. indica*, V-2007 (1♀); Fazenda Monte Alegre, *M. sapientum*, IX-2007 (1♀), *C. nucifera*, IX-2007 (1♀), I-2008 (1♀), V-2008 (2♀), *A. muricata*, I-2008 (2♀); Fazenda Nossa Senhora Auxiliadora, *T. ca-*

cao, VII-2007 (1♀), *T. grandiflorum*, XI-2007 (1♀); Fazenda Terra Nova, *C. nucifera*, V-2007 (1♀), IX-2007 (4♀, 1♂); Sítio Agrotropical, *C. nucifera*, III-2007 (2♀); Sítio Sabino, *A. squamosa* VIII-2007 (2♀).

**Female.** Eleven specimens measured. Dorsal shield 394 (348–425) long, 313 (282–351) wide, *j1* 23 (20–26), *j3* 34 (32–37), *j4* 5 (4–6), *j5* 5 (4–6), *j6* 6 (5–7), *J2* 7 (6–8), *J5* 8 (7–9), *z2* 6 (5–7), *z4* 6 (5–7), *z5* 5 (4–7), *Z1* 7 (6–7), *Z4* 141 (128–149), *Z5* 192 (184–200), *s4* 149 (140–157), *S2* 7 (6–8), *S4* 6 (5–7), *S5* 6 (5–8), *r3* 6 (6–7), *R1* 7 (6–8); distances between *St1–St3* 52 (50–55), *St2–St2* 77 (75–79) and *St5–St5* 103 (95–111); ventrianal shield 121 (111–130) long, 160 (152–165) wide at level of *ZV2* and 115 (110–120) wide at level of anus; movable cheliceral digit 34 (32–36) long, with 4 teeth; fixed cheliceral digit 27 (25–28) long, with 11 teeth; calyx of spermatheca 13 (12–15) long; *Sge I* 63 (58–68), *Sge II* 38 (34–43), *Sge III* 66 (60–71), *Sti III* 31 (29–35), *Sge IV* 138 (130–145), *Sti IV* 82 (76–87), *St IV* 43 (37–47).

**Male.** Five specimens measured. Dorsal shield 326 (315–340) long, 243 (220–269) wide, *j1* 24 (23–25), *j3* 35 (31–39), *j4* 5 (4–6), *j5* 5 (4–6), *j6* 7 (6–7), *J2* 6 (5–7), *J5* 7 (6–8), *z2* 6 (5–7), *z4* 6 (6–7), *z5* 5 (5–6), *Z1* 6 (5–7), *Z4* 109 (105–118), *Z5* 182 (176–187), *s4* 106 (100–110), *S2* 7 (6–7), *S4* 6 (5–7), *S5* 6 (5–7), *r3* 6 (6–7), *R1* 6 (6–7); ventrianal shield 142 (141–145) long and 191 (186–197) wide at anterior corners; shaft of spermatodactyl 20 (20–22) long; *Sge I* 47 (45–49), *Sge II* 33 (30–35), *Sge III* 44 (42–47), *Sti III* 28 (25–30), *Sge IV* 95 (92–98), *Sti IV* 65 (60–70), *St IV* 44 (42–48).

**Remarks.** Measurements of the specimens collected are similar to those of the original description, except for the width of dorsal shield (reported as 147 for the female holotype, probably a mistake). This is the first description of a male of this species.

### *Iphiseiodes setillus* Gondim Jr. & Moraes

*Iphiseiodes setillus* Gondim Jr. & Moraes, 2001: 75.

*Iphiseiodes setillus*: Moraes et al. 2004: 91.

*Iphiseiodes setillus*: Chant and McMurtry 2007: 98.

**Specimens examined.** UESC, *E. oleracea*, VII-2007 (1♀), *T. cacao*, VII-2007 (1♀), I-2008 (2♀), IV-2008 (2♀); Fazenda Bela Vista, *T. cacao*, XI-2007 (1♀); Fazenda Frupical, *T. cacao*, VII-2007 (1♀).

**Female.** Six specimens measured. Dorsal shield 312 (256–364) long and 224 (212–246) wide, *j1* 8 (6–8), *j3* 15 (14–16), *j4* 15 (14–15), *j5* 16, *j6* 17 (15–18), *J2* 17 (17–18), *J5* 9 (7–9), *z2* 11, *z4* 16 (15–17), *z5* 14 (12–15), *Z1* 17 (16–18), *Z4* 18 (16–19), *Z5* 25 (21–27), *s4* 15 (14–16), *S2* 17 (16–18), *S4* 17 (15–18), *S5* 16 (14–17), *r3* 11 (10–11), *R1* 11 (11–12); distances between *St1–St3* 41 (40–42), *St2–St2* 60 (58–61) and *St5–St5* 80 (76–84); ventrianal shield 77 (75–79) long, 95 (86–98) wide at level of *ZV2* and 75 (73–80) wide at level of anus; movable cheliceral digit 22 (21–23) long, with 2 teeth; fixed cheliceral digit 19 (18–20) long, with 6 teeth; calyx

of spermatheca 10 (8–11) long; *Sge I* 8 (7–8), *Sge II* 11 (11–12), *Sge III* 13 (13–14), *Sti III* 14 (13–15), *Sge IV* 13 (13–14), *Sti IV* 14 (14–15), *St IV* 20 (19–20).

**Remarks.** Measurements of specimens collected are similar to those of the original description. This is the first record of this species in Bahia.

### *Iphiseiodes zuluagai* Denmark & Muma

*Iphiseiodes zuluagai* Denmark & Muma, 1972: 23.

*Iphiseiodes zuluagai*: Moraes et al. 2004: 91.

*Iphiseiodes zuluagai*: Chant and McMurtry 2007: 98.

**Specimens examined.** UESC, *C. nucifera*, VII-2007 (6♀), I-2008 (1♂), IV-2008 (1♀), *C. papaya*, V-2007 (2♀), *T. grandiflorum*, VIII-2007 (2♀), *E. guineensis*, VII-2007 (1♀), I-2008 (1♀); CEPLAC, *C. nucifera*, fruit, XII-2008 (1♀); Fazenda Barra, *A. occidentale*, VIII-2007 (1♀), *A. muricata*, XI-2007 (1♀); Fazenda Bela Vista, *T. cacao*, IV-2007 (3♀, 2♂), XI-2007 (2♀); Fazenda Formiga, *A. muricata*, VIII-2007 (1♀), *C. nucifera*, VIII-2007 (1♀); Fazenda Liberdade, *E. oleracea*, V-2007 (1♀), *T. cacao*, V-2007 (3♀); Fazenda Monte Alegre, *C. nucifera*, I-2008 (2♀, 1♂), V-2008 (2♀, 2♂); Fazenda Nossa Senhora Auxiliadora, *P. caimito*, IV-2007 (2♀); Fazenda Terra Nova, *C. nucifera*, IX-2007 (1♂); Sítio Agrotropical, *C. nucifera*, III-2007 (1♀, 1♂); Sítio Sabino, *M. indica*, VIII-2007 (5♀).

**Female.** Thirteen specimens measured. Dorsal shield 353 (307–397) long, 291 (256–333) wide, *j1* 21 (15–25), *j3* 29 (20–35), *j4* 2 (2–3), *j5* 3 (2–3), *j6* 3 (3–4), *J2* 3 (3–4), *J5* 4 (3–4), *z2* 3 (2–3), *z4* 3 (2–3), *z5* 3 (2–3), *Z1* 3 (2–4), *Z4* 4 (3–5), *Z5* 131 (105–144), *s4* 109 (87–122), *S2* 3 (3–4), *S4* 3 (2–4), *S5* 3 (2–4), *r3* 6 (4–7), *R1* 4 (4–5); distances between *St1–St3* 49 (41–53), *St2–St2* 77 (70–82) and *St5–St5* 104 (89–115); ventrianal shield 99 (82–110) long, 120 (96–133) wide at level of *ZV2* and 88 (70–112) wide at level of anus; movable cheliceral digit 36 (34–37) long; fixed cheliceral digit 29 (25–31) long; calyx of spermatheca 7 (5–8) long; *Sge I* 51 (40–60), *Sge II* 33 (26–40), *Sge III* 51 (39–59), *Sti III* 29 (23–32), *Sge IV* 93 (63–107), *Sti IV* 62 (48–72), *St IV* 38 (30–44).

**Male.** Eight specimens measured. Dorsal shield 287 (269–312) long, 223 (205–256) wide, *j1* 21 (15–25), *j3* 37 (32–40), *j4* 2 (2–3), *j5* 3 (2–3), *j6* 3 (3–4), *J2* 4 (3–4), *J5* 4 (3–5), *z2* 3 (2–4), *z4* 3 (2–3), *z5* 3 (2–3), *Z1* 3 (3–4), *Z4* 4 (3–6), *Z5* 93 (77–108), *s4* 82 (64–92), *S2* 4 (3–4), *S4* 3 (2–5), *S5* 5 (3–7), *r3* 7 (6–8), *R1* 5 (3–5); ventrianal shield 110 (100–121) long and 169 (148–187) wide at anterior corners; shaft of spermatodactyl 16 (15–18) long; *Sge I* 41 (31–47), *Sge II* 30 (24–33), *Sge III* 40 (29–46), *Sti III* 26 (20–33), *Sge IV* 65 (52–72), *Sti IV* 49 (38–56), *St IV* 35 (26–46).

**Remarks.** Measurements of the specimens collected are similar to those of the original description, except for the longer *r3* and *R1* (2 for the holotype). The calyx of the spermatheca is shorter than reported by Lofego et al. (2009) for specimens from São Paulo state [14 (12–15)]. Measurements of male specimens fit the measurements of the allotype male and those of Lofego et al. (2004).

***Paraamblyseius multicircularis* Gondim Jr. & Moraes**

*Paraamblyseius multicircularis* Gondim Jr. & Moraes, 2001: 79.

*Paraamblyseius multicircularis*: Moraes et al. 2004: 158.

*Paraamblyseius multicircularis*: Chant and McMurtry 2007: 103.

**Specimens examined.** UESC, *E. oleracea*, I-2008 (1♀), *T. cacao*, VII-2007 (3♀), I-2008 (1♀, 1♂), IV-2008 (2♀).

**Female.** Seven specimens measured. Dorsal shield 331 (302–360) long, 248 (225–270) wide, *j1* 13 (13–15), *j3* 24 (22–25), *j4* 30 (27–33), *j5* 38 (34–42), *j6* 38 (35–42), *J2* 42 (38–44), *J5* 13 (12–14), *z2* 29 (27–31), *z4* 36 (33–41), *z5* 28 (24–30), *Z1* 45 (43–49), *Z4* 44 (40–46), *Z5* 29 (27–31), *s4* 42 (37–46), *S2* 33 (30–35), *S4* 33 (31–35), *S5* 24 (23–26), *r3* 13, *RI* 17 (16–18); distances between *St1–St3* 32 (30–34), *St2–St2* 64 (62–67) and *St5–St5* 100 (95–104); ventrianal shield 110 (103–117) long, 191 (182–201) wide at level of *ZV2* and 117 (110–125) wide at level of anus; movable cheliceral digit 20 (19–21) long, with 2 teeth; fixed cheliceral digit 20 (19–21) long; calyx of spermatheca 18 (17–20) long.

**Male.** One specimen measured. Dorsal shield 218 long, 179 wide, *j1* 13, *j3* 23, *j4* 26, *j5* 30, *j6* 28, *J2* 30, *J5* 8, *z2* 18, *z4* 30, *z5* 21, *Z1* 37, *Z4* 38, *Z5* 23, *s4* 41, *S2* 31, *S4* 24, *S5* 18, *r3* 11, *RI* 12; ventrianal shield 92 long and 153 wide at anterior corners; shaft of spermatodactyl 20 long.

**Remarks.** Measurements of the females collected are similar to those of the original description, except for longer dorsal shield (263 for the holotype). This is the first description of a male of this species and the first record of this genus in Bahia.

***Paraphytoseius orientalis* (Narayanan, Kaur & Ghai)**

*Typhlodromus (Amblyseius) orientalis* Narayanam et al., 1960: 394.

*Paraphytoseius orientalis*: Moraes et al. 2004: 162.

*Paraphytoseius orientalis*: Chant and McMurtry 2007: 53.

**Specimens examined.** Fazenda Nossa Senhora Auxiliadora, *P. guajava*, XI-2007 (2♀).

**Female.** Two specimens measured. Dorsal shield 270, 282 long, 159 wide, *j1* 31, 38, *j3* 80, 88, *j4* 4, 5, *j5* 3, 5, *j6* 6, *J5* 4, *z2* 8, 9, *z4* 8, *z5* 4, 5, *Z1* 6, 7, *Z4* 69, 76, *Z5* 106, 110, *s4* 123, 127, *r3* 40, 47, *RI* 25, 26; distances between *St1–St3* 64, *St2–St2* 64 and *St5–St5* 80; ventrianal shield 85, 107 long, 55 wide at level of *ZV2* and 51 wide at level of anus; movable cheliceral digit 29, 31 long, with 2 teeth; fixed cheliceral digit 24, 25 long, with 11 teeth; calyx of spermatheca 4, 5 long; *Sge IV* 26, 31, *Sti IV* 36, 37, *St IV* 43, 46.

**Remarks.** Only measurements of the dorsal shield and of the longer setae were given in the original description. The specimens collected are slightly smaller and concurrently have slightly shorter setae than the holotype. Measurements of the specimens

collected are similar to those of the original description of its junior synonym, *Paraphytoseius multidentatus* Swirski & Shechter, 1961, except for the longer Z5 (76–91 in the latter). They also are similar to the measurements provided by Lofego et al. (2009).

### ***Phytoscutus sexpilis* Muma**

*Phytoscutus sexpilis* Muma, 1961: 275.

*Phytoscutus sexpilis*: Moraes et al. 2004: 166.

*Phytoscutus sexpilis*: Chant and McMurtry 2007: 101.

**Specimens examined.** Fazenda Nossa Senhora Auxiliadora, *P. guajava*, VII-2007 (1♀).

**Female.** One specimen measured. Dorsal shield 350 long, 330 wide, *j1* 15, *j3* 33, *j4* 12, *j6* 12, *J5* 9, *z2* 11, *z4* 14, *z5* 10, *Z1* 15, *Z4* 192, *Z5* 276, *s4* 184, *S4* 9, *S5* 9, *r3* 14, *R1* 16; distances between *St1–St3* 48, *St2–St2* 57 and *St5–St5* 87; ventrianal shield 137 long, 165 wide at level of ZV2 and 128 wide at level of anus; movable cheliceral digit 25 long; fixed cheliceral digit 17 long; calyx of spermatheca 15 long; *Sge IV* 84, *Sti IV* 69.

**Remarks.** Measurements of the specimen collected fit the redescription of the holotype given by Yoshida-Shaul and Chant (1997), except for the shorter *j1*, *J5*, *z4*, *Z1*, *S4* and *S5* and the longer *j4* and *r3* [respectively, 21, 13, 22, 26, 13, 14, 9 and 10]. Gondim Jr. and Moraes (2001) reported specimens from São Paulo state to have longer *z2*, *z4* and *S5* [respectively, 16, 26 and 12].

### ***Proprioseiopsis ovatus* (Garman)**

*Amblyseiopsis ovatus* Garman, 1958: 78.

*Proprioseiopsis ovatus*: Moraes et al. 2004: 184.

*Proprioseiopsis ovatus*: Chant and McMurtry 2007: 89.

**Specimens examined.** UESC, *M. sapientum*, IV-2008 (1♀); Fazenda Terra Nova, *E. oleracea*, I-2008 (1♀); Sítio Sabino, *P. edulis*, XI-2007 (1♀).

**Female.** Three specimens measured. Dorsal shield 366 (360–372) long, 306 (287–320) wide, *j1* 31 (30–32), *j3* 69 (63–72), *j4* 7, *j5* 4 (3–5), *j6* 12 (10–13), *J5* 9, *z2* 42 (32–50), *z4* 22 (21–23), *z5* 8, *Z1* 22 (20–24), *Z4* 110 (101–115), *Z5* 84 (79–90), *s4* 102 (92–112), *S2* 22 (21–23), *S4* 12 (11–12), *S5* 11 (10–11), *r3* 20 (20–21), *R1* 11 (10–12); distances between *St1–St3* 58 (57–58), *St2–St2* 74 (72–76) and *St5–St5* 91 (90–92); ventrianal shield 114 (113–115) long, 111 (110–112) wide at level of ZV2 and 91 (85–96) wide at level of anus; movable cheliceral digit 32 (30–34) long; fixed cheliceral digit 31 (30–32) long; calyx of spermatheca 15 (12–15) long; *Sge II* 18 (17–18), *Sge III* 30 (29–31), *Sti III* 26 (25–27), *Sge IV* 61 (60–62), *Sti IV* 41 (35–45), *St IV* 89 (88–90).



**Remarks.** Measurements of the specimens collected fit the redescription of the holotype given by Moraes and McMurtry (1983), except for the shorter *S4* and longer width of dorsal shield, *J5* and *Sge IV* [respectively, 23, 252, 4 and 48 in the latter]. The measurements fit the redescription given by Lofego et al. (2009) and Ferla et al. (2011) for Brazilian specimens.

### ***Proprioiseiopsis dominigos* (El-Banhawy)**

*Amblyseius dominigos* El-Banhawy, 1984: 130.

*Proprioiseiopsis dominigos*: Moraes et al. 2004: 175.

*Proprioiseiopsis dominigos*: Chant and McMurtry 2007: 89.

**Specimens examined.** Fazenda Monte Alegre, *M. sapientum*, IX-2007 (1♀), *P. guajava*, I-2008 (1♀, 1♂); Sítio Agrotropical, *C. nucifera*, III-2007 (1♀).

**Female.** Three specimens measured. Dorsal shield 397 long, 307 wide, *j1* 35, *j3* 106 (100–114), *j4* 3, *j5* 3, *j6* 4 (3–4), *J5* 6 (5–7), *z2* 32 (31–33), *z4* 54 (50–57), *z5* 3, *Z1* 6 (5–7), *Z4* 128 (123–131), *Z5* 120 (114–125), *s4* 123 (118–127), *S2* 8 (6–10), *S4* 8 (7–9), *S5* 5 (5–6), *r3* 28 (27–30), *R1* 9 (8–9); distances between *St1–St3* 54 (52–56), *St2–St2* 74 (71–75) and *St5–St5* 130 (129–130); ventrianal shield 117 (110–122) long, 138 (127–145) wide at level of *ZV2* and 113 (110–117) wide at level of anus; movable cheliceral digit 45 long; fixed cheliceral digit 40 long; calyx of spermatheca 22 (20–25) long; *Sge I* 31, *Sge II* 25, *Sge III* 31 (30–32), *Sti III* 29 (28–30), *Sge IV* 54 (52–56), *Sti IV* 37 (36–37), *St IV* 51 (46–56).

**Male.** One specimen measured. Dorsal shield 333 long, 230 wide, *j1* 30, *j3* 82, *j4* 2, *j5* 3, *j6* 3, *J5* 4, *z2* 28, *z4* 43, *z5* 3, *Z1* 5, *Z4* 97, *Z5* 88, *s4* 90, *S2* 9, *S4* 7, *S5* 6, *r3* 17, *R1* 8; ventrianal shield 148 long and 183 wide at anterior corners; shaft of spermatodactyl 19 long; *Sge I* 22, *Sge II* 20, *Sge III* 21, *Sti III* 23, *Sge IV* 40, *Sti IV* 27, *St IV* 44.

**Remarks.** Measurements of the specimens collected are similar to those of the original description. They also agree with those of Moraes et al. (2013), except for the longer *z2*, *z4* and *S4* [respectively, 21, 37 and 5 in the latter]. This is the first record of this species in Bahia.

### ***Proprioiseiopsis neotropicus* (Ehara)**

*Amblyseius neotropicus* Ehara, 1966: 133.

*Proprioiseiopsis neotropicus*: Moraes et al. 2004: 183.

*Proprioiseiopsis neotropicus*: Chant and McMurtry 2007: 89.

**Specimens examined.** UESC, *C. nucifera*, I-2008 (1♀); Fazenda Barra, *T. cacao*, XI-2007 (2♀); Fazenda Liberdade, *T. cacao*, V-2007 (1♀); Fazenda Monte Alegre, *M. sapientum*, I-2008 (1♀); Fazenda Terra Nova, *D. kaki*, V-2007 (1♀).

**Female.** Four specimens measured. Dorsal shield 424 (376–453) long, 331 (300–371) wide, *j1* 34 (31–36), *j3* 47 (46–48), *j4* 6 (4–7), *j5* 5 (5–6), *j6* 7 (6–8), *J5* 5 (5–6), *z2* 26 (22–29), *z4* 21 (17–25), *z5* 6 (5–7), *Z1* 8 (7–8), *Z4* 120 (112–130), *Z5* 106 (98–116), *s4* 119 (107–126), *S2* 8 (7–8), *S4* 8 (7–8), *S5* 8 (7–9), *r3* 24 (21–26), *RI* 13 (10–17); distances between *St1–St3* 66 (65–66), *St2–St2* 85 (82–88) and *St5–St5* 109 (103–117); ventrianal shield 114 (108–120) long, 115 (105–122) wide at level of *ZV2* and 94 (91–97) wide at level of anus; movable cheliceral digit 40 (38–43) long, with 3 teeth; fixed cheliceral digit 33 (32–35) long; calyx of spermatheca 22 (20–25) long; *Sge I* 34 (32–39), *Sge II* 35 (33–40), *Sge III* 36 (34–38), *Sti III* 32 (31–34), *Sge IV* 79 (72–82), *Sti IV* 57 (50–65), *St IV* 69 (65–73).

**Remarks.** Measurements of the specimens collected are similar to those of the original description, except for the longer *j3* and *r3* (respectively 32 and 15 in the latter). They also agree with those of Lofego et al. (2004), except for the longer *z4* (13 in the latter). This is the first record of this species in Bahia.

### ***Proprioiseiopsis pentagonalis* (Moraes & Mesa)**

*Amblyseius pentagonalis* Moraes et al., 1991: 127.

*Proprioiseiopsis pentagonalis*: Moraes et al. 2004: 186.

*Proprioiseiopsis pentagonalis*: Chant and McMurtry 2007: 89.

**Specimens examined.** Fazenda Monte Alegre, *T. cacao*, I-2008 (1♀).

**Female.** One specimen measured. Dorsal shield 328 long, 207 wide, *j1* 24, *j3* 23, *j4* 6, *j5* 6, *j6* 7, *J5* 10, *z2* 10, *z4* 11, *z5* 6, *Z1* 9, *Z4* 102, *Z5* 164, *s4* 74, *S2* 10, *S4* 10, *S5* 8, *r3* 18, *RI* 11; distances between *St1–St3* 63, *St2–St2* 69 and *St5–St5* 61; ventrianal shield 108 long, 85 wide at level of *ZV2* and 71 wide at level of anus; movable cheliceral digit 35 long; fixed cheliceral digit 30 long; calyx of spermatheca 50 long; *Sge II* 25, *Sge III* 22, *Sti III* 17, *Sge IV* 58, *Sti IV* 37, *St IV* 52.

**Remarks.** Measurements of the specimen collected are similar to those of the original description and of Moraes et al. (2013) for specimens from São Paulo state. This is the first record of this species in Bahia.

### ***Typhlodromalus peregrinus* (Muma)**

*Typhlodromus peregrinus* Muma, 1955: 270.

*Typhlodromalus peregrinus*: Moraes et al. 2004: 202.

*Typhlodromalus peregrinus*: Chant and McMurtry 2007: 111.

**Specimens examined.** CEPLAC, *C. nucifera*, fruits, XII-2008 (1♀), XII-2009 (1♀); Fazenda Barra, *T. cacao*, XI-2007 (1♀); Fazenda Bela Vista, *P. guajava*, IV-2007 (1♀), XI-2007 (1♀); Fazenda Kamuí, *C. nucifera*, XI-2007 (1♀); Fazenda Nossa Senhora Auxiliadora, *C. papaya*, VII-2007 (1♀).

**Female.** Five specimens measured. Dorsal shield 351 (335–399) long, 226 (192–302) wide, *j1* 28 (27–29), *j3* 33 (31–37), *j4* 14 (12–15), *j5* 14 (11–15), *j6* 18 (15–20), *J2* 18 (14–21), *J5* 9 (8–10), *z2* 20 (18–22), *z4* 28 (27–30), *z5* 14 (11–15), *Z1* 25 (21–27), *Z4* 46 (41–52), *Z5* 64 (58–67), *s4* 41 (37–44), *S2* 30 (26–36), *S4* 25 (19–27), *S5* 12 (9–14), *r3* 20 (18–21), *R1* 16 (15–17); distances between *St1*–*St3* 64 (62–65), *St2*–*St2* 61 (60–63) and *St5*–*St5* 72 (69–74); ventrianal shield 109 (102–114) long, 64 (62–66) wide at level of *ZV2* and 61 (59–63) wide at level of anus; movable cheliceral digit 33 (30–35) long, with 3 teeth; fixed cheliceral digit 28 (25–30) long; calyx of spermatheca 22 (15–24) long; *Sge I* 15 (10–18), *Sge II* 18 (15–21), *Sge III* 28 (25–30), *Sti III* 18 (16–19), *Sge IV* 44 (39–47), *Sti IV* 21 (17–29), *St IV* 62 (58–70).

**Remarks.** Setal measurements were not given in the original description. Measurements of the specimens collected are similar to those of McMurtry (1983), except for the longer *Z1* (16–17 in the latter); they also agree with the redescription given by Moraes et al. (2013).

### *Typhlodromips mangleae* De Leon

*Typhlodromips mangleae* De Leon, 1967: 28.

*Typhlodromips mangleae*: Moraes et al. 2004: 217.

*Typhlodromips mangleae*: Chant and McMurtry 2007: 63.

**Specimens examined.** Sítio São Jorge, *C. nucifera*, XI-2007 (1♀).

**Female.** One specimen measured. Dorsal shield 343 long and 220 wide, *j1* 18, *j3* 20, *j4* 10, *j5* 10, *j6* 11, *J2* 12, *J5* 9, *z2* 11, *z4* 10, *z5* 10, *Z1* 12, *Z4* 36, *Z5* 73, *s4* 22, *S2* 12, *S4* 10, *S5* 8, *r3* 14, *R1* 14; distances between *St1*–*St3* 56, *St2*–*St2* 62 and *St5*–*St5* 63; ventrianal shield 115 long, 87 wide at level of *ZV2* and 85 wide at level of anus; movable cheliceral digit 27 long; fixed cheliceral digit 23 long; calyx of spermatheca 5 long; *Sge I* 27, *Sge II* 26, *Sge III* 31, *Sti III* 24, *Sge IV* 43, *Sti IV* 36, *St IV* 53.

**Remarks.** Measurements of the specimens collected are similar to those of the original description, except for shorter calyx of spermatheca (9 in the holotype). Lofego et al. (2004, 2009) reported a slightly longer *Z4* [respectively, 39 (37–41) and 38 (35–45)]. Measurements of the specimens collected are similar to those of Gondim Jr. and Moraes (2001). This is the first record of this species in Bahia.

### *Typhlodromips theobromae* Souza, Oliveira & Gondim Jr.

*Typhlodromips theobromae*: Souza et al. 2010: 49.

**Specimens examined.** UESC, *T. cacao*, VII-2007, IV-2008 (3♀, 2♂); Fazenda Monte Alegre, *T. cacao*, V-2007 (1♀), *M. indica*, V-2008 (2♀, 3♂); Fazenda Terra Nova, *T. cacao*, IX-2007, I-2008 (4♀).

**Female.** Ten specimens measured. Dorsal shield 301 (288–312) long, 211 (194–246) wide; *j1* 19 (17–21), *j3* 23 (17–26), *j4* 8 (6–10), *j5* 8 (6–10), *j6* 10 (8–12), *J2* 10 (8–15), *J5* 7 (7–8), *z2* 10 (8–12), *z4* 18 (16–21), *z5* 7 (6–9), *Z1* 13 (8–17), *Z4* 45 (43–47), *Z5* 56 (46–60), *s4* 23 (20–25), *S2* 15 (11–19), *S4* 10 (6–16), *S5* 9 (6–11), *r3* 11 (10–12), *R1* 9 (7–10); distances between *St1–St3* 58 (55–60), *St2–St2* 72 (68–74) and *St5–St5* 65 (61–65); ventrianal shield 88 (82–93) long, 91 (86–93) wide at level of *ZV2* and 74 (70–79) wide at level of anus; movable cheliceral digit 39 (38–40) long, with 3 teeth; fixed cheliceral digit 30 long, with 15 teeth; calyx of spermatheca 9 (7–10) long; *Sge II* 14 (12–16), *Sge III* 15 (14–18), *Sti III* 14 (12–15), *Sge IV* 31 (27–35), *Sti IV* 15(12–18), *St IV* 28 (25–30).

**Male.** Five specimens measured; dorsal shield 229 (212–241) long and 160 (152–176) wide; *j1* 16 (14–18), *j3* 22 (19–24), *j4* 9 (8–9), *j5* 8 (8–9), *j6* 10 (9–10), *J2* 8 (8–9), *J5* 6 (5–6), *z2* 10 (8–11), *z4* 19 (17–22), *z5* 7 (7–8), *Z1* 9 (8–10), *Z4* 33 (31–35), *Z5* 36 (33–39), *s4* 21 (20–22), *S2* 13 (12–14), *S4* 9 (8–10), *S5* 7 (6–7), *r3* 12 (11–12), *R1* 8 (7–10); ventrianal shield 91 (87–96) long and 132 (128–138) wide at anterior corners; shaft of spermatodactyl 15 long; *Sge I* 13 (12–14), *Sge II* 12 (12–13), *Sge III* 12 (11–13), *Sti III* 12 (11–13), *Sge IV* 19 (18–20), *Sti IV* 14 (12–16), *St IV* 24 (23–25).

**Remarks.** Measurements of the specimens collected are similar to those of the original description.

## Phytoseiinae Berlese

### *Phytoseius latinus* El-Banhawy

*Phytoseius latinus* El-Banhawy, 1984: 141.

*Phytoseius latinus*: Moraes et al. 2004: 217.

*Phytoseius latinus*: Chant and McMurtry 2007: 129.

**Specimens examined.** Fazenda Bela Vista, *P. guajava*, IV-2007 (7♀, 6♂), VII-2007 (4♀, 2♂), VIII-2007 (1♀), XI-2007 (1♀).

**Female.** Five specimens measured. Dorsal shield 272 (256–302) long, 135 (128–148) wide; *j1* 19 (18–20), *j3* 32, *j4* 7 (7–8), *j5* 7, *j6* 9 (8–10), *J2* 9 (8–10), *J5* 7 (6–8), *z2* 10 (9–10), *z3* 37 (35–38), *z4* 11 (10–11), *z5* 7 (6–7), *Z4* 47 (45–50), *Z5* 52 (50–55), *s4* 51 (48–55), *s6* 59 (56–62), *r3* 36 (33–37), *R1* 14 (13–16); distances between *St1–St3* 62 (60–64), *St2–St2* 64 (63–65) and *St5–St5* 53 (52–55); ventrianal shield 90 (85–94) long, 51 (48–55) wide at level of *ZV2* and 45 (40–49) wide at level of anus; movable cheliceral digit 29 (28–30) long; fixed cheliceral digit 25 (24–26) long; calyx of spermatheca 15 (15–16) long; *Sge IV* 29 (27–30), *Sti IV* 28 (27–29), *St IV* 29 (27–30).

**Male.** Two specimens measured. Dorsal shield 228, 236 long, 115, 123 wide, *j1* 15, 16, *j3* 28, 29, *j4* 7, 8, *j5* 6, 8, *j6* 7, 9, *J2* 7, *J5* 6, *z2* 8, 10, *z3* 31, 32, *z4* 9, 10, *z5* 6, 7, *Z4* 32, 33, *Z5* 27, 30, *s4* 37, 38, *s6* 43, 45, *r3* 28, 30, *R1* 10, 11; ventrianal shield

88 long and 126 wide at anterior corners; shaft of spermatodactyl 12, 14 long; *Sge IV* 17, 18, *Sti IV* 16, *St IV* 21, 25.

**Remarks.** The females collected differ from the original description by having longer *j3* and *Z4* and shorter *j4*, *J5* and *z4* (respectively 24, 36, 10, 17 and 15 in the holotype). This is the first record of this species in Bahia.

### ***Phytoseius woodburyi* De Leon**

*Phytoseius (Phytoseius) woodburyi* De Leon, 1965: 130.

*Phytoseius woodburyi*: Moraes et al. 2004: 246.

*Phytoseius woodburyi*: Chant and McMurtry 2007: 131.

**Specimens examined.** Fazenda Bela Vista, *C. nucifera*, XI-2007 (1♀), *P. guajava* VII-2007 (11♀).

**Female.** Five specimens measured. Dorsal shield 279 (265–310) long, 145 (140–150) wide, *j1* 29 (28–31), *j3* 33 (33–35), *j4* 5 (5–6), *j5* 5 (5–6), *j6* 5 (5–6), *J5* 6 (5–8), *z2* 13 (13–15), *z3* 30 (28–33), *z4* 12 (10–13), *z5* 6 (5–8), *Z4* 87 (85–91), *Z5* 73 (68–78), *s4* 118 (110–125), *s6* 78 (73–82), *r3* 43 (40–46); distances between *St1–St3* 55 (53–58), *St2–St2* 58 (55–60) and *St5–St5* 58 (55–61); ventrianal shield 77 (65–99) long, 32 (28–45) wide at level of *ZV2* and 39 (35–47) wide at level of anus; movable cheliceral digit 21 (18–25) long; fixed cheliceral digit 21 (20–23) long; calyx of spermatheca 8 (6–10) long; *Sge IV* 8 (8–9), *Sti IV* 49 (46–52), *St IV* 27 (24–30).

**Remarks.** Measurements of the specimens collected are similar to those of the original description, except for the longer *St IV* (20 in the holotype). Specimens collected in this study have the ventrianal shield longer than reported by Gondim Jr. and Moraes (2001) for specimens from São Paulo state [53 (50–58) in the latter]. This is the first record of this species in Bahia.

### **Typhlodrominae Wainstein**

#### ***Cocoseius elsalvador* Denmark & Andrews**

*Cocoseius elsalvador* Denmark & Andrews, 1981: 155.

*Cocoseius elsalvador*: Moraes et al. 2004: 263.

*Cocoseius elsalvador*: Chant and McMurtry 2007: 132.

**Specimens examined.** Fazenda Terra Nova, *T. cacao*, IX-2007 (1♀), *C. nucifera*, IX-2007 (2♀); Sítio São Jorge, *C. nucifera*, XI-2007 (2♀).

**Female.** Five specimens measured. Dorsal shield 302 (279–315) long, 166 (154–192) wide; *j1* 27 (25–28), *j3* 54 (51–56), *j4* 34 (32–37), *j5* 52 (50–55), *j6* 66 (62–70), *J2* 63 (58–67), *J5* 28 (22–31), *z2* 21 (20–21), *z3* 31 (30–32), *z4* 73 (68–77), *z5* 34

(29–37), *z6* 93 (88–98), *Z4* 74 (68–79), *Z5* 77 (73–81), *s4* 79 (77–82), *S4* 75 (69–80), *r3* 42 (37–46), *R1* 72 (67–75); distances between *St1–St3* 55 (54–56), *St2–St2* 60 (57–63) and *St5–St5* 59 (57–62); ventrianal shield 70 (63–75) long, 60 (50–64) wide at level of *ZV2* and 59 (55–63) wide at level of anus; movable cheliceral digit 26 (25–26) long; fixed cheliceral digit 25 (25–26) long; calyx of spermatheca 21 (20–21) long; *Sge IV* 49 (46–52), *Sti IV* 37 (34–40), *St IV* 76 (73–79).

**Remarks.** Measurements of the specimens collected are similar to those of the original description, except for the longer *Sge IV* and *Sti IV* (respectively 39 and 25 in the holotype). Gondim Jr. and Moraes (2001) reported specimens from Pernambuco state to have longer *z2* [27 (25–29)].

### ***Cocoseius palmarum* Gondim Jr., Moraes & McMurtry**

*Cocoseius palmarum* Gondim Jr. et al., 2000: 1226.

*Cocoseius palmarum*: Moraes et al. 2004: 263.

*Cocoseius palmarum*: Chant and McMurtry 2007: 134.

**Specimens examined.** UESC, *C. nucifera*, VII-2007 (3♀), I-2008 (1♀), IV-2008 (1♀); Fazenda Formiga, *C. nucifera*, VIII-2007 (1♀); Fazenda Terra Nova, *C. nucifera*, IX-2007 (1♀), I-2008 (2♀).

**Female.** Eight specimens measured. Dorsal shield 270 (265–275) long, 182 (161–189) wide, *j1* 25 (23–27), *j3* 41 (38–45), *j4* 54 (52–59), *j5* 61 (57–64), *j6* 68 (65–70), *J2* 68 (65–72), *J5* 12 (11–13), *z2* 26 (21–28), *z3* 34 (31–36), *z4* 69 (65–75), *z5* 34 (31–38), *z6* 88 (85–92), *Z4* 73 (70–75), *Z5* 69 (67–71), *s4* 76 (72–81), *S4* 64 (60–68), *r3* 57 (52–65), *R1* 54 (50–59); distances between *St1–St3* 48 (46–50), *St2–St2* 55 (53–57) and *St5–St5* 63 (60–66); ventrianal shield 93 (88–94) long, 70 (67–73) wide at level of *ZV2* and 63 (60–65) wide at level of anus; movable cheliceral digit 25 (23–26) long, with 1 teeth; fixed cheliceral digit 24 (22–25) long, with 3 teeth; calyx of spermatheca 16 (15–18) long; *Sge IV* 37 (36–40), *Sti IV* 26 (25–28), *St IV* 53 (50–56).

**Remarks.** Measurements of the specimens collected are similar to those of the original description, except for a slightly shorter calyx of spermatheca [21 (20–23) in the original description]. This is the first record of this species in Bahia.

### ***Leonseius regularis* (De Leon)**

*Typhloseiopsis regularis* De Leon, 1965: 123.

*Leonseius regularis*: Moraes et al. 2004: 275.

*Leonseius regularis*: Chant and McMurtry 2007: 161.

**Specimens examined.** UESC, *T. cacao*, VII-2007 (7♀), I-2008 (5♀), *E. guineensis*, VII-2007 (1♂), *A. integrifolia*, VII-2007 (2♀); Fazenda Bela Vista, *T. cacao*, IV-2007

(5♀, 1♂), XI-2007 (8♀), *P. guajava* VII-2007 (1♀), *S. mombin*, IV-2007 (1♀), *C. nucifera*, IV-2007 (1♀), XI-2007 (1♀), *A. muricata*, IV-2007 (4♀), *A. integrifolia*, IV-2007 (1♀), *G. americana*, IV-2007 (3♀); Fazenda Monte Alegre, *T. cacao*, V-2007 (6♀, 1♂), *C. nucifera*, IX-2007 (2♀); Fazenda Nossa Senhora Auxiliadora, *T. cacao*, VII-2007 (2♀); Fazenda Terra Nova, *T. cacao*, I-2008 (1♀, 1♂); Sítio Sabino, *M. indica*, VIII-2007 (1♀), *S. malaccense*, VIII-2007 (1♀).

**Female.** Twelve specimens measured. Dorsal shield 366 (333–397) long, 257 (212–282) wide, *j1* 28 (25–30), *j3* 38 (35–42), *j4* 4 (3–5), *j5* 4 (3–5), *j6* 5 (5–6), *J2* 6 (5–8), *J5* 8 (7–9), *z2* 4 (3–5), *z3* 11 (9–14), *z4* 6 (5–7), *z5* 4 (3–5), *Z4* 101 (85–111), *Z5* 276 (262–292), *s4* 89 (71–98), *s6* 8 (6–12), *S5* 6 (5–7), *r3* 10 (8–12), *R1* 9 (7–11); distances between *St1–St3* 61 (59–66), *St2–St2* 70 (67–73) and *St5–St5* 73 (70–76); ventrianal shield 115 (98–128) long, 63 (55–70) wide at level of *ZV2* and 64 (55–70) wide at level of anus; movable cheliceral digit 32 (30–33) long, with 4 teeth; fixed cheliceral digit 28 (25–30) long; calyx of spermatheca 18 (17–18) long; *Sge I* 54 (50–57), *Sge II* 40 (37–43), *Sge III* 51 (47–55), *Sti III* 38 (33–44), *Sge IV* 119 (107–128), *Sti IV* 68 (55–82), *St IV* 54 (48–60).

**Male.** Four specimens measured. Dorsal shield 258 (243–266) long, 171 (166–179) wide, *j1* 22 (20–24), *j3* 32 (30–34), *j4* 5 (4–6), *j5* 5 (4–6), *j6* 6 (5–7), *J2* 6 (4–8), *J5* 7 (6–8), *z2* 4 (3–5), *z3* 10 (8–12), *z4* 5 (4–7), *z5* 4 (3–5), *Z4* 63 (62–64), *Z5* 193 (185–202), *s4* 51 (48–55), *s6* 8 (6–10), *S5* 6 (5–8), *r3* 10 (8–11), *R1* 7 (7–8); ventrianal shield 111 (107–115) long and 141 (138–146) wide at anterior corners; shaft of spermatodactyl 18 (17–18) long; *Sge I* 36 (35–37), *Sge II* 30 (28–32), *Sge III* 38 (36–40), *Sti III* 29 (28–31), *Sge IV* 68 (66–70), *Sti IV* 42 (40–45), *St IV* 44 (42–48).

**Remarks.** Measurements of the specimens collected are similar to those of the original description, except for the shorter *s4* (105 in the holotype). They also agree with those reported by Moraes et al. (2013). This is the first record of this genus in Bahia.

### *Metaseiulus ferlai* Moraes, McMurtry & Lopes

*Metaseiulus* (*Metaseiulus*) *ferlai* Moraes et al., 2006: 352.

*Metaseiulus ferlai*: Chant and McMurtry 2007: 174.

**Specimens examined.** Fazenda Barra, *A. occidentale*, VIII-2007 (1♀).

**Female.** One specimen measured. Dorsal shield 350 long, 189 wide, *j1* 21, *j3* 21, *j4* 14, *j5* 14, *j6* 17, *J2* 20, *J5* 9, *z2* 17, *z3* 18, *z4* 20, *z5* 15, *Z4* 30, *Z5* 29, *s4* 21, *s6* 24, *S2* 24, *S5* 21, *r3* 21, *R1* 21; distances between *St1–St3* 63, *St2–St2* 66 and *St5–St5* 58; ventrianal shield 116 long, 87 wide at level of *ZV2* and 85 wide at level of anus; movable cheliceral digit 31 long; fixed cheliceral digit 29 long; calyx of spermatheca 25 long.

**Remarks.** Measurements of the specimens collected fit the original description. This is the first record of this genus in Bahia.

***Typhlodromina subtropica* Muma & Denmark**

*Typhlodromina subtropica* Muma & Denmark, 1969: 412.

*Typhlodromina subtropica*: Moraes et al. 2004: 305.

*Typhlodromina subtropica*: Chant and McMurtry 2007: 169.

**Specimens examined.** Fazenda Terra Nova, *D. kaki*, V-2007 (1♀).

**Female.** One specimen measured. Dorsal shield 353 long, 262 wide, *j1* 22, *j4* 35, *j5* 31, *J2* 46, *J5* 13, *z2* 32, *z3* 37, *z4* 43, *z5* 35, *Z4* 57, *Z5* 51, *s4* 43, *s6* 55, *S5* 54, *r3* 30, *R1* 21; distances between *St1–St3* 56, *St2–St2* 61 and *St5–St5* 77; ventrianal shield 125 long, 90 wide at level of *ZV2* and 72 wide at level of anus; movable cheliceral digit 28 long; fixed cheliceral digit 22 long; calyx of spermatheca 20 long.

**Remarks.** Measurements of the specimens collected fit the redescription of the holotype given by Moraes and McMurtry (1983), except for the longer *Z5* (42 in the holotype).

***Typhlodromus (Anthoseius) transvaalensis* (Nesbitt)**

*Kampimodramus transvaalensis* Nesbitt, 1951: 55.

*Typhlodromus (Anthoseius) transvaalensis*: Moraes et al. 2004: 355.

*Typhlodromus (Anthoseius) transvaalensis*: Chant and McMurtry 2007: 169.

**Specimens examined.** CEPLAC, *C. nucifera*, fruit, VII-2008 (1♀).

**Female.** One specimen measured. Dorsal shield 369 long, 205 wide, *j1* 30, *j3* 38, *j5* 30, *J2* 42, *J5* 9, *z2* 22, *z3* 40, *z4* 42, *z5* 26, *Z4* 51, *Z5* 60, *s4* 44, *s6* 48, *S2* 52, *S4* 53, *S5* 9, *r3* 34, *R1* 37; distances between *St1–St3* 64, *St2–St2* 62 and *St5–St5* 71; ventrianal shield 120 long, 69 wide at level of *ZV2* and 66 wide at level of anus; movable cheliceral digit 30 long; fixed cheliceral digit 29 long; *Sge IV* 22, *Sti IV* 30, *St IV* 46.

**Remarks.** Measurements of the specimen collected agree with the redescription of the holotype given by Schicha (1981), except the longer *j1* (23 in the holotype). They are similar to measurements given by Ferla et al. (2011), except for the shorter *j3* (48 in the latter).

**Key to the phytoseiid species collected in the present work (females)**

- |   |   |    |
|---|---|----|
| 1 | Setae <i>z3</i> and/ or <i>s6</i> present .....   | 2  |
| – | Setae <i>z3</i> and <i>s6</i> absent .....  | 3  |
| 2 | Setae <i>Z1</i> , <i>S2</i> , <i>S4</i> and <i>S5</i> absent; seta <i>r3</i> inserted on dorsal shield .....                            | 23 |
| – | At last one of the setae <i>Z1</i> , <i>S2</i> , <i>S4</i> or <i>S5</i> present; seta <i>r3</i> inserted on unsclerotised cuticle ..... | 24 |
| 3 | Posterior margin of sternal shield lightly sclerotized, often indistinct, with a posteromedian projection .....                         | 4  |



- Posterior margin of sternal shield distinct, without a posteromedian projection..... **5**
- 4 Ratio  $s4:Z1 > 3$ ; seta  $Z4$  not as long as distance between its base and base of  $S4$ ; dorsal shield mostly smooth, with few anterolateral striae ..... **Amblydromalus maniboti** (Moraes)
- Ratio  $s4:Z1 < 3$ ; seta  $Z4$  longer than distance between its base and base of  $S4$ ; dorsal shield reticulated..... **Typhlodromalus peregrinus** (Muma)
- 5 Setae  $J2$ ,  $S2$ ,  $S4$  and  $S5$  absent; lateral margin of dorsal shield with distinct incision at level of  $s4$ ; setae  $j3$ ,  $s4$ ,  $Z4$  and  $Z5$  thickened, strongly serrate, inserted on tubercles ... **Paraphytoseius orientalis** (Narayanan, Kaur & Ghai)
- Setae  $J2$  and  $S2$  present or absent,  $S4$  and  $S5$  present; lateral margin of dorsal shield without distinct incision at level of  $s4$ ; setae  $j3$ ,  $s4$ ,  $Z4$  and  $Z5$  not markedly thickened, smooth to lightly serrate, not inserted on tubercles..... **6**
- 6 Ratio  $s4:Z1 < 4$ ; body never red or dark brown; seta  $J2$  present ..... **7**
- Ratio  $s4:Z1 > 4$  (except for *Iphiseiodes setillus* and *Paraamblyseius multicircularis*); some species heavily sclerotized, body red or dark brown; seta  $J2$  present or absent ..... **8**
- 7 Calyx of spermatheca dish-shaped; seta  $Z4$  not as long as distance between its base and base of  $S5$ ; *Sge IV* sharp-tipped... **Typhlodromips manglae** De Leon
- Calyx of spermatheca short-tubular; seta  $Z4$  at least as long as distance between its base and base of  $S5$ ; *Sge IV* with tiny knob..... **Typhlodromips theobromae** Souza, Oliveira & Gondim Jr.
- 8 All shields generally strongly sclerotized; sternal shield usually wider than long; ventrianal shield usually as wide as long or wider than long; setae  $z2$  and/or  $z4$  rarely short/ minute ..... **9**
- All shields lightly sclerotized; sternal shield about as wide as long; ventrianal shield longer than wide; setae  $z2$  and  $z4$  usually short/ minute..... **18**
- 9 Seta  $J2$  present (if absent, then  $j5$  and  $S2$  also absent) ..... **10**
- Seta  $J2$  absent;  $j5$  and  $S2$  present..... **15**
- 10 Setae  $j5$ ,  $J2$  and  $S2$  absent ..... **Phytoscutus sexpilis** Muma
- Setae  $j5$ ,  $J2$  and  $S2$  present ..... **11**
- 11 Dorsal shield without marked circular ornamentation or reticulation; macrosetae present on legs I-IV ..... **12**
- Dorsal shield with marked circular ornamentation or reticulate; macrosetae, if present, only on basitarsus IV ..... **14**
- 12 With one pair of enlarged metapodal plates; seta  $Z4$  longer than distance between its base and base of  $Z5$  ..... **Iphiseiodes metapodalis** (El-Banhawy)
- With 2 pairs of metapodal plates, none distinctly large; seta  $Z4$  shorter than distance between its base and base of  $Z5$ ..... **13**
- 13 Except for  $j1$ ,  $j3$ ,  $s4$  and  $Z5$ , dorsal shield setae short/ minute; setae  $s4$  and  $Z5$  considerably longer than others; setae  $Z5$ , *Sge IV* and *Sti IV* knobbed..... **Iphiseiodes zuluagai** Denmark & Muma

- Dorsal shield setae of medium lengths, none considerably longer than others; setae *Z5*, *Sge IV* and *Sti IV* sharp-tipped .....  
.....***Iphiseiodes setillus* Gondim Jr. & Moraes**
- 14 Dorsal, genital and ventrianal shields with circular ornamentation; dorsal shield setae of medium length, none considerably longer than others; seta *ZV3* absent; with a pair of enlarged metapodal plates; leg macrosetae absent .....  
.....***Paraamblyseius multicircularis* Gondim Jr. & Moraes**
- Dorsal, genital and ventrianal shields reticulate; some dorsal shield setae much longer than others; seta *ZV3* present; with 2 pairs of metapodal plates, none distinctly enlarged; *St IV* present .....  
.....***Arrenoseius urquharti* (Yoshida-Shaul & Chant)**
- 15 Calyx of spermatheca tubular, flared near vesicle and inflate near atrium, longer than 40  $\mu\text{m}$  .....***Proprioseiopsis pentagonalis* (Moraes & Mesa)**
- Calyx of spermatheca saccular or funnel-shaped, shorter than 30  $\mu\text{m}$ ..... **16**
- 16 Seta *j3* at most 1.2 times longer than distance between their bases; calyx saccular .....***Proprioseiopsis neotropicus* (Ehara)**
- Seta *j3* at least 1.5 times longer than distance between their bases; calyx funnel-shaped ..... **17**
- 17 Seta *j3* longer than 100  $\mu\text{m}$ ; seta *Z5* longer than 110  $\mu\text{m}$ ; seta *z2* shorter than *z4*; calyx longer than 20  $\mu\text{m}$ ..... ***Proprioseiopsis dominigos* (El-Banhawy)**
- Seta *j3* shorter than 75  $\mu\text{m}$ ; seta *Z5* shorter than 100  $\mu\text{m}$ ; seta *z2* longer than *z4*; calyx shorter than 20  $\mu\text{m}$ ..... ***Proprioseiopsis ovatus* (Garman)**
- 18 Ventral shield separated from anal shield...***Amblyseius perditus* Chant & Baker**
- Ventral and anal shields fused, constituting a ventrianal shield..... **19**
- 19 Calyx of spermatheca shorter than 13  $\mu\text{m}$ ; less than 4 times as long as width at median length..... **20**
- Calyx of spermatheca longer than 14  $\mu\text{m}$ ; over 5 times as long as width at median length..... **22**
- 20 Seta *Z4* not as long as distance between its base and base of *Z5*; setae *s4* and *Z4* respectively 41–44 and 30–33  $\mu\text{m}$  long; calyx cup-shaped, 6  $\mu\text{m}$  long, increasing progressively in diameter toward the base .....  
..... ***Amblyseius impeltatus* Denmark & Muma**
- Zeta *Z4* longer than distance between its base and base of *Z5*; setae *s4* and *Z4* respectively longer than 55 and 50  $\mu\text{m}$ ; calyx short-tubular, at least 8  $\mu\text{m}$  long, somewhat constricted in the middle..... **21**
- 21 Setae *j1*, *j3*, *s4*, *Z4* and *Z5* respectively 23–26, 35–39, 56–61, 60–66 and 130–155  $\mu\text{m}$  long.....***Amblyseius igarassuensis* Gondim Jr. & Moraes**
- Setae *j1*, *j3*, *s4*, *Z4* and *Z5* respectively 33–42, 42–55, 91–120, 115–148 and 223–307  $\mu\text{m}$  long..... ***Amblyseius operculatus* De Leon**
- 22 Setae *s4*, *Z4* e *Z5* respectively 105–111, 120–143 and 271–315  $\mu\text{m}$  long; atrium nodular, distinct; major duct narrower than calyx.....  
.....***Amblyseius aerialis* (Muma)**

- Setae *s4*, *Z4* e *Z5* respectively 90–92, 100–115 and 227–246 µm long; atrium small, indistinct, incorporated into base of calyx; major duct approximately of the same diameter as calyx ..... ***Amblyseius tamatavensis* Blommers**
- 23 Setae *J2* and *R1* present; seta *s4* shorter than *s6*; insertions of setae *Z4* and *Z5* distinctly separated; each macroseta of leg IV with a tiny knob, none markedly shorter than others ..... ***Phytoseius latinus* El-Banhawy**
- Setae *J2* and *R1* absent; seta *s4* longer than *s6*; insertions of setae *Z4* and *Z5* approximate; macrosetae on leg IV broadly clavate, *Sge IV* markedly shorter than others..... ***Phytoseius woodburyi* (De Leon)**
- 24 Seta *S5* absent; ventrianal shield with 1 or 3 pairs of pre-anal setae; dorsal shield setae not distally swollen ..... **25**
- Setae *S5* present; ventrianal shield with 4 pairs of pre-anal setae (if with 3 pairs, then most dorsal shield setae distally swollen) ..... **26**
- 25 Dorsal shield lightly reticulate; ventrianal shield with one pair of preanal setae; seta *ZV3* present ..... ***Cocoseius elsalvador* Denmark & Andrews**
- Dorsal shield smooth, except for a few anterolateral striae; ventrianal shield with 3 pairs of preanal setae; seta *ZV3* absent.....
- ..... ***Cocoseius palmarum* Gondim Jr., Moraes & McMurtry**
- 26 Seta *S4* present; ventrianal shield with 3 pairs of preanal setae .....  
..... ***Typhlodromus transvaalensis* (Nesbitt)**
- Seta *S4* absent; ventrianal shield with 4 pairs of preanal setae..... **27**
- 27 Dorsal setae, except *j1*, *j3*, *s4*, *Z4* and *Z5*, short/ minute; setae *s4*, *Z4* and *Z5* greatly longer than other dorsal setae; setae *JV4* and *ZV3* present .....  
..... ***Leonseius regularis* (De Leon)**
- Pattern of dorsal setae lengths not as above; seta *JV4* absent; seta *ZV3* present or absent ..... **28**
- 28 Seta *S2* absent, seta *ZV3* present; seta *R1* inserted on unsclerotized cuticle; seta *s6* over twice as long as *R1*; seta *S5* longer than distance between its base and base of *Z5* ..... ***Typhlodromina subtropica* Muma & Denmark**
- Seta *S2* present, seta *ZV3* absent; seta *R1* on dorsal shield; seta *s6* less than twice as long as *R1*; seta *S5* not as long as distance between its base and base of *Z5*..... ***Metaseiulus ferlai* Moraes, McMurtry & Lopes**

## Discussion

Fifty-one phytoseiid species have been reported from Bahia (Bonato et al. 1999; Denmark and Muma 1973; Farias et al. 1981; Fiaboe et al. 2004, 2007; Lawson-Balagbo et al. 2008; Lofego et al. 2000, 2013; Moraes et al. 1993, 1994, 1997; Moraes and Denmark 1999; Moraes and McMurtry 1983; Noronha et al. 1997; Noronha and Moraes 1989; Oliveira et al. 2007; Souza et al. 2010, 2012). In the present study, fifteen species are reported for the first time in that state, raising the number of known species to sixty-six.

By far most of the species and of the specimens collected belong to Amblyseiinae (72 and 81%, respectively), followed by the Typhlodrominae (21 and 13%) and the Phytoseiinae (7 and 6%). Similar patterns were summarized by Castro and Moraes (2010) for similar surveys conducted in the Atlantic Forest of southeastern Brazil. There was a general trend for less specific phytoseiid species, i.e., those found on larger number of host plants, to be most abundant. Most of the phytoseiid species was found on a single or few host species. The largest numbers of phytoseiids on *C. nucifera*, *T. cacao* and *P. guajava* suggest that the microhabitat on the leaves of these plants favor these predators, but should not be taken to indicate the preference of these mites for those plants, given that the collecting effort was not the same on all plant species. These higher numbers could be due to the fact that these plants were among the most common in the localities where the study was conducted.

The most diverse genus in the present study was *Amblyseius*, as also found by Lawson-Balagbo et al. (2008) on coconut palm in the coastal region of Bahia. *Amblyseius operculatus* was the most abundant species and the species found in the largest number of plants. In total, the total number of specimens of this species was higher than the sum of the second and third most common species, and they were found in every month of the year, except (probably by chance) in February.

A noticeable absence in this study was mites belonging to the genus *Euseius* Wainstein. Although species of this genus have been reported as diverse and numerous in surveys conducted on different crops in the inland semiarid region of Bahia (Moraes et al. 1993; Moraes and McMurtry 1983), they were not found in the present study nor in previous surveys conducted in the southern coastal region of Bahia (A.R. Oliveira, personal observation) on different plant species.

Total annual rainfall in the semiarid region in the inland of Bahia ranges between 700 and 1,300 mm (Moraes et al. 1993; Moraes and McMurtry 1983), whereas in southern coast it is approximately 1,700 mm, with no pronounced dry season (Almeida and Valle 2010). Daud and Feres (2005) reported significant correlations between the population levels of *Euseius citrifolius* Denmark & Muma and rainfall (negative) or pollen abundance (positive). Pollen is known to constitute an important part of the diet of *Euseius* species (McMurtry et al. 2013). Thus, the apparent absence (or scarcity) of *Euseius* species in the present work could be related to the high rainfall in the southern coastal region and low pollen availability in the tropical fruit trees plantations surveyed. Species of this genus were not rare in a similar survey conducted in the coast of São Paulo state (Castro and Moraes 2010), where rainfall is quite similar to that reported in the southern coastal region of Bahia (Climate-Data.org 2015). This apparent discrepancy could be related to the fact that in that study these species were only found on plants of spontaneous growth, which could be protected at a certain level from the direct effect of rainfall. Those were not sampled in the present study.

The results of this study may contribute to the determination of future research themes, to subsidize future implementation of the use of phytoseiids as biological control agents in the region where the study was conducted. A next step in this trajectory

could involve studies under controlled laboratory conditions to evaluate the interactions between the most common predators found and the most common pest species.

## Acknowledgements

To Carlos H.W. Flechtmann (ESALQ, USP) for his invaluable support throughout the conduction of this work. To Afonso L.G.E. de Freitas for his help in collecting and mounting the mites. To the owners of the properties for allowing the collections. To CNPq (Brazilian National Council for Scientific and Technological Development) for the MSc scholarship granted to IVS (563094/2008-7). To CAPES (Coordination for the Improvement of Higher Education Personnel) for the PNPd grant to PSA. To the two anonymous reviewers for their valuable comments and suggestions which improved the manuscript. MGCG JR and GJM are CNPq researchers (301096/2009-0 and 302406/2009-3, respectively). This work was funded by the State of Bahia Research Foundation (FAPESB 7736/2006).

## References

- Almeida AAF, Valle RR (2010) Cacao: ecophysiology of growth and production. In: Da Matta FM (Ed.) *Ecophysiology of Tropical Tree Crops*. Nova Science Publishers Inc., Hauppauge, 37–70.
- Blommers L (1974) Species of the genus *Amblyseius* Berlese, 1914, from Tamatave, East Madagascar (Acarina: Phytoseiidae). *Bulletin Zoologisch Museum Universiteit van Amsterdam* 3: 143–155.
- Bonato O, Noronha ACS, Moraes GJ (1999) Distribution of *Amblyseius manihoti* (Acari, Phytoseiidae) on manioc and development of sampling plan. *Journal of Applied Entomology-Zeitschrift Fur Angewandte Entomologie* 123: 541–546. doi: 10.1046/j.1439-0418.1999.00421.x
- Castro TMMG, Moraes GJ (2010) Diversity of phytoseiid mites (Acari: Mesostigmata: Phytoseiidae) in the Atlantic Forest of São Paulo. *Systematics and Biodiversity* 8: 301–307. doi: 10.1080/14772001003801375
- Chant DA, Baker EW (1965) The Phytoseiidae (Acarina) of Central America. *Memoirs of the Entomological Society of Canada* 41: 1–56. doi: 10.4039/entm9741fv
- Chant DA, McMurtry JA (2007) *Illustrated Keys and Diagnoses for the Genera and Subgenera of the Phytoseiidae of the World (Acari: Mesostigmata)*. Indira Publishing House, Michigan, 219 pp.
- Chant DA, Yoshida-Shaul E (1989) Adult dorsal setal patterns in the family Phytoseiidae (Acari: Gamasina). *International Journal of Acarology* 15: 219–233. doi:10.1080/01647958908683852
- Chant DA, Yoshida-Shaul E (1991) Adult ventral setal patterns in the family Phytoseiidae (Acari: Gamasina). *International Journal of Acarology* 17: 187–199. doi: 10.1080/01647959108683906

- Climate-Data.org: Dados Climáticos para Cidades Mundiais. <http://pt.climate-data.org>
- Daud RD, Feres RJF (2005) Diversidade e flutuação populacional de ácaros (Acari) em *Mabea fistulifera* Mart. (Euphorbiaceae) de dois fragmentos de mata estacional semidecídua em São José do Rio Preto, SP. Neotropical Entomology 34: 191–201. doi: 10.1590/S1519-566X2005000200007
- De Leon D (1965) Phytoseiid mites from Puerto Rico with descriptions of new species (Acarina: Mesostigmata). The Florida Entomologist 48: 121–131. doi: 10.2307/3493102
- De Leon D (1967) Some mites of the Caribbean Area. Part I. Acarina on plants in Trinidad, West Indies. Allen Press Inc., Lawrence, 66 pp.
- Demite PR, Moraes GJ, McMurtry JA, Denmark HA, Castilho RC (2014) Phytoseiidae Database. [www.lea.esalq.usp.br/phytoseiidae](http://www.lea.esalq.usp.br/phytoseiidae) [accessed 15 Feb 2015]
- Denmark HA, Andrews KL (1981) Plant associated Phytoseiidae of El Salvador, Central America (Acarina: Mesostigmata). The Florida Entomologist 64: 147–158. doi: 10.2307/3494606
- Denmark HA, Muma MH (1972) Some Phytoseiidae of Colombia (Acarina: Phytoseiidae). The Florida Entomologist 55: 19–29. doi: 10.2307/3493192
- Denmark HA, Muma MH (1973) Phytoseiidae mites of Brazil (Acarina, Phytoseiidae). Revista Brasileira de Biologia 33: 235–276.
- Denmark HA, Muma MH (1989) A revision of the genus *Amblyseius* Berlese, 1914 (Acari: Phytoseiidae). Occasional Papers of the Florida State Collection of Arthropods 4: 1–149.
- Ehara S (1966) Some mites associated with plants in the State of São Paulo, Brazil, with a list of plant mites of South America. Japanese Journal of Zoology 15: 129–149.
- El-Banhawy EM (1984) Description of some phytoseiid mites from Brazil (Acarina: Phytoseiidae). Acarologia 25: 125–144.
- Farias ARN, Flechtmann CHW, Moraes GJ, McMurtry JA (1981) Predadores do ácaro verde da mandioca, no nordeste do Brasil. Pesquisa Agropecuária Brasileira 16: 313–317.
- Ferla NJ, Johann L, Botton M, Majolo F, Klock CL (2011) Phytoseiid mites associated with vineyards in the Rio Grande do Sul State, Brazil Zootaxa 2976: 15–31.
- Fiaboe KKM, Gondim Jr. MGC, Moraes GJ, Ogol CKPO, Knapp M (2007) Surveys for natural enemies of the tomato red spider mite *Tetranychus evansi* (Acari: Tetranychidae) in northeastern and southeastern Brazil. Zootaxa: 33–58.
- Fiaboe KKM, Moraes GJ, Gondim Jr. MGC (2004) A new genus and a new species of phytoseiid mite (Acari: Phytoseiidae) from northeastern Brazil. Zootaxa: 1–4.
- Garman P (1958) New species belonging to the genera *Amblyseius* and *Amblyseiopsis* with keys to *Amblyseius*, *Amblyseiopsis* and *Phytoseiulus*. Annals of the Entomological Society of America 51: 69–79. doi: 10.1093/aesa/51.1.69
- Gondim Jr. MGC, Moraes GJ (2001) Phytoseiid mites (Acari: Phytoseiidae) associated with palm trees (Arecaceae) in Brazil. Systematic and Applied Acarology 6: 65–94.
- Gondim Jr. MGC, Moraes GJ, McMurtry JA (2000) A New Species of *Cocoseius* (Acari: Phytoseiidae) from Brazil and Redefinition of the Genus. Annals of the Entomological Society of America 93: 1226–1229. doi: 10.1603/0013-8746(2000)093[1226:ansoca]2.0.co;2
- Hoy MA (2011) The Phytoseiidae: Effective Natural Enemies. In: Agricultural acarology: introduction to integrated mite management. CRC Press, Boca Raton, Florida, 159–177. doi: 10.1201/b10909-16

- Lawson-Balagbo LM, Gondim Jr. MGC, Moraes GJ, Hanna R, Schausberger P (2008) Exploration of the acarine fauna on coconut palm in Brazil with emphasis on *Aceria guerreronis* (Acari : Eriophyidae) and its natural enemies. *Bulletin of Entomological Research* 98: 83–96. doi: 10.1017/s0007485307005421
- Lindquist EE, Evans GO (1965) Taxonomic Concepts in the Ascidae, with a Modified Setal Nomenclature for the Idiosoma of the Gamasina (Acarina: Mesostigmata). *Memoirs of the Entomological Society of Canada* 97: 5–66. doi: 10.4039/entm9747fv
- Lofego AC, Demite PR, Kishimoto RG, Moraes GJ (2009) Phytoseiid mites on grasses in Brazil (Acari: Phytoseiidae). *Zootaxa* 2240: 41–59.
- Lofego AC, Moraes GJ, Castro LAS (2004) Phytoseiid mites (Acari: Phytoseiidae) on Myrtaceae in the State of São Paulo, Brazil. *Zootaxa* 516: 1–18. doi: 10.1080/00222933.2011.596950
- Lofego AC, Moraes GJ, McMurtry JA (2000) Three new species of phytoseiid mites (Acari: Phytoseiidae) from Brazil. *Anais da Sociedade Entomológica do Brasil* 29: 461–467. doi: 10.1590/S0301-80592000000300008
- Lofego AC, Rezende JM, Verona RLC, Feres RJF (2013) Mites (Acari) associated with three species of the genus *Jatropha* (Euphorbiaceae) in Brazil, with emphasis on *Jatropha curcas*. 2013: 13. doi: 10.11158/saa.18.4.10
- McMurtry JA (1983) Phytoseiid mites from Guatemala, with description of two new species and redefinition of the genera *Euseius*, *Typhlodromus* and the *Typhlodromus occidentalis* species group (Acari: Mesostigmata). *International Journal of Acarology* 25: 249–272.
- McMurtry JA, Moraes GJ, Sourassou NF (2013) Revision of the lifestyles of phytoseiid mites (Acari: Phytoseiidae) and implications for biological control strategies. *Systematic & Applied Acarology* 18: 297–320. doi: 10.11158/saa.18.4.1
- Moraes GJ, Alencar JA, Lima JLS, Yaninek JS, Delalibera Jr. I (1993) Alternative plant habitats for common phytoseiid predators of the cassava green mite (Acari: Phytoseiidae, Tetranychidae) in northeast Brazil. *Experimental and Applied Acarology* 17: 77–90.
- Moraes GJ, Barbosa MFC, Castro TMMG (2013) Phytoseiidae (Acari: Mesostigmata) from natural ecosystems in the State of São Paulo, Brazil. *Zootaxa* 3700: 47. doi: 10.11646/zootaxa.3700.3.1
- Moraes GJ, Denmark HA (1999) The genus *Proprioseius* Chant (Acari: Phytoseiidae), with descriptions of two new species from Brazil. *Systematic and Applied Acarology* 4: 97–102. doi: 10.1007/BF00156945
- Moraes GJ, Flechtmann CHW (2008) *Manual de Acarologia, Acarologia Básica e Ácaros de Plantas Cultivadas no Brasil*. Holos Ribeirão Preto, 308 pp.
- Moraes GJ, McMurtry JA (1983) Phytoseiid mites (Acarina) of northeastern Brazil with descriptions of four new species. *International Journal of Acarology* 9: 131–148. doi: 10.1080/01647958308683326
- Moraes GJ, McMurtry JA, Denmark HA, Campos CB (2004) A revised catalog of the mite family Phytoseiidae. *Zootaxa* 434: 1–494.
- Moraes GJ, McMurtry JA, Lopes PC (2006) Redefinition of *Metaseiulus* Muma (Acari: Phytoseiidae) and description of a new species from Brazil. *International Journal of Acarology* 32: 351–354. doi: 10.1080/01647950608684481

- Moraes GJ, Melo EL, Gondim Jr. MGC (1997) Description of a new species of phytoseiid mite from Northeastern Brazil and redescription of *Neoseiulus gracilis* (Acari: Phytoseiidae). The Florida Entomologist 80: 319–324. doi: 10.2307/3495765
- Moraes GJ, Mesa NC, Braun A (1991) Some phytoseiid mites of Latin America (Acari: Phytoseiidae). International Journal of Acarology 17: 117–139. doi: 10.1080/01647959108683892
- Moraes GJ, Mesa NC, Braun A, Melo EL (1994) Definition of the *Amblyseius limonicus* species group (Acari: Phytoseiidae), with descriptions of two new species and new records. International Journal of Acarology 20: 209–217. doi: 10.1080/01647959408684019
- Muma MH (1955) Phytoseiidae (Acarina) associated with citrus in Florida. Annals of the Entomological Society of America 48: 262–272. doi: 10.1093/aesa/48.4.262
- Muma MH (1961) Subfamilies, genera, and species of Phytoseiidae (Acarina: Mesostigmata). Bulletin of the Florida State Museum Biological Sciences 5: 267–302.
- Muma MH, Demark HA, De Leon D (1970) Phytoseiidae of Florida. Arthropods of Florida and Neighboring Land Areas, 6. Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Gainesville, 150.
- Muma MH, Denmark HA (1969) The *conspicua* species-group of *Typhlodromina* Muma, 1961. Annals of the Entomological Society of America 62: 406–413. doi: 10.1093/aesa/62.2.406
- Narayanam ES, Kaur RB, Ghai S (1960) Importance of some taxonomic characters in the family Phytoseiidae Berl., 1916, (predatory mites) with new records and descriptions of species. Proceedings of the National Institute of Science of India 26B: 384–394.
- Nesbitt HHJ (1951) A taxonomic study of the Phytoseiidae (Family Laelaptidae) predaceous upon Tetranychidae of economic importance. Zoologische Verhandelingen 12: 64 pp. + 32 plates.
- Noronha ACS, Carvalho JEB, Caldas RC (1997) Ácaros em citros nas condições de Tabuleiros Costeiros. Revista Brasileira de Fruticultura 19: 373–376.
- Noronha ACS, Moraes GJ (1989) Flutuação populacional do ácaro verde da mandioca e seus predadores fitoseídeos (Acari: Tetranychidae, Phytoseiidae) em Cruz das Almas - Bahia. Revista Brasileira de Mandioca 8: 31–39.
- Oliveira VS, Noronha ACS, Argolo PS, Carvalho JEB (2007) Acarofauna em pomares cítricos nos municípios de Inhambupe e Rio Real no Estado da Bahia. Magistra 19: 257–261.
- Rowell HJ, Chant DA, Hansell RIC (1978) The determination of setal homologies and setal patterns on the dorsal shield in the family Phytoseiidae (Acarina: Mesostigmata). The Canadian Entomologist 110: 859–876. doi: 10.4039/ent110859-8
- Santos-Serejo JA, Dantas JLL, Sampaio CV, Coelho YS (2009) Fruticultura Tropical: espécies regionais e exóticas. Embrapa Informação Tecnológica, Brasília, 509 pp.
- Schicha E (1981) A new species of *Amblyseius* (Acari: Phytoseiidae) from Australia compared with ten closely related species from Asia, America & Africa. International Journal of Acarology 7: 203–216. doi: 10.1080/01647958108683262
- Souza IV, Gondim Jr. MGC, Ramos ALR, Santos EA, Ferraz MIF, Oliveira AR (2012) Population dynamics of *Aceria guerreronis* (Acari: Eriophyidae) and other mites associated with coconut fruits in Una, state of Bahia, northeastern Brazil. Experimental and Applied Acarology 58: 221–233. doi: 10.1007/s10493-012-9576-3



- Souza IV, Oliveira AR, Gondim Jr. MGC (2010) A new species of the genus *Typhlodromips* De Leon (Acari: Phytoseiidae) from the state of Bahia, Brazil. *International Journal of Acarology* 36: 49–52. doi: 10.1080/01647950903508490
- Swirski E, Shechter R (1961) Some phytoseiid mites (Acarina: Phytoseiidae) of Hong-Kong, with a description of a new genus and seven new species. *The Israel Journal of Agricultural Research* 11: 97–117.
- Yoshida-Shaul E, Chant DA (1988) Descriptions of two unusual new species in the genus *Amblyseius* Berlese (Acari: Phytoseiidae). *Canadian Journal of Zoology* 66: 2053–2056. doi: 10.1139/z88-302
- Yoshida-Shaul E, Chant DA (1997) A world review of the genus *Phytoscutus* Muma (Phytoseiidae: Acari). *Acarologia* 38: 219–238.