

# A new species of *Isoperla* (Insecta, Plecoptera) from the Karawanken, with considerations on the Southern Limestone Alps as centers of endemism

Wolfram Graf<sup>1</sup>, Martin Konar<sup>2</sup>, Dávid Murányi<sup>3</sup>,  
Kirill Márk Orci<sup>4</sup>, Simon Vitecek<sup>5</sup>

**1** Institute of Hydrobiology and Aquatic Ecosystem Management, University of Natural Resources and Applied Life Sciences, Max Emanuel-Strasse 17, A-1180 Vienna, Austria **2** Amt der Kärntner Landesregierung, Abteilung 8 (Kompetenzzentrum Umwelt, Wasser und Naturschutz), Kärntner Institut für Seenforschung, 9020 Klagenfurt am Wörthersee, Kirchengasse 43, Austria **3** Department of Zoology, Hungarian Natural History Museum, Baross u. 13, H-1088 Budapest, Hungary **4** MTA-ELTE-MTM Ecology Research Group, Pázmány Péter s. 1/C, H-1117 Budapest, Hungary **5** Department of Limnology & Bio-Oceanography, Faculty of Life Sciences, University of Vienna, Althanstrasse 14, A-1090 Vienna, Austria

Corresponding author: *Wolfram Graf* (wolfram.graf@boku.ac.at)

---

Academic editor: *M. Gottardo* | Received 27 August 2014 | Accepted 30 September 2014 | Published 20 October 2014

---

<http://zoobank.org/5B6A06E7-A1A6-4B58-B14B-6E7905DD7EF2>

---

**Citation:** Graf W, Konar M, Murányi D, Orci KM, Vitecek S (2014) A new species of *Isoperla* (Insecta, Plecoptera) from the Karawanken, with considerations on the Southern Limestone Alps as centers of endemism. ZooKeys 448: 27–36. doi: 10.3897/zookeys.448.8509

---

## Abstract

A new species of the genus *Isoperla* (Plecoptera, Perlodidae), belonging to the *oxylepis* species-group is described, and the male mating call is characterized. Its range falls within a small region of the Southern Limestone Alps which is well known to be one endemism-centre of aquatic insects.

## Keywords

*Isoperla*, new species, endemism, Austria, Slovenia, Southern Alps

## Introduction

The genus *Isoperla* consists of about 150 species (DeWalt et al. 2011, Baumann and Lee 2009, Murányi 2011, Szczytko and Stewart 1979, Zwick and Surenkhorloo 2005) and covers the Holarctic and Oriental regions. In Europe 56 species are known so far (Graf et al. 2009, Murányi 2011), of which ten occur in Austria (Graf 2010). *Isoperla* is a morphologically difficult genus, especially the *grammatica* and *tripartita* species groups that both exhibit high variability, and requires further resolution. A synthesis of zoogeographical, morphological, molecular, and possibly behavioural data will be required to get full knowledge on the diversity of this highly interesting genus.

Recently a series of specimens were collected from the Karawanken Alps in southern Austria and the nearby Kamnik Alps in northern Slovenia deviating from all hitherto known species. In this paper we provide morphological descriptions of males, females and the larva. Additionally we illustrate drumming signals of one male.

## Material and methods

Adult specimens were collected using sweep nets, larvae were collected by handpicking from cobbles (mesolithal), the dominant substrate type. Collected specimens were stored in 70% ethanol. Morphological characteristics of male terminalia were examined in KOH-treated, cleared specimens. Comparative material from the authors' collections enabled the identification of the new species.

Vibratory signal recordings were made using a small, dynamic speaker (SAL YD78) as a vibration transducer. The speaker was connected to the microphone input socket of a solid state, digital recorder (Zoom H4n). The examined specimen was placed on the diaphragm of the speaker. To prevent the specimen from escaping the speaker was covered by a sheet of hobby glass. During the recordings ambient air temperature was measured using a P 300W thermometer. Vibration recordings were analysed and oscillograms produced using the software Adobe Audition 1.5 (Adobe Systems Incorporated, San Jose, California, USA). Drumming signal terminology follows Stewart and Sandberg (2006) and Murányi et al. (2014).


## Results

### *Isoperla claudiae* Graf & Konar, sp. n.

<http://zoobank.org/50F79ECE-AD68-4DD1-BD7F-643D25205189>

Figs 1–3

**Type material.** Holotype: 1 male, Austria, Carinthia, Dolintschitschach brook south-east of Feistritz ob Bleiburg (46°32'6"N, 14°45'52"E), 600m a.s.l., 30.5.2014, leg.



**Figure 1.** *Isoperla claudiae* sp. n. **A** habitus **B** colouration of the head of *I. claudiae* **C** ventral view of the male abdomen with extruded penis **D** ventral view of the female abdomen.

W. Graf; Paratypes: 3 males, 2 females, same place, date and collector. The holotype is deposited at the Linzer Landesmuseum, Linz, Austria, paratypes are stored in the first author's collection.

Other material. 1 male (drumming call examined), 1 female (HNHM: PLP4333), Slovenia, Upper Carniola, Kamnik municipality, Kamnik Alps, small forest brook S of Podvolovljek Pass (46°16.250'N 14°41.325'E), 980m a.s.l., 09.07.2013, leg. D. Murányi, I. Sivec.

**Type locality.** Austria, Carinthia, Feistriz ob Bleiburg, Dolintschitschach brook.

**Etymology.** The species is named in honour of the second author's wife Claudia.

**Diagnosis.** An *Isoperla* exhibiting the following combination of characters: (1) a small medial penial armature in the form of an equilateral triangle, lacking lateral penial armatures; (2) yellow head and pronotum with a small horseshoe-like brown marking connecting the ocelli.

**Description.** Medium-sized species, macropterous. Body length: holotype 10.5 mm, allotypes 11–12 mm; forewing length: holotype 12 mm, paratypes 12–14 mm. Primary colouration yellow, head and pronotum mostly yellow with dark brown


markings; pilosity short. Primary colouration of the head yellow, with a dark horse-shoe-like brown patch connecting the three ocelli (Fig. 1A, B). Occiput with indistinct rugosities but with brown patches laterally. Eyes normal sized. Scape brown, pedicel and the following antennomeres brown; palpi greyish to light brown. Pronotum yellow with a delicate brownish marking at the posterior margin, trapezoidal, edges angled; rugosities hardly visible and yellow. Anterior part of the mesonotum yellow, remaining portions brown; metanotum medially dark brown, laterally and anterior of the insertion of wings whitish. Wings yellowish, particularly the anterior half; venation mostly whitish to yellow, costa and apical part of radii brown. Ventral surface of thorax pale, meso- and metabasisternum inconspicuous, furcasternites and furcal pits pale. Femora brown dorsally and yellow ventrally. Tibiae brownish dorsally, pale ventrally; tarsi brown.

Male abdomen (Fig. 1C): 1<sup>st</sup> to 7<sup>th</sup> tergite dorsally brown (with some tiny pale spots) with increasing laterally whitish areas towards the apex, 8<sup>th</sup> to 10<sup>th</sup> tergite mostly yellowish with small brown medial patches and medially interrupted anterior stripes up to T9, T10 pale without markings. Laterally and ventrally all segments whitish to yellow, lacking dark markings. Pilosity on segment posterior ends short and inconspicuous. Ventral lobe of sternite VIII yellow, slightly longer than wide, its posterior margin strongly convex with long marginal pilosity. Sternite IX yellowish. Paraprocts brown, regularly curved in caudal with blunt tips; cerci light brown, apically dark brown.

Penis (Fig. 2): Divided into four lobes and a basal section in extruded position. Medial penial armature located on the medial lobe adjacent to the ventral lobe, lateral penial armatures lacking. The medial penial armature resembles an equilateral triangle of 130  $\mu\text{m}$  width and 97  $\mu\text{m}$  length formed by slightly brownish coloured scales that are relatively blunt and short and vary in length (4.98–6.26  $\mu\text{m}$ ). The median basal area is sparsely covered by shorter scales. The medial penial armatures are connected distally by a narrow band of colourless scales with an area densely covered by smaller triangular scales. Similar scales are located proximal to the medial armatures. Their length varies between 4.4 and 7.2  $\mu\text{m}$ . With the exception of the medial armatures the central area of the ventral penis is bald. Lateral portions of lateral lobes covered by dense scales similar to the ones on medial and ventral lobes, being denser at the connection to the ventral lobe.


Female abdomen (Fig. 1D): 1<sup>st</sup> to 7<sup>th</sup> tergite dorsally brown with increasing laterally whitish areas towards the apex, 8<sup>th</sup> to 10<sup>th</sup> tergite mostly yellowish with small brown medial patches. Laterally and ventrally all segments entirely whitish with dark markings reduced to delicate brownish lines at the posterior end of sternites. Subgenital plate covers most of sternite VIII width and half of sternite IX length, posterior margin rounded semicircularly. Sternite X and paraprocts yellowish; cerci generally brownish, the first segment being pale.

Larva (Fig. 3): Body length of the matured larva: 13 mm. General colour brown but with pale markings. Pilosity dense, pronotal, posterior tergal and cercal fringes short and acute; swimming hairs lacking. Head dark brown with two yellow spots




**Figure 2.** Penis of *Isoperla claudiae* sp. n. **A** ventral view of the extruded penis **B** medial penial armature, scale bar 50  $\mu\text{m}$  **C** medial penial armature, scale bar 50  $\mu\text{m}$  **D** scales of the medial penial armature, scale bar 20  $\mu\text{m}$  **E** scales found caudally the medial penial armature, scale bar 20  $\mu\text{m}$  **F** medial penial armature of *I. orobica*, scale bar 200  $\mu\text{m}$ . Photographs **A–E** by W. Lechthaler, Vienna.

anterior to the M-line, two posterior to the M-line, one around the median ocellus and one laterally to the each posterior ocellus. Two large pale spots laterally on the occipit (Fig. 3A). M-line distinct, tentorial callosities hardly visible; eyes normal



**Figure 3.** Larval characters of *I. claudiae* sp. n. **A** mature larva of *I. claudiae* sp. n. in dorsal view, head and pronotum **B** ventral view of the abdomen end of *I. claudiae* sp. n. **C** lacinia of *I. claudiae* sp. n.

sized. Scape and pedicel pale, the following antennomeres light brown; palpi yellowish, mouthparts light brown. Lacinia triangular, with 6 strong setae beneath the two apical teeth, thin hairs present all along the inner margin; galea with scattered setae on the whole surface (Fig. 3C). Pronotum rectangular with rounded corners, twice as wide as long, brown but with a narrow medial pale stripe along the medial suture and a marbled impression due to several medial pale areas, lateral parts uniformly brown, margins laterally pale. Mesonotum and metanotum mostly brown but with a pale, marmoreal pattern; wingpads brownish. Ventral surface of thorax pale, furcasternites and furcal pits inconspicuous. Legs uniformly pale. Abdominal tergites brown with a pair of roundish pale spots laterally to a median, darker area. The spots are increasing in size towards the entirely pale last tergite. Ventral surface of abdomen pale brown, the distal segments darker. Paraprocts brown; cerci light brown with dense circumferential rows of bristles of varying length at the end of each segment.



**Figure 4.** Oscillograms showing the drumming call of an *I. claudiae* sp. n. male (ambient air temperature 24.2 °C). **A** oscillogram showing rhythm and amplitude variation patterns of a call **B** a faster oscillogram of two bi-beats from the second half of the call **C** variation of interval duration between bi-beats during the call presented in **A**. Inter-beat intervals were measured from the amplitude peak of one bi-beat to the amplitude peak of the next bi-beat (measured on the same call presented in **A**).

**Ecology and distribution.** The species was collected in a small spring-brook at 535 m a.s.l. in the Karawanken, and a small forest brook at 980 m a.s.l. in the Kamnik Alps (Southern Limestone Alps).

**Preliminary description of the male drumming call.** Since only one signal from a single male could be recorded we cannot give any information on the variation range of the signal parameters in this species. The aim of this preliminary description is only to report the basic features of the signal, but even that should be treated with some caution since we cannot be sure whether or not the recorded signal shows some deviant features.

As it is observable in (Fig. 4A) the male call is a sequence of bi-beats. After an initial crescendo the peak amplitude of bi-beats fluctuate around a constant value. In bi-beats the first beat is of lower amplitude (missing in the low amplitude initial part of the call and sometimes hardly detectable even in the main part of the signal), the second one is of higher amplitude and followed by a long, decaying wave train (Fig. 4B). Inter beat interval within bi-beats varied between 8–20 ms during the call. The interval between bi-beats (or single beats at the initial part) gradually increased during the sequence (except for a short initial part of the sequence where inter beat interval decreases) varying between 230–350 ms (Fig. 4C).

## Discussion

### Relationships

The new species can be attributed to the *oxylepis* species-group sensu Murányi (2011), currently comprising *I. oxylepis oxylepis* (Despax), *I. oxylepis balcanica* Raušer, *I. bosnica* Aubert, *I. orobica* Ravizza and *I. submontana* Raušer. These species develop similarly shaped medial penial armatures and scales of penial armatures, lack real lateral armatures, and develop dense, uncoloured scales on each lobe.

*Isoperla claudiae* sp. n. is most similar to *I. orobica*, a species restricted to the southwestern Alps, but can be easily distinguished from the latter species as the scales of the medial penial armature are shorter in *I. claudiae* sp. n., a higher density of uncoloured scales on the penis in *I. claudiae* sp. n., as well as yellow, hardly visible rugosities of the pronotum in *I. claudiae* sp. n.

The male drumming call of *I. claudiae* sp. n. is clearly different from the drumming call of *I. oxylepis*, which is the only species of the *I. oxylepis* species-group, where published information regarding the vibratory signals is available (Rupprecht 1969, 1983). Amongst the European species of *Isoperla* the male call of *I. claudiae* sp. n. is most similar to that of *I. rivulorum* (Pictet) (Rupprecht 1969, Tierno de Figueroa and Sánchez-Ortega 1999, Tierno de Figueroa et al. 2000, 2002, 2011, Tierno de Figueroa and Luzón-Ortega 2002, Luzón-Ortega et al. 2010), but the beat group repetition period seems to be longer in this species (230–350 ms, 24.2 °C, Fig. 4C) than in *I. rivulorum* (Luzón-Ortega et al. 2010) reported 103–163 ms at 20 °C), and *I. rivulorum* frequently produces 3 beats per beat group.

### The Southern Limestone Alps as centers of endemism

The southern slopes of the Alps from the Ligurian Prealps in the southwest to the Pohorje Mountains in the east are densely covered by microendemic species. Concentrations of endemic species in the south and south-eastern Alps are well known among Trichoptera species (Malicky 1983, 2000); regarding Plecoptera *Leuctra dylani* Graf, *L. juliettae* Vinçon & Graf, *L. muranyii* Vinçon & Graf and *Protonemura bipartita* Consiglio are restricted to small areas from the Bergamo prealps to the Lessinian Alps; the apterous *L. istenicae* Sivec and *Siphonoperla ottomoogi* Graf are nested as microendemics in southeastern refugia referred to as the Steirische Randgebirge.

The western alpine slopes (Biellesse, Graian and Cottian Alps) are another area of alpine endemism where a high diversity within the genus *Leuctra* is found (Ravizza Dematteis and Ravizza 1988; Vinçon and Ravizza 1998), and their distribution patterns are associated with the presence of nunataks during the Würm glaciation (Ravizza and Ravizza Dematteis 1993, 1994). The hitherto known range of *Isoperla claudiae* sp. n. fits well in this hot-spot of biodiversity and supports the Dinodal theory of Malicky (1983, 2000), which suggests glacial species-specific refugia

within the Alps based on distribution patterns of endemic caddisfly species. Most of these microendemic species are stenoeious elements of springs and small streams in medium altitudes.

## Acknowledgements

The authors acknowledge support from the BioFresh EU project–Biodiversity of Freshwater Ecosystems: Status, Trends, Pressures and Conservation Priorities (contract No. 226874). The bioacoustic research in this study was supported by the Hungarian Scientific Research Fund (OTKA K 81929). This study further was morally supported by the Austrian Science Fund (FWF) (project number P23687-B17).

## References

- Baumann RW, Lee JJ (2009) Two interesting new species of *Isoperla* from Northern California (Plecoptera: Perlodidae). *Illiesia* 5: 1–10.
- DeWalt RE, Neu-Becker U, Steuber G (2011) Plecoptera Species File Online. Version 1.1/4.0, <http://Plecoptera.SpeciesFile.org>
- Graf W, Lorenz AW, Tierno de Figueroa JM, Lücke S, López-Rodríguez MJ, Davies C (2009) Distribution and Ecological Preferences of European Freshwater Organisms - Volume 2: Plecoptera. In: Schmidt-Kloiber A, Hering D (Eds) *Distribution and Ecological Preferences of European Freshwater Organisms*. Pensoft Publishers, Sofia-Moscow, 262 pp.
- Graf W (2010) Aktualisierte Check-Liste der Steinfliegen (Insecta: Plecoptera) Österreichs. *Lauterbornia, Dinkelscherben* 71: 175–183.
- Luzón-Ortega JM, López-Rodríguez MJ, Tierno de Figueroa JM (2010) Confirmation of the presence of *Isoperla rivulorum* (Pictet, 1841) (Plecoptera, Perlodidae) and first data of the male drumming call in the Iberian Peninsula. *Boletín de la Asociación Española de Entomología* 34(3/4): 441–443.
- Malicky H (1983) Chlorological patterns and biome types of European Trichoptera and other freshwater insects. *Archiv für Hydrobiologie* 96: 223–244.
- Malicky H (2000) Arealodynamik und Biomgrundtypen am Beispiel der Köcherfliegen (Trichoptera). *Entomologica Basiliensia* 22: 235–259.
- Murányi D (2011) Balkanian species of the genus *Isoperla* Banks, 1906 (Plecoptera: Perlodidae). *Zootaxa* 3049: 1–46.
- Murányi D, Gamboa M, Orci KM (2014) *Zwicknia* gen. n. for the *Capnia bifrons* species group, with descriptions of three new species based on morphology, drumming signals and molecular genetics, and a synopsis of the West Palaearctic and Nearctic genera of Capniidae (Plecoptera). *Zootaxa* 3812: 001–082.
- Ravizza Dematteis E, Ravizza C (1988) Les Plécopteres de la vallée supérieure du Po (Alpes Cottiennes). *Notes faunistiques et écologiques. Annales de Limnologie* 24: 243–360. doi: 10.1051/limn/1988021

- Ravizza C, Ravizza Dematteis E (1993) Zoogeographical aspects of the Plecoptera population of the Biellese mountains (Pennine Alps, Italy). *Bollettino della Società Entomologica Italiana*, Genova 125: 6–22.
- Ravizza C, Ravizza Dematteis E (1994) *Leuctra vinconi aubertorum*, a new subspecies of *Leuctra* from the Ticino canton, Switzerland (Plecoptera, Leuctridae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 67: 37–41.
- Rupprecht R (1969) Zur Artspesifität der Trommelsignale der Plecopteren (Insecta). *Oikos* 20: 26–33. doi: 10.2307/3543742
- Rupprecht R (1983) Kommunikationssignale von Arten der Gattung *Isoperla* (Plecoptera) und deren Eignung zur Abgrenzung von Arten. *Verhandlungen der Deutschen zoologischen Gesellschaft* 76: 198.
- Szczytko SW, Stewart KW (1979) The genus *Isoperla* (Plecoptera) of Western North America; holomorphology and systematics, and a new stonefly genus *Cascadoperla*. *Memoirs of the Entomological Society of America* 32: 1–120.
- Vinçon G, Ravizza C (1998) Les Leuctridé (Plecoptera, Leuctridae) des Alpes. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 71: 285–342.
- Stewart KW, Sandberg JB (2006) Vibratory communication and mate searching behaviour in stoneflies. In: Drosopoulos S, Claridge MF (Eds) *Insect sounds and communication: Physiology, behaviour, ecology and evolution*. Taylor & Francis, CRC Press, Boca Raton, 179–186.
- Tierno de Figueroa JM, Sánchez-Ortega A (1999) The male drumming call of *Isoperla nevada* Aubert, 1952 (Plecoptera, Perlodidae). *Aquatic Insects* 21: 33–38. doi: 10.1076/aqin.21.1.33.4543
- Tierno de Figueroa JM, Luzón-Ortega JM, Sánchez-Ortega A (2000) Male calling, mating and oviposition in *Isoperla curtata* (Plecoptera, Perlodidae). *European Journal of Entomology* 97: 171–175. doi: 10.14411/eje.2000.032
- Tierno de Figueroa JM, Luzón-Ortega JM (2002) The drumming call of *Isoperla bipartita* Aubert, 1962 (Plecoptera, Perlodidae). *Aquatic Insects* 24: 87–90. doi: 10.1076/aqin.24.2.87.4901
- Tierno de Figueroa JM, Luzón-Ortega JM, Stewart KW (2002) The drumming of *Isoperla pallida* Aubert and *Guadalgenus franzi* (Aubert) (Plecoptera, Perlodidae), and review and evolutionary considerations of Southern Iberian Peninsula Perlodidae. *Zoological Science* 19: 871–875. doi: 10.2108/zsj.19.871
- Tierno de Figueroa JM, Gaetani B, Luzón-Ortega JM, López-Rodríguez MJ, Fochetti R (2011) On the identity of *Isoperla curtata* (Plecoptera: Perlodidae): behavioural and molecular approaches show the existence of two separate species. *Zootaxa* 3000: 49–58.
- Zwick P, Surenkhorloo P (2005) The Mongolian species of *Isoperlinae* (Plecoptera: Perlodidae). *Acta Zoologica Academiae Scientiarum Hungaricae* 51: 253–277.