

Checklist of the families Chaoboridae, Dixidae, Thaumaleidae, Psychodidae and Ptychopteridae (Diptera) of Finland

Jukka Salmela¹, Lauri Paasivirta², Gunnar M. Kvifte³

1 *Metsähallitus, Natural Heritage Services, P.O. Box 8016, FI-96101 Rovaniemi, Finland* **2** *Ruuhikoskenkatu 17 B 5, 24240 Salo, Finland* **3** *Department of Limnology, University of Kassel, Heinrich-Plett-Str. 40, 34132 Kassel-Oberzwehren, Germany*

Corresponding author: *Jukka Salmela* (jukka.e.salmela@gmail.com)

Academic editor: *J. Kahanpää* | Received 17 March 2014 | Accepted 22 May 2014 | Published 19 September 2014

<http://zoobank.org/87CA3FF8-F041-48E7-8981-40A10BACC998>

Citation: Salmela J, Paasivirta L, Kvifte GM (2014) Checklist of the families Chaoboridae, Dixidae, Thaumaleidae, Psychodidae and Ptychopteridae (Diptera) of Finland. In: Kahanpää J, Salmela J (Eds) Checklist of the Diptera of Finland. ZooKeys 441: 37–46. doi: 10.3897/zookeys.441.7532

Abstract

A checklist of the families Chaoboridae, Dixidae, Thaumaleidae, Psychodidae and Ptychopteridae (Diptera) recorded from Finland is given. Four species, *Dixella dyari* Garret, 1924 (Dixidae), *Threticus tridactylis* (Kincaid, 1899), *Panimerus albifacies* (Tonnoir, 1919) and *P. przhiboroi* Wagner, 2005 (Psychodidae) are reported for the first time from Finland.

Keywords

Finland, Diptera, species list, biodiversity, faunistics

Introduction

Psychodidae or moth flies are an intermediately diverse family of nematocerous flies, comprising over 3000 species world-wide (Pape et al. 2011). Its taxonomy is still very unstable, and multiple conflicting classifications exist (Duckhouse 1987, Vaillant 1990, Ježek and van Harten 2005). The nomenclature used in the Finnish check-list follows the

Norwegian one (Kvifte et al. 2011). For differences between the taxonomy used herein and that used in previous works on Finnish Psychodidae (e.g. Hackman 1980, Salmela 2003, 2005, Salmela et al. 2007, Autio and Salmela 2010) see Kvifte et al. (2011). Two species, *Pneumia nubila* and *Pneumia palustris*, listed in the previous check-lists by Hackman (1980) and Wagner (2013) have not been confirmed through study of material. Of these, *P. palustris* have been suggested to be synonymous with *Pneumia gracilis* (Eaton, 1893) (Withers 1989). Both these species have been linked to some taxonomic confusion in the past, and thus the record of *P. nubila* may represent *Pneumia trivialis*. The present check-list contains the first Finnish records of the species *Threticus tridactilis* (Kincaid, 1899), *Panimerus albifacies* (Tonnoir, 1919) and *P. przhiboroi* Wagner, 2005 (see Notes).

Finnish Psychodidae are very well studied compared to other Nordic countries, however new species are still being discovered at a regular rate. Most of these are new country records, but in recent years the fauna has also yielded some species new to science (Salmela et al. 2012, Kvifte unpubl.). This trend is likely to continue. Probably between twelve and thirty further species can be expected to be found in the Finnish fauna. There are no good identification sources available for Psychodids, as all works suffer from taxonomic or nomenclatural errors and/or insufficient descriptions. The best works relevant to the Finnish fauna are Vaillant (1971-1983), Withers (1989) and Wagner (1997c, genera only).

Phantom midges (Chaoboridae) are close relatives to mosquitoes (Culicidae) and meniscus midges (Dixidae) (Saether 1970, 1997). There are around 50 known species of phantom midges in the world and 15 in the palearctic region, of which five are holarctic species. In Finland, there are eight species in three genera (Hirvenoja 1960, Hackman 1980, in which the species in the genus *Mochlonyx* were still doubtful).

The only strictly lacustrine species in Finland is *Chaoborus flavicans*. The other species live in small fishless lakes, ponds and springs where their predators are mainly dragon flies, dytiscid beetles and frogs. *Chaoborus nyblaei* and *Cryophila lapponica* are found from small fishless ponds on fells and ponds created by melting water. However, in 2013 larvae of *C. nyblaei* were found from the boreal zone, NE Lapland (Savukoski) from a small but permanent pond. *Chaoborus pallidus* lives in eutrophic lakes and small lakes in southern and central Finland (Hirvenoja 1960, 1963, Paasivirta 2002). However, there have not been any systematic surveys of the fauna across the biogeographic provinces and the conservation statuses of the species have not been evaluated. Finnish species can be identified by using Saether (1997, 2002).

Meniscus midges (Dixidae) belong to infraorder Culicomorpha. There are nearly 200 known species of which 53 occur in the Palaearctic region. In Finland two genera and 16 species are present (Hackman 1980, Salmela 2003, Salmela et al. 2007). One species (*Dixella dyari* Garret, 1924) is reported here for the first time from Finland (see Notes). Larvae of meniscus midges dwell in or slightly above water meniscus, having a characteristic U-shaped position (Wagner 1997a). Members of the genus *Dixella* are associated with lentic or slow flowing waters, such as ponds, lake shores and bog pools. Species of the genus *Dixa* are lotic, dwelling in running waters of varying size. Two Finnish species, *Dixa dilatata* and *D. submaculata* prefer springs and spring brooks. The Finnish fauna is generally rather well known, although three species have hitherto

Table 1. Number of species by family.

Family	Number of species in			Level of knowledge
	World (Pape et al. 2011)	Europe	Finland	
Psychodomorpha:				
Psychodidae	2958	493	61–63	average
Culicomorpha:				
Chaoboridae	54	9	8	good
Dixidae	186	32	16	average
Thaumaleidae	182	76	1	good
Ptychopteromorpha:				
Ptychopteridae	69 (Wagner et al. 2008)	14	7	good

been collected from single localities. One northern *Dixa* species close to *D. dilatata* is still undescribed. A majority of the species can be identified by using Martini (1928), Peus (1934, 1936), Vaillant (1967) and Disney (1999).

Trickle midges (Thaumaleidae) is a small family belonging to infraorder Culicomorpha. Globally, nearly 200 species are known of which 76 are European. Central European mountains harbor a rich assemblage of trickle midges (Wagner 2002), at least if compared to the poor diversity (5 spp.) in Fennoscandian area (Wagner 1997b, Andersen et al. 2013). Only one species, *Thaumalea truncata*, has been hitherto recorded from Finland (see the discussion in Andersen et al. 2013). Larvae of trickle midges dwell on hygropetric substrates or splash zones of running waters (Wagner 1997b). *Thaumalea truncata* occurs mainly in northern Finland, both boreal and subarctic ecoregions, and adult specimens have been caught around springs and headwater streams (Salmela 2003, Salmela 2008).

Phantom crane flies (Ptychopteridae) belong to infraorder Ptychopteromorpha, being a sister group of Psychodomorpha and Culicomorpha (Wiegmann et al. 2011). Ptychopteridae is a small family with over 69 species in the world and 14 in Europe. Seven species are known from Finland, all belonging to the genus *Ptychoptera*. Only one Finnish species, *Ptychoptera minuta*, is widespread, locally abundant and almost ubiquitous. Other species are much less frequent, occurring in peatlands, headwater streams, springs and swamps. Adult *Ptychoptera* superficially resemble Tipuloidea crane flies, being slender and having long legs. Larvae of *Ptychoptera* live in shallow aquatic habitats, usually in soft substrates (Andersson 1997). Finnish species can be identified by using Andersson (1997) and Krzeminski (1986).

Checklist

Nematocera Dumeril, 1805

infraorder Psychodomorpha Hennig, 1968

PSYCHODIDAE Newman, 1834

PSYCHODINAE Newman, 1834

tribe Pericomaini Enderlein, 1935

PARABAZARELLA Vaillant, 1983

Parabazarella subneglecta (Tonnoir, 1922)

BERDENIELLA Vaillant, 1976

Berdeniella freyi (Berdén, 1954)

CLYTOCERUS Eaton, 1904

sg. **Boreoclytocerus** Duckhouse, 1978

Clytocerus ocellaris (Meigen, 1818)

Clytocerus rivosus (Tonnoir, 1919)

Clytocerus tetracorniculatus Wagner, 1977

PERICOMA Walker, 1856

sg. **Pachypericoma** Vaillant, 1978

Pericoma blandula Eaton, 1893

Pericoma nielseni Kvifte, 2010

= *formosa* Nielsen, 1964 (preocc. Meunier, 1905)

sg. **Pericoma** Walker, 1856

Pericoma rivularis Berdén, 1954

PNEUMIA Enderlein, 1935

Pneumia borealis (Berdén, 1954)

Pneumia bucegica (Vaillant, 1981)

Pneumia mutua (Eaton, 1893)

? *Pneumia nubila* (Meigen, 1818)

? *Pneumia palustris* (Meigen, 1818)

Pneumia pilularia (Tonnoir, 1940)

Pneumia stammeri (Jung, 1954)

Pneumia trivialis (Eaton, 1893)

Pneumia ussurica (Wagner, 1994)

ULOMYIA Walker, 1856

Ulomyia cognata (Eaton, 1893)

Ulomyia fuliginosa (Meigen, 1804)

tribe Psychodini Newman, 1834

PHILOSEPEDON Eaton, 1904

Philosepedon balkanicus Krek, 1970

Philosepedon humeralis (Meigen, 1818)

Philosepedon soljani Krek, 1971

PSYCHODA Latreille, 1796

Psychoda albipennis Zetterstedt, 1850

= *parthenogenetica* Tonnoir, 1940

Psychoda alternata Say, 1824

Psychoda brevicornis Tonnoir, 1940

Psychoda buxtoni Withers, 1988

Psychoda cinerea Banks, 1894

- Psychoda crassipennis* Tonnoir, 1940
Psychoda cultella Salmela, Kvifte & More, 2012
Psychoda erminea Eaton, 1898
Psychoda gemina (Eaton, 1904)
Psychoda grisescens Tonnoir, 1922
Psychoda itoco Tokunaga & Komyo, 1955
Psychoda lativentris Berdén, 1952
Psychoda lobata Tonnoir, 1940
Psychoda minuta Banks, 1894
Psychoda phalaenoides Linnaeus, 1758
Psychoda satchelli Quate, 1955
Psychoda setigera Tonnoir, 1922
Psychoda trinodulosa Tonnoir, 1922
Psychoda uniformata Haseman, 1907
THRETICUS Eaton, 1904
Threticus tridactilis (Kincaid, 1899)
TRICHOPSYCHODA Tonnoir, 1922
Trichopsychoda hirtella (Tonnoir, 1919)
tribe Maruinini Enderlein, 1937
LOBULOSA Szabo, 1960
Lobulosa pollex (Berdén, 1954)
TONNOIRIELLA Vaillant, 1971
Tonnoiriella nigricauda (Tonnoir, 1919)
tribe Mormiini Enderlein, 1937
MORMIA Enderlein, 1935
Mormia niesiolowskii Wagner, 1985
Mormia strobli (Ježek, 1986)
tribe Paramormiini Enderlein, 1937
PARAJUNGIELLA Vaillant, 1972
Parajungiella consors (Eaton, 1893)
Parajungiella ellisi (Withers, 1987)
Parajungiella pseudolongicornis Wagner, 1975
LEPISEODINA Enderlein, 1937
Lepiseodina rothschildi (Eaton, 1912)
PANIMERUS Eaton, 1913
sg. Panimerus Eaton, 1913
Panimerus albomaculatus (Wahlgren, 1904)
Panimerus notabilis (Eaton, 1893)
sg. Psycmera Ježek, 1984
Panimerus albifacies (Tonnoir, 1919)
Panimerus intellegus (Jung, 1956)
Panimerus przhiboroi Wagner, 2005

PARAMORMIA Enderlein, 1935*Paramormia polyscoidea* (Krek, 1970)*Paramormia ustulata* (Walker, 1856)**PERIPSYCHODA** Enderlein, 1935*Peripsychoda auriculata* (Curtis, 1839)*Peripsychoda fusca* (Macquart, 1826)**TELMATOSCOPIUS** Eaton, 1904*Telmatoscopus advena* (Eaton, 1893)*Telmatoscopus similis* Tonnoir, 1922

SYCORACINAE Rondani, 1856

SYCORAX Haliday in Curtis, 1839*Sycorax silacea* Haliday in Curtis, 1839

infraorder Culicomorpha Hennig, 1948

superfamily Culicoidea Meigen, 1818

CHAOBORIDAE Newman, 1834**CHAOBORUS** Lichtenstein, 1800sg. *Chaoborus* Lichtenstein, 1800*Chaoborus crystallinus* (De Geer, 1776)*Chaoborus flavicans* (Meigen, 1830)*Chaoborus obscuripes* (van der Wulp, 1859)sg. *Peusomyia* Sæther, 1970*Chaoborus pallidus* (Fabricius, 1794)sg. *Schadanophasma* Dyar & Shannon, 1924*Chaoborus nyblaei* (Zetterstedt, 1838)**CRYOPHILA** Edwards, 1930*Cryophila lapponica* (Martini, 1928)**MOCHLONYX** Loew, 1844*Mochlonyx fuliginosus* Felt, 1905= *M. martinii* Edwards, 1930= *M. velutinus* of authors*Mochlonyx velutinus* (Ruthe, 1831)= *M. culiciformis* (De Geer, 1776) preocc.**DIXIDAE** Schiner, 1868**DIXA** Meigen, 1818*Dixa dilatata* Strobl, 1900*Dixa nebulosa* Meigen, 1830*Dixa puberula* Loew, 1849*Dixa submaculata* Edwards, 1920**DIXELLA** Dyar & Shannon, 1924*Dixella aestivalis* (Meigen, 1818)*Dixella amphibia* (De Geer, 1776)

Dixella autumnalis (Meigen, 1838)
Dixella borealis (Martini, 1928)
Dixella dyari (Garret, 1924)
Dixella filicornis (Edwards, 1926)
Dixella hyperborea (Bergroth, 1889)
Dixella laeta (Loew, 1849)
Dixella naevia (Peus, 1934)
Dixella nigra (Staeger, 1840)
Dixella obscura (Loew, 1849)
Dixella serotina (Meigen, 1818)

THAUMALEIDAE Bezzi, 1913

THAUMALEA Ruthe, 1831

Thaumalea truncata Edwards, 1929

infraorder Ptychopteromorpha Wood & Borkent, 1986
 superfamily Ptychopteroidea Osten Sacken, 1862

PTYCHOPTERIDAE Osten Sacken, 1862

PTYCHOPTERA Meigen, 1803

sg. Paraptychoptera Tonnoir, 1919

Ptychoptera lacustris Meigen, 1830

Ptychoptera paludosa Meigen, 1804

sg. Ptychoptera Meigen, 1803

Ptychoptera albimana (Fabricius, 1787)

Ptychoptera contaminata (Linnaeus, 1758)

Ptychoptera hugoi Tjeder, 1968

Ptychoptera minuta Tonnoir, 1919

Ptychoptera scutellaris Meigen, 1818

Excluded species

Dixa maculata Meigen, 1818. There is one female specimen collected from Russia, Kantalaks (=Kandalaksha) that belongs to *Dixella borealis*.

Notes

Threticus tridactylis: Ostrobothnia borealis pars borealis, Rovaniemi, Nammalikkokivalo, 13.6.–11.7.2004, J. Salmela leg.; Regio kuusamoensis, Taivalkoski, Poikaloukusanoja, 1.6.–1.7.2006, J. Salmela leg.; Regio kuusamoensis, Taivalkoski, Syväoja 1.6.-1.7.2006, J. Salmela leg.

- Panimerus albifacies***: Regio aboensis, Turku, Pomponrahka-Isosuo mire conservation area, 1.8.–28.9.2011, J. Salmela & E. Nummela leg.
- Panimerus przhiboroi***: Ostrobothnia borealis pars borealis, Tervola, Ruuttulampi conservation area, 4.6.–2.7.2012 J. Salmela leg. (New for European fauna)
- Psychoda albipennis*** Zetterstedt has been confirmed to occur in Finland by Bo-W. Svensson (female specimen from Tavastia australis, South Finland, Tampere, 2009, J. Salmela leg.).
- Dixella dyari***: Lapponia enontekiensis, Enontekiö, Pallas-Yllästunturi National Park, Lumikero E, 25.8.2010, J. Salmela leg.

References

- Andersen T, Hagenlund LK, Håland Ø, Kvifte GM, Skartveit J (2013) New records of lower Diptera (“Nematocera”) from Finnmark. Norwegian Journal of Entomology 60: 182–189.
- Andersson H (1997) Diptera, Ptychopteridae, Phantom Crane Flies. In: Nilsson A (Ed) The Aquatic Insects of North Europe 2: 193–207.
- Autio O, Salmela J (2010) The semi-aquatic fly fauna (Diptera) of wetlands of the Åland Islands. Memoranda Societatis pro Fauna et Flora Fennica 86: 43–53.
- Disney RHL (1999) British Dixidae (Meniscus midges) and Thaumaleidae (Trickle midges): Keys with ecological notes. Freshwater Biological Association, Scientific Publication No. 56.
- Duckhouse DA (1987) A revision of Afrotropical *Setomima*, elucidation of their genealogical relationships and descriptions of other Afrotropical Psychodinae (Diptera: Psychodidae). Annals of the Natal Museum 28: 231–281.
- Hirvenoja M (1960) Ökologische Studien über die Wasserinsekten in Riihimäki (Südfinnland). I. Chaoborinae (Dipt., Culicidae). Annales entomologici fennici 26: 31–44.
- Hirvenoja M (1963) Ein neuer Fund von *Chaoborus (Sayomyia) pallidus* Fabr. (Dipt., Culicidae) aus Finland. Annales entomologici fennici 29: 283.
- Ježek J (1983) Contribution to the taxonomy of the genus *Logima* Eat. (Diptera, Psychodidae). Acta entomologica musei nationalis pragrae 41: 213–259.
- Ježek J, van Harten A (2005) Further new taxa and little-known species of non-biting moth flies (Diptera, Psychodidae) from Yemen. Acta entomologica musei nationalis pragrae 45: 199–220.
- Krzeminski W (1986) Ptychopteridae of Poland (Diptera, Nematocera). Polskie Pismo Entomologiczne 56: 105–131.
- Kvifte GM, Håland Ø, Andersen T (2011) A revised checklist of Norwegian moth flies (Diptera, Psychodidae). Norwegian Journal of Entomology 58: 180–188.
- Martini E (1928) Dixinae und Corethrinae des Museums zu Helsingfors. Notulae entomologicae 8: 33–35.
- Paasivirta L (2002) Surviaissääsket, sulkasääsket ja hyttyset. In: Salokannel J (Ed) Iidesjärven hyönteisselvitys, Diamina 11: 25–27.
- Pape T, Blagoderov V, Mostovski MB (2011) Order Diptera Linnaeus, 1758. In: Zhang Z-Q (Ed) Animal biodiversity: An outline of higher-level classification and survey of

- taxonomic richness. *Zootaxa* 3148: 222–229. <http://www.mapress.com/zootaxa/2011/flzt03148p229.pdf>
- Peus F (1934) Dixiden und Culiciden aus Lettland. *Notulae entomologicae* 14: 69–78.
- Peus F (1936) Zur Kenntnis der Dixiden-Fauna Nordeuropas (Dipt., Nematoc.). *Norsk Entomologisk Tidsskrift* 4: 117–127.
- Saether OA (1970) Nearctic and Palaearctic Chaoborus (Diptera: Chaoboridae). *Bulletin of the Fisheries Research Board of Canada* 174: 1–57.
- Saether OA (1997) Diptera, Chaoboridae, Phantom midges. In: Nilsson A (Ed) *The Aquatic Insects of North Europe* 2: 149–161.
- Saether OA (2002) Insecta: Diptera: Chaoboridae. *Süßwasserfauna von Mitteleuropa* 21: 3–38.
- Salmela J (2003) New records of the families Psychodidae, Dixidae and Thaumaleidae (Diptera, Nematocera) from Finland. *Studia dipterologica* 10: 143–152.
- Salmela J (2005) New moth flies for eastern Fennoscandia (Diptera: Psychodidae). *Sahlbergia* 10: 1–3.
- Salmela J (2008) Semiaquatic fly (Diptera, Nematocera) fauna of fens, springs, headwater streams and alpine wetlands in the northern boreal ecoregion, Finland. *w-album* 6: 3–63. <http://org.utu.fi/harrastus/TEKS/w-album/608.pdf>
- Salmela J, Autio O, Ilmonen J (2007) A survey on the nematoceran (Diptera) communities of southern Finnish wetlands. *Memoranda Societatis Pro Fauna et Flora Fennica* 83: 33–47.
- Salmela J, Kvifte GM, More A (2012) Description of a new *Psychoda* Latreille species from Fennoscandia (Diptera: Psychodidae). *Zootaxa* 3313: 34–43.
- Vaillant F (1967) Two Dixid flies from near the arctic circle in Sweden (Dipt.). *Opuscula entomologica* 32: 284–288.
- Vaillant F (1971–1983) Psychodidae–Psychodinae. In: Lindner E (Ed) *Die Fliegen Der Palearktischen Region*. Lieferungen 287, 291, 292, 305, 310, 313, 317, 320, 326, 328. E. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart, 258 pp.
- Vaillant F (1990) Propositions pour une révision de la classification des Diptères Psychodidae Psychodinae. *Bulletin de la Société vaudoise des sciences naturelles* 80: 141–163.
- Wagner R (1997a) Diptera Dixidae, Meniscus midges. In: Nilsson A (Ed) *Aquatic Insects of North Europe*. Volume 2. Odonata–Diptera. Apollo Books, Strenstrup, 145–148.
- Wagner R (1997b) Diptera Thaumaleidae. In: Nilsson A (Ed) *Aquatic Insects of North Europe*. Volume 2. Odonata–Diptera. Apollo Books, Strenstrup, 187–191.
- Wagner R (1997c) Diptera Psychodidae, Moth flies. In: Nilsson A (Ed) *Aquatic Insects of North Europe*. Volume 2. Odonata–Diptera. Apollo Books, Strenstrup, 133–144.
- Wagner R (2002) Insecta: Diptera: Thaumaleidae. *Süßwasserfauna von Mitteleuropa* 21: 41–110.
- Wagner R (2013) Fauna Europaea: Psychodidae. In: Beuk P, Pape T (2013) *Fauna Europaea: Diptera, Psychodidae*. Fauna Europaea version 1.0, <http://www.faunaeur.org>
- Wagner R, Barták M, Borkent A, Courtney G, Goddeeris B, Haenni J-P, Knutson L, Pont A, Rotheray GE, Rozkošný R, Sinclair B, Woodley N, Zatwarnicki T, Zwick P (2007) Global diversity of dipteran families (Insecta Diptera) in freshwater (excluding Simuliidae, Culicidae, Chironomidae, Tipulidae and Tabanidae). *Hydrobiologia* 595(1): 489–519. doi: 10.1007/s10750-007-9127-9

- Wiegmann BM, Trautwein MD, Winkler IS, Barr NB, Kim J-W, Lambkin C, Bertone MA, Cassel BK, Bayless KM, Heimberg AM, Wheeler BM, Peterson KJ, Pape T, Sinclair BJ, Skevington JH, Blagoderov V, Caravas J, Kutty SN, Schmidt-Otto U, Kampmeier GE, Thompson FC, Grimaldi DA, Beckenbach AT, Courtney GW, Friedrich M, Meier R, Yeats DK (2011) Episodic radiations in the fly tree of life. *Proceedings of the National Academy of Sciences of the United States of America* 108: 5690–5695. doi: 10.1073/pnas.1012675108
- Withers P (1989) Moth flies. *Diptera: Psychodidae. Dipterists Digest* 4: 1–83.