

First report on C-banding, fluorochrome staining and NOR location in holocentric chromosomes of *Elasmolomus (Aphanus) sordidus* Fabricius, 1787 (Heteroptera, Rhyparochromidae)

Vikas Suman¹, Harbhajan Kaur²

1 Department of Entomology, Y.S. Parmar University of Horticulture and Forestry, Nauni, Solan 173 230, Himachal Pradesh, India **2** Department of Zoology and Environmental Sciences, Punjabi University, Patiala 147 002, Punjab, India

Corresponding author: Vikas Suman (viks_suman@yahoo.co.in)

Academic editor: S. Grozeva | Received 7 November 2012 | Accepted 9 February 2013 | Published 30 July 2013

Citation: Suman V, Kaur H (2013) First report on C-banding, fluorochrome staining and NOR location in holocentric chromosomes of *Elasmolomus (Aphanus) sordidus* Fabricius, 1787 (Heteroptera, Rhyparochromidae). In: Popov A, Grozeva S, Simov N, Tasheva E (Eds) Advances in Hemipterology. ZooKeys 319: 283–291. doi: 10.3897/zookeys.319.4265

Abstract

In spite of various cytogenetic works on suborder Heteroptera, the chromosome organization, function and its evolution in this group is far from being fully understood. Cytologically, the family Rhyparochromidae constitutes a heterogeneous group differing in chromosome numbers. This family possesses XY sex mechanism in the majority of the species with few exceptions. In the present work, multiple banding techniques viz., C-banding, base-specific fluorochromes (DAPI/CMA₃) and silver nitrate staining have been used to cytologically characterize the chromosomes of the seed plant pest *Elasmolomus (Aphanus) sordidus* Fabricius, 1787 having $2n=12=8A+2m+XY$. One pair of the autosomes was large while three others were of almost equal size. At diplotene, C-banding technique revealed, that three autosomal bivalents show terminal constitutive heterochromatic bands while one medium sized bivalent was euchromatic. Microchromosomes (m-chromosomes) were positively heteropycnotic. After DAPI and CMA₃ staining, all the autosomal bivalents showed equal fluorescence, except CMA₃ positive signals, observed at both telomeric heterochromatic regions of one medium sized autosomal bivalent. Silver nitrate staining further revealed that this chromosome pair carries Nucleolar Organizer Regions (NORs) at the location of CMA₃ positive signals. The X chromosome showed a thick C-band, positive to both DAPI /CMA₃ while Y, otherwise C-negative, was weakly positive to DAPI and negative to CMA₃, m-chromosomes were DAPI bright and CMA₃ dull.

Keywords

C-banding, DAPI, CMA₃, NOR location

Introduction

Heteroptera is a large cosmopolitan suborder comprising about 42,300 known species (Henry 2009). The species of Heteroptera are distributed into 7 infraorders and a total of 24 superfamilies worldwide (Schuh and Slater 1995). Lygaeidae, Rhyparochromidae Pyrrhocoridae, Coreidae, Pentatomidae, Reduviidae and Miridae are some of the major families, each having its individual economic importance (Schaefer and Panizzi 2000). Rhyparochromidae (seed bugs) were considered by most workers to be a subfamily within the Lygaeidae until revision by Henry (1997) who recognized them at the family level. Rhyparochromids are mostly ground dwellers, living in the shadow vegetation and feeding primarily on seeds. *Elasmolomus (Aphanus) sordidus* is a serious pest, occurring on pods left drying in the fields and in stores. Groundnuts and sesame pods infested by this insect have shrivelled kernels. Like other heteropterans, Rhyparochromidae are characterized by holokinetic chromosomes and post reductional division of sex chromosomes, as well by presence of m-chromosomes and XY sex mechanism in all the species with few exceptions (Ueshima 1979).

In the present contribution, cytological characterization of *E. (Aphanus) sordidus*, reported as *Aphanus sordidus* having chromosomal complement $2n=12=8A+2m+XY$ (Parshad 1957), has been done using different banding techniques. The amount and location (C-banding) and composition (AT/GC base richness) of heterochromatin have been studied. Further silver banding was employed to locate the position and number of nucleolar organiser regions (NORs). The application of CMA₃/DAPI banding revealed correspondence between NORs (r-DNA sites) and GC rich domains.

Material and methods

Adult males of *E. sordidus* (9 specimens) were collected from fields of sesame and groundnut plants in Punjab (India). Insects were dissected remove the gonads and air dried slides were prepared. Aged air dried slides were used for C-banding after Kaur et al. (2010). To study the localization of NORs, silver staining was done using one step method with a protective colloidal developer (gelatine and formic acid) (Howell and Black 1980). To reveal the base composition of C-heterochromatin, two fluorochromes: AT sequence specific DAPI (4-6' Diamidino-2-phenylindole) and GC sequence specific CMA₃ (chromomycin A₃), were applied, following the protocol suggested by Manicardi and Gautam (1994). Well-spread stages were photographed under the microscope Nikon-Optiphot-2. Slides stained with fluorochrome dyes DAPI/CMA₃ were studied and photographed under Nikon fluorescent microscope using UV filter (for DAPI) and BV (for CMA₃).

Results

The chromosomal complement consisted of twelve elements. Of these, eight were autosomes, two were m-chromosomes, while two of different sizes were sex chromosomes, large X and small Y respectively. The chromosomal complement was confirmed as $2n=12=8A+2m+XY$.

C-banding

At diplotene, three bivalents showed terminal C-bands while one was euchromatic. The X chromosome showed thick C-band covering almost two thirds of the chromosome while Y was C-negative; m-chromosomes were slightly C-positive (Figs 1, 2).

DAPI/CMA₃ staining

All the four autosomal bivalents showed equal fluorescence with both DAPI and CMA₃ (Figs 3, 4). However, one of the medium sized autosomal bivalents showed bright CMA₃ signals at both ends, which correspond to NORs (Fig. 4). The X was positive to both DAPI/CMA₃ while Y was weak to DAPI and negative to CMA₃; m-chromosomes were DAPI bright and CMA₃ dull (Figs 5, 6).

Silver staining

NORs were found to be associated with both ends of a medium-sized autosomal bivalent (Figs 7, 8).

Discussion

E. sordidus is a pest of pod crops, mainly groundnut and sesame in India. Parshad (1957) was first to study its standard chromosomal complement ($2n=12=8A+2m+XY$) and male meiosis of this species (as *Aphanus sordidus*). The same chromosomal complement has been observed by the present authors. In the present study, C-banding, silver staining and DNA sequence-specific staining have been used to reveal the distribution and constitution of constitutive heterochromatin and also to find the correspondence between NORs and GC-rich regions.




Figure 1–8. C-banding (1, 2) 1, 2 Diplotene stages showing distribution of C-bands. Arrows showing heterochromatic chromosomes while arrowhead showing single euchromatic chromosome. **Sequence-specific banding (3–6)** 3 Diplotene stage with DAPI 4 Diplotene stage with localized CMA₃ signals on one autosomal bivalent (shown by arrows) 5 Late diplotene stage with DAPI 6 Late diplotene stage with CMA₃. **Silver banding (7, 8)** 7, 8 Diplotene stages showing location of NORs (shown by arrows) and nucleolar bodies (N). Bar=0.01 mm.

C-banding

Terminal C-bands have been observed in three autosomal pairs of *E. sordidus*. In Heteroptera, the terminal C-bands are of wide occurrence. This kind of C-band location has been reported in *Antiteuchus mixtus* (Fabricius, 1787) (Pentatomidae) by Lanzone and Souza (2006), in *Dieuches uniguttatus* (Thumb, 1822) and *D. insignis* (Distant, 1918) (Rhyparochromidae) by Kaur et al. (2010). Usually, telomeric bands are absent, if interstitial blocks are present in a chromosome. This is reported in one or two chromosomes of *Nezara viridula* Linnaeus, 1758 (Pentatomidae) and *Triatoma patagonica* Del-Ponte, 1929 (Reduviidae) by Camacho et al. (1985) and Panzera et al. (1997) respectively.

One of the autosomal bivalent in *E. sordidus* was found to be euchromatic. A similar condition is observed in *Nezara icterica* (Horvath, 1889) (Pentatomidae) by Dey and Wangdi (1990), in *Dieuches coloratus* Distant, 1909 (Rhyparochromidae) by Kaur et al. (2010) and in *Neophysopelta schlanbuschi* Ahmad & Abbas, 1987 (Largidae) by Suman et al. (2012).

The X chromosome is almost (2/3) completely C-positive and this condition has been earlier reported in Pentatomidae by Camacho et al. (1985), in Tingidae by Grozeva and Nokkala (2001) and in Nabidae by Grozeva et al. (2004), whereas, the Y chromosome, is C-negative. This condition is not uncommon in Heteroptera and has been reported previously in some species belonging to Coreidae, Pentatomidae and Tingidae (Muramoto 1980, Camacho et al. 1985, Dey and Wangdi 1990, Grozeva and Nokkala 2001).

Microchromosomes were originally described by Wilson (1905); since then they have been discovered in many heteropteran families, including Rhyparochromidae. Microchromosomes are C-positive in *E. sordidus*. Similar observation have been made in *Leptoglossus impictus* (Stål, 1860) and *Phthia picta* (Drury, 1773) (Coreidae) by Bressa et al. (2005) and in *Dieuches uniguttatus* and *D. insignis* (Rhyparochromidae) by Kaur et al. (2010). Microchromosomes are DAPI bright and CMA₃ dull. Similar set of observations have been previously made by Kaur et al. (2010) in *Dieuches uniguttatus* and *D. insignis* (Rhyparochromidae). Information on chromatin composition of m-chromosomes is still very poor and their genetic constitution is not fully known.

DAPI/CMA₃ staining

The use of DNA binding fluorochromes having different base specificities allows a better characterization of heterochromatic regions in terms of their relative enrichment with AT or GC base pairs. In Heteroptera, still there is little information on heterochromatin base composition. The bright fluorescence after DAPI and CMA₃ staining observed in *E. sordidus* indicates that the constitutive heterochromatic regions possess interspersed AT and GC repeats. Similar observations have been made in *Edessa mediatibunda* (Fabricius, 1974) and *E. rufomarginata* (De Geer, 1773) (Pentatomidae) by

Rebagliati et al. (2003), in *Antiteuchus mixtus*, *A. macraspis* (Perty, 1834), *A. sepulcralis* (Fabricius, 1803) (Pentatomidae) by Lanzone and Souza (2006) and in *Arachnocoris trinitatus* Bergroth, 1916 (Nabidae) by Kuznetsova et al. (2007).

After silver banding and fluorochrome staining, the localization of CMA₃ positive bands in NOR regions on medium sized autosomal bivalent was revealed. It was confirmed that ribosomal genes are GC rich. This correspondence of CMA₃ signals with NORs have also been reported for several true bug species at interstitial or terminal positions either on autosomes or sex chromosomes by Gonzalez-Garcia et al. (1996), Papeschi et al. (2003), Rebagliati et al. (2003), and Grozeva et al. (2004). However, NORs do not always show GC base richness as is reported in *Carlisis wahlbergi* Stål, 1858 (Coreidae) by Fossey and Liebenberg (1995).

A common feature of the sex chromosomes of Heteroptera is that they demonstrate bright fluorescence after both DAPI and CMA₃ during the meiotic prophase (Rebagliati et al. 2003). In the present study, the X chromosome showed fluorescence after both DAPI and CMA₃. Similar observations have been also made in *Cimex emarginatus* Simov, Ivanova & Schunger, 2006 by Grozeva and Nokkala (2002), *Cimex lectularius* (Cimicidae) by Grozeva et al. (2010), in *Edessa mediatubunda* and *E. rufomarginata* (Pentatomidae) by Rebagliati et al. (2003), in *Athaumastus haematicus* (Stål, 1860), *Leptoglossus impictus* and *Phibia picta* (Coreidae), *Jadera sanguinolenta* (Fabricius, 1775) (Rhopalidae) by Bressa et al. (2005), in *Antiteuchus mixtus*, *A. macraspis* and *A. sepulcralis* (Pentatomidae) by Lanzone and Souza (2006). In the present study, however, the Y chromosome is C-negative, but DAPI positive and CMA₃ negative. Similar observations have been made in *Triatoma vitticeps* (Stål, 1859) (Reduviidae) by Severi-Aguiar et al. (2006).

Silver staining

The silver impregnation stains not only the NORs but also the nucleolus at specific points of some chromosomes (Castanhole et al. 2008). In the present study, NORs were found to present on terminal regions of one of medium sized autosomal pairs of *E. sordidus*, like in *Nysius californicus* Stål, 1859 (Lygaeidae) (Souza et al. 2007), and in *Arachnocoris trinitatus* Bergroth, 1916 (Nabidae) (Kuznetsova et al. 2007). However, in Belostomatidae, NORs have been reported on either autosomes, on sex chromosomes or on both autosomes and sex chromosomes (Papeschi and Bidau 1985).

Conclusion

Till date, very few Rhyparochromid species have been analysed cytologically based on banding techniques. The present study was able to reveal some cytogenetic characters which were used as markers for better knowledge of chromosome organization and the identification of separate chromosomes in *E. sordidus*. Much more information about true

bug chromosomes could be obtained if new molecular cytogenetic techniques involving FISH (fluorescence *in situ* hybridization) mapping of chromosomes are used (Grozeva et al. 2011 and references therein, Kuznetsova et al. 2012 and references therein).

Acknowledgements

The authors thank the Department of Science and Technology, New Delhi, India for providing financial support (Grant Number 96915) to carry out the present study.

References

- Bressa MJ, Larramendy ML, Papeschi AG (2005) Heterochromatin characterization in five species of Heteroptera. *Genetica* 124: 307–317. doi: 10.1007/s10709-005-4524-3
- Camacho JPM, Belda J, Cabrero J (1985) Meiotic behaviour of holocentric chromosomes of *Nezara viridula* (Insecta: Heteroptera) analysed by C-banding and Silver impregnation. *Canadian Journal of Genetics and Cytology* 27: 490–497.
- Castanhole MMU, Pereira LLV, Souza HV, Bicudo HEMC, Costa LAA, Itoyama MM (2008) Heteropicnotic chromatin and nucleolar activity in meiosis and spermiogenesis of *Limnogonus aduncus* (Heteroptera, Gerridae): a stained nucleolar organizing region that can serve as a model for studying chromosome behaviour. *Genetics and Molecular Research* 7(4): 1398–1407. doi: 10.4238/vol7-4gmr527
- Dey SK, Wangdi T (1990) Banding patterns of the holocentric chromosomes in some species of Heteroptera. *Cytologia* 55: 181–186. doi: 10.1508/cytologia.55.181
- Fossey A, Liebenberg H (1995) Meiosis and nucleolar structures in the stink bug *Carlisia wahlbergi* Stål (Coreidae: Heteroptera). *Cytobios* 81: 7–15.
- Gonzalez-Garcia JM, Antonio C, Siya JA, Rufas JS (1996) Meiosis in holocentric chromosomes: kinetic activity is randomly restricted to the chromatid ends of sex univalents in *Graphosoma italicum* (Heteroptera). *Chromosome Research* 4: 124–132. doi: 10.1007/BF02259705
- Grozeva SM, Nokkala S (2001) Chromosome numbers, sex determining systems, and patterns of the C-heterochromatin distribution in 13 species of lace bugs (Heteroptera, Tingidae). *Folia Biologica (Kraków)* 49: 29–41.
- Grozeva SM, Nokkala S (2002) Achiasmatic male meiosis in *Cimex* sp. (Heteroptera, Cimicidae). *Caryologia* 55: 189–192. doi: 10.1080/00087114.2002.10589276
- Grozeva S, Kuznetsova VG, Nokkala S (2004) Patterns of chromosome banding in four nabid species (Heteroptera, Cimicomorpha, Nabidae) with high chromosome number karyotypes. *Hereditas* 140(2): 99–104. doi: 10.1111/j.1601-5223.2004.01782.x
- Grozeva S, Kuznetsova V, Anokhin B (2010) Bed bug cytogenetics: karyotype, sex chromosome system, FISH mapping of 18S rDNA, and male meiosis in *Cimex lectularius* Linnaeus, 1758 (Heteroptera: Cimicidae). *Comparative Cytogenetics* 4: 151–160. doi: 10.3897/compcytogen.v4i2.36

- Grozeva S, Kuznetsova VG, Anokhin BA (2011) Karyotypes, male meiosis and comparative FISH mapping of 18S ribosomal DNA and telomeric (TTAGG)_n repeat in eight species of true bugs (Hemiptera, Heteroptera). *Comparative Cytogenetics* 5(4): 355–374. doi: 10.3897/CompCytogen.v5i4.2307
- Henry TJ (1997) Phylogenetic analysis of family groups within the infraorder Pentatomorpha (Hemiptera:Heteroptera) with emphasis on Lygaeoidea. *Annals of Entomology Society America* 90: 275–301.
- Henry TJ (2009) Biodiversity of Heteroptera. In: Footitt R, Alder P (Eds) *Insect Biodiversity Science and Society*. Chichester, Blackwell Publishers, Chichester, 223–263. doi: 10.1002/9781444308211.ch10
- Howell M, Black DA (1980) Controlled silver staining of nucleolus organizer regions with protective colloidal developer: 1-step method. *Experientia* 36: 104–105. doi: 10.1007/BF01953855
- Kaur H, Suman V, Kaur R (2010) A first report on C- banding and Fluorescent banding in species of *Dieuches* (Rhyparochrominae: Lygaeidae: Heteroptera). *Entomological Research* 40(1): 1–7. doi: 10.1111/j.1748-5967.2009.00255.x
- Kuznetsova VG, Grozeva S, Sewlal JN, Nokkala S (2007) Cytogenetic characterization of the endemic of Trinidad, *Arachnocoris trinitatus* Bergroth: the first data for the tribe Arachnocorini (Heteroptera: Cimicomorpha: Nabidae). *Folia Biologica (Kraków)* 55(1–2): 17–26. doi: 10.3409/173491607780006344
- Kuznetsova VG, Grozeva SM, Anokhin BA (2012) The first finding of (TTAGG)_n telomeric repeat in chromosomes of true bugs (Heteroptera, Belostomatidae). *Comparative Cytogenetics* 6(4): 341–346. doi: 10.3897/CompCytogen.v6i4.4058
- Lanzone C, Souza M (2006) C-banding, fluorescent staining and NOR location in holokinetic chromosomes of bugs of Neotropical genus *Antiteuchus* (Heteroptera: Pentatomidae: Discocephalinae). *European Journal of Entomology* 103: 239–243.
- Manicardi GC, Gautam DC (1994) Cytogenetic investigations on the holokinetic chromosomes of *Tetraneurella akinirei* (Sasaki) (Homoptera, Pemphigidae). *Caryologia* 47: 159–165.
- Muramoto N (1980) A study of the C-banded chromosomes in some species of Heteropteran insects. *Proceedings of Japan Academy* 56 (B): 125–130.
- Panzer F, Horhos S, Pereira J, Cestau R, Canale D, Diotaiuti L, Dujardin JP, Perez R (1997) Genetic variability and geographic differentiation among three species of Triatominae bugs (Hemiptera, Reduviidae). *American Journal of Tropical Medicine and Hygiene* 57: 732–739.
- Papeschi AG, Bidau CJ (1985) Chromosome complement and male meiosis in four species of *Belostoma latreille* (Belostomatidae, Heteroptera). *Brazilian Journal of Genetics* 8: 249–261.
- Papeschi AG, Mola LM, Bressa MJ, Greizerstein EJV, Poggio L (2003) Behaviour of ring bivalents in holocentric systems: alternate sites of spindle attachment in *Pachylis argentinus* and *Nezara viridula* (Heteroptera). *Chromosome Research* 11: 725–733. doi: 10.1023/B:CHRO.0000005740.56221.03
- Parshad R (1957) Chromosome number and sex mechanism in twenty species of the Indian Heteroptera. *Current Science* 26: 125.

- Rebagliati PJ, Papeschi AG, Mola LM (2003) Meiosis and fluorescent banding in *Edessa med-
itabunda* and *E. rufomarginata* (Heteroptera: Pentatomidae: Edessinae). *European Journal
of Entomology* 100: 11–18.
- Schaefer CW, Panizzi AR (2000) *Heteroptera of Economic Importance*. CRC press Boca Ra-
ton, London, New York, Washington DC, 828 pp.
- Schuh RT, Slater JA (1995) *True Bugs of the World (Hemiptera: Heteroptera)*, Classification
and Natural History. Ithaca: Cornell University Press XII, 336 pp.
- Severi-Aguiar GD, Lourenco LB, Bicudo HEMC, Azeredo-Oliveira MTV (2006) Meiosis as-
pects and Nucleolar activity in *Triatoma vitticeps* (Triatominae, Heteroptera). *Genetica*
126: 141–151. doi: 10.1007/s10709-005-1443-2
- Suman V, Kaur H, Singh D, Kaur R (2012) Species-specific sex chromosome behaviour and
banding patterns in three Largid species (Heteroptera). *Chromosome Science* 15 (1–2):
31–37.
- Souza H V, Bicudo HEMC, Itoyama MM (2007) Study of chromosomal and nucleolar aspects
in testes of *Nysius californicus* (Heteroptera-Lygaeidae). *Genetics and Molecular Research*
6 (1): 33–40.
- Ueshima N (1979) Hemiptera II: Heteroptera. In: John B (Eds) *Animal Cytogenetics*. Berlin-
Stuttgart, 113pp.
- Wilson EB (1905) Studies of chromosome. II: The paired microchromosomes, idiochromo-
somes and heterotropic chromosomes in Hemiptera. *Journal of Experimental Zoology*
2: 507–545. doi: 10.1002/jez.1400020405