

A new species of *Haplothrips* from southern Iran (Thysanoptera, Phlaeothripidae)

Kambiz Minaei^{1†}, Maryam Aleosfoor^{1‡}

¹ Department of Plant Protection, College of Agriculture, Shiraz University, Shiraz, Iran

† [urn:lsid:zoobank.org/author:11F6AD27-C98E-43C4-A10F-5E19156799C2](https://doi.org/urn:lsid:zoobank.org/author:11F6AD27-C98E-43C4-A10F-5E19156799C2)

‡ [urn:lsid:zoobank.org/author:84EDF5CC-EE8D-4803-B93C-D8D0677EE2E3](https://doi.org/urn:lsid:zoobank.org/author:84EDF5CC-EE8D-4803-B93C-D8D0677EE2E3)

Corresponding author: Kambiz Minaei (kminaei@shirazu.ac.ir)

Academic editor: L. Mound | Received 2 December 2012 | Accepted 3 February 2013 | Published 6 March 2013

[urn:lsid:zoobank.org/pub:4566E772-15EA-41FD-BB89-11C49F5D6AF7](https://doi.org/urn:lsid:zoobank.org/pub:4566E772-15EA-41FD-BB89-11C49F5D6AF7)

Citation: Minaei K, Aleosfoor M (2013) A new species of *Haplothrips* from southern Iran (Thysanoptera, Phlaeothripidae). ZooKeys 275: 91–99. doi: 10.3897/zookeys.275.4433

Abstract

Haplothrips herajius sp. n. is described from leaves and flowers of a species of *Suaeda* in the south of Fars Province, Iran. This is the second Iranian species of *Haplothrips* with the unusual character state of extra setae on the metanotum. Information on variation in color and structure of the new species is provided. The similarities and host plant associations of this new species and *H. kermanensis* are discussed, as both are phytophagous on species of Chenopodiaceae.

Keywords

Haplothrips, new species, phytophagous, *Suaeda* sp.

Introduction

Traditionally and still widely accepted, the known species of thrips are placed in a single order, the Thysanoptera, within which two suborders are recognized, the Terebrantia and Tubulifera (Mound et al. 1980, Mound and Morris 2007, Buckman et al. 2013). The suborder Tubulifera comprises a single family, the Phlaeothripidae with about 3500 described species (Mound 2013) classified into two subfamilies. In the more speciose of the subfamilies, the Phlaeothripinae, *Haplothrips* Amyot &

Serville, with 226 species worldwide, is the second largest genus, exceeded in number of described species only by *Liothrips* Uzel with 250 species (Mound 2013). Most species in this genus are Eurasian with just three described from South America (Mound and Zapater 2003, see also Goldarazena et al. 2012) and some world-wide in distribution (Pitkin 1976).

Among 20 genera of Phlaeothripidae recorded from Iran so far (Minaei 2013), the genus *Haplothrips* is considered to be the richest in this country (Minaei and Mound 2008). Apart from some species that are predators on other arthropods (Putman 1965, Bailey and Caon 1986, Palmer and Mound 1990, zur Strassen 1995, Kakimoto et al. 2006, Okajima 2006) most species in this genus live on two plant families, Asteraceae and Poaceae, with a few species found on plants in other families. One of these is *H. kermanensis* zur Strassen which was described from Iran based on specimens collected on *Haloxylon* sp. (Chenopodiaceae) (zur Strassen 1975), and this appears to be a specific host for this thrips (Minaei and Mound 2008). The objective of this paper is to describe a new species of *Haplothrips* collected on another chenopod species, *Suaeda* sp. These two thrips species are very similar in structure as discussed below.

Materials and methods

The new species discussed below was collected by beating leaves and flowers of *Suaeda* sp. (Chenopodiaceae) onto a plastic tray. The specimens were removed with a fine brush into a collecting vial containing 90% ethyl alcohol. They were then mounted onto slides in Canada balsam using a form of the protocol given by Mound and Kibby (1998). The line drawings were sketched using a drawing attachment. Terminology follows Mound and Minaei (2007) and Minaei and Mound (2008). The holotype and other specimens studied here are deposited in the collection of the Department of Plant Protection, College of Agriculture, Shiraz University, Shiraz, Iran. A few paratypes are deposited in the Australian National Insect Collection, Canberra and the Natural History Museum, London. The following abbreviations are used for pronotal setae: am—anteromarginals; aa—anteroangulars; ml—midlaterals; epim—epimerals; pa—posteroangulars.

Taxonomy

Haplothrips herajius sp. n.

urn:lsid:zoobank.org:act:FFCC20A9-BDC2-4700-B252-C87F008BA82B

http://species-id.net/wiki/Haplothrips_herajius

Type material. Holotype female, Iran, Fars Province, Mohr, Heraj village; *Suaeda* sp. (leaves), 31.iii.2012. (Mohsen Abdolahi); Paratypes: 58 females, 11 males taken with holotype; 14 females, 3 males, same place, *Suaeda* sp. (flowers), 21. ix. 2012.

Description. Female macroptera. Body brown (paler in summer forms), all tarsi, fore tibiae in distal half, distal apex of mid and hind tibiae are yellow; antennal segments I–II brown but the color of remaining segments variable depending on collecting date (III–VI yellow, VII–VIII yellow-brownish in summer forms; III yellow, IV–VIII yellow-brownish, gradually darker brown in spring forms); fore wing pale except for basal area; major body setae as well as sub-basal wing setae pale but tergite setae and anal setae slightly shaded at base.

Antennae 8-segmented, segment III with two, IV with four sensoria, VII slightly constricted at base, VIII short and broad at base (Fig. 1). Head a little longer than wide with maxillary stylets 0.2–0.3 of head width apart, retracted anterior to post ocular setae; post ocular setae blunt or capitate, extending to posterior margin of eye (Fig. 2). Cheeks weakly rounded. Maxillary bridge well developed. Mouth cone rounded.


Pronotum transverse, without sculpture lines except close to posterior margin; notopleural sutures complete; five pairs of developed setae present: am, aa, ml, epim and pa, all blunt or capitate (Fig. 2); prosternum with paired basantra and ferna as well as a spinasternum, ferna broad (Fig. 3). Mesonotum transversely weakly reticulate, with no microtrichia, lateral setae well developed, weakly capitate (Fig. 4). Mesopresternum eroded medially (Fig. 3). Metanotum reticulate, with no microtrichia, median setae slender and acute, arise on posterior half of sclerite, with 2–4 small setae on anterior half (Fig. 4). Fore tarsal tooth conspicuous (Fig. 2). Fore wing constricted medially (Fig. 6), sub-basal setae S1, S2 and S3 blunt or capitate, their bases arranged in a triangle (Fig. 5), 2–7 duplicated cilia present (Fig. 6).

Pelta triangular, weakly reticulate (Fig. 7). Tergite II–VII with wing-retaining setae, anterior pair weaker than posterior one, these being weakest on tergite II; tergites II–VII with a few lines of sculpture and 3–5 discal setae lateral to two pairs of developed wing-retaining setae; marginal setae S1 and S2 on tergites VII–IX long and finely pointed, S2 on other tergites finely pointed but S1 usually blunt, rarely finely pointed and sometimes variable on different tergites, tending to be more pointed on posterior than anterior tergites. Tergite VII with two campaniform sensilla not close to each other, separated by at least 0.1 width of tergite, with four micro-setae laterally; tergite VIII campaniform sensilla further apart, more than two times as those on tergite VIII, three to four micro-setae between sensilla or sometimes in front of them (Fig. 8). Tube short, about twice as long as basal width (Fig. 9); anal setae usually longer than tube.


Measurements. (holotype female, in microns). Body distended length 1845. Head, length 190; median width 180; postocular setae 35. Pronotum, length 35; width 68; major setae am 34, aa 33, ml 26, epim 51, pa 43. Fore wing length 700; sub-basal wing setae 41, 50, 68. Tergite IX setae S1 95, S2 85. Tube length 108; basal width 58. Antennal segments III–VIII length 38, 47, 43, 41, 33, 21.

Male macroptera. Color and structure similar to female. Sternites with no pore plates; tergite IX setae S2 short and stout (Fig. 10). Pseudovirga spoon shaped at apex (Fig. 11).


Diagnosis. The reticulation on the mesonotum and metanotum of *H. herajius* (Fig. 4) is unique among Iranian *Haplothrips* as well as for most other *Haplothrips* species. In other species of *Haplothrips* recorded from Iran, this reticulation is weakly



Figures 1–9. *Haplothrips herajius* sp. n. Female. **1** Antenna **2** Head and pronotum **3** Prosternum and meso-presternum **4** Mesonotum and metanotum **5** Sub basal wing setae **6** Forewing **7** Pelta **8** Tergites VII–VIII **9** Tergite IX and tube.



Figures 10–11. *Haplothrips herajius* sp. n. Male **10** Tergite IX and tube **11** Pseudovirga.



Figures 12–14. *Haplothrips kermanensis*. Female **12** Sub basal wing setae **13** Mesonotum and metanotum. Male **14** Pseudovirga.

developed or absent. The new species is very close to *H. kermanensis*. Both species have extra setae on metanotum (Figs 4, 13) that are not seen in other Iranian species of *Haplothrips*. Moreover, in both species the basal wing setae are arranged in a triangle (Figs 5, 12) (this arrangement in *H. kermanensis* was not reported by Minaei and Mound (2008), and the apex of the mid and hind tibiae are pale, also a conspicuous fore tarsal tooth is present in both species. However, the number of small setae anterior

to the median pair of metanotal setae in the new species is variable, 2–4 (rarely 0, 5 or 6), whereas available specimens of *H. kermanensis* all consistently have one pair. In addition, am setae on the pronotum in *H. herajius* are blunt or capitate in contrast to *H. kermanensis* in which they are pointed. Furthermore, fore wing sub-basal setae S3 in the new species is blunt compared with weakly pointed in *H. kermanensis*. Males of the two species are clearly different in genitalia: spoon shaped in *H. herajius* but rod shaped in *H. kermanensis* (Figs 11, 14).

Variability. Color of body and antennal segments varies among specimens, being paler in summer specimens compared with specimens collected in early spring. The fore tarsal tooth is conspicuous, but variable from small to large among male specimens. Maxillary stylets are retracted to postocular setae but rarely are low in the head and not reaching the postocular setae. Moreover, in a few specimens, the pronotal am setae are not developed.

Etymology. Heraj is a village of Mohr city in the south of Fars Province, south of Iran which is located 300 km south of Shiraz, the capital of Fars Province.

Discussion

The presence on two separate chenopod species of two *Haplothrips* species that share unusual character states, as discussed above, is interesting. The large number of collected specimens of *H. herajius* on both leaves (early spring) and flowers (late summer) of *Suaeda* sp. suggest that the new species is phytophagous, and apparently this plant species is a specific host for the new species in that area. Recently Minaei et al. (2012) described another thrips species, *Ankothrips zayandicus* (Melanthripidae), from the same plant in Isfahan Province, central Iran.

Considering that most species of Chenopodiaceae bloom in summer months, the specimens studied here were collected at two seasons: in early spring on leaves and in late summer on flowers. The color of body and antennae differs between specimens collected at these two seasons as mentioned above. This difference is remarkable because the effect of temperature on body color during development was not noticed in any species of *Haplothrips* so far. However, in onion thrips, *Thrips tabaci*, a well-known pest of thripid family, experiments showed low temperatures during pupal development induce dark adult body color (Murai and Toda 2002). Similarly, in Australia an endemic and very common thrips, plague thrips, *Thrips imaginis*, is commonly dark after winter but pale yellow during summer (Mound and Masumoto 2005).

Bhatti et al. (2009) in their book on the Thysanoptera of Iran listed 29 species of *Haplothrips*, among them two species (*Haplothrips bagnalli*, *Haplothrips* nr. *bagrolis*) reported by Manzari (2004) from Iranian islands. However, only one *Haplothrips* (*H. bagnalli*) was reported by him (the other two species were thripids), although towards the end of the report he introduces, obscurely, a comparison between *H.* nr. *bagrolis* and *H. ganglbaueri*. The somewhat cursory report was published in an informal newsletter and, given its nature; the presence of both species in Iran requires further verifi-

cation. Similarly, the report of three other species, *caespitis*, *minutus* and *rabinovitchi* by Bagheri and Alavi (2007) needs to be confirmed (Minaei and Mound 2008, Bhatti et al. 2009). Moreover, Minaei and Mound (2010) demonstrated that *H. cerealis* does not occur in Iran, the reported occurrence of this species being a misidentification of *H. tritici*. So with the addition of the species described here, the confirmed Iranian species in *Haplothrips* comprise 24 species i.e. about 12% of the total Thysanoptera fauna in this country. Similarly this genus comprises about 8% of Thysanoptera fauna in Britain (Collins 2010). In contrast, the species composition of the Australian insect fauna (southern Hemisphere) is very different and the species of *Haplothrips* comprise about 3% of the known thrips fauna of that continent (Mound and Minaei 2007).

Acknowledgement

We are much indebted to Mr Mohsen Abdolahi and Ms Fatemeh Mohammadzadeh, BSc students of Plant Protection Department, Shiraz University, for collecting the materials of the new species described in this paper and sketching the illustrations respectively. Dr Ahmad Reza Khosravi, Department of Biology, Shiraz University, identified the plant on which the thrips were collected. The manuscript was improved through the advice and criticisms kindly provided by Dr Alice Wells (Australian Biological Resources Study, Canberra, Australia) and two anonymous reviewers. Open access to this paper was supported by the Encyclopedia of Life (EOL) Open Access Support Project (EOASP).

References

- Bagheri S, Alavi J (2007) New records of 20 species Thysanoptera on forest and rangeland plants for Iran and Khuzestan province. Proceedings of the 17th Iranian Plant Protection Congress; Tehran, Karaj I: 78.
- Bailey P, Caon G (1986) Predation on two-spotted mite, *Tetranychus urticae* Koch (Acarina: Tetranychidae) by *Haplothrips victoriensis* Bagnall (Thysanoptera: Phlaeothripidae) and *Stethorus nigripes* Kapur (Coleoptera: Coccinellidae) on seed lucerne crops in South Australia. Australian Journal of Zoology 34: 515–525. doi: 10.1071/ZO9860515
- Bhatti JS, Alavi J, zur Strassen R, Telmadarraiy Z (2009) Thysanoptera in Iran 1938–2007. An Overview. Part 1. Thrips 7: 1–82.
- Buckman RS, Mound LA, Whiting MF (2013) Phylogeny of Thrips (Insecta: Thysanoptera) based on five molecular loci. Systematic Entomology 38: 123–133. doi: 10.1111/j.1365-3113.2012.00650.x
- Collins DW (2010) Thysanoptera of Great Britain: a revised and updated checklist. Zootaxa 2412: 21–41.
- Goldarazena A, Gattesco F, Atencio R, Korytowski C (2012) An updated checklist of the Thysanoptera of Panama with comments on host associations. Check List 8: 1232–1247. Available from: <http://200.144.120.19/pdf/SL070-12.pdf>

- Kakimoto K, Inoue H, Hinomoto N, Noda T, Hirano K, Kashio T, Kusigemati K, Okajima S (2006) Potential of *Haplothrips brevitubus* (Karny) (Thysanoptera: Phlaeothripidae) as a predator of mulberry thrips *Pseudodendrothrips mori* (Niwa) (Thysanoptera: Thripidae). *Biological Control* 37: 314–319. doi: 10.1016/j.biocontrol.2006.01.014
- Manzari S (2004) Report of three species of thrips from Iranian islands in Persian Gulf. *Newsletter of Entomological Society of Iran* 21: 2 (in Persian).
- Minaei K (2013) The *Phlaeothrips*-lineage of fungus feeding thrips in Iran with a new species of *Hindsiothrips*. *Zootaxa* 3599: 279–290. doi: 10.11646/zootaxa.3599.3.5
- Minaei K, Haftbaradaran F, Mound LA (2012) A new *Ankothrips* species (Thysanoptera: Melanthripidae) from Iran with unusually short setae. *Zootaxa* 3552: 37–42. Available from: <http://www.mapress.com/zootaxa/2012/f/zt03552p042.pdf>
- Minaei K, Mound LA (2010) Taxonomic problems in character state interpretation: variation in the wheat thrips *Haplothrips tritici* (Kurdjumov) (Thysanoptera: Phlaeothripidae) in Iran. *Deutsche Entomologische Zeitschrift* 57: 233–241. doi: 10.1002/mmnd.201000020
- Minaei K, Mound LA (2008) The Thysanoptera Haplothripini (Phlaeothripidae) of Iran. *Journal of Natural History* 42: 2617–2658. Available from: <http://www.tandfonline.com/doi/pdf/10.1080/00222930802354159>, doi: 10.1080/00222930802354159
- Mound LA (2013) Thysanoptera (Thrips) of the world- a checklist. Available from: <http://www.ento.csiro.au/thysanoptera/worldthrips.html>
- Mound LA, Heming BS, Palmer JM (1980) Phylogenetic relationships between the families of recent Thysanoptera. *Zoological Journal of the Linnean Society of London* 69: 111–141. doi: 10.1111/j.1096-3642.1980.tb01934.x
- Mound LA, Kibby G (1998) *Thysanoptera: an identification guide*. Second edition. – CAB International Institute of Entomology and British Museum (Natural History), London 70 pp.
- Mound LA, Masumoto M (2005) The genus *Thrips* (Thysanoptera, Thripidae) in Australia, New Caledonia and New Zealand. *Zootaxa* 1020: 1–64. Available from: <http://www.mapress.com/zootaxa/2005f/zt01020p064.pdf>
- Mound LA, Minaei K (2007) Australian thrips of the *Haplothrips* lineage (Insecta: Thysanoptera). *Journal of Natural History* 41: 2919–2978. Available from: <http://www.tandfonline.com/doi/abs/10.1080/00222930701783219>, doi: 10.1080/00222930701783219
- Mound LA, Morris DC (2007) The insect order Thysanoptera: classification versus systematics. *Zootaxa* 1668: 395–411. Available from: <http://www.mapress.com/zootaxa/2007f/zt01668p411.pdf>
- Mound LA, Zapater MC (2003) South American *Haplothrips* species (Thysanoptera: Phlaeothripidae), with a new species of biological control interest to Australia against weedy *Heliotropium amplexicaule* (Boraginaceae). *Neotropical Entomology* 32: 437–442. Available from: <http://www.scielo.br/pdf/ne/v32n3/18757.pdf>, doi: 10.1590/S1519-566X2003000300009
- Murai T, Toda S (2002) Variation of *Thrips tabaci* in color and size. In: Marullo R and Mound LA (Eds) *Thrips and Tospoviruses: Proceedings of the 7th international symposium on thysanoptera*. Australian National. Insect Collection, Canberra, 377–378. Available from: <http://www.ento.csiro.au/thysanoptera/Symposium/Section10/58-Murai-Toda.pdf>
- Okajima S (2006) The suborder Tubulifera (Thysanoptera). *Insects of Japan* 2: 1–720.

- Palmer JM, Mound LA (1990) Thysanoptera In: Rosen D (Ed) The armoured scale insects, their biology, natural enemies and control. Amsterdam (The Netherlands), Elsevier, 67–75.
- Pitkin BR (1976) A revision of the Indian species of *Haplothrips* and related genera (Thysanoptera: Phlaeothripidae). Bulletin of the British Museum (Natural History) (Entomology) 34: 223–280.
- Putman WL (1965) The predaceous thrips *Haplothrips faurei* Hood (Thysanoptera: Phlaeothripidae) in Ontario peach orchards. Canadian Entomologists 97: 1208–1221. doi: 10.4039/Ent971208-11
- zur Strassen R (1975) Eine neue Haloxylon-bewohnende *Haplothrips*-Art aus dem südlichen Iran (Thysanoptera, Phlaeothripidae). Entomologische Zeitschrift 85: 137–143.
- zur Strassen R (1995) Bionomial data of some predacious thrips. In: Parker BL (Ed) Thrips biology and management. New York, Plenum Press, 325–328.