

A new species of sardine, *Sardinella pacifica* from the Philippines (Teleostei, Clupeiformes, Clupeidae)

Harutaka Hata¹, Hiroyuki Motomura²

1 The United Graduate School of Agricultural Sciences, Kagoshima University, 1–21–24 Korimoto, Kagoshima 890–0065, Japan **2** The Kagoshima University Museum, 1–21–30 Korimoto, Kagoshima 890–0065, Japan

Corresponding author: Hata Harutaka (k2795502@kadai.jp)

Academic editor: Nina Bogutskaya | Received 19 October 2018 | Accepted 11 January 2019 | Published 11 March 2019

<http://zoobank.org/8F6BD1CD-B01B-4BA5-BD29-677E7A161E8B>

Citation: Hata H, Motomura H (2019) A new species of sardine, *Sardinella pacifica* from the Philippines (Teleostei, Clupeiformes, Clupeidae). ZooKeys 829: 75–83. <https://doi.org/10.3897/zookeys.829.30688>

Abstract

A new sardine, *Sardinella pacifica* **sp. n.**, is described on the basis of 21 specimens collected from the Philippines. The new species closely resembles *Sardinella fimbriata* (Valenciennes, 1847), both species having lateral scales with centrally discontinuous striae, a dark spot on the dorsal-fin origin, more than 70 lower gill rakers on the first gill arch, the pelvic fin with eight rays, and 17 or 18 prepelvic and 12 or 13 postpelvic scutes. However, the new species is distinguished from the latter by lower counts of lateral scales, pseudobranchial filaments, and postpelvic scutes (38–41, 14–19 and 12–13, respectively vs. 44–46, 19–22 and 13–14), and a shorter lower jaw (10.4–11.6% of standard length vs. 11.1–12.2%). *Sardinella pacifica* **sp. n.** is known only from the Philippines, whereas *S. fimbriata* is restricted to the Indian Ocean, although previously considered to be an Indo-West Pacific species, distributed from India to the Philippines.

Keywords

morphology, *Sardinella fimbriata*, Southeast Asia, taxonomy

Introduction

Sardinella Valenciennes, 1847, an Indo-Pacific and Atlantic genus of marine, brackish and/or fresh water sardines (Clupeidae), comprises 22 valid species (Whitehead 1985, Stern et al. 2016). Many species, including the endemic fresh water species *Sardinella tawilis* (Herre, 1927), occur in the Philippines (Fowler 1941, Rau and Rau 1980,

Whitehead 1985, Conlu 1986, Munroe et al. 1999, Willette et al. 2011a, b, Willette and Santos 2012, Stern et al. 2016, Hata 2017a, b), some being an important fisheries resource (locally named “tambam”, “tuloy”, and “tunsoy”) (Rau and Rau 1980, Conlu 1986). A recently described species, *Sardinella goni* Stern, Rinkevich & Goren, 2016 was based on specimens collected from Boracay Island, the Philippines.

During a revisionary study of *Sardinella*, 21 specimens of a clupeid fish from the Philippines were found to be characterized by a unique combination of scales with centrally discontinuous striae, a dark spot on the dorsal-fin origin, and low counts of lateral scales in the longitudinal series and pseudobranchial filaments. They are described herein as a new species of *Sardinella*.

Materials and methods

Counts and proportional measurements followed Hubbs and Lagler (1947) with additions as in Kimura et al. (2009). All measurements were made with digital calipers to the nearest 0.01 mm. Standard length is abbreviated as SL. Institutional codes follow Sabaj (2016).

Sardinella pacifica sp. n.

<http://zoobank.org/30675329-0FBF-45F3-ACF5-30D6C6669C39>

Figures 1–2, Table 1

Sardinella fimbriata: Fowler 1941: 609 (Bacon, Manilla and Aparri, Philippines); Chan 1965 (in part): 14 (Philippines); Rau and Rau 1980: 203 (Philippines); Whitehead 1985 (in part): 98, unnumbered fig. (Philippines); Conlu 1986: 45, fig. 20 (Alabat Island; Appari, Cagayan; Bacon, Sorsogon; Bauang, La Union; Calapan, Mindoro; Cavite, Cavite; Davao Gulf; Estancia, Iloilo; Malolos, Bulacan; Manila Bay; Margosatubig, Zamboanga; Nasugbu, Batangas; Ragay Gulf, Quezon; San Miguel Bay, Camarines Sur; Samar, Philippines); Munroe et al. 1999 (in part): 1814, unnumbered fig. (Philippines); Luceño et al. 2013: 30, fig. 2 (Butuan, Dipolog, and Pagadian, Mindanao Island, Philippines); Stern et al. 2016 (in part): 9, fig. 2 (b), fig. 4 (b) (Manilla, Philippines). (**non** Valenciennes)

Holotype. BMNH 1985.4.12.1, 105.1 mm SL, Manila Harbor, Manila Bay, Luzon Island, Philippines.

Paratypes. 20 specimens, 90.2–105.9 mm SL, all from the Philippines. BMNH 1960.4.7.52, 90.2 mm SL, Palawan Island; BMNH 1985.4.12.2, 98.7 mm SL, Manila Harbor, Manila Bay, Luzon Island; CAS 38365, 105.9 mm SL, Manila Bay, Luzon Island; CAS 51909, 96.5 mm SL, Manila Bay, Luzon Island; CAS 52501, 98.4 mm SL, Manila Bay, Luzon Island; CAS 59712, 100.3 mm SL, Bacon, Sorsogon, Luzon Island; CAS-SU 28568, 96.5 mm SL, Alabat Island; CAS-SU 28569, 101.3 mm SL,




Figure 1. Holotype of *Sardinella pacifica* sp. n., BMNH 1985.4.12.1, 105.1 mm SL, Manila Bay, Luzon Island, Philippines.

Manila Bay, Luzon Island; CAS-SU 29920, 2 specimens, 97.6–103.3 mm SL, Manilla Bay, Luzon Island; CAS-SU 32915, 2 specimens, 95.7–97.8 mm SL, Manila Bay, Luzon Island; KAUM–I. 125000, 95.9 mm SL, Manila Bay, Luzon Island; USNM 56232, 94.5 mm SL, USNM 56233, 92.2 mm SL, Bacon, Sorsogon, Luzon Island; USNM 72197, 92.9 mm SL, Manila, Luzon Island; USNM 177667, 2 specimens, 93.4–96.7 mm SL; USNM 403460, 95.9 mm SL, Navatos, Manila, Luzon Island; USNM 427789, 94.9 mm SL, Catbalogan, Samar Island, Visayas.

Diagnosis. A species of *Sardinella* with the following combination of characters: caudal fin with black posterior margin; lateral body scales with centrally discontinuous vertical striae, and few perforations and pores posteriorly; 38–41 (modally 38) lateral scales in longitudinal series; body scales deciduous; black spot on dorsal-fin origin; pelvic fin with one unbranched and seven branched rays; gill rakers 40–53 (43) in upper series on 1st gill arch, 71–84 (72) in lower series, 112–137 (118) in total; gill rakers 40–56 (42) in upper series on 2nd gill arch, 70–94 (79) in lower series, 112–148 (115) in total; gill rakers 37–52 (42) in upper series on 3rd gill arch, 57–75 (60) in lower series, 95–127 (99) in total; gill rakers 31–43 (35) in upper series on 4th gill arch, 44–63 (48) in lower, 78–106 (80) in total; gill rakers 30–43 (34) on hind face of 3rd gill arch; 17 or 18 (18) + 12 or 13 (13) = 29–31 (30) scutes on ventral edge of body; anal fin with 18–21 (20) rays; lower jaw rather short, 10.4–11.6% of SL.

Description. Counts and measurements, expressed as percentages of SL, are given in Table 1. Data for the holotype are presented first, followed by paratype data in parentheses. Body oblong, compressed, deepest at dorsal-fin origin. Dorsal profile of body elevated from snout tip to dorsal-fin origin, thereafter decreasing to uppermost point of caudal-fin base. Ventral profile of body curved downward from lower-jaw tip to pelvic-fin insertion, thereafter rounded to ventralmost point of caudal-fin base. Abdomen from isthmus to anus with 30 (29–31) scutes. Predorsal scutes absent. Anteriormost point of pectoral-fin insertion anterior to posteriormost point of opercle. Upper, posterior and ventral margins of pectoral fin nearly linear. Posterior tip of pectoral fin pointed. Pectoral-fin axillary scale present. Posteriormost dorsal-fin ray not filamentous. Anteriormost point of pelvic-fin insertion located directly below origin of 8th (7th–10th) dorsal-fin ray. Posterior tip of depressed pelvic fin reaching between a vertical through posterior end of dorsal-fin base and anus. Pelvic-fin axillary scale

Table 1. Counts and measurements of specimens of *Sardinella pacifica* sp. n. and *S. fimbriata*.

	<i>Sardinella pacifica</i> sp. n.			<i>Sardinella fimbriata</i>		
	Holotype	Paratypes		Lectotype	Non-types	
	Manilla Bay, Philippines BMNH 1985.4.12.1	Philippines n = 20	Modes	Malabar, India MNHN 3227	Indian Ocean n = 16	Modes
Standard length (SL; mm)	105.1	90.2–105.9		118.2	89.7–123.6	
Counts						
Dorsal-fin rays (unbranched)	4	4–5	4	4	4	4
Dorsal-fin rays (branched)	15	14–16	14	15	14–16	15
Anal-fin rays (unbranched)	3	3	3	3	3	3
Anal-fin rays (branched)	17	15–18	17	16	15–19	17
Pectoral-fin rays (unbranched)	1	1	1	1	1	1
Pectoral-fin rays (branched)	13	12–15	14	14	13–16	14
Pelvic-fin rays (unbranched)	1	1	1	1	1	1
Pelvic-fin rays (branched)	7	7	7	7	7	7
Caudal-fin rays (upper+ lower)	10 + 9	10 + 9	10 + 9	10 + 9	10 + 9	10 + 9
Gill rakers on 1st gill arch (upper)	44	40–53	43	49	40–49	49
Gill rakers on 1st gill arch (lower)	72	71–84	72	74	71–79	78
Gill rakers on 1st gill arch (total)	116	112–137	118	123	112–127	121
Gill rakers on 2nd gill arch (upper)	44	40–56	42	47	40–53	48
Gill rakers on 2nd gill arch (lower)	79	70–94	79	87	75–95	87
Gill rakers on 2nd gill arch (total)	123	112–148	115	134	115–146	123
Gill rakers on 3rd gill arch (upper)	43	37–52	42	49	37–50	45
Gill rakers on 3rd gill arch (lower)	61	57–75	60	69	60–82	75
Gill rakers on 3rd gill arch (total)	104	95–127	99	118	100–131	122
Gill rakers on 4th gill arch (upper)	35	31–43	35	39	30–40	36
Gill rakers on 4th gill arch (lower)	49	44–63	48	51	43–53	48
Gill rakers on 4th gill arch (total)	84	78–106	80	90	74–93	90
Gill rakers on posterior face of 3rd gill arch	32	30–43	34	36	31–39	36
Prepelvic scutes	18	17–18	18	18	17–18	18
Postpelvic scutes	12	12–13	13	14	13–14	14
Total scutes	30	29–31	30	32	31–32	32
Lateral scales in longitudinal series	41	38–41	38	45	44–46	45
Pseudobranchial filaments	18	14–19	18	21	19–22	21
Measurements (%SL)			Means			Means
Head Length	24.9	23.1–26.8	25.3	26.5	25.0–28.5	26.5
Body depth	31.5	28.3–36.9	31.3	33.1	28.8–32.3	31.0
Pre-dorsal-fin length	42.3	41.9–46.1	44.3	44.9	43.9–46.7	45.2
Snout tip to pectoral-fin insertion	25.1	24.2–27.9	26.3	27.9	25.2–27.9	26.8
Snout tip to pelvic-fin insertion	51.4	48.3–55.1	51.7	53.4	50.5–53.1	51.7
Pre-anal-fin length	76.4	72.9–79.3	77.3	77.9	75.2–77.7	76.7
Dorsal-fin base length	16.1	13.9–16.5	15.2	13.4	13.4–16.6	14.9
Anal-fin base length	15.7	13.9–17.6	16.0	16.0	13.7–17.5	16.1
Caudal-peduncle length	9.5	7.7–10.4	9.0	9.1	8.1–10.2	9.4
Caudal-peduncle depth	9.5	9.1–11.1	10.0	10.5	9.5–10.0	9.8
Dorsal-fin origin to pectoral-fin insertion	31.8	28.4–34.2	32.1	33.9	30.6–34.1	32.1
Dorsal-fin origin to pelvic-fin insertion	30.5	27.4–35.9	30.5	32.3	28.0–31.4	30.1
Dorsal-fin origin to anal-fin origin	42.1	39.9–45.0	43.0	43.0	39.5–43.5	41.4
Pectoral-fin insertion to pelvic-fin insertion	27.9	23.5–9.5	26.7	27.0	25.0–27.4	25.9
Pelvic-fin insertion to anal-fin origin	30.5	26.1–31.8	28.8	27.7	26.1–29.7	27.4
Pectoral-fin length	broken	18.2–20.8	19.6	18.8	17.4–19.7	18.7
Pelvic-fin length	10.7	10.3–11.9	11.2	11.0	9.5–11.6	10.9
Interorbital width	4.2	4.0–5.0	4.5	4.6	3.8–5.0	4.5
Postorbital length	12.4	10.4–13.7	12.0	12.6	11.2–13.9	12.8
Upper-jaw length	9.6	9.3–10.9	10.0	10.7	9.5–11.0	10.4
Mandible length	10.7	10.4–11.6	11.0	11.6	11.1–12.2	11.5
1st unbranched dorsal-fin ray length	1.5	1.1–3.0	1.8	0.9	1.4–3.5	2.1

	<i>Sardinella pacifica</i> sp. n.			<i>Sardinella fimbriata</i>		
	Holotype Manilla Bay, Philippines BMNH 1985.4.12.1	Paratypes Philippines n = 20	Means	Lectotype Malabar, India MNHN 3227	Non-types Indian Ocean n = 16	Means
2nd unbranched dorsal-fin ray length	7.2	2.5–9.0	5.2	2.7	3.5–7.5	5.1
3rd unbranched dorsal-fin ray length	11.8	6.2–13.7	9.5	7.6	7.4–12.5	9.4
1st unbranched anal-fin ray length	1.0	0.8–1.9	1.3	broken	0.6–1.8	1.3
2nd unbranched anal-fin ray length	broken	2.3–5.3	3.6	broken	1.9–3.9	2.8
3rd unbranched anal-fin ray length	broken	4.9–6.1	5.6	5.7	4.6–6.5	5.5
1st pectoral-fin ray length	broken	16.9–19.9	18.5	broken	16.5–19.4	18.0
1st pelvic-fin ray length	10.6	10.3–11.9	11.1	11.0	10.4–11.6	10.9

present. Anal-fin origin posterior to vertical through posteriormost point of dorsal-fin base. Two posteriormost anal-fin rays enlarged. Caudal fin forked. Posterior tips of caudal-fin lobes pointed. Anus on ventral midline, slightly anterior to anal-fin origin, posterior to midpoint of body. Scales cycloid, thin, deciduous, except for robust ventral scutes. Scales on lateral body surface with several centrally discontinuous vertical striae, few perforations and pores posteriorly (Fig. 2). Bases of dorsal and anal fins with low scaly sheaths. Predorsal scales paired. No elongate, wing-like scales present beneath normal paired scales. No scales on head and fins, except for a broad triangular sheath of scales on caudal fin. Mouth terminal, small, posterior tip of maxilla not reaching vertical through anterior margin of iris. Premaxilla and hypomaxilla without teeth. Ventral margin of maxilla toothed. Lower jaw with several conical teeth anteriorly. Posterior ramus of lower jaw elevated. Second supramaxilla symmetrical. Orbit elliptical, eye and iris round. Eyes covered with adipose eyelid posteriorly. Interorbital space flat. Nostrils close to each other, anterior to orbit. Eight (8–10) on top of head. No lateral line. Gill rakers long, slender, with small asperities on anterior surface. Pseudobranchial filaments present. Gill opening with two fleshy outgrowths on posterior margin and a large papilla on ventral margin. Posterior margins of preopercle and opercle smooth.

Color of preserved specimens. Body dark brown dorsally, elsewhere yellowish silver. Black spot on dorsal-fin origin. Melanophores scattered on upper part of dorsal fin and first pectoral-fin ray. Posterior margin of caudal fin dark. Fresh coloration shown in Luceño et al. (2013) (as *S. fimbriata*).

Distribution. Currently known only from the Philippines.

Etymology. The specific name *pacifica* (in reference to the Pacific Ocean) is given to distinguish the species from *S. fimbriata*, with which it had been confused, and which is now considered to be restricted to the Indian Ocean.

Remarks. The new species is assignable to the genus *Sardinella*, defined by Whitehead (1985) and Munroe et al. (1999), due to its compressed body, abdomen covered with prominently keeled scutes, paired predorsal scales, a symmetrical second supramaxilla, toothless hypo-maxilla, two posteriormost anal-fin rays enlarged, the dorsal fin without filamentous rays, and two fleshy outgrowths on the hind margin of the gill opening. It most closely resembles *Sardinella fimbriata* (Valenciennes, 1847) (Fig. 3), sharing centrally discontinuous striae on the lateral body scales, a dark spot on the dor-



Figure 2. Photograph of a stained scale, collected from mid-body below the dorsal fin, of *Sardinella pacifica* sp. n. (BMNH 1985.4.12.1, 105.1 mm SL, Manila Bay, Luzon Island, Philippines).



Figure 3. Lectotype of *Sardinella fimbriata*, MNHN 3227, 118.2 mm SL, Malabar, India.

sal-fin origin, more than 70 lower gill rakers on the first gill arch, eight pelvic-fin rays, 15–18 branched anal-fin rays, and 17 or 18 prepelvic scutes (Whitehead 1985, Munroe et al. 1999, Stern et al. 2016). However, *S. pacifica* can be distinguished from the latter by lower counts of lateral scales in the longitudinal series (38–41 vs. 44–46 in *S. fimbriata*; Table 1), pseudobranchial filaments (14–19 vs. 19–22; Table 1; Fig. 4A) and postpelvic scutes (12 or 13 vs. 13 or 14; Table 1), and a shorter lower jaw (10.4–11.6% SL vs. 11.1–12.2%; Table 1; Fig. 4B). Moreover, the deciduous body scales of the new species are distinctively diagnostic, the body scales of *S. fimbriata* being non-deciduous. Although *S. fimbriata* has been regarded as an Indo-West Pacific species, distributed from India to the Philippines (Whitehead 1985, Munroe et al. 1999, Stern et al. 2016), no Pacific region specimens of *S. fimbriata* appear to have been collected (see comparative materials), and the species is judged herein to be an Indian Ocean endemic.




Figure 4. Relationships of (A) pseudobranchial filament numbers, and (B) lower-jaw length (as % of standard length) to SL in *Sardinella pacifica* sp. n. [solid circles (solid star = holotype)] and *S. fimbriata* [open triangles (open star = lectotype)]

Comparative material examined. *Sardinella fimbriata* (Valenciennes, 1847) (17 specimens, 89.7–123.6 mm SL): BMNH 1889.2.1.1778, 112.0 mm SL, Madras, India; BMNH 1889.2.1.1915–1916, 1 of 2 specimens, 97.9 mm SL, Orissa, India; BMNH 1889.2.1.1917, 99.5 mm SL, Akyab, Burma; CAS 41433, 2 specimens, 89.7–93.9 mm SL, Calicut, India; CAS 41434, 2 specimens, 94.2–97.2 mm SL, Ernakulam, Cochin, India; CAS 41435, 119.3 mm SL, Madras, India; MNHN 3227, lectotype of *Spratella fimbriata*, 118.2 mm SL, Malabar, India; USNM 276446, 121.8 mm SL, Cochin, Kerala, India; USNM 276447, 105.6 mm SL, Kovalam, Trivandrum, India; USNM 276449, 2 specimens, 108.7–114.6 mm SL, Vizhinam, Trivandrum, Kerala, India; USNM 276450, 4 specimens, 111.4–123.6 mm SL, Calicut, Kerala, India.

Acknowledgements

We thank O Crimmen and J Maclaine (BMNH), D Catania (CAS), R Causse and P Pruvost (MNHN), and J Williams, K Murphy, S Raredon and D Pitassy (USNM), for opportunities to examine specimens of *Sardinella*. We also thank Y Haraguchi and other volunteers, and students of KAUM for their curatorial assistance, and G Hardy (Ngunguru, New Zealand) for reading the manuscript and providing help

with English. This study was supported in part by a Grant-in-Aid from the Japan Society for the Promotion of Science for JSPS Fellows (DC2: 29-6652); the Sasakawa Scientific Research Grant from the Japan Science Society (28-745); JSPS KAKENHI Grant Numbers JP19770067, JP26241027, JP24370041, JP23580259, and JP26450265; the “Biological Properties of Biodiversity Hotspots in Japan” project of the National Museum of Nature and Science, Tsukuba, Japan; “Establishment of Research and Education Network on Biodiversity and Its Conservation in the Satsunan Islands” project of Kagoshima University adopted by the Ministry of Education, Culture, Sports, Science and Technology, Japan; and the “Island Research” project by Kagoshima University.

References

- Chan WL (1965) A systematic revision of the Indo-Pacific clupeid fishes of the genus *Sardinella* (family Clupeidae). Japanese Journal of Ichthyology 13 (1/3): 1–39.
- Conlu PV (1986) Guide to Philippine flora and fauna. Volume IX, fishes. Natural Resources Management Center Ministry of Natural Resources, Manila and University of the Philippines, Manila, 495 pp.
- Fowler HW (1941) Contributions to the biology of the Philippine archipelago and adjacent regions. The fishes of the groups Elasmobranchii, Holocephali, Isospondyli, and Ostariophysi obtained by the United States Bureau of Fisheries steamer “Albatross” in 1907 to 1910, chiefly in the Philippine Islands and adjacent seas. Bulletin of the United States National Museum No. 100 13: 1–879.
- Hata H (2017a) *Sardinella gibbosa* (Bleeker 1849). In: Motomura H, Alama UB, Muto N, Babaran RP, Ishikawa S (Eds) Commercial and bycatch market fishes of Panay Island, Republic of the Philippines, The Kagoshima University Museum, Kagoshima, University of the Philippines Visayas, Iloilo and Research Institute for Humanity and Nature, Kyoto, 41.
- Hata H (2017b) *Sardinella* cf. *longiceps* Valenciennes 1847. In: Motomura H, Alama UB, Muto N, Babaran RP, Ishikawa S (Eds) Commercial and bycatch market fishes of Panay Island, Republic of the Philippines, The Kagoshima University Museum, Kagoshima, University of the Philippines Visayas, Iloilo and Research Institute for Humanity and Nature, Kyoto, 42.
- Herre AWCT (1927) Four new fishes from Lake Taal (Bombon). Philippine Journal of Science 34(3): 273–279. [pls. 1–3]
- Hubbs CL, Lagler JF (1947) Fishes of the Great Lakes region. Bulletin of Cranbrook Institute of Sciences 26: 1–186.
- Kimura S, Hori K, Shibukawa K (2009) A new anchovy, *Stolephorus teguhi* (Clupeiformes: Engraulidae), from North Sulawesi, Indonesia. Ichthyological Research 56: 292–295. <https://doi.org/10.1007/s10228-009-0103-4>
- Luceño AJM, Torres MAJ, Tabugo SR, Demayo CG (2013) Describing the body shapes of three populations of *Sardinella fimbriata* (Valenciennes, 1847) from Mindanao Island, Philippines using relative warp analysis. Annals of Biological Research 4(10): 29–39.

- Munroe TA, Wongratana T, Nizinski, MS (1999) Order Clupeiformes. Clupeidae. Herrings (also, sardines, sprat, pilchard, and menhades). In: Carpenter KE, Niem VH (Eds) FAO Species Identification Guide for Fishery Purposes. The Living Marine Resources of the Western Central Pacific, Vol. 3. Batoid Fishes, Chimaeras and Bony Fishes Part 1 (Elopidae to Linophrynidae), FAO, Rome, 1775–1821.
- Rau N, Rau A (1980) Commercial fishes of the Philippines. German Agency for Technical Cooperation, Eschborn, 623 pp.
- Sabaj MH (2016) Standard symbolic codes for institutional resource collections in herpetology and ichthyology: an online reference. Version 6.5 (16 August 2016). <http://www.asih.org/resources/standard-symbolic-codes-institutionalresource-collections-herpetology-ichthyology> [1 March 2018]
- Stern N, Rinkevich B, Goren M (2016) Integrative approach revises the frequently misidentified species of *Sardinella* (Clupeidae) of the Indo-West Pacific Ocean. *Journal of Fish Biology* 89(5): 2282–2305. <https://doi.org/10.1111/jfb.13114>
- Valenciennes A (1847) Chapitre VIII. Du genre *Spratella* (*Spratella*). In: Cuvier G, Valenciennes A (Eds) Histoire naturelle des poissons. Tome vingtième. Livre vingt et unième. De la famille des Clupéoïdes. vol 20, P Bertrand, Paris, 356–361.
- Whitehead PJP (1985) FAO species catalogue Vol. 7. Clupeoid Fishes of the World (Suborder Clupeoidei). An Annotated and Illustrated Catalogue of the Herrings, Sardines, Pilchards, Sprats, Shads, Anchovies and Wolf-herrings. Part 1, Chirocentridae, Clupeidae and Pristigasteridae. FAO Fisheries Synopsis 125, 7(1): 1–303.
- Willette DA, Bognot EDC, Mutia MT, Santos MD (2011a) Biology and ecology of sardines in the Philippines; a review. Bureau of Fisheries and Aquatic Resources National Fisheries Research and Development Institute Technical Paper Series 13(1): 1–18.
- Willette DA, Santos MD (2012) Correcting widespread misidentifications of the highly abundant and commercially important sardine species *Sardinella lemuru* in the Philippines. *Journal of Applied Ichthyology* 2012: 1–5.
- Willette DA, Santos MD, Aragon MA (2011b) First report of the Taiwan sardinella *Sardinella hualiensis* (Clupeiformes: Clupeidae) in the Philippines. *Journal of Fish Biology* 79: 2087–2094. <https://doi.org/10.1111/j.1095-8649.2011.03133.x>