

A new species of *Scelidopetalon* Delkeskamp (Coleoptera, Erotylidae) from China with a key to world species of the genus

Cong-Chao Dai^{1,†}, Mei-Jun Zhao^{1,‡}

¹ Department of Biology, College of Life and Environmental Sciences, Shanghai Normal University, Shanghai, 200234, P. R. China

[†] [urn:lsid:zoobank.org/author:F296F7D8-8259-4A69-841C-C2F74F0A54E7](https://doi.org/urn:lsid:zoobank.org/author:F296F7D8-8259-4A69-841C-C2F74F0A54E7)

[‡] [urn:lsid:zoobank.org/author:FBE4FA1F-5BCF-405E-98C1-4F86894643EA](https://doi.org/urn:lsid:zoobank.org/author:FBE4FA1F-5BCF-405E-98C1-4F86894643EA)

Corresponding author: Mei-Jun Zhao (mjzhao@shnu.edu.cn)

Academic editor: Michael Thomas | Received 12 March 2013 | Accepted 15 July 2013 | Published 19 July 2013

[urn:lsid:zoobank.org/pub:AA71D289-CF05-4C7D-A7A4-E5A9E99E4FC6](https://doi.org/urn:lsid:zoobank.org/pub:AA71D289-CF05-4C7D-A7A4-E5A9E99E4FC6)

Citation: Dai C-C, Zhao M-J (2013) A new species of *Scelidopetalon* Delkeskamp (Coleoptera, Erotylidae) from China with a key to world species of the genus. ZooKeys 317: 81–87. doi: 10.3897/zookeys.317.5112

Abstract

A new species *Scelidopetalon biwenxuanii* sp. n. is described from China, representing the first occurrence of the genus in Hainan province. A key to the world species of this genus is provided.

Keywords

Coleoptera, Erotylidae, *Scelidopetalon*, *Pseudamblyopus* identification key, new record genus, new species, China

Introduction

The subfamily Erotylinae includes colorful fungus-feeding beetles commonly called “pleasing fungus beetles”. They are worldwide in distribution with over 2500 described species. In general, species of the tribe Tritomini are characterized by an antennal club of 3 to 5 antennomeres, dilated maxillary palpi, closed procoxal cavities (with prosternal-proepimeral suture located at the midpoint posterior to the procoxae), meso-metasternal suture having a broadened dicondylic connection, and pseudotetramerous tarsi (tarsomere IV small and hidden by an expanded tarsomere III). Larvae and adults feed on larger basidiomycete fungi (e.g. mushrooms, polypore or bracket fungi, etc.).

The name “pleasing fungus beetle” is likely derived from the large size and colorful patterns of many species.

While examining Burmese specimens of *Amblyopus* Lacordaire, Gorham (1896) established *Petaloscelis* based on its small and finely faceted eyes (Fig. 1), which are large and coarsely faceted in *Amblyopus* (Fig. 2). Arrow (1925, 1926) described four additional species in *Petaloscelis*. While examining the African species, Arrow (1945) noted that Gorham confused African species with an Indian species and considered *Petaloscelis* a synonym of *Amblyscelis* Gorham. Delkeskamp (1957) thought *Petaloscelis* and *Amblyscelis* were distinct based on eye characters as mentioned above and on differences in tibial dilation. Because the name *Petaloscelis* Gorham was a junior homonym of *Petaloscelis* Bergoth (see Chûjô and Chûjô 1990), Delkeskamp (1957) proposed the name *Scelidopetalon* for the species of *Petaloscelis* Gorham with *Petaloscelis instabilis* Gorham, 1896, as the type.

Araki (1941) described *Pseudamblyopus* for the Japanese species *Amblyopus palmipes* Lewis. Araki noted *Pseudamblyopus* is easily distinguished from *Amblyopus* by the small and finely faceted eyes and mentioned this genus differs from *Petaloscelis* in the antennal club structure. But neither Arrow nor Delkeskamp were aware of Araki’s work. Thus, *Scelidopetalon* and *Pseudamblyopus* were both separated from *Amblyopus* based on the same characters.

Because no specimens of *Pseudamblyopus* are available for study, we are not treating *Scelidopetalon* as a synonym to *Pseudamblyopus*. This should be considered in future revisionary work.

Previously, a total of six species have been reported for *Scelidopetalon* and two species for *Pseudamblyopus* worldwide.

Scelidopetalon instabilis (Gorham 1896) (Burma, Viet-Nam)

Scelidopetalon similis (Arrow 1925) (Assam Valley)

Scelidopetalon varicolor (Arrow 1925) (India)

Scelidopetalon solidus (Arrow 1925) (India)

Scelidopetalon fasciatus (Arrow 1926) (Sumatra, N. Borneo)

Scelidopetalon arrowi Delkeskamp (1957) (Singapore)


Pseudamblyopus palmipes (Lewis 1889) (Japan)

Pseudamblyopus similis (Lewis 1887) (Far East, Japan)


In this work, one new species of *Scelidopetalon* is described and illustrated: *Scelidopetalon biwenxuani* sp. n. from Hainan Province, China.

Material and methods

To examine the genitalia, the abdominal segments were detached from the body after softening in hot water. The genitalia, together with other dissected parts, were mounted in Euparal (Chroma Gesellschaft Schmidt, Koengen, Germany) on plastic slides. Photos of sexual characters were taken with a FUJIFILM X10 camera attached to an



Figures 1–2. Head of 1 *Scelidopetalon biwenxuani* and 2 *Amblyopus vittatus* in dorsal view. Scale = 0.5 mm.



Figures 3–4. Habitus of *Scelidopetalon biwenxuani* in dorsal and ventral view. Scale = 2 mm.

***Scelidopetalon biwenxuani* Dai & Zhao, sp. n.**

urn:lsid:zoobank.org:act:B6AA8AE6-86AE-4B7D-A172-CC01ED471933

http://species-id.net/wiki/Scelidopetalon_biwenxuani

Figs 1, 3–4, 5–9

Type material. Holotype: CHINA: Hainan Prov.: 1♀, Jianfengling N.R., Mingfeng-gu Valley, 18°44'N, 108°50'E, alt. 1000 m, 15.V.2011, Bi Wen-Xuan leg. (SNUC).

Description. Body (Fig. 3, 4) oval, convex, shining; length: 6.50 mm; width: 3.79 mm. Body black; legs, palpi and antennae reddish-brown. Elytron black, with red at basal third.

Head (Fig. 1) width between eyes = 5.5 times eye diameter in dorsal view; punctation coarse, separated by 0.5–1.0 puncture diameters laterally and 2–4 puncture diameters medially; stridulatory files not evident. Antennae (Fig. 5) short, not extending behind posterior border of pronotum; antennomere III about 2.0 times as long as IV; antennomeres IV to VIII short; antennomeres IX to XI broad and transverse; relative lengths of antennomeres II–XI: 9.0: 14.0: 7.0: 7.0: 7.0: 7.0: 6.5: 9.0: 9.0: 8.0. Maxillary terminal palpomeres trapezoidal, 1.67 times wider than long. Mentum broad with anterior projection, almost pentagonal, 1.75 times wider than long.

Pronotum transverse, convex above, widest at base (pl/pw = 0.70); anterior angles weakly projecting; lateral margins gently curved toward eyes. Pronotum distinctly punctured medially, finely and closely punctured laterally.

Prosternum (Fig. 7) strongly punctured, the front margin is produced to a short point in the middle, prosternal lines short, not extending in front of procoxal cavities; Mesosternum coarsely punctured. Metasternum coarsely punctured at the sides and almost smooth in the middle, with distinct coxal lines. Abdomen fairly strongly and closely, its sides coarsely, punctured, with distinct coxal lines on first ventrite nearly attaining posterior margin.

Scutellum pentagonal, finely and sparsely punctured.

Legs short, tibiae triangular, with the extremities very broad and hollowed for the tarsi.


Elytra widest at middle, then gradually narrowing to apex; each with 9 punctate lines.

Female genitalia (Fig. 9) with gonostyli fringed apically with some elongate setae; female spermatheca (Fig. 6) with capsule almost egg-shaped.

Distribution. China (Hainan Province).

Diagnosis. The new species can be distinguished from other species in this genus by the black pronotum and red markings of the elytra.

Etymology. This species is named in honor of Mr. Wen-Xuan Bi, collector of the new species.



Figures 5–9. *Scelidopetalon biwenxuani*. **5** antenna **6** spermatheca **7** prosternum **8** tibia and tarsus **9** female genitalia. Scales = 0.5 mm (**5**, **7**, **9**), Scales = 0.1 mm (**6**), Scales = 1.0 mm (**9**).

Acknowledgements

We thank Paul E. Skelley, Florida State Collection of Arthropods, USA and Richard A.B. Leschen, New Zealand for all of their assistance providing methods and information for this study. We thank Wen-xuan, Bi, the collector of this new species for providing specimens for this work. The research was supported by the National Natural Science Foundation of China (No. 31101659 and No. 31172134), the Innovation Program of Shanghai Municipal Education Commission (No. 12YZ077) and by Shanghai Normal University (DZL125).

References

- Araki H (1941) Description of a new genus of Erotylidae (Coleoptera) from Japan proper. Studies on the Erotylidae from Japanese Empire (1). Transactions of the Natural History Society of Formosa 31: 364–366.
- Arrow GJ (1925) Coleoptera. Clavicornia. Erotylidae, Languriidae, and Endomychidae. The Fauna of British India, including Ceylon and Burma. London: Taylor and Francis, xv + 416 pp., 1 pl., 1 map.
- Arrow GJ (1926) Fauna Sumatrensis. Endomychidae and Erotylidae. Entomologische Mitteilungen 15: 248–262.
- Arrow GJ (1945) Systematic notes on a few genera of Erotylid Coleoptera. Proceedings of the Royal Entomological Society of London (B) 14: 117–118.
- Chûjô M, Chûjô M (1990) A Catalogue of the Erotylidae (Coleoptera) from the Old World (excl. the Ethiopian Region) III. Esakia 29: 1–67.
- Delkeskamp H (1957) Neue Erotyliden aus dem Britischen Museum. 21. Beitrag zur Kenntnis der Erotyliden (Coleoptera). Bulletin of the British Museum (Natural History) (Entomology) 6: 79–90.
- Gorham HS (1896) Viaggio di Leonardo Fea in Birmania e regioni vicine. LXIX. Languriidae, Erotylidae and Endomychidae. Annali del Museo Civico di Storia Naturale di Genova 36: 257–302.
- Lewis G (1887) A list of fifty Erotylidae from Japan, including thirty-five new species and four new genera. The Annals and Magazine of Natural History (5) 20: 53–73. doi: 10.1080/00222938709460010
- Lewis G (1889) On a species of *Amblyopus* (Erotylidae) from Japan. The Entomologist's Monthly Magazine 25: 397–398.