

Two new species of the *Stenochinus amplus* species-group from China (Coleoptera, Tenebrionidae, Stenochiini)

Cai-Xia Yuan^{1,2}, Guo-Dong Ren¹

1 College of Life Sciences, Hebei University, Baoding, 071002, P. R. China **2** College of Life Sciences, Yan'an University, Yan'an 716000, P. R. China

Corresponding author: Guo-Dong Ren (gdren@hbu.edu.cn)

Academic editor: C. Majka | Received 22 March 2014 | Accepted 21 May 2014 | Published 17 June 2014

<http://zoobank.org/9A62D665-9B33-481A-97AA-3505854B65D7>

Citation: Yuan C-X, Ren G-D (2014) Two new species of the *Stenochinus amplus* species-group from China (Coleoptera, Tenebrionidae, Stenochiini). ZooKeys 416: 67–76. doi: 10.3897/zookeys.416.7568

Abstract

Two new species of the *Stenochinus amplus* species-group are described, *S. apiciconcavus* **sp. n.** (CHINA: Shaanxi) and *S. xinyicus* **sp. n.** (CHINA: Guangdong). Also, some new distribution data are provided for *S. cylindricus* (Gebien, 1914), and a key to the seven species of the *S. amplus* species-group from China is given.

Keywords

Tenebrionidae, Stenochiini, *Stenochinus*, new species, China

Introduction

The genus *Stenochinus* was established by Motschulsky (1860) for *S. reticulatus*. Contributions to the taxonomy of this genus have been made by several workers, such as Marseul (1876), Fairmaire (1896), Gebien (1911, 1914, 1925), Gravely (1915), Pic (1921, 1922, 1923, 1926, 1927, 1958), Kulzer (1954), Nakane (1956, 1985), Ardoin (1969), Kaszab (1978, 1980a, 1980b), Shibata (1980), Merkl (1992), Masumoto and Akita (2002), Ren and Yang (2004), Schawaller (2006), Löbl et al. (2008) and Masumoto et al. (2013), this group currently includes 43 species and subspecies, which are mainly distributed in the Oriental Region. The species of Taiwan and Japan were divided into three species-groups by Shibata (1980), one of them, the *Stenochinus amplus* species-group has so far 12 species

(Shibata 1980, Masumoto and Akita 2002, Ren and Yang 2004, Schawaller 2006, Masumoto et al. 2013). In the present study, two new species of this group from China are described. Besides, a key to the Chinese species of the *Stenochinus amplus* species-group is provided, including new distribution data for *S. cylindricus* (Gebien, 1914).

Material and methods

Specimens were examined under a Nikon (SMZ800) dissecting microscope. Measurements were taken and photographs captured using a Leica (M205 A) dissecting microscope. The habitus photos were taken with a Canon (EOS 5D mark II) camera.

The measurements were as follows: body length: length of the body from the anterior edge of the clypeus to elytral apex with the head in its natural position; body width: length of the maximal elytral width; pronotal length: length of the pronotum along the midline; elytral length: length of the elytra from the base of the scutellum to the elytral apex along the suture. All measurements are given in millimeters.

The following codens of the collections are used:

MHBU Museum of Hebei University, Baoding, China;

SYSU Research Institute of Entomology, Sun Yat-Sen University, Guangzhou, China;

IZCAS Institute of Zoology, Chinese Academy of Sciences, Beijing, China.

Taxonomy

Stenochinus amplus species-group

Diagnosis. Body covered with distinct scale-like hairs, mandibles subtruncate, apical projections of pronotum not or hardly reaching anterior margins of eyes, tibiae with rows of suberect hairs along inner margins, male genitalia widened basally.

Distribution. China (including Taiwan), Japan, Vietnam, India, Nepal, Sri Lanka.

Key to the species of *Stenochinus amplus* species-group in China

- | | | |
|---|--|---|
| 1 | Elytra distinctly convex, oval. Brachypterous species..... | 2 |
| – | Elytra weakly convex, subcylindrical (Figs 17–20). Winged species | 4 |
| 2 | Anterior margin of pronotum distinctly emarginate..... | 3 |
| – | Anterior margin of pronotum without emargination | |
| | <i>S. mysticus</i> Masumoto, Akita & Lee | |
| 3 | Body slender, gradually widened in basal one-fourth part of elytra..... | |
| | <i>S. akiyamai</i> Masumoto, Akita & Lee | |
| – | Body stouter, subparallel-sided in basal one-fourth part of elytra | |
| | <i>S. amplus</i> (Gebien) | |

- 4 Pronotum moderately projecting forward (Figs 1–3, 5) 5
 – Pronotum strongly projecting forward (Figs 4, 6) ***S. cylindricus* (Gebien)**
 5 Anterior margin of pronotum without emargination (Figs 2–4). Scutellum subquadrate 6
 – Anterior margin of pronotum distinctly emarginate (Fig 1). Scutellum subrectangular ***S. apiciconcavus* sp. n.**
 6 Pronotum weakly reflexed at anterior margin, scale-like hairs on pronotum longer and wider than those on elytra ***S. xinyicus* sp. n.**
 – Pronotum not reflexed at anterior margin, scale-like hairs on pronotum nearly same as on elytra ***S. leprosus* (Ren & Yang)**

***Stenochinus apiciconcavus* sp. n.**

<http://zoobank.org/3BD90C95-FA8E-4919-AF90-196DA9F0CF1E>


Figs 1, 7, 11–12, 17

Type material. Holotype: ♂, China, Shaanxi, Nanzheng county, Beiba town, 20.vi.2005, Yi-Bin Ba leg. (MHBUS).

Diagnosis. This new species is similar to *S. cylindricus* (Gebien, 1914), but can be distinguished from the latter by the pronotum moderately projecting anteriorly, with emargination in middle of anterior margin; scutellum subrectangular; maxillary palpomere IV strongly expanded (in *S. cylindricus*, pronotum strongly projecting anteriorly, without emargination at anterior margin (Figs 4, 6); scutellum subquadrate; maxillary palpomere IV moderately expanded). It also resembles *S. wakoi* Masumoto & Akita, 2002, but differs in the following characters: antennomeres VIII–XI wider than long; pronotum with emargination in middle of anterior margin; scutellum subrectangular, glabrous (in *S. wakoi*, antennomeres VI–XI wider than long; pronotum without emargination at anterior margin; scutellum subquadrate, covered with scale-like hairs on both sides).

Etymology. The specific name is derived from the Latin *apici* [apex] + *concauus* [concave], a reference to the pronotum with an emargination in middle of anterior margin.

Description. Male (Fig. 17). Body length 10.3 mm, elongate, subcylindrical. Head, elytra and legs dark reddish brown, antennae reddish brown, pronotum dark brown. Scale-like hairs on the surface pale golden. Head transversely subelliptical, surface densely punctate; clypeus truncate at anterior margin, clypeogenal suture grooved, frontoclypeal suture invisible; genae weakly raised, with rounded outer margins; frons wide, distance between eyes 2.56 times as wide as transverse diameter of an eye in dorsal view. Eyes medium-sized, weakly protruding, each side with a groove along inner and posterior margins. Antennae (Fig. 7) clavate, antennomeres VIII–XI each wider than long, antennomeres XI ovate, ratio of the length of antennomeres II–XI as 0.12: 0.18: 0.09: 0.09: 0.07: 0.08: 0.07: 0.10: 0.08: 0.22. Maxillary palpomere IV strongly expanded. Pronotum (Fig. 1) 1.13 times as long as wide, widest in middle; anterior margin with a distinct emargination in middle; posterior margin weakly bisinuate, with deep emar-



Figures 1–6. Pronotum. 1 *S. apiconcavus* sp. n. 2–3, 5 *S. xinyicus* sp. n. (2, 5 ♂ 3 ♀) 4, 6 *S. cylindricus*. Scale bars = 0.5 mm.

gination in middle; both sides steeply inclined downward, lateral margins sinuate before posterior angles; anterior angles acute and directed anteriorad, posterior angles acute and directed postero-laterad; disc subelliptically projecting in anterior parts, this projecting

area distinctly impressed, surface roughly and deeply punctate, punctures often fused with one another. Scutellum subrectangular, glabrous. Elytra 2.5 times as long as wide, widest at apical 1/3, 2.76 times as long as and 1.25 times as wide as pronotum; dorsum convex but flattened in lateral view; disc with rows of subquadrate punctures, which are larger and deeper anteriorly, each puncture with a granule on each lateral margin; intervals somewhat transversely wrinkled, weakly ridged in lateral parts, scale-like hairs on the intervals shorter than those on pronotum. Ventral side covered with dense punctures and scale-like hairs, which distinctly shorter than those on pronotum. Legs relatively short, ratio of the lengths of metatarsomeres I–IV as 0.45: 0.24: 0.17: 0.69. Male genitalia (Figs 11–12) curved in middle in lateral view, 1.94 mm long, 0.43 mm wide; apicale 0.71 mm long, weakly curved in lateral view.

Female: Unknown.

***Stenochinus xinyicus* sp. n.**

<http://zoobank.org/D193ED7A-DE68-42F1-9BDE-F748463B88A3>


Figs 2–3, 5, 8–9, 13–14, 18–19

Type material. Holotype: ♂, China, Guangdong, Xinyi City, Dawuling, 1.vi.2002, Cheng Chen leg. (MHBUS). Paratypes: 3 ♀♀, same locality as holotype, 31.v.2002, Zhong-Ping Deng leg.; 1.vi.2002, Cui-Feng Li leg.; 6.vii.1988, Feng-Long Jia leg. (MHBUS); 2 ♀♀, same locality as holotype, 31.v.2002, Bing-Lan Zhang leg.; 7.vii.1988, He-Xiang Zhou leg. (SYSU).

Diagnosis. This new species is similar to *S. cylindricus*, but can be distinguished from the latter by antennomeres VII–XI wider than long, pronotum moderately projecting anteriorly (Fig. 5), shape of male genitalia different (Figs 13–16) (in *S. cylindricus*, antennomeres VIII–XI wider than long (Fig. 10), pronotum strongly projecting anteriorly (Fig. 6)).

Etymology. The specific name is derived from the type locality of this species.

Description. Male (Fig. 18). Body length 11.0 mm, elongate, subcylindrical. Head, elytra, legs reddish brown, pronotum brown, antennae and mouthparts yellowish brown; scale-like hairs surface pale golden. Head transversely subelliptical, surface densely punctate; clypeus transverse, weakly convex in middle, clypeogenal suture grooved, frontoclypeal suture invisible; genae weakly raised; frons wide, distance between eyes 2.53 times as wide as the transverse diameter of an eye in dorsal view. Eyes medium-sized, weakly protruding, with groove along inner and posterior margins. Antennae (Fig. 8) clavate, antennomeres VII–XI wider than long, antennomeres XI oval, length ratio of antennomeres II–XI as 0.16: 0.17: 0.12: 0.10: 0.09: 0.09: 0.10: 0.13: 0.10: 0.23. Maxillary palpomere IV moderately expanded. Pronotum (Fig. 2) 1.14 times as long as wide, widest shortly before middle; anterior margin weakly reflexed; posterior margin weakly bisinuate, with deep emargination in middle; both sides steeply inclined downward, lateral margins sinuate before posterior angles; anterior angles acute and directed anteriorly, posterior angles obtuse and



Figures 7–16. 7–10 Antennae 11–16 Male genitalia in dorsal view and lateral view 7, 11–12 *S. apiciconcavus* sp. n. 8–9, 13–14 *S. xinyicus* sp. n. (8 ♂ 9 ♀) 10, 15–16 *S. cylindricus*. Scale bars = 0.5 mm

directed posteriorly; disc subelliptically projected in anterior parts, projecting area weakly impressed, surface roughly and deeply punctate, punctures often fused with another. Scutellum subquadrate, glabrous. Elytra 2.51 times as long as wide, widest at apical 1/3, 2.80 times as long as and 1.11 times as wide as pronotum; dorsum convex but flattened in lateral view; disc with rows of large, deep and subquadrate



Figures 17–20. Habitus. 17 *S. apiconcavus* sp. n. ♂ 18–19 *S. xinyicus* sp. n. (18 ♂ 19 ♀) 20 *S. cylindricus* ♂.

punctures, each puncture with a granule on each lateral margin; intervals somewhat transversely wrinkled, scale-like hairs shorter and narrower than those on pronotum. Ventral side covered with dense punctures and scale-like hairs, which shorter than those on pronotum. Legs relatively short, length ratio of metatarsomeres I–IV as 0.40: 0.33: 0.21: 0.67. Male genitalia (Figs 13–14) strongly curved in middle in lateral view, 2.29 mm long, 0.43 mm wide; apicale 0.77 mm long, weakly curved in lateral view.

Female (Fig. 19): Body 11.0–12.5 mm, dark brown. Antennae (Fig. 9) thicker, length ratio of antennomeres II–XI as 0.12: 0.14: 0.11: 0.10: 0.07: 0.08: 0.09: 0.12: 0.11: 0.26. Pronotum (Fig. 3) 1.10 times as long as wide. Elytra 2.42 times as long as wide.

***Stenochinus cylindricus* (Gebien, 1914)**

Figs 4, 6, 10, 15–16, 20

Dicraeosia cylindricus Gebien, 1914: 9.

Material examined. 2 ♂♂, 5 unsexed, China, Taiwan, Hualien county, Xiulin township, Tailuge, 60–500 m, 13.v.2010, Wen-Yi Zhou leg. (MHBUS); 1 unsexed, China, Taiwan, Taitung city, Lianren town, Shouka, 450 m, 16.vi.2010, Wen-Yi Zhou leg. (MHBUS).

Distribution. South China: Taiwan.

***Stenochinus leprosus* (Ren & Yang, 2004)**

Dicraeosia leprosus Ren & Yang, 2004: 317.

Type material examined. Holotype: ♂, China, Guangxi, Fangcheng, Fulong, 500 m, 24.v.1999, Da-Jun Liu leg. (IZCAS).

Distribution. South China: Guangxi.

Acknowledgments

We thank very much Dr. Wolfgang Schawaller (Staatliches Museum für Naturkunde, Stuttgart), Dr. Kimio Masumoto (Tokyo, Japan) and Dr. Ottó Merkl (Hungarian Natural History Museum, Budapest) for helpful comments on this manuscript, also, Dr. O. Merkl provided the important paper. Many thanks are also due to Dr. Yu-Xia Yang (College of Life Sciences, Hebei University), Dr. Ji-Shan Xu (College of Agriculture and Biology, Dali University) and Dr. J. MacDermott (USA) for kindly helping to review the manuscript. Two anonymous reviewers provided helpful comments on the manuscript. This work is financially supported by the National Natural Foundation of China (No. 31093430) and the High-level university construction projects of Shaanxi Province (No. 2012SXTS03).

References

- Ardoin P (1969) Contributions à la connaissance de la faune entomologique de la Côte d'Ivoire (J. Decelle, 1961–1964) XXVII – Coleoptera Tenebrionidae. *Annales du Musée Royal de l'Afrique Centrale, Sciences Zoologiques* 175: 139–285.
- Fairmaire L (1896) Hétéromères de l'Inde recueillis par M. Andrewes. *Annales de la Société Entomologique de Belgique* 40: 6–62.
- Gebien H (1911) Tenebrionidae Ill, IV. Pars 28, 37. In: Schenkling S (Ed) *Coleopterorum Catalogus*. Vol. 18. W. Junk, Berlin, 355–740.
- Gebien H (1914) H. Sauter's Formosa-Ausbeute. Tenebrionidae (Coleopt.). *Archiv für Naturgeschichte A* 79(9) [1913]: 1–60.
- Gravely FH (1915) XLII. Coleoptera, IX: Tenebrionidae. In: Zoological results of the Abor Expedition, 1911–12. XLII. Coleoptera IX: Tenebrionidae. *Records of the Indian Museum* 8 [1912–1922]: 519–536.
- Kaszab Z (1978) Vier neue Tenebrioniden aus Nordbengal (Coleoptera). *Folia Entomologica Hungarica* (N. S.) 31(2): 173–176.
- Kaszab Z (1980a) Neue Tenebrioniden (Coleoptera) aus Sri Lanka. 1. *Acta Zoologica Academiae Scientiarum Hungaricae* 26(1–3): 123–196.
- Kaszab Z (1980b) Angaben zur Kenntnis der Tenebrioniden Nordvietnams (Coleoptera). *Annales Historico-Naturales Musei Nationalis Hungarici* 72: 169–221.
- Gebien H (1925) Die Tenebrioniden (Coleoptera) des indomalayischen Gebietes, unter Berücksichtigung der benachbarten Faunen, III. *The Philippine Journal of Science*, Manila 26(4): 535–577.
- Kulzer H (1954) Achter Beitrag zur Kenntnis der Tenebrioniden (Col.). *Entomologische Arbeiten aus dem Museum G. Frey* 5: 20–73.
- Löbl I, Merkl O, Ando K, Bouchard P, Lillig M, Masumoto K, Schawaller W (2008) Tenebrionidae. In: Löbl I, Smetana A (Eds) *Catalogue of Palaearctic Coleoptera*. Vol. 5. Apollo Books, Stenstrup, 105–352.
- Marseul SA de (1876) Coléoptères du Japon recueillis par M. Georges Lewis. Énumération des Hétéromères avec la description des espèces nouvelles. *Annales de la Société Entomologique de France* (5)6: 93–142.
- Masumoto K, Akita K (2002) New or little-known tenebrionid species from Japan (Part 3). *Entomological Review of Japan* 57(2): 147–153.
- Masumoto K, Akita K, Lee CF (2013) New tenebrionid (Coleoptera) beetles from Taiwan (8) Descriptions of six new species and new occurrence records of four species from Taiwan. *Elytra*, Tokyo, New Series 3(2): 255–273.
- Merkl O (1992) Tenebrionidae (Coleoptera) from Laos and Vietnam, with reclassification of Old World “*Doliema*”. *Acta Zoologica Academiae Scientiarum Hungaricae* 38(3–4): 261–280.
- Motschulsky V de (1860) Insectes des Indes orientales, et de contrées analogues. *Études Entomologiques* 8[1859]: 25–118.
- Nakane T (1956) New or little-known Coleoptera from Japan and its adjacent regions. XIII. *The Science Reports of the Saikyo University* (A) 2(3): 164–174.

- Nakane T (1985) New or little-known Coleoptera from Japan and its adjacent regions. XXXVIII. *Fragmenta Coleopterologica* 38–40: 162–164.
- Pic M (1921) Nouveautés diverses. *Mélanges Exotico-Entomologiques* 34: 1–33.
- Pic M (1922) Coléoptères exotiques en partie nouveaux. *L'Échange, Revue Linnéenne* 38: 31–32.
- Pic M (1923) Nouveautés diverses. *Mélanges Exotico-Entomologiques* 39: 1–32.
- Pic M (1926) Nouveautés diverses. *Mélanges Exotico-Entomologiques* 45: 1–32.
- Pic M (1927) Coléoptères de L'Indochine. *Mélanges Exotico-Entomologiques* 49: 1–36.
- Pic M (1958) La réserve naturelle intégrale du Mont Nimba, fascicule IV. *Coleopteres Lycides*. VIII. *Coleopteres nouveaux* (Ptinidae, Tenebrionidae, Oedemeridae, Anthicidae, Meloidae, Lampyridae, Cleridae, Helodidae, Chrysomelidae). *Mémoire de l'Institut Français d'Afrique Noire* 53: 199–200.
- Ren GD, Yang XJ (2004) *Coleoptera: Tenebrionidae*. In: Yang XK (Ed) *Insects from Mt. Shiwandashan Area of Guangxi*. China Forestry Publishing House, Beijing, 668 pp. [in Chinese, English title]
- Schawaller W (2006) New species, new records and new synonyms of Tenebrionidae (Insecta: Coleoptera) from Nepal. *Veröffentlichungen des Naturkundemuseums Erfurt* 25: 475–482.
- Shibata T (1980) Notes on Tenebrionidae from Taiwan and Japan, III. (Coleoptera). *Entomological Review of Japan* 34(1–2): 63–74.