

A new species and new records of the leafhopper genus *Taperus* Li & Wang, 1994 (Hemiptera, Cicadellidae, Evacanthinae) from China

Yu-jian Li^{1,2,†}, Zi-zhong Li^{1,2,‡}

1 Institute of Entomology, Guizhou University, Guiyang, Guizhou Province 550025, China **2** The Provincial Key Laboratory for Agricultural Pest Management of Mountainous Region, Guizhou University, Guiyang, Guizhou Province 550025, China

† [urn:lsid:zoobank.org:author:533BDC07-7555-4C51-916C-FB76C091FC9B](https://doi.org/urn:lsid:zoobank.org:author:533BDC07-7555-4C51-916C-FB76C091FC9B)

‡ [urn:lsid:zoobank.org:author:9BA8A6EF-F7C3-41F8-AD7D-485FB93859F2](https://doi.org/urn:lsid:zoobank.org:author:9BA8A6EF-F7C3-41F8-AD7D-485FB93859F2)

Corresponding author: Zi-zhong Li (yujian528@163.com, lizizhong38@163.com)

Academic editor: Mike Wilson | Received 9 April 2011 | Accepted 19 May 2011 | Published 25 July 2011

[urn:lsid:zoobank.org:pub:66EF85AD-9939-4AC4-9DCC-754D5BA4306B](https://doi.org/urn:lsid:zoobank.org:pub:66EF85AD-9939-4AC4-9DCC-754D5BA4306B)

Citation: Li Y, Li Z (2011) A new species and new records of the leafhopper genus *Taperus* Li & Wang, 1994 (Hemiptera, Cicadellidae, Evacanthinae) from China. ZooKeys 120: 1–8. doi: 10.3897/zookeys.120.1373

Abstract

The paper deals with the species of the Oriental leafhopper genus *Taperus* Li & Wang. A new species, *T. daozenensis* **sp. n.**, from Guizhou Province, China is described and new records for other Chinese species are given together with a key for their separation. The type specimens of the new species are deposited in the Institute of Entomology, Guizhou University (GUGC).

Keywords

Auchenorrhyncha, morphology, identification, China

Introduction

The leafhopper genus *Taperus* was established by Li and Wang in 1994 with *Taperus fasciatus* Li & Wang as its type species, at same time Li and Wang described two more species, *T. albivittatus* and *T. apicalis*. Subsequently, three new species were described in the genus by Cai and Shen (1997, 1999) and Li and Wang (2001). Zhang et al. (2010)

reviewed the genus and described three new species from China in addition to transferring one species from *Onukia* to *Taperus* and three species from *Taperus* to the genus *Convexana* Li.

A new species from Guizhou Province, China is here described and illustrated. A key to all known species of *Taperus* is given and new records for other Chinese species of the genus are provided. Currently eight species are included in the genus.

The specimens used in this study are deposited in the Institute of Entomology, Guizhou University, Guiyang, China (GUGC).

Taxonomy

Taperus Li & Wang

<http://species-id.net/wiki/Taperus>

Taperus Li and Wang 1994: 374

Type species. *Taperus fasciatus* Li & Wang.

Diagnosis. *Taperus* is similar to *Onukia* Matsumura, it differs from this genus in having the median longitudinal carina of vertex very weak, nearly indistinct; area between median carina and submarginal carina of vertex nearly flat rather than concave and male pygofer with rows or groups of stout spine-like macrosetae marginally.

For detailed generic description see Zhang et al. (2010).

Key to species of the genus *Taperus* Li & Wang based on male adults

- 1 Aedeagal shaft with a triangular posterior process in lateral view; foot-like apical process of style with apex very long and slender 2
- Aedeagal shaft not or slightly developed posteriorly near base in lateral view; foot-like apical process of style with apex short and robust..... 4
- 2 Male pygofer longer than wide, with apical half distinctly narrowed in lateral view 3
- Male pygofer nearly discoidal in lateral view *T. apicalis* Li & Wang
- 3 Male pygofer with a row of long spine-like macrosetae on dorsal margin *T. bannaensis* Zhang, Zhang & Wei
- Male pygofer with short thick macrosetae at apex *T. fasciatus* Li & Wang
- 4 Male pygofer with apex truncate in lateral view 5
- Male pygofer distinctly narrowed apically with apex rounded in lateral view...6
- 5 Male pygofer somewhat rectangular in lateral view; style elongate..... *T. flavifrons* (Matsumura)
- Male pygofer nearly quadrangular in lateral view; style short *T. quadragulatus* Zhang, Zhang & Wei
- 6 Apex of aedeagus with processes..... 7

- Apex of aedeagus without processes
..... *T. luchunensis* Zhang, Zhang & Wei
- 7 Apex of aedeagus with three spine-like processes on dorsal margin and four pairs of spine-like process on ventral margin *T. albivittatus* Li & Wang
- Apex of aedeagus with two retrorse spine-like processes on dorsal margin and one big spine-like process on dorsal margin subapically (Figs 4, 5).....
..... *T. daozenensis* sp. n.

***Taperus albivittatus* Li & Wang**

http://species-id.net/wiki/Taperus_albivittatus

Taperus albivittatus Li and Wang, 1994: 375; 1996: 115; Zhang et al. 2010: 39

Distribution. China (Sichuan).

Hosts. Unknown.

Material examined. 1♂ (Holotype): CHINA, Sichuan: Emeishan, Wanniansi, 3 August 1991, coll. Li Zizhong.

Notice. After reexamining the holotype of this species, we found the count of spine-like processes on the apex of the aedeagus was wrong in the original description, we corrected it in above keys.

***Taperus apicalis* Li & Wang**

http://species-id.net/wiki/Taperus_apicalis

Taperus apicalis Li and Wang, 1994: 377; 1996: 116; Zhang et al. 2010: 39

Distribution. China (Guizhou).

Hosts. Unknown.

Material examined. 1♂ (Holotype): CHINA, Guizhou: Shuicheng, 30 September 1987, coll. Li Zizhong.

***Taperus fasciatus* Li & Wang**

http://species-id.net/wiki/Taperus_fasciatus

Taperus fasciatus Li and Wang, 1994: 378; 1996: 117; Zhang et al. 2010: 39

Distribution. China (Guizhou, Hainan, Guangxi, Sichuan, Fujian, Shaanxi, Hunan, Zhejiang, Jiangxi); Vietnam.

Hosts. Unknown.

Material examined. 1♂ (Holotype): CHINA, Guizhou: Daozhen, 18 September 1988, coll. Li Zizhong; 20♂♂, 17♀♀: CHINA, Guizhou: Leigongshan, Xiaodanjiang, 13–14 September 2005, coll. Li Zizhong & Zhang Bin; 6♂♂7♀♀: CHINA, Guizhou: Kuankuoshui, Chachang, 10–17 August 2010, coll. Dai Renhuai, Li Hu & Fan Zhihua; 16♂♂7♀♀: CHINA, Guizhou: Kuankuoshui, Chachang, 14–17 August 2010, coll. Yu Xiaofei; 3♂♂: CHINA, Guizhou: Kuankuoshui, Chachang, 12 August 2010, coll. Li Yujian; 2♂♂: CHINA, Hainan: Jianfengling, 10–12 July 2007, coll. Li Yujian; 2♂♂: CHINA, Guangxi: Huaping, 11 June 1994, coll. Du Yuzhou; 1♂: CHINA, Zhejiang: Longquan, 17 June 1980, coll. Tong Xuesong; 3♂♂: CHINA, Sichuan: Baishuihe, 27 August 2007, coll. Xing Jichun.

***Taperus flavifrons* (Matsumura)**

http://species-id.net/wiki/Taperus_flavifrons

Onukia flavifrons Matsumura, 1912: 45

Taperus flavifrons (Matsumura), Zhang et al. 2010: 39

Distribution. China (Hainan, Taiwan); Japan.

Hosts. Unknown.

Material examined. 2♂♂: CHINA, Hainan: Wuzhishan, 13–15 July 2007, coll. Zhang Bin; 1♂: CHINA, Hainan: Wuzhishan, 13–15 July 2007, coll. Li Yujian; 1♂: CHINA, Hainan: Diaoluoshan, 16–18 July 2007, coll. Zhang Bin; 1♀: CHINA, Hainan: Wuzhishan, 13–15 July 2007, coll. Zhang Hui; 1♀: CHINA, Hainan: Wuzhishan, 13–15 July 2007, coll. Song Yuehua; 1♀: CHINA, Hainan: Diaoluoshan, 16–18 July 2007, coll. Zhang Hui; 1♀: CHINA, Hainan: Diaoluoshan, 16–18 July 2007, coll. Zhang Bin.

***Taperus bannaensis* Zhang, Zhang & Wei**

http://species-id.net/wiki/Taperus_bannaensis

Taperus bannaensis Zhang et al. 2010: 44

Distribution. China (Yunnan, Hainan).

Hosts. Unknown.

Material examined. 1♂: CHINA, Hainan: Limuling, 23 May 1997, coll. Yang Maofa.

***Taperus quadragulatus* Zhang, Zhang & Wei**

http://species-id.net/wiki/Taperus_quadragulatus

Taperus quadragulatus Zhang et al. 2010: 44

Distribution. China (Hunan, Guizhou).

Hosts. Unknown.

***Taperus luchunensis* Zhang, Zhang & Wei**

http://species-id.net/wiki/Taperus_luchunensis

Taperus luchunensis Zhang et al. 2010: 45

Distribution. China (Guizhou, Hubei, Yunnan).

Hosts. Unknown.

Material examined. 4♂♂, 5♀♀: CHINA, Guizhou: Leigongshan, Xiaodanjiang, 13-14 September 2005, coll. Li Zizhong & Zhang Bin; 1♂: CHINA, Hubei: Shennongjia, 13 August 2004, coll. Peng Jingyang.

***Taperus daozenensis* sp. n.**

urn:lsid:zoobank.org:act:35578DE4-6CF7-4389-B007-F306E375AE90

http://species-id.net/wiki/Taperus_daozenensis

Figs 1–7

Measurements. ♂: body length: 6.0 mm; head width (incl. eyes): 1.1 mm; head length: 0.8 mm.

Vertex, pronotum and scutellum dark brown; pale median longitudinal band yellow, extending from apex of vertex to posterior margin of pronotum (Figs 1, 2). Face nearly pale yellow (Figs 1, 2). Forewing (Figs 1, 2) dark brown, with nearly white and transparent plaque in middle of costal area.

Male pygofer (Fig. 3) nearly triangular in lateral view, row of long macrosetae at end of dorsal and ventral margin. Apex of foot-like apical process of style very long and slender, about one third length of style (Fig. 6). Aedeagal shaft, recurved dorsally, expanded distally in lateral view, broadly rounded apically with pair of retrorse spine-like processes on dorsal margin subapically and one big spine-like process on dorsal margin near base (Figs 4, 5).

Other characteristics are as shown in Figs 1–7.

Type Material. Holotype: ♂, China: Guizhou: Daozhen, 17–22 August 2004, coll. Yang Maofa; Paratypes: 2♂♂, China: Guizhou: Daozhen, 17–22 August 2004, coll. Yang Maofa.


Etymology. The species is named after the locality of the type specimens, Daozhen county.

Remarks. This species resembles *T. albivittatus* Li & Wang, from which it is distinguished by: aedeagus with pair of subapical retrorse spine-like processes dorsally and



Figures 1–2. *Taperus daozhenensis* sp. n. 1–2 Male, dorsal view and lateral view.

one big spine-like process on dorsal margin near base. Apex of aedeagus of *T. albivittatus* has three spine-like processes on dorsal margin and four pairs of spine-like process on ventral margin.



Figures 3–7. *Taperus Daozhenensis* sp. n. **3** Male pygofer, lateral view **4** Aedeagus, lateral view **5** Aedeagus, ventral view **6** style and connective **7** Subgenital plate.

Acknowledgements

We thank Professor Yang Maofa, Guizhou University, for his gift of the holotype. We also thank Dr. Mike Wilson (National Museum Wales, Subject editor of Zookeys), Mr. M. D. Webb (Natural History Museum, London) and anonymous reviewer for reading the manuscript and making some suggestions. This study is supported by the National Natural Science Foundation of China (Grant No. 31093430) and the Ministry of Science and Technology of the People's Republic of China (MOST grant No. 2006FY110500).

References

- Cai P, Shen XC (1997) Four new species of Evacanthinae (Homoptera: Cicadellidae) from China. *Entomotaxonomia* 19: 246–251.
- Cai P, Shen XC (1999) Nine leafhopper new species of Baotianman. In: Shen, Pei (Ed) *Fauna of insects Henan Province of China Vol. 4*. China Agriculture Press, Beijing, 23–25 pp.
- Dietrich CH (2004) Phylogeny of the leafhopper subfamily Evacanthinae with a review of Neotropical species and notes on related groups (Hemiptera: Membracoidea: Cicadellidae). *Systematic Entomology* 29: 455–487. doi:10.1111/j.0307-6970.2004.00250.x
- Li ZZ (1994) A new genus and three new species of Evacanthini from China (Homoptera: Cicadellidae). *Acta Zootaxonomica Sinica* 19: 465–470.
- Li ZZ, Wang LM (1994) A new genus and three new species of the tribe Evacanthini (Insecta: Homoptera: Cicadellidae) with a key to the genera and a list of species occurring in China. *Journal of Natural History* 28: 373–382. doi:10.1080/00222939400770161
- Li ZZ, Wang LM (2001) Three new species of Evacanthinae from Yunnan, China (Homoptera: Cicadellidae). *Zoological Research* 22: 387–391.
- Huang KW (1992) Taxonomy of Evacanthini of Taiwan (Homoptera: Cicadellidae: Cicadellinae). *Bulletin of National Museum of Natural Science* 3: 159–185.
- Matsumura S (1912) *Die Cicadinen Japans II*. *Annotationes Zoologicae Japonenses* 8: 15–51.
- Zhang YL (1990) *A Taxonomic Study of Chinese Cicadellidae (Homoptera)*. Tianze Eldonejo, Yangling, 218 pp.
- Zhang XM, Zhang YL, Wei C (2010) Review of the leafhopper genus *Taperus* Li & Wang (Hemiptera: Cicadellidae: Evacanthinae) from China, with description of three new species. *Zootaxa* 2721: 39–46.