

# A new species of *Corydalus* Latreille from Venezuela (Megaloptera, Corydalidae)

Atilano Contreras-Ramos<sup>1†</sup>, Klaus von der Dunk<sup>2‡</sup>

**1** Instituto de Biología, UNAM, Depto. de Zoología, Apdo. Postal 70-153, 04510 México, D.F., México

**2** Kreis Nürnberger Entomologen, Ringstraße 62, 91334 Hemhofen, Germany

† [urn:lsid:zoobank.org:author:1EC6004F-2261-415D-AC90-46D1B7344255](https://doi.org/urn:lsid:zoobank.org:author:1EC6004F-2261-415D-AC90-46D1B7344255)

‡ [urn:lsid:zoobank.org:author:DDE65F45-DFDC-4B13-A38F-CC0F9B9DCE89](https://doi.org/urn:lsid:zoobank.org:author:DDE65F45-DFDC-4B13-A38F-CC0F9B9DCE89)

Corresponding author: Atilano Contreras-Ramos ([acontreras@ibiologia.unam.mx](mailto:acontreras@ibiologia.unam.mx))

---

Academic editor: Terry Erwin | Received 26 June 2010 | Accepted 25 October 2010 | Published 10 November 2010

---

[urn:lsid:zoobank.org:pub:DC0630A9-00B2-4B9A-AE48-CD06697586B2](https://doi.org/urn:lsid:zoobank.org:pub:DC0630A9-00B2-4B9A-AE48-CD06697586B2)

---

**Citation:** Contreras-Ramos A, Dunk K von der (2010) A new species of *Corydalus* Latreille from Venezuela (Megaloptera, Corydalidae). ZooKeys 67: 11–19. doi: 10.3897/zookeys.67.702

---

## Abstract

A new species of dobsonfly, *Corydalus wanningeri*, from Venezuela, is described and illustrated. It superficially resembles *C. neblinensis* Contreras-Ramos, with a uniform reddish coloration of body and wings. Yet, because of male genitalic structure it might be closely related to *C. crossi* Contreras-Ramos. Specimens were collected from a rain forest transitional zone between the Orinoco lowlands and the Gran Sabana plateau, in Bolívar state. This is the 15<sup>th</sup> species of *Corydalus* to be recorded from Venezuela, rendering this the country with the highest number of documented *Corydalus* species. A key to the sexually dimorphic, long-mandibled Venezuelan species of the genus is provided.

## Keywords

Dobsonfly, taxonomy, biodiversity, South America, key

## Introduction

The dobsonfly genus *Corydalus* Latreille was revised nearly a decade ago (Contreras-Ramos 1998), with 30 species recognized. Four species from Venezuela were later add-

ed to the genus (Contreras-Ramos 2002), and a 35<sup>th</sup> species is herein described, also from Venezuela. *Corydalus* is the most species rich genus of New World dobsonflies, a monophyletic lineage also including *Chloronia* Banks and *Platyneuromus* Weele. Most *Corydalus* species, 27 (77%), are South American only, and three are South and Central American. Fifteen species (43%) have been recorded from Venezuela, of which seven (20%), or possibly eight, are known only from this country (Table 1). Brazil rates second, with 11 species recorded (31%), of which only three (9%), or possibly four, are uniquely recorded for the country. These numbers indicate a pattern of high species richness and strong endemism for dobsonflies in Venezuela.

Late last year the authors established contact. From images of a collection of Neotropical dobsonflies, a series of *Corydalus* specimens stood out as potentially new, a supposition corroborated upon specimen examination. Specimens of the new species had been collected by Professor Rupert Wanninger of Bavaria, where he is now a retired teacher of arts and sports. Prof. Wanninger is also a recognized amateur entomologist, deeply versed in Natural History, with extensive experience in breeding of exotic beetles. For years, he has motivated young people in an interest and respect for Nature. This contribution is in homage to Prof. Wanninger's lifetime as an educator and insect lover.

## Methods

Isolated single specimens from a single site were collected by Prof. Wanninger, amounting to about 40 collections over a more than a 10 year span. All specimens were collected using mercury vapor light. The collecting site is located adjacent to the NE limit of Parque Nacional Canaima, in a portion of a winding road known as La Escalera (Spanish for ladder), highway 10, between Piedra de la Virgen and Danto Falls, around Km 110–112, at 1,000 m of elevation. A large communications antenna is a landmark for the collecting site. Highway 10 connects the Orinoco lowlands with the Gran Sabana plateau in the south. La Escalera is a humid slope covered with rain forest, with several brooks and waterfalls, potential habitat for the hellgrammites. Collections by Prof. Wanninger from nearby sites at higher and lower elevations did not produce any more specimens of the new species. Specimens were dissected and observed using standard techniques (Contreras-Ramos 1998).

Specimens will be deposited at Colección Nacional de Insectos, Instituto de Biología, UNAM, Mexico City (CNIN-UNAM), Museo del Instituto de Zoología Agrícola, Universidad Central de Venezuela, Maracay (MIZA), Zoologische Staatssammlung München, Bavaria, Germany (ZSM), and Prof. Wanninger's private collection (RW). The identification key herein provided applies to species with males whose mandibles are elongate, with reduced dentition. Species with males having short, female-like mandibles may be identified with Contreras-Ramos (2002).

**Table 1.** Species of *Corydalus* Latreille recorded from Venezuela (Contreras-Ramos 1999, 2005).

Species	Distribution
<i>Corydalus affinis</i> Burmeister, 1839	Argentina, Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Paraguay, Peru, Venezuela
<i>Corydalus armatus</i> Hagen, 1861	Argentina, Bolivia, Colombia, Ecuador, Peru, Venezuela
<i>Corydalus arpi</i> Navás, 1936†	Brazil, Venezuela
<i>Corydalus batesii</i> MacLachlan, 1868	Bolivia, Brazil, Colombia, Ecuador, French Guiana, Guyana, Peru, Suriname, Venezuela
<i>Corydalus clavijoi</i> Contreras-Ramos, 2002	Venezuela
<i>Corydalus crossi</i> Contreras-Ramos, 2002	Venezuela
<i>Corydalus flavicornis</i> Stitz, 1914	Colombia, Costa Rica, Ecuador, El Salvador, Guatemala, Honduras, Panama, Peru, Venezuela
<i>Corydalus flinti</i> Contreras-Ramos, 1998†	Venezuela
<i>Corydalus hayashii</i> Contreras-Ramos, 2002†	Venezuela
<i>Corydalus hecate</i> MacLachlan, 1866†	Brazil, Peru, Venezuela‡
<i>Corydalus mayri</i> Contreras-Ramos, 2002†	Venezuela
<i>Corydalus neblinensis</i> Contreras-Ramos, 1998	Venezuela
<i>Corydalus nubilus</i> Erichson, 1848	Brazil, French Guiana, Guyana, Venezuela
<i>Corydalus peruvianus</i> Davis, 1903	Argentina, Bolivia, Colombia, Costa Rica, Ecuador, Guatemala, Mexico, Panama, Peru, Venezuela
<i>Corydalus tessellatus</i> Stitz, 1914	Colombia‡, Venezuela
<i>Corydalus wanningeri</i> sp. n.	Venezuela

†Male mandible short, female-like, with discrete dentition; ‡Doubtful record.

## Taxonomy

### *Corydalus wanningeri* Contreras-Ramos & Von der Dunk, sp. n.

urn:lsid:zoobank.org:act:D8CD9346-1591-4C3D-8898-A5145867FB18

Figures 1–8

**Etymology.** Named after Prof. Rupert Wanninger, amateur and outreach entomologist from Donaustauf, Bavaria, Germany, collector of the type series.

**Type material.** Holotype, male, VENEZUELA: Bolívar, Escalera Km 110, el. 1,000 m, 26.viii.1994, leg. Rupert Wanninger [Head width 11.8 mm, mandible length 29.5 mm, antenna length 65.3 mm, forewing length 80.4 mm] (CNIN-UNAM). Paratypes: VENEZUELA, [Bolívar], Guyana, Km 120, 1997, 1 male [genitalia dissected] (MIZA); [Bolívar], Escalera, [Km 110, 1,000 m], 25.viii. 1999, [R. Wanninger], 1 female [genitalia dissected] (CNIN-UNAM); Bolívar, Escalera, Km 110,



**Figure 1.** *Corydalus wanningeri* sp. n. Male holotype, habitus.

1,000 m, 2.ii.2001, leg. R. Wanninger, 1 female (ZSM); [Bolívar, Escalera, Km 110, no date, R. Wanninger], 1 male (ZSM), 1 female (RW).

**Diagnosis.** Head and pronotum are reddish-brown, ferruginous (Figs 1–3), thus resembling in color species such as *Corydalus cephalotes* Rambur and *C. hecate* MacLachlan, both with monomorphic mandibles (males with short mandibles with discrete dentition), or *C. batesii* MacLachlan and *C. holzenthali* Contreras-Ramos, with sexually dimorphic mandibles (males with elongate mandibles with reduced dentition). However, both color of body and wings most closely resembles *C. neblinensis* Contreras-Ramos (Contreras-Ramos 1998, figs 124–127). In both species, antennae are paler than head and wings are pale reddish brown, unpatterned. However, in the new species ninth gonostyli are distinct (Figs 4, 5), with a narrowed apex (subclavate, unmodified in *C. neblinensis*, Contreras-Ramos 1998, figs 26A, 26B). Tenth sternite lobes (Figs 5, 6) are sclerotized, close to each other, convergent, and bluntly pointed (semimembranous, widely separated, and papilliform in *C. neblinensis*, Contreras-Ramos 1998, fig. 26C). Females may be distinguished by the unpatterned reddish color and by a mandibular dentitional arrangement with an inner prepedal concavity and moderately separated first and second teeth (Figs 3, 8), similar to *C. nubilus* and *C. crossi* (Contreras-Ramos 1998, fig. 27I; Contreras-Ramos 2002, fig. 27).

**Phylogenetic position.** Despite a close superficial resemblance to *C. neblinensis*, *C. wanningeri* sp. n., does not appear to be closely related to that or other kindred species, such as *C. batesii* or *C. holzenthali*. The new species appears to be closest to *C. crossi* Contreras-Ramos, because of the structure of the 10<sup>th</sup> sternite and a basal protrusion of the 9<sup>th</sup> gono-



**Figure 2.** *Corydalus wanningeri* sp. n. Female paratype, habitus.

stylus. If so, the new species would be basal among species with elongate mandibles. At least, *C. wanningeri*, new species, should be basal with respect to species with a subclavate 9<sup>th</sup> gonostylus and non-incurvate 10<sup>th</sup> tergite, and so would lay out of Unnamed Group 1 (Contreras-Ramos 1998, table 36). However, a certain phylogenetic position of the new species would be known only after a formal phylogenetic analysis. Both *C. crossi* and *C. wanningeri*, new species, share a Guayana Shield affinity.

**Adult male.** Head width 11.8–12.8 mm (average 12.3,  $n = 3$ ), mandible length 29.5–30.3 mm (average 29.9 mm,  $n = 3$ ), antenna length 64.5–65.3 mm (average 64.9 mm,  $n = 2$ ), forewing length 77.7–80.4 mm (average 79.5 mm,  $n = 3$ ), antenna length/forewing length 0.81–0.83. Color uniform dark reddish-brown. Head dark reddish-brown, unpatterned, mandible elongate with reduced dentition (Fig. 1). Clypeal margin thinly black, lateral projections moderately developed, flat to slightly concave, median projection shallowly incised (Fig. 7). Antenna 87–89-segmented, filiform, scape and flagellum pale brown, tip infusate. Maxilla blackish, 4-segmented palp brown. Labial palp 3-segmented, pale brown, last segment elongate.

Pronotum dark reddish-brown, unpatterned. Forewing pale reddish-brown, semitranslucent, unpatterned; veins reddish-yellow, except basal half of Sc and R infusate;  $M_{1+2}$  3-branched (variably 4-branched),  $M_{3+4}$  a single vein. Hindwing pale-reddish, semitranslucent, basal 1/4 of R infusate.

Genitalia (Figs 4–6). Ninth tergum subquadrate, V-shaped internal inflection reaching midlength of tergum. Anal tubercle without lateral sclerites. Tenth tergites slightly longer than ninth tergum, digitiform; basal 1/3 wide, roundly concave (Fig. 4). Ninth gonostylus subclavate, about 4/5 as long as 10<sup>th</sup> tergite, with narrowed digitiform apex



**Figure 3.** *Corydalus wanningeri*, sp. n. Female paratype, head and pronotum (dorsal).


(Fig. 5). Ninth sternum subquadrate, semimembranous, posterolateral lobes moderately developed (Fig. 5). Membrane between 9<sup>th</sup> and 10<sup>th</sup> sternites with thickened wrinkled portion. Tenth sternite moderately convex, anteromedian margin slightly convex; anterolateral projections moderately developed, blunt; lobes well sclerotized, elongate-trianguloid, apically convergent (Fig. 6). Pregenital sacs apparently absent.

**Adult female.** Forewing length 65.1–80.6 mm (average 74.1 mm,  $n = 3$ ). Color of body and wings similar to male (Figs 2, 3). Mandible with base dark reddish-brown, rest blackish; shape similar to *C. nubilus* Erichson and *C. crossi*, basal preapical tooth moderately separated from second (Fig. 8; Contreras-Ramos 1998, fig. 27I; Contreras-Ramos 2002, fig. 27). Antenna filiform, pale brown.

Terminalia non distinct. Sternal pouch between 6<sup>th</sup> and 7<sup>th</sup> abdominal segments well developed. Eighth sternum moderately sclerotized, discontinuous with pleural area, posterior margin mesally semimembranous, concave.

**Key to long-mandibled males of known *Corydalus* species from Venezuela (modified from Contreras-Ramos 2002).**

1. Ninth sternum modified, subattenuate and more sclerotized posteromesally (Contreras-Ramos 1998, fig. 27B) or with posteromedian projection (Contreras-Ramos 1998, fig. 31B) ..... **2**
- Ninth sternum unmodified, subquadrate (Fig. 5; Contreras-Ramos 1998, figs 2B, 4B) ..... **4**



Figures 4–5. 4 *Corydalus wanningeri*, sp. n. 4 male genitalia (dorsal) 5 male genitalia (ventral).





Figure 6. *Corydalus wanningeri*, sp. n. Male tenth sternite.

- 2. Ninth sternum subattenuate, noticeably more sclerotized posteromesally (Contreras-Ramos 1998, fig. 27B) ..... ***C. nubilus* Erichson**
- Ninth sternum with posteromedian projection ..... **3**
- 3. Posteromedian projection of 9<sup>th</sup> sternum large (nearly as long as sternum), thumblike; 9<sup>th</sup> gonostylus unguiform (Contreras-Ramos 1998, fig. 31B) ..... ***C. tessellatus* Stitz**
- Posteromedian projection of 9<sup>th</sup> sternum small (about 1/2 as long as sternum), narrow; 9<sup>th</sup> gonostylus tubular, with conspicuous preapical claw (Contreras-Ramos 2002, fig. 17) ..... ***C. clavijoi* Contreras-Ramos**
- 4. Ninth gonostylus elongate, somewhat flattened or tubular (Contreras-Ramos 1998, fig. 2B) ..... **5**
- Ninth gonostylus subclavate (Fig. 5; Contreras-Ramos 1998, figs 4B, 7B, 17B) ..... **6**



**Figure 7.** *Corydalus wanningeri*, sp. n. Male clypeal margin (dorsal).



**Figure 8.** *Corydalus wanningeri*, sp. n. Female mandible (right, dorsal).

5. Ninth gonostylus and 10<sup>th</sup> tergite slender, subequal in length and shape (Contreras-Ramos 1998, fig. 2A) ..... ***C. affinis* Burmeister**  
 – Ninth gonostylus narrower and noticeably shorter than 10<sup>th</sup> tergite (Contreras-Ramos 2002, fig. 23) ..... ***C. crossi* Contreras-Ramos**
6. Head and pronotum reddish brown; 10<sup>th</sup> tergite apex without incurvation (Fig. 5), although it may be enlarged (Contreras-Ramos 1998, Figs 7E, 26E) ....7  
 – Head and pronotum yellowish to greenish brown; 10<sup>th</sup> tergite with well developed apical incurvation (Contreras-Ramos 1998, Figs 4A, 17B) ..... **9**
7. Forewing contrastingly patterned (Contreras-Ramos 1998, fig. 58) .....  
 ..... ***C. batesii* MacLachlan**  
 – Forewing not so contrastingly patterned ..... **8**
8. Forewing pale, clear, nearly translucent, few subtle small white spots (Contreras-Ramos 1998, Figs 124–126); 9<sup>th</sup> gonostylus unmodified, 10<sup>th</sup> sternite lobes papiliform, separated (Contreras-Ramos 1998, Figs 26B, 26C) .....  
***C. neblinensis* Contreras-Ramos**  
 – Forewing rather opaque, uniformly pale reddish (Fig. 1); 9<sup>th</sup> gonostylus with narrowed digitiform apex, 10<sup>th</sup> sternite lobes elongate-trianguloid, close to each other (Figs 5, 6) ..... ***C. wanningeri* sp. n.**
9. Antenna conspicuously subserrate, sinuate (Contreras-Ramos 1998, fig. 17F); 10<sup>th</sup> sternite with anteromedian projection (Contreras-Ramos 1998, fig. 17C) ..... ***C. flavicornis* Stitz**  
 – Antenna slightly subserrate; 10<sup>th</sup> sternite without anteromedian projection (Contreras-Ramos 1998, fig. 4C) ..... **10**

10. Antenna, including scape and pedicel, pale to dark brown, apically infusate (Contreras-Ramos 1998, Figs 43, 44, 48); 10<sup>th</sup> sternite lobes typically subequal in width and length, less than half length of lobe surpassing posterior edge of 10<sup>th</sup> sternite (Contreras-Ramos 1998, fig. 4C); pregenital sacs well developed, conspicuous (Contreras-Ramos 1998, fig. 4F)..... ***C. armatus* Hagen**
- Antenna, including scape and pedicel, yellow to yellowish green, up to distal 1/3 infusate (Contreras-Ramos 1998, Figs 139–141); 10<sup>th</sup> sternite lobes typically about twice as long as wide, about half of lobe surpassing posterior edge of 10<sup>th</sup> sternite (Contreras-Ramos 1998, fig. 29C); pregenital sacs apparently absent, inconspicuous ..... ***C. peruvianus* Davis**

## Acknowledgments

Help with line drawings by Jorge A. González Martínez is greatly appreciated. Comments from David E. Bowles, Missouri State University, and an anonymous reviewer, improved a previous draft of this paper. Institutional support to ACR from Biblioteca (Georgina Ortega), Jefatura de Zoología (Patricia Escalante), Secretaría Técnica (Noemí Chávez), and Dirección (Tila M. Pérez) is gratefully appreciated.

## References

- Contreras-Ramos A (1998) Systematics of the dobsonfly genus *Corydalus* (Megaloptera: Corydalidae). Thomas Say Publications in Entomology: Monographs, Entomological Society of America, Lanham, 360 pp.
- Contreras-Ramos A (1999) List of species of Neotropical Megaloptera (Neuropterida). Proceedings of the Entomological Society of Washington 101: 274-284.
- Contreras-Ramos A (2002) Six new species of dobsonflies from Venezuela (Megaloptera: Corydalidae: Corydalinae). Aquatic Insects 24: 55-75.
- Contreras-Ramos A (2005) Recent accounts on the systematics and biogeography of Neotropical Megaloptera (Corydalidae, Sialidae). Annali del Museo Civico di Storia Naturale di Ferrara 8: 67-72.

