

Host insect species of *Ophiocordyceps sinensis*: a review

Xiao-Liang Wang^{1,2}, Yi-Jian Yao¹

1 State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, P.O. Box 2714, Beijing 100101, P.R. China **2** Graduate University of the Chinese Academy of Sciences, Beijing 100049, P.R. China

Corresponding author: Yi-Jian Yao (yaoyj@sun.im.ac.cn)

Academic editor: E. van Nieuwerkerken | Received 14 December 2010 | Accepted 12 August 2011 | Published 8 September 2011

Citation: Wang X-L, Yao Y-J (2011) Host insect species of *Ophiocordyceps sinensis*: a review. ZooKeys 127: 43–59. doi: 10.3897/zookeys.127.802

Abstract

Ophiocordyceps sinensis (\equiv *Cordyceps sinensis*) is one of the most valued medicinal fungi in China, used for its invigorating effects in strengthening the body and restoring energy. The fungus parasitizes larvae of moths and converts them into sclerotia from which the fungus fruiting body grows. Since the late 1950s, considerable effort has been devoted to the study of host insects related to the fungus. In the present paper, the research history of insect species associated with *O. sinensis* is briefly reviewed and an extensive literature survey is presented. Ninety-one insect names, spanning 13 genera, related to host insects of *O. sinensis* are investigated. The relationships between the reported insect species and *O. sinensis* are analyzed. Fifty-seven of these are considered as recognizable potential host species of the fungus distributed throughout the Tibetan Plateau, whilst eight are considered as indeterminate hosts and 26 as non-hosts. Among the names of recognizable potential host insects, three are invalid (*nomen nudum*) and require further study. This work provides basic information for management of the insect resources and for the conservation and sustainable use of *O. sinensis*.

Keywords

Cordyceps, Fungi, Hepialidae, host insects, *Ophiocordyceps*

Introduction

Ophiocordyceps sinensis (Berk.) G.H. Sung, J.M. Sung, Hywel-Jones & Spatafora is an ascomycete fungus, which is also known as the Chinese Caterpillar Fungus or “Dong Chong Xia Cao” (winter worm, summer grass) in Chinese, or “Hia Tsao Tong

Tchong” and “Hea Tsaon Tsong Chung” in early English translations (Pegler et al. 1994). The fungus parasitizes larvae of moths belonging to the order Lepidoptera, especially *Hepialus/Thitarodes*. The infected larva is converted into a sclerotium covered by the intact exoskeleton of the insect to withstand the winter, which is regarded as “winter worm”. In the late spring or summer of the next year, a clavate stroma of the fungus grows from the sclerotium and emerged from the ground appearing as a herb, which is regarded as “summer grass” (Pegler et al. 1994, Yao 2004). As a valued Chinese herb and tonic, *O. sinensis* has a long history of use and a high reputation of value both in China and abroad. In Traditional Chinese Medicine (TCM), the fungus is believed to nourish the lungs and kidneys (Wu 1757). It has also been shown in recent studies to have multiple pharmacological effects, including immunomodulating (Wu et al. 2006), hypocholesterolemic (Koh et al. 2003), hypoglycemic (Zhang et al. 2006), anti-tumor (Wu et al. 2005), anti-oxidation (Dong and Yao 2008) and anti-aging (Ji et al. 2009) activities.

The natural product of *O. sinensis* for medicinal use is actually a combination of the fungus and an insect larva. The fungus parasitizes underground dwelling larvae of moths and converts them into sclerotia, from which the fruiting body of the fungus grows (Pegler et al. 1994, Wang 1995, Yao 2004). *Ophiocordyceps sinensis* is endemic to the Tibetan Plateau, with a distribution covering five provinces in China, i.e., Gansu, Qinghai, Sichuan, Tibet and Yunnan. It may be found in alpine meadow and shrub habitat from an altitude of 3000 m up to the snow-line (Wang 1995, Yao 2004). The natural production of the fungus is limited owing to its strict host-specificity, confined geographic distribution and over exploitation by humans in recent decades. It is therefore currently listed as an endangered species under the second class of state protection (State Forestry Administration and Ministry of Agriculture 1999).

Insect host species of *O. sinensis* belong to the family Hepialidae (Lepidoptera) (Chu et al. 2004). Since the late 1950s, much effort has been devoted to study the insect species related to the fungus in China. In 1958, researchers from the Institute of Zoology, Chinese Academy of Sciences, began their investigation in some parts of Qinghai and Sichuan provinces. The first report on host insects of *O. sinensis* in China was on *Hepialus armoricanus* Oberthür (= *Thitarodes armoricanus* Oberthür, Chu 1965) and then followed by studies of biological characteristics of the moth (Chen et al. 1973). The research on Hepialidae diversity and taxonomy grew rapidly in China during the 1980s, leading to a series of publications of new taxa, including four new genera, 71 new species and one subspecies (see Chu and Wang 1985a, b; Liang et al. 1988; Wang 1990; Wu 1992; Li et al. 1993; Fu et al. 1991, 2002; Yang 1993, 1994; Yang et al. 1991a, 1992a, 1995; Liang 1995; Yang and Jiang 1995; Shen and Zhou 1997; Yan 2000; Wang et al. 2001; Chu et al. 2004; Zhang et al. 2007; Tu et al. 2009). A number of attempts have been made to summarize the insect species associated with *O. sinensis* and various numbers of host species were recorded, e.g., five by Yin (1987); eleven by Yin et al. (1991) without a name list; two, 22 and 23 by Jiang (1989, 1991, 2001); 22 by Chen and Jin

(1992); 20 by Long (1992); 37 by Liu et al. (1995); 19 by Li (1996); 38 by Dong and Luo (1996); 31 by Wang et al. (1996) and 37 by Yang (1998). Recently, Chu et al. (2004) published the volume on Hepialidae and Epiplemlidae in the Fauna Sinica, in which seven genera and 82 species or subspecies of Hepialidae in China were listed, and 14 species in six genera were believed to be hosts of *O. sinensis* but the names were not given in that list. However, in two other recent publications (Liu et al. 2005, 2006), 66 and 69 insect names were listed respectively as the hosts of the fungus. However, all these accounts, except Chu et al. (2004), provided only the number or a list of insect names without any relevant information to determine whether they are hosts of the fungus or not. Therefore, the number of insect host species of *O. sinensis* and the relationship between those insects and the fungus remain unclear. To clarify this situation, an extensive survey of the literature on the host of *O. sinensis* was carried out to gather all the insect names related to the fungus in the literature and to analyze the relationship between the insect species and the fungus. The results of this work are reported here.

Recently, a global inventory of the suborder Exoporia, comprising Mnesarchaeoidea and Hepialoidea, was presented by Nielsen et al. (2000), in which the systematic position of many taxa was checked and adjusted. Nielsen et al.'s classification system for Hepialidae is adopted in this study.

Methods

Based upon an exhaustive literature search, a total of 4793 publications related to *Cordyceps/Ophiocordyceps* and *Hepialus/Thitarodes*, in either English or Chinese, were gathered. Those publications relevant to host insects of *O. sinensis*, including reports on taxonomy, checklists, fauna, biological characteristics, ecology and geographical distribution were examined for information about these insects. All the insect names associated with *O. sinensis* were assessed based on the following criteria to determine their relationship with the fungus. Taxa which met both of the following requirements were considered as recognizable potential insect host species of *O. sinensis*: (1) The distribution areas of the insect overlapped that of *O. sinensis*, which was determined on the basis of field collections made by this research group during the years 2000–2010, examination of herbarium specimens, and another exhaustive literature analysis carried out in this laboratory (Li et al. in press). (2) The insect was reported from an altitude above 3000 m on the Tibetan Plateau. However, stem-boring insects were excluded as hosts of the fungus, even if they were hepialid and distributed above 3000 m within the distribution areas of *O. sinensis*, because the fungus infects only subterranean root-boring insects. Species of root-borers lacking altitude information were considered as indeterminate hosts of *O. sinensis* requiring further confirmation, despite the overlap of distribution areas with *O. sinensis*. Species falling in both of the following circumstances were deemed not to be host insects of *O. sinensis*: the distribution of the insect was outside that of *O. sinensis* and below an altitude of 3000 m.

Results

A total of 91 names in 13 genera of Hepialidae were found in the literature search. They are listed in alphabetical order in Table 1, together with geographic distribution, altitude, main references and the relationship with *O. sinensis* as determined by this study. Insect names used in the references, if different from that in Nielsen et al. (2000), are also given. There are 67 names in the references being combined in different genera by Nielsen et al. (2000) and a total of 71 species were originally described from China. Twenty four species described in the literature were not included in Nielsen et al. (2000).

Fifty-seven species are considered here as recognizable potential host insects of *O. sinensis*, whilst eight as indeterminate hosts and 26 as non-hosts. The recorded altitude ranges of the recognized potential host insects were found to vary from 2800 to 5100 m. The distribution areas of these species covered 26 provinces in China and more than 12 other countries. Three of the recognizable potential host names are invalid (*nomen nudum*).

Table 1. Potential insect hosts of *Ophiocordyceps sinensis*.

Insect name	Geographic distribution	Altitude (m)	Main references	Status of host insect [†]	Different name in the main references
<i>Bipecilus yunnanensis</i> Chu & Wang, 1985	Yunnan Province: Lijiang County [‡]	3200	Chu and Wang 1985a, Nielsen 1988	P	
<i>Bipecilus zhejiangensis</i> Wang, 2001 [§]	Zhejiang Province: Anji County [‡] ; Fujian Province	—	Wang et al. 2001, Huang 2006, Wu 2007	N	
<i>Endoclita anhuiensis</i> (Chu & Wang, 1985)	Anhui Province: Yuexi County [‡]	—	Chu and Wang 1985b, Chu et al. 2004	N	<i>Phassus anhuiensis</i> Chu & Wang, 1985
<i>Endoclita davidi</i> (Poujade, 1886)	Sichuan Province: Baoxing and Danba Counties; Fujian and Guangxi Provinces	3600	Chu and Wang 1985b, Yang 1998, Chu et al. 2004	P	<i>Hepialus davidi</i> Poujade, 1886; <i>Phassus giganodus</i> Chu & Wang, 1985
<i>Endoclita excrescens</i> (Butler, 1877)	Sichuan Province: Yingjing County; Anhui, Hebei, Heilongjiang, He'nan, Jilin, Liaoning, Shandong and Shanxi Provinces; Inner Mongolia Autonomous Region; Japan	—	Chu et al. 2004	N	<i>Phassus excrescens</i> (Butler, 1877); <i>Phassus camphorae</i> Sasaki, 1908
<i>Endoclita fujianodus</i> (Chu & Wang, 1985)	Fujian Province [‡]	—	Chu and Wang 1985b, Chu et al. 2004	N	<i>Phassus fujianodus</i> Chu & Wang, 1985
<i>Endoclita jingdongensis</i> (Chu & Wang, 1985)	Yunnan Province: Jingdong County [‡] , Xishuangbanna Prefecture [‡]	—	Chu and Wang 1985b, Chu et al. 2004	N	<i>Phassus jingdongensis</i> Chu & Wang, 1985

Insect name	Geographic distribution	Altitude (m)	Main references	Status of host insect [†]	Different name in the main references
<i>Endoclita nodus</i> (Chu & Wang, 1985) ^l	Anhui Province: Yuexi County [‡] ; Guangxi, Guizhou, Hainan, Hu'nan, Jiangxi and Zhejiang Provinces	—	Chu and Wang 1985b, Chu et al. 2004	N	<i>Phassus nodus</i> Chu & Wang, 1985
<i>Endoclita signifer</i> (Walker, 1856) ^l	Hu'nan Province	—	Chu and Wang 1985b, Chu et al. 2004	N	<i>Phassus humanensis</i> Chu & Wang, 1985
<i>Endoclita sinensis</i> (Moore, 1877) ^l	Fujian, Guangdong, Guangxi, Hainan, Hebei, He'nan, Hubei, Hu'nan, Jiangxi, Shandong, Shanxi, Sichuan, Yunnan and Zhejiang Provinces; Shanghai Municipality; D.P.R. Korea; India; Japan; Sri Lanka	—	Chu et al. 2004	N	<i>Phassus sinensis</i> Moore, 1877; <i>Phassus herzi</i> Fixsen, 1887
<i>Endoclita xizangensis</i> (Chu & Wang, 1985) ^l	Tibet Autonomous Region: Nyalam County [‡]	—	Chu and Wang 1985b, Wang et al. 1996, Chu et al. 2004	N	<i>Phassus xizangensis</i> Chu & Wang, 1985
<i>Endoclita yunnanensis</i> (Chu & Wang, 1985) ^l	Yunnan Province: Jinghong Municipality [‡] ; Guangdong and Hainan Provinces	—	Chu and Wang 1985b, Chu et al. 2004	N	<i>Phassus yunnanensis</i> Chu & Wang, 1985
<i>Gazoryctra ganna</i> (Hübner, [1808])	Qinghai Province: Zadoi County; Heilongjiang Province; Inner Mongolia Autonomous Region; Northern Europe; Russia	3900 [‡]	Chu and Wang 1985a, Wang et al. 1996, Yang 1998, Chu et al. 2004, Karsholt and Nieuwerkerken 2010	P	<i>Hepialus ganna</i> (Hübner, [1808])
<i>Gazoryctra macilentus</i> (Eversmann, 1851)	Hebei and Heilongjiang Provinces; Inner Mongolia Autonomous Region; Eastern Siberia; Mongolia	340–1300	Wang et al. 1996, Yang 1998, Chu et al. 2004	N	<i>Hepialus macilentus</i> Eversmann, 1851
<i>Hepialiscus jiangbeiensis</i> Chu & Wang, 2004 [§]	Chongqing Municipality [‡]	—	Chu et al. 2004	N	
<i>Hepialiscus ledongensis</i> Chu & Wang, 2004 [§]	Hainan Province: Ledong County [‡]	—	Chu et al. 2004	N	
<i>Hepialiscus nepalensis</i> (Walker, 1856)	Tibet Autonomous Region: Nyalam County; India; Nepal; Sikkim	—	Chu and Wang 1985a, Wang et al. 1996, Chu et al. 2004	I	<i>Hepialiscus flavus</i> Chu & Wang, 1985
<i>Hepialus bibelteus</i> Shen & Zhou, 1997 [§]	Yunnan Province: Deqên County [‡]	4500	Shen and Zhou 1997, Chu et al. 2004	P	
<i>Hepialus biruensis</i> Fu, 2002 [§]	Tibet Autonomous Region: Biru County [‡]	4400–4700	Fu et al. 2002, Chu et al. 2004	P	
<i>Hepialus dinggyeensis</i> Chu & Wang, 2004 [§]	Tibet Autonomous Region: Dinggyê County [‡]	—	Chu et al. 2004	I	

Insect name	Geographic distribution	Altitude (m)	Main references	Status of host insect [†]	Different name in the main references
<i>Hepialus gangcaensis</i> Chu & Wang, 2004 [§]	Qinghai Province: Gangca County [‡]	3195 [¶]	Chu et al. 2004	P	
<i>Hepialus guidera</i> Yan, 2001 [§]	Qinghai Province: Guide County	3400–3600	Yan 2001a, Li et al. 2002, Li and Li 2004	P, IN	<i>Hepialus guidera</i> Yan, 2001
<i>Hepialus hainanensis</i> Chu & Wang, 2004 [§]	Hainan Province: Ledong County [‡]	—	Chu et al. 2004	N	
<i>Hepialus humuli</i> (Linnaeus, 1758)	Heilongjiang Province; Europe and Siberia	—	Chu et al. 2004, Karsholt and Nieukerken 2010	N	
<i>Hepialus lagii</i> Yan, 2001 [§]	Qinghai Province: Guide County	3400–3600	Yan 2001b; Yan 2001c, Li et al. 2002, Li and Li 2004, Zhang et al. 2009	P, IN	
<i>Hepialus latitegumenus</i> Shen & Zhou, 1997 [§]	Yunnan Province: Deqên County [‡]	4500	Shen and Zhou 1997, Chu et al. 2004	P	
<i>Hepialus maquensis</i> Chu & Wang, 2004 [§]	Gansu Province: Maqu County [‡]	3300 [¶]	Chu et al. 2004	P	
<i>Hepialus namensis</i> Chu & Wang, 2004 [§]	Tibet Autonomous Region: Damxung County [‡]	4200 [¶]	Chu et al. 2004	P	
<i>Hepialus namlinensis</i> Chu & Wang, 2004 [§]	Tibet Autonomous Region: Namling County [‡]	3704 [¶]	Chu et al. 2004	P	
<i>Hepialus pui</i> Zhang, Gu & Liu, 2007 [§]	Tibet Autonomous Region: Nyingchi County [‡]	4100–5000	Zhang et al. 2007	P	
<i>Hepialus xiaojinensis</i> Tu, Ma & Zhang 2009 [§]	Sichuan Province: Xiaojin [‡] and Jinchuan County [‡]	3500–4800	Tu et al. 2009	P	
<i>Hepialus xingazeensis</i> Chu & Wang, 2004 [§]	Tibet Autonomous Region: Xigazê Prefecture [‡]	—	Chu et al. 2004	I	
<i>Hepialus yadongensis</i> Chu & Wang, 2004 [§]	Tibet Autonomous Region: Yadong County [‡]	—	Chu et al. 2004	I	
<i>Hepialus yongshengensis</i> Chu & Wang, 2004 [§]	Yunnan Province: Yongsheng County [‡]	—	Chu et al. 2004	I	
<i>Hepialus zadoiensis</i> Chu & Wang, 2004 [§]	Qinghai Province: Zadoi County [‡]	3900 [¶]	Chu et al. 2004	P	
<i>Magnificus jiuizhiensis</i> Yan, 2000 [§]	Qinghai Province: Jigzhi County [‡]	3800–3900	Yan 2000	P	
<i>Magnificus zhiduoensis</i> Yan, 2000 [§]	Qinghai Province: Zhidoi County [‡]	4400–4600	Yan 2000	P	
<i>Napialus chenzhouensis</i> Chu & Wang, 2004 [§]	Hu'nan Province: Chenzhou City [‡] ; Shanghai Municipality	—	Chu et al. 2004, Chen and Wang 2006	N	
<i>Napialus chongqingensis</i> Wu, 1992	Chongqing Municipality [‡]	—	Wu 1992, Chu et al. 2004	N	
<i>Napialus hunanensis</i> Chu & Wang, 1985	Hu'nan Provinces: Changsha City [‡] ; Guangdong, Guangxi, Hainan, and Jiangxi Provinces	—	Chu and Wang 1985a, Wang et al. 1996, Chu et al. 2004	N	

Insect name	Geographic distribution	Altitude (m)	Main references	Status of host insect [†]	Different name in the main references
<i>Napialus jiangxiensis</i> Chu & Wang, 2004 [§]	Jiangxi Province: Taihe County [‡]	—	Chu et al. 2004	N	
<i>Palpifer sexnotatus</i> (Moore, 1879) ^l	Sichuan and Taiwan Provinces; Kashmir; India; Sri Lanka; Japan	—	Chu et al. 2004	N	
<i>Parahepialiscus borneensis</i> (Pfitzner in Pfitzner & Gaede, 1933)	Hu'nan Province; Malaysia	—	Chu et al. 2004	N	<i>Hepialiscus borneensis</i> Pfitzner, 1933
<i>Pharmacis carna</i> ([Denis & Schiffermüller], 1775)	Sichuan Province: Luhuo County; Europe	3050 [¶]	Chu et al. 2004, Karsholt and Nieukerken 2010	P	<i>Hepialus carna</i> ([Denis & Schiffermüller], 1775)
<i>Pharmacis fusconebulosa</i> (De Geer, 1778)	Sichuan Province: Kangding District; Europe; Russia	3500 [¶]	Chu et al. 2004, Yu 2004, Karsholt and Nieukerken 2010	P	<i>Hepialus fusconebulosa</i> (De Geer, 1778); <i>Hepialus gallicus</i> Lederer, 1852
<i>Pharmacis pyrenaicus</i> (Donzel, 1838)	Sichuan Province: Dègè County; Southwest Europe	3880 [¶]	Chu et al. 2004, Karsholt and Nieukerken 2010	P	<i>Hepialus alticola</i> Oberthür, 1881
<i>Sthenopsis regius</i> (Staudinger, 1896) ^l	—	—	Yin 1987	N	<i>Phassus regius</i> (Staudinger, 1896)
<i>Sthenopsis roseus</i> (Oberthür, 1911) ^l	Hubei Province	—	Chu and Wang 1985b, Chu et al. 2004	N	<i>Phassus miniatus</i> Chu & Wang, 1985
<i>Thitarodes albipictus</i> (Yang, 1993)	Yunnan Province: Deqên County [‡]	4500–4800	Wang et al. 1996, Yang 1993	P	<i>Hepialus albipictus</i> Yang, 1993
<i>Thitarodes altaicola</i> (Wang, 1990)	Xinjiang Uygur Autonomous Region [‡]	1300–1800	Wang 1990, Yang 1998, Chu et al. 2004	N	<i>Hepialus altaicola</i> Wang, 1990
<i>Thitarodes anomopterus</i> (Yang, 1994)	Yunnan Province: Jianchuan [‡] and Lijiang Counties [‡]	2800–3100	Yang 1994, Yang 1998	P	<i>Hepialus anomopterus</i> Yang, 1994
<i>Thitarodes armoricanus</i> (Oberthür, 1909)	Gansu, Qinghai, Sichuan and Yunnan Provinces; Tibet Autonomous Region; Xinjiang Uygur Autonomous Region	3600–5000	Chu 1965, Chen et al. 1973, Yang et al. 1987, Chu et al. 2004	P	<i>Hepialus armoricanus</i> Oberthür, 1909
<i>Thitarodes baimaensis</i> (Liang in Liang et al., 1988)	Yunnan Province: Deqên County [‡]	4500–4900	Liang et al. 1988, Yang 1998	P	<i>Hepialus baimaensis</i> Liang, 1988
<i>Thitarodes baqingensis</i> (Yang & Jiang, 1995)	Tibet Autonomous Region: Baqên County [‡]	4600–4800	Yang and Jiang 1995	P	<i>Hepialus baqingensis</i> Yang and Jiang, 1995

Insect name	Geographic distribution	Altitude (m)	Main references	Status of host insect [†]	Different name in the main references
<i>Thitarodes callinivalis</i> (Liang, 1995)	Tibet Autonomous Region; Yunnan Province : Deqên County [‡]	4300–4600	Liang 1995, Yang 1998	P	<i>Hepialus callinivalis</i> Liang, 1995
<i>Thitarodes cingulatus</i> (Yang & Zhang in Yang et al., 1995)	Gansu Province: Wenxian County [‡]	3200–3800	Yang et al. 1995, Yang 1998	P	<i>Hepialus cingulatus</i> Yang & Zhang, 1995
<i>Thitarodes damxungensis</i> (Yang in Yang & Jiang, 1995)	Tibet Autonomous Region: Damxung County [‡]	4500–4680	Yang and Jiang 1995	P	<i>Hepialus damxungensis</i> Yang, 1995
<i>Thitarodes deqinensis</i> (Liang in Liang et al., 1988)	Yunnan Province: Deqên County	4200–4700	Liang et al. 1988, Yang et al. 1992b	P	<i>Hepialus deqinensis</i> Liang, 1988
<i>Thitarodes dongyuensis</i> (Liang in Yang et al., 1992)	Tibet Autonomous Region: Markam County; Yunnan Province: Deqên County	4000–4700	Yang et al. 1992b, Yang et al. 1996, Hu and Zha 2010	P, IN	<i>Hepialus dongyuensis</i> Liang in Yang et al., 1992
<i>Thitarodes ferrugineus</i> (Li, Yang & Shen, 1993)	Yunnan Province: Deqên County [‡]	4200–4700	Yang et al. 1992b, Li et al. 1993, Chu et al. 2004	P	<i>Hepialus ferrugineus</i> Li, Yang & Shen, 1993
<i>Thitarodes gonggaensis</i> (Fu & Huang in Fu et al., 1991)	Sichuan Province: Kangding County [‡]	3800–4400	Fu et al. 1991, Yang 1998, Chu et al. 2004	P	<i>Hepialus gonggaensis</i> Fu & Huang, 1991
<i>Thitarodes jialangensis</i> (Yang, 1994)	Tibet Autonomous Region: Zogang County [‡]	4000–4600	Yang 1994, Chu et al. 2004	P	<i>Hepialus jialangensis</i> Yang, 1994
<i>Thitarodes jianchuanensis</i> (Yang, 1994)	Yunnan Province: Jianchuan County [‡]	2900–3500	Yang 1994, Yang 1998	P	<i>Hepialus jianchuanensis</i> Yang, 1994
<i>Thitarodes jinshaensis</i> (Yang, 1993)	Yunnan Province: Deqên County [‡]	4600	Yang 1993, Chu et al. 2004	P	<i>Hepialus jinshaensis</i> Yang, 1993
<i>Thitarodes kangdingensis</i> (Chu & Wang, 1985)	Sichuan Province: Kangding County [‡]	3600–4500	Chu and Wang 1985a, Yang et al. 1991b, Chu et al. 2004	P	<i>Hepialus kangdingensis</i> Chu & Wang, 1985
<i>Thitarodes kangdingroides</i> (Chu & Wang, 1985)	Sichuan Province: Kangding County [‡]	4200	Chu and Wang 1985a, Yang 1998, Chu et al. 2004	P	<i>Hepialus kangdingroides</i> Chu & Wang, 1985
<i>Thitarodes lijiangensis</i> (Chu & Wang, 1985)	Yunnan Province: Lijiang County [‡]	3500–4400	Chu and Wang 1985a, Yang 1998, Chu et al. 2004	P	<i>Hepialus lijiangensis</i> Chu & Wang, 1985
<i>Thitarodes litangensis</i> (Liang, 1995)	Sichuan Province: Litang [‡] and Batang County; Tibet Autonomous Region	4300–4700	Liang 1995, Yang 1996, Yang 1998	P	<i>Hepialus litangensis</i> Liang, 1995
<i>Thitarodes luquensis</i> (Yang & Yang in Yang et al., 1995)	Gansu Province: Luqu County [‡]	4276–4300	Yang et al. 1995, Yang 1998	P	<i>Hepialus luquensis</i> Yang & Yang, 1995

Insect name	Geographic distribution	Altitude (m)	Main references	Status of host insect [†]	Different name in the main references
<i>Thitarodes markamensis</i> (Yang, Li & Shen, 1992)	Tibet Autonomous Region: Markam County [‡] ; Yunnan Province: Deqên County	4500–4900	Yang et al. 1992a,b, Yang 1998	P	<i>Hepialus markamensis</i> Yang, Li & Shen, 1992
<i>Thitarodes meiliensis</i> (Liang in Liang et al., 1988)	Yunnan Province: Deqên County [‡]	3650–4700	Liang et al. 1988, Wang et al. 1996, Yang 1998	P	<i>Hepialus meiliensis</i> Liang, 1988
<i>Thitarodes menyuanicus</i> (Chu & Wang, 1985)	Gansu Province: Jishishan County; Qinghai Province: Hualong, Menyuan [‡] and Tongren Counties	—	Chu and Wang 1985a, Ma et al. 1995, Yang 1998	I	<i>Hepialus menyuanicus</i> Chu & Wang, 1985
<i>Thitarodes nebulosus</i> (Alpheraky, 1889)	Qinghai Province: Yushu Prefecture; Tibet Autonomous Region: Amdo [‡] and Damxung Counties [‡] , Nagqu Prefecture	4500	Yin 1987, Yang 1998, Chu et al. 2004	P	<i>Hepialus nebulosus</i> Alpheraky, 1889
<i>Thitarodes oblifurcus</i> (Chu & Wang, 1985)	Qinghai Province: Yushu Prefecture [‡] ; Sichuan Province: Kangding County	4000–4500	Chu and Wang, 1985a, Gao et al. 1992, Yang 1998	P	<i>Hepialus oblifurcus</i> Chu & Wang, 1985
<i>Thitarodes pratensis</i> (Yang, Li & Shen, 1992)	Yunnan Province: Deqên County [‡]	4350	Yang et al. 1992a	P	<i>Hepialus pratensis</i> Yang, Li & Shen, 1992
<i>Thitarodes renzhiensis</i> (Yang in Yang et al., 1991)	Yunnan Province: Deqên County [‡]	3880–5100	Yang et al. 1991a, Wang et al. 1996, Yang 1998	P	<i>Hepialus renzhiensis</i> Yang, 1991
<i>Thitarodes sichuanus</i> (Chu & Wang, 1985)	Sichuan Province [‡] : Aba Prefecture, Emei and Kangding Counties; Chongqing Municipality	3600–3800	Chu and Wang 1985a, Wang et al. 1996, Yang 1998, Chu et al. 2004	P	<i>Hepialus sichuanus</i> Chu & Wang, 1985
<i>Thitarodes varians</i> (Staudinger, 1896)	Sichuan Province: Batang County; Tibet Autonomous Region: Qamdo County	4500	Yin 1987, Yang 1998	P	<i>Hepialus varians</i> Staudinger, 1896
<i>Thitarodes xizangensis</i> (Chu & Wang, 1985)	Tibet Autonomous Region: Nyalam County [‡]	2200	Chu and Wang 1985a, Wang et al. 1996	N	<i>Forkalus xizangensis</i> Chu & Wang, 1985
<i>Thitarodes xunhuaensis</i> (Yang & Yang in Yang et al., 1995)	Qinghai Province: Xunhua County [‡]	3800	Yang et al. 1995, Yang 1998	P	<i>Hepialus xunhuaensis</i> Yang & Yang, 1995
<i>Thitarodes yeriensis</i> (Liang, 1995)	Yunnan Province: Deqên County [‡]	4500–4700	Liang 1995, Yang 1998	P	<i>Hepialus yeriensis</i> Liang, 1995
<i>Thitarodes yulongensis</i> (Liang, 1988)	Yunnan Province: Lijiang County [‡]	4150–4500	Liang et al. 1988, Wang et al. 1996, Yang 1998	P	<i>Hepialus yulongensis</i> Liang, 1988
<i>Thitarodes yunlongensis</i> (Chu & Wang, 1985)	Yunnan Province: Yunlong [‡] and Dali Counties; Hainan Province	3600–4200	Chu and Wang 1985a, Wang et al. 1996, Yang 1998, Chu et al. 2004	P	<i>Hepialus yunlongensis</i> Chu & Wang, 1985

Insect name	Geographic distribution	Altitude (m)	Main references	Status of host insect [†]	Different name in the main references
<i>Thitarodes yunnanensis</i> (Yang, Li & Shen, 1992)	Yunnan Province: Jianchuan [‡] , Lanping [‡] , Lijiang [‡] and Weixi Counties	3600–4100	Yang et al. 1992a,b, Wang et al. 1996, Yang 1998	P	<i>Hepialus yunnanensis</i> Yang, Li & Shen, 1992
<i>Thitarodes yushuensis</i> (Chu & Wang, 1985)	Qinghai Province: Yushu Prefecture [‡] , Batang, Chindu and Zadoi Counties; Gansu Province	4500–4900	Chu and Wang, 1985a, Yang et al. 1991b, Yang 1998, Ma et al. 1995	P	<i>Hepialus yushuensis</i> Chu & Wang, 1985
<i>Thitarodes zaliensis</i> (Yang, 1994)	Tibet Autonomous Region: Markam County [‡]	4600–4900	Yang 1994, Yang 1998	P	<i>Hepialus zaliensis</i> Yang, 1994
<i>Thitarodes zhangmoensis</i> (Chu & Wang, 1985)	Tibet Autonomous Region: Nyalam County [‡]	2200	Chu and Wang 1985a, Wang et al. 1996	N	<i>Hepialus zhangmoensis</i> Chu & Wang, 1985
<i>Thitarodes zhayuensis</i> (Chu & Wang, 1985)	Tibet Autonomous Region: Zayü [‡] and Markam Counties; Yunnan Province: Deqên and Gongshan County	4200–4400	Chu and Wang 1985a, Yang et al. 1987, Yang 1998	P	<i>Hepialus zhayuensis</i> Chu & Wang, 1985
<i>Thitarodes zhongzhiensis</i> (Liang, 1995)	Yunnan Province: Deqên County [‡]	4000–4600	Liang 1995, Wang et al. 1996	P	<i>Hepialus zhongzhiensis</i> Liang, 1995
<i>Triodia nubifer</i> (Lederer, 1853)	Sichuan Province: Kangding Prefecture; Central Asia	—	Chu et al. 2004	I	<i>Hepialus nubifer</i> Lederer, 1853
<i>Triodia sylvina</i> (Linnaeus, 1761)	Sichuan Province: Kangding County; Central Asia; Central Europe and Northern Europe	—	Chu and Wang 1985a, Chu et al. 2004, Karsholt and Nieuwerkerken 2010	I	<i>Hepialiscus sylvinus</i> (Linnaeus, 1761)

[†] The status of host insect of *O. sinensis* determined in this study: I = indeterminate host, N = non-host, IN = invalid name, P = potential host;

[‡] Type-locality;

[§] Names not included in Nielsen et al. (2000);

[†] Stem-borers;

^{*} The lowest altitude of the reported locality in China.

Discussion

Through an extensive literature survey, all the Hepialidae species reported from China were listed and analyzed using detailed information on their geographic distribution, altitude and nomenclature. The relationships between the insect species and *O. sinensis* were clarified based on available information. The data provided here serve as a foundation for further investigations on the conservation biology of this endangered fungal species and its insect hosts.

Species in different genera of Chinese hepialids can be divided into two categories according to the feeding strategy of the larvae (Chu and Wang 1985a, b; Chu et al. 2004): stem-borers (12 taxa) and root-borers (79 taxa). The stromata of *O. sinensis* are produced directly on the dead larvae of hepialids which were tunneling under the ground (Wang 1995, Yao 2004, Sung et al. 2007), and the host larvae of the fungus feed on plant roots underground (Chen et al. 1973, Shen et al. 1983, Wang 1995, Yao 2004). Therefore, the stem-borers, including nine in *Endoclita*, one in *Palpifer* and two in *Sthenopis* (Table 1), apparently can not be hosts of *O. sinensis*. The remaining 79 taxa found in this survey were categorized as potential hosts, indeterminate hosts, or non-hosts assessed based on the criteria described in the methods. Fourteen of the 79 root-borers were ruled out as hosts of *O. sinensis*, including 1 *Bipectilus*, 1 *Gazoryctra*, 2 *Hepialiscus*, 2 *Hepialus*, 4 *Napialus*, 1 *Parahepialiscus* and 3 *Thitarodes* species (Table 1), because they have not been reported from the distribution area of *O. sinensis* and were found below the elevation of 3000 m, either far away from the Tibetan Plateau (12 species), e.g., *Bipectilus zhejiangensis* from Zhejiang Province, *Hepialus hainanensis* from Hainan Province, etc., or on the Plateau (two species), e.g., *Thitarodes xizangensis* and *T. zhangmoensis*, which were found in Zhangmu Town in Tibet Autonomous Region, where the altitude range is from 1700 to 2400 m (People's Government of Tibet Autonomous Region, 2011) and no evidence for the occurrence of *O. sinensis* has been found (Li et al. in press).

Eight species, including 1 *Hepialiscus*, 4 *Hepialus*, 1 *Thitarodes* and 2 *Triodia* species (Table 1), are considered as indeterminate hosts of *O. sinensis*. While the distribution ranges of these species are within that of *O. sinensis*, they lack an altitude record and require further confirmation before being considered as potential hosts of *O. sinensis*, e.g., *Hepialus yadongensis*, *Triodia sylvina*, etc.

Fifty-seven taxa are recognized as potential hosts of *O. sinensis*, including 1 *Bipectilus*, 1 *Endoclita*, 1 *Gazoryctra*, 12 *Hepialus*, 2 *Magnificus*, 3 *Pharmacis* and 37 *Thitarodes* species (Table 1). The distribution ranges of these insects overlap that of *O. sinensis*. Altitude information for these insects was reported in three ways in the literature: (1) The altitude range of the insect was reported unambiguously above 3000 m, e.g., *Thi-*

tarodes baimaensis, *T. meiliensis*, etc. (37 species). Among these, the lowest altitude of 3200 m was reported for *T. cingulatus* (Yang 1998). (2) The altitude range of the species was not specified, but the types were collected at an elevation above 3000 m, e.g., *T. baqingensis*, *Magnificus jiuzhiensis*, etc. (11 species). The lowest altitude of the type locality is at 3200 m for *B. yunnanensis* (Chu and Wang 1985a). (3) There is no data reported on the altitude range for the species or the type specimen, but the altitude of the recorded localities of the moth were above 3000 m, e.g., *Hepialus gangcaensis*, *Pharmacis carna*, etc. (nine species). The lowest altitude for the locality of this group is 3050 m for *P. carna* (Chu et al. 2004) in Luhuo County, Sichuan Province, where the occurrence *O. sinensis* was confirmed (Li et al. in press).

Three names of the recognizable potential host insects are invalid (*nomen nudum*) because no full description of the species was published in the literature, although the names appeared several times in various publications (Table 1). Among them, *Thitarodes dongyuensis* was described by Yang (1992) as '*Hepialus dongyuensis*' and deemed as a *nomen nudum* in Nielsen et al. (2000), while *Hepialus guidera* and *H. lagii* were described by Yan (2001a, b) and recognized as *nomen nudum* in the present study. Further study is required to describe these species in full.

Species of *Hepialus* and *Phassus* described from China after 1984 have been transferred to *Thitarodes* and *Endoclita* respectively by Nielsen et al. (2000). Most of these species were described on the male genitalia and occasionally venation of one or very few individuals but not all morphological characteristics of the adult (Nielsen et al. 2000). However, the structure of the valve on male genitalia was still employed recently as the sole basis for classification in the revision of Chinese *Hepialus* by Zou et al. (2010). Further, disparate and incongruent regional taxonomies were regarded as developing rapidly for the Chinese Hepialidae (Nielsen et al. 2000), but the situation has not been changed much. As seen in this study, 24 names listed in Table 1 were not included in Nielsen et al. (2000). Two of them were described pre-2000 and apparently missed by Nielsen and his colleagues, while the remaining 22 were newly described after the year 2000 (Table 1). It seems that further study, especially robust phylogenetic hypotheses from molecular data, of these taxa is required to clarify their taxonomic status and generic placement.

Natural production of *O. sinensis* has been declining significantly over the last few decades while the market demands on the fungus have increased sharply in recent years. Clarification of the host insects of *O. sinensis* will provide basic information for management of the insect resources and for the conservation and sustainable use of the fungus. This work has gathered the available information on the host insects of *O. sinensis* and will lay a foundation for further studies of the relationship between the fungus and its hosts, especially their co-evolution (an ongoing research project based on DNA sequence analyses in this laboratory), and also for the cultivation of this valuable fungus for massive production.

Acknowledgements

This work is supported by the National Science and Technology Supporting Projects operated by the Ministry of Science and Technology of the People's Republic of China (2007BAI32B03), the Key Research Project of Innovation Programmes (KSCX2-YW-G-076, KSCX2-YW-G-074-04, KSCX2-SW-101C) and the scheme of Introduction of Overseas Outstanding Talents operated by the Chinese Academy of Sciences, and the National Science Funds for Distinguished Young Scholars from the National Natural Science Foundation of China (30025002).

References

- Chen LG, Wang ZY (2006) A new record species of *Napialus* (Lepidoptera: Hepialidae) in Shanghai. In: Cheng ZM (Eds) Science and Technology Innovation and Green Plant Protection. Agricultural Science and Technology Press, Beijing, 303–305. [In Chinese]
- Chen SJ, Jin SY (1992) Research survey on hosts of *Cordyceps sinensis* in China. Lishizhen Medicine and Materia Medica Research 3: 37–39. [In Chinese]
- Chen TL, Tang JJ, Mao JL (1973) A preliminary study on the biology of the 'insect herb', *Hepialus armoricanus* Oberthür. Acta Entomologica Sinica 16 (2): 198–202. [In Chinese]
- Chu HF (1965) The host insect of Chinese 'insect herb', *Hepialus armoricanus* Oberthür. Acta Entomologica Sinica 14 (6): 620–621. [In Chinese]
- Chu HF, Wang LY (1985a) 'Insect-herb' versus Hepialids with descriptions of new genera and new species of Chinese Hepialidae. Sinozoologia 3: 121–134. [In Chinese with English abstract]
- Chu HF, Wang LY (1985b) On the stem-borers of Chinese hepialids (Lepidoptera: Hepialidae). Acta Entomologica Sinica 28 (3): 293–301. [In Chinese with English abstract]
- Chu HF, Wang LY, Han HX (2004) Fauna Sinica, volume 38, Lepidoptera: Hepialidae, Epiplemlidae. Science Press, Beijing, 291 pp. [In Chinese with English abstract]
- Dong CH, Yao YJ (2008) In vitro evaluation of antioxidant activities of aqueous extracts from natural and cultured mycelia of *Cordyceps sinensis*. LWT-Food Science and Technology 41 (4): 669–677. doi:10.1016/j.lwt.2007.05.002.
- Dong YC, Luo DQ (1996) Research on *Cordyceps sinensis* in China. Territory & Natural Resources Study (3): 47–51. [In Chinese]
- Fu SQ, Huang TF, Chen SJ, Yin DH, Luo QM, Zha X, Zhama CR, Sun JH (2002) A new species of *Hepialus* (Lepidoptera: Hepialidae). Acta Entomologica Sinica 45 Suppl: 56–57. [In Chinese with English abstract]
- Fu SQ, Huang TF, Luo QM (1991) A new species of *Hepialus* (Lepidoptera: Hepialidae). Acta Entomologica Sinica 34 (3): 362–363. [In Chinese with English abstract]

- Gao ZX, Chen J, Zhao ZH, Song HM (1992) Study on the main host swiftmoth, *Hepialus oblifurcus* Chu et Wang, of caterpillar fungus, *Cordyceps sinensis* (Berkeley) Sacc., in Kangding. *Acta Entomologica Sinica* 35 (3): 317–321. [In Chinese with English abstract]
- Hu P, Zha LS (2010) Records of Chinese Edible Insects. *Journal of Anhui Agricultural Sciences* 38 (1): 522–526. [In Chinese with English abstract]
- Huang JC (2006) Preliminary study on spatial distribution pattern of *Bipectilus zhejiangensis* larva. *Journal of Fujian Forestry Science and Technology* 33 (4): 138–141. [In Chinese with English abstract]
- Ji DB, Ye J, Li CL, Wang YH, Zhao J, Cai SQ (2009) Antiaging effect of *Cordyceps sinensis* extract. *Phytotherapy Research* 23 (1): 116–122. doi: 10.1002/ptr.2576.
- Jiang SJ (1989) Exploitation of host insects of *Cordyceps* fungi. *Edible Fungi* (4): 5–6. [In Chinese]
- Jiang SJ (1991) *Cordyceps* fungi and their host insects in China. *Edible Fungi of China* 10 (2): 35–37. [In Chinese]
- Jiang SJ (2001) Host insect resources of *Cordyceps sinensis* and *Cordyceps militaris*. *Special Economic Animal and Plant* 4 (5): 15. [In Chinese]
- Karsholt O, Nieukerken EJ van (eds.) (2010) *Lepidoptera, Moths*. Fauna Europaea version 2.4, <http://www.faunaeur.org>
- Koh JH, Kim JM, Chang UJ, Suh HJ (2003) Hypocholesterolemic effect of hot-water extract from mycelia of *Cordyceps sinensis*. *Biological & Pharmaceutical Bulletin* 26: 84–87. doi: 10.1248/bpb.26.84.
- Li CD, Yang DR, Shen FR (1993) A new species of the genus *Hepialus* from Yunnan, China (Lepidoptera: Hepialidae). *Acta Entomologica Sinica* 36 (4): 495–496. [In Chinese with English abstract]
- Li JM, Li FA (2004) Research progress in *Cordyceps sinensis*. *Qinghai Medical Journal* 34 (8): 56–61. [In Chinese]
- Li Y, Wang XL, Jiao L, Jiang Y, Li H, Jiang SP, Lhosumtseiring N, Fu SZ, Dong CH, Zhan Y, Yao YJ (in press) A survey of geographic distribution of *Ophiocordyceps sinensis*. *The Journal of Microbiology*.
- Li YL (1996) Progress in the research on *Cordyceps sinensis*. *Chinese Qinghai Journal of Animal and Veterinary Sciences* 26 (6): 37–39. [In Chinese]
- Li YL, Liu X, Xu HF, Wang HS, Yan L (2002) Progress and trends in the research on *Cordyceps sinensis* in Qinghai. *Qinghai Science and Technology* 9 (4): 15–17. [In Chinese]
- Liang XC, Yang DR, Shen FR, Long YC, Yang YX (1988) Four new species of the genus *Hepialus* (Ghost Moth) from Yunnan, China. *Zoological Research* 9 (4): 419–425. [In Chinese with English abstract]
- Liang XC (1995) Four new species of the genus *Hepialus* (Ghost Moth) from Yunnan, Sichuan of China (Lepidoptera: Hepialidae). *Zoological Research* 16 (3): 207–212. [In Chinese with English abstract]
- Liu F, Wu XL, Yin DH, Chen SJ, Zeng W (2005) Progress in biological studies on the host insects of *Cordyceps sinensis*. *Chongqing Journal of Research on Chinese Drugs and Herbs* (1): 45–52. [In Chinese]

- Liu F, Wu XL, Yin DH, Chen SJ, Zeng W (2006) Progress in the research on species and distribution of host insects of *Cordyceps sinensis*. *Chongqing Journal of Research on Chinese Drugs and Herbs* (1): 47–50. [In Chinese]
- Liu LP, Long QX, Zhou CQ (1995) The host insect of the *Cordyceps sinensis* fungus-Hepialids. *Natural Enemies of Insects* 17 (4): 184–190. [In Chinese with English abstract]
- Long QX (1992) Research on *Cordyceps sinensis* and its utilization. *Natural Enemies of Insects* 14 (2): 96–100. [In Chinese]
- Ma QL, Wang Z, Ma FQ (1995) Species identification of the host insects of *Cordyceps sinensis* in Gansu. *Gansu Agricultural Science and Technology* (12): 35. [In Chinese]
- Nielsen ES (1988) The peculiar Asian ghost moth genus *Bipectilus* Chu & Wang: taxonomy and systematic position (Lepidoptera: Hepialidae s.str.). *Systematic Entomology* 13 (2): 171–195. doi: 10.1111/j.1365-3113.1988.tb00240.x.
- Nielsen ES, Robinson GS, Wagner DL (2000) Ghost-moths of the world: a global inventory and bibliography of the Exoporia (Mnesarchaeoidea and Hepialoidea) (Lepidoptera). *Journal of Natural History* 34 (6): 823–878. doi: 10.1080/002229300299282.
- Pegler DN, Yao YJ, Li Y (1994) The Chinese ‘Caterpillar Fungus’. *Mycologist* 8 (1): 3–5.
- People’s Government of Tibet Autonomous Region (2011) <http://www.xznjw.gov.cn/?action-model-name-rkz-itemid-117>. [In Chinese]
- Shen FR, Zhou YS (1997) Two new species of the genus *Hepialus* from Yunnan, China (Lepidoptera: Hepialidae). *Acta Entomologica Sinica* 40 (2): 198–201. [In Chinese with English abstract]
- Shen NY, Zeng L, Zhang XC, Zhou ZR, San Z, Wei SL (1983) Study on feeding habits of *Hepialus armoricanus* larva. *Special Wild Economic Animal and Plant Research* (3): 23–24. [In Chinese]
- State Forestry Administration and Ministry of Agriculture, Order No. 4. 1999. The list of the wild plants under the state emphasized protection. http://www.gov.cn/gongbao/content/2000/content_60072.htm. [See also: CITES Management Authority of China and China Customers (2000) Announcement of Adjustment for Catalogue of Exporting and Importing Wild Animals and Plants. Beijing: CITES Management Authority of China and China Customers.] [In Chinese]
- Sung GH, Hywel-Jones NL, Sung JM, Luangsa-ard JJ, Shrestha B, Spatafora JW (2007) Phylogenetic classification of *Cordyceps* and the clavicipitaceous fungi. *Studies in Mycology* 57: 5–59. doi: 10.3114/sim.2007.57.01.
- Tu YQ, Ma KS, Zhang DL (2009) A New Species of the Genus *Hepialus* (Lepidoptera: Hepialidae) from China. *Entomotaxonomia* 31 (2): 123–126. [In Chinese with English abstract]
- Wang GD (1995) *Cordyceps* species: ecology, cultivation and application. Science and Technology Reference Press, Beijing, 307 pp. [In Chinese]
- Wang LY (1990) A new species of *Hepialus* (Lepidoptera: Hepialidae). *Sinozoologia* 7: 173–174. [In Chinese with English abstract]
- Wang LY, Zheng JJ, Chen JY (2001) A new species of *Bipectilus* from China (Lepidoptera: Hepialidae). *Acta Entomologica Sinica* 44 (3): 348–349. [In Chinese with English abstract]

- Wang W, Zhong YC, Zhou SN, Chen CT (1996) Studies on *Cordyceps sinensis* I. Confirmations of Chinese name and the insect-hosts. Supplement to the Journal of Sun Yatsen University (2): 50–54. [In Chinese with English abstract]
- Wu WW (1992) A new species of the genus *Napialus* (Lepidoptera: Hepialidae) from Chongqing, China. Entomotaxonomia 14 (1): 55–56. [In Chinese with English abstract]
- Wu Y, Sun H, Qin F, Pan Y, Sun C (2006) Effect of various extracts and a polysaccharide from the edible mycelia of *Cordyceps sinensis* on cellular and humoral immune response against ovalbumin in mice. Phytotherapy Research 20 (8): 646–652. doi: 10.1002/ptr.1921.
- Wu YL (1757, Reproduced in 1958) Renewed Materia Medica. Science, Technology and Health Press, Shanghai, 377 pp. [In Chinese]
- Wu YL, Ishurd O, Sun CR, Pan YJ (2005) Structure analysis and antitumor activity of (1→3)-beta-d-glucans (cordyglucans) from the mycelia of *Cordyceps sinensis*. Planta Medica 71 (4): 381–384. doi: 10.1055/s-2005-864111.
- Wu ZC (2007) Relations between the occurrence of *Bipectilus zhejiangensis* and the environment. Entomological Journal of East China 16 (2): 92–95. [In Chinese with English abstract]
- Yan L (2000) A new genus and two new species of the Hepialidae (Lepidoptera) from Qinghai, China. Journal of Qinghai University 18 (5): 1–6. [In Chinese with English abstract]
- Yan L (2001a) Relation between quantity of population and ecology factors in *Hepialus guidera* Yan. Qinghai Prataculture 10 (1): 1–3. [In Chinese with English abstract]
- Yan L (2001b) The taxonomic feature of *Hepialus lagii* larva (Lepidoptera: Hepialidae) from Qinghai. Journal of Qinghai University 19 (1): 5–8. [In Chinese with English abstract]
- Yan L (2001c) Dynamic for population system of *Hepialus guidera* Yan in Qinghai Province. Jiangsu Agricultural Sciences (5): 66–68. [In Chinese]
- Yang DR (1993) Two new species of the genus *Hepialus* from Yunnan, China (Lepidoptera: Hepialidae). Acta Zootaxonomica Sinica 18 (2): 184–187. [In Chinese with English abstract]
- Yang DR (1994) Four new species of the genus *Hepialus* from Yunnan and Xizang, China (Lepidoptera: Hepialidae). Zoological Research 15 (3): 5–11. [In Chinese with English abstract]
- Yang DR (1998) *Cordyceps sinensis*. In: Yang GH (Eds) Utilization and Industrialization of Insect Resources in China. China Agriculture Press, Beijing, 68–80. [In Chinese]
- Yang DR, Jiang CP (1995) Two new species of the genus *Hepialus* (Lepidoptera: Hepialidae) from north Xizang, China. Entomotaxonomia 17 (3): 215–218. [In Chinese with English abstract]
- Yang DR, Li CD, Shen FR (1992a) Three new species of the genus *Hepialus* from Yunnan and Xizang, China (Lepidoptera: Hepialidae). Zoological Research 13 (3): 245–250. [In Chinese with English abstract]
- Yang DR, Li CD, Shu C, Yang YX (1992b) Relationship between alpine meadow vegetation and *Hepialus* moths distribution. Southwest China Journal of Agricultural Sciences 5 (2): 68–73. [In Chinese with English abstract]

- Yang DR, Li CD, Shu C, Yang YX (1996) Studies on the Chinese species of the genus *Hepialus* and their geographical distribution. *Entomologica Sinica* 39 (4): 413–422. [In Chinese with English abstract]
- Yang DR, Long YC, Shen FR, Yang YX, Yu RQ, Lu Z, Chun S, Sina DJ (1987) Research on the ecology of Yunnan Hepialids I. Regional and ecogeographical distribution. *Zoological Research* 8 (1): 1–11. [In Chinese with English abstract]
- Yang DR, Shen FR, Yang YX, Liang XC, Dong DZ, Chun S, Sina DJ (1991a) Biological studies and notes on a new species of the genus *Hepialus* from Yunnan, China. *Acta Entomologica Sinica* 34 (2): 218–224. [In Chinese with English abstract]
- Yang DR, Shu C, Li CD, Shen FR, Yang YX (1991b) Comparative study of 4 Hepialids and their fungal pathogens. In: Botanical Society of China (Eds) *Study and Application of Entomogenous Fungi in China*. China Agricultural Science and Technology Press, Beijing, 69–73. [In Chinese with English abstract]
- Yang DR, Yang YX, Zhang SY (1995) Three new species of the genus *Hepialus* from Qinghai and Gansu, China (Lepidoptera: Hepialidae). *Acta Entomologica Sinica* 38 (3): 359–362. [In Chinese with English abstract]
- Yao YJ (2004) Conservation and rational use of the natural resources of *Cordyceps sinensis*. *Science News* (15): 28–29. [In Chinese]
- Yin DH, Zeng W, Ye DJ (1991) Research progress in host insects of *Cordyceps sinensis*. *Special Wild Economic Animal and Plant Research* (3): 28–31. [In Chinese]
- Yin XC (1987) Hosts and geographical distribution of the genus *Cordyceps* (Fr.) Link. In: Northwest Institute of Plateau Biology (Eds) *Acta Biologica Plateau Sinica* 6. Science Press, Beijing, 1–24. [In Chinese]
- Yu YX (2004) Studies on artificial culture of *Cordyceps sinensis*. *Journal of Fungal Research* 2 (2): 42–46. [In Chinese with English abstract]
- Zhang GQ, Huang YD, Bian Y, Wong JH, Ng TB, Wang HX (2006) Hypoglycemic activity of the fungi *Cordyceps militaris*, *Cordyceps sinensis*, *Tricholoma mongolicum* and *Omphalia lapidescens* in streptozotocin-induced diabetic rats. *Applied Microbiology and Biotechnology* 72 (6): 1152–1156. doi: 10.1007/s00253-006-0411-9.
- Zhang GR, Gu DX, Liu X (2007) A new species of *Hepialus* (Lepidoptera, Hepialidae) from China. *Acta Zootaxonomica Sinica* 32 (2): 473–476. [In Chinese with English abstract]
- Zhang ZH, Li YL, Xu HF, Ma SL (2009) Hatch Rate Experiment on Egg of *Hepialus lagii* in Qinghai. *Special Wild Economic Animal and Plant Research* 31 (1): 19–22. [In Chinese with English abstract]
- Zou ZW, Liu X, Zhang GR (2010) Revision of taxonomic system of the genus *Hepialus* (Lepidoptera, Hepialidae) currently adopted in China. *Journal of Hunan University of Science & Technology (Natural Science Edition)* 25 (1): 114–120. [In Chinese with English abstract]