

First record of *Scolopendrellopsis* from China with the description of a new species (Myriapoda, Symphyla)

Ya-Li Jin¹, Yun Bu¹

¹ Natural History Research Center, Shanghai Natural History Museum, Shanghai Science & Technology Museum, Shanghai, 200041, China

Corresponding author: Yun Bu (buy@sstm.org.cn)

Academic editor: Pavel Stoev | Received 12 June 2018 | Accepted 16 August 2018 | Published 10 October 2018

<http://zoobank.org/8737EB89-629A-4DA2-B998-5F46F2C5E4B5>

Citation: Jin Y-L, Bu Y (2018) First record of *Scolopendrellopsis* from China with the description of a new species (Myriapoda, Symphyla). ZooKeys 789: 103–113. <https://doi.org/10.3897/zookeys.789.27356>

Abstract

The genus *Scolopendrellopsis* Bagnall, 1913 is recorded from China for the first time and *Scolopendrellopsis glabrus* sp. n. is described and illustrated. The new species is characterized by the short central rod on head, third tergite complete, four kinds of sensory organs present on antenna, and the cerci rather short and covered with a low number of straight setae.

Keywords

antennal sensory organ, chaetotaxy, taxonomy, tergal process, Tömösváry organ

Introduction

There are 204 symphylian species known in the world to date (Szucsich and Scheller 2011; Domínguez Camacho and Vandenspiegel 2012; Bu and Jin 2018); however, only few publications deal with those from Asia. Hansen firstly described five species of Symphyla from Southeast Asia (Hansen 1903). After that several species were described from India (Scheller 1971), Indonesia (Scheller 1988), USSR (Scheller and Golovatch 1982), Russian Far East (Scheller and Mikhaljova 2000) and Iran (Scheller et al. 2011). Symphyla is poorly studied in China with only *Hanseniella caldaria* from Zhejiang province and *Geophilella orientalis* from Hebei province recorded (Zhang and Wang 1992; Bu and Jin 2018). Three genera, *Scutigerebella*, *Scolopendrelloides*, and *Symphylella*,

were also mentioned for China, but without determined species recorded (Zhang and Wang 1992). During our ecological survey of soil animals of Zhejiang, Jiangsu, and Hainan provinces in recent years, many symphylans were obtained. Among them, one new species of *Scolopendrellopsis* was identified and is described in the present paper.

Materials and methods

Most specimens were collected during a project for soil animal survey of Gutian Mountain of Zhejiang Province during the years 2012 to 2013; others were collected in Jiangsu province and Hainan province recently. All were extracted by means of the Tullgren funnels from soil and humus samples and preserved in 75% ethanol. They were mounted under slides using Hoyer's solution and dried in an oven at 60 °C. Observations were made with a phase contrast microscope (Leica DM 2500). Photographs were taken by a digital camera installed on the microscope (Leica DMC 4500). Line drawings were drawn using a drawing tube. All specimens are deposited in the collections of Shanghai Natural History Museum (SNHM) and Shanghai Entomological Museum (SEM), Shanghai, China.

Taxonomy

Family Scolopendrellidae Bagnall, 1913

Genus *Scolopendrellopsis* Bagnall, 1913, new record

Diagnosis. Habitus slender. First pair of legs present, 3-segmented and with claws, not more than one-half length of the following pairs. Trunk with 16 or 17 tergites and most of tergites with a pair of posterior processes, without any striped band between each pair of processes on tergites, some tergites transversely divided.

Distribution. The genus *Scolopendrellopsis* includes fifteen species and is subcosmopolitan, widely distributed in Palaearctic, Nearctic, Neotropical, Ethiopian, Oriental, and Australian regions (Szucsich and Scheller 2011). It is newly recorded from China in this paper.

Scolopendrellopsis glabrus sp. n.

<http://zoobank.org/95E5B444-5DEF-49CB-A699-E9730BD69528>

Figs 1–3. Tables 1–3

Diagnosis. *Scolopendrellopsis glabrus* sp. n. is characterized by the short central rod on head, 3rd tergite not divided and with only weak middle indentation, rod-like sensory

organs with setae surrounded on dorsal side of 3rd–17th antennal segments, cavity-shaped organs on dorsal side of subapical 5–6 antennal segments, mushroom-shaped organs at lateral side of subapical 4–7 segments and bladder-shaped organs on subapical 3–6 antennal segments, first pair of legs longer than the tarsus of the last pair of legs, cerci short and covered with a low number of straight setae.

Material examined. *Holotype*, female (slide no. ZJ-GTS-SY2012017) (SNHM), China, Zhejiang Province, Gutian Mountain, extracted from soil samples in broad-leaved forest, Alt. 1000 m, 29°15'N, 118°06'E, 11-IV-2012, coll. Y. Bu et al. *Paratypes*, 2 female (slides nos. ZJ-GTS-SY2012010, ZJ-GTS-SY2012016) (SNHM), same date as holotype; 1 female (slide no. ZJ-GTS-SY2012051) (SEM), *ibidem*, 14-X-2012; 2 females (slides nos. ZJ-GTS-SY2012055, ZJ-GTS-2012060) (SNHM), *ibidem*, 17-XI-2012; 1 female (slide no. ZJ-GTS-SY2013015) (SNHM), *ibidem*, 24-IV-2013; 1 male (slide no. JS-WX-SY2017001) (SNHM), China, Jiangsu Province, Wuxi, Daji Mountain, extracted from soil samples in bamboo forest, Alt. 5 m, 31°32'N, 120°12'E, 9-X-2017, coll. Y. Bu. **Other material** (SNHM): 8 juveniles with 8–10 pairs of legs (slides nos. ZJ-GTS-SY2012002, ZJ-GTS-SY2012004, ZJ-GTS-SY2012006, ZJ-GTS-SY2012012–ZJ-GTS-SY2012015, ZJ-GTS-SY2012019), same data as holotype; 1 juvenile with 10 pairs of legs (slide no. ZJ-GTS-SY2012023), *ibidem*, 19-VI-2012, coll. Y. Bu et al; 6 juveniles with 8–11 pairs of legs (slides nos. ZJ-GTS-SY2012028–ZJ-GTS-SY2012032, ZJ-GTS-SY2012039), *ibidem*, 15-VII-2012, coll. Y. Bu et al; 2 juveniles with 9 and 10 pairs of legs (slides nos. ZJ-GTS-SY2012046, ZJ-GTS-SY2012051), *ibidem*, 14-X-2012, coll. Y. Bu et al; 2 juveniles with 10 and 9 pairs of legs respectively (slides nos. ZJ-GTS-SY2012052, ZJ-GTS-SY2012055), *ibidem*, 17-XI-2012, coll. Y. Bu et al; 3 juveniles with 8–10 pairs of legs (slides nos. ZJ-GTS-SY2012064–ZJ-GTS-SY2012066), *ibidem*, 12-XII-2012, coll. Y. Bu et al; 1 juvenile with 8 pairs of legs (slide no. ZJ-GTS-SY2013004), *ibidem*, 23-II-2013, coll. Y. Bu et al; 6 juveniles with 8–10 pairs of legs (slides nos. ZJ-GTS-SY2013005, ZJ-GTS-SY2013009, ZJ-GTS-SY2013011, ZJ-GTS-SY2013013–ZJ-GTS-SY2013015), *ibidem*, 27-III-2013, coll. Y. Bu et al; 1 juvenile with 9 pairs of legs (slide no. HN-SY-SY2017001), China, Hainan Province, Sanya, Yalong bay tropical paradise forest park, extracted from soil samples in bamboo forest, Alt. 67 m, 18°15' N, 109°37'E, 22-III-2017, coll. Y. Bu.

Description. Adult body 1.57 mm long in average (1.45–1.65 mm, n = 8), holotype 1.65 mm (Figure 1A). *Head* longer than wide, length 145–175 µm, width 133–170 µm, with widest part a little behind the middle on a level with the points of articulation of mandibles. Central rod distinct and with anterior part absent, length 45–49 µm, approximately one-third of head. Dorsal side of head covered with sparse setae of different length, longest setae (12–17 µm) located at the anterior part of head, approx. 3.0 times as long as central ones (4–5 µm). Cuticle around Tömösváry organ and anterolateral part of head with rather coarse granulation. Central and posterior part of head with dense pubescence (Figs 1B, 3A).




Figure 1. *Scolopendrellopsis glabrus* sp. n. (Holotype) **A** habitus **B** head, dorsal view **C** right antenna, 3th–6th segments, dorsal view **D** right antenna, 12th–16th segments, ventral view **E** right antenna, 11th–16th segments, dorsal view **F** left Tömösváry organ **G** stylus on base of 6th leg (arrow indicated) **H** stylus on base of 11th leg (arrow indicated) **I** first pair of legs **J** 3rd leg and coxal sac **K** 9th leg and coxal sacs **L** cerci, dorsal view. ro-rod-like sensory organs with surrounded setae, co-cavity-shaped organ, mo-mushroom-shaped organ, bo-bladder-shaped organ. Scale bars: 100 μ m (**A**), 20 μ m (**B**–**L**).

Tömösváry organ oval, maximum diameter 17.0–22.5 μ m, somewhat shorter than the greatest diameter of 3rd antennal segment (20–23 μ m), opening at front position, with diameter (4–5 μ m) approx. one-fourth of 3rd segment of antennae (Figs 1F, 3A).

Mandible with eleven teeth and divided into two parts by a gap, with five anterior and six posterior teeth respectively. First maxilla has two lobes, inner lobe with four




Figure 2. *Scolopendrellopsis glabrus* sp. n. (Holotype) **A** first and second maxilla **B** 1st and 2nd tergite **C** 3rd tergite **D** 4th tergite **E** 6th tergite, left side **F** 8th tergite, left side **G** 9th tergite, left side **H** 12th tergite, left side **I** 14th tergite, left side **J** 15th tergite, left side. Scale bars: 20 μ m.

hook-shaped teeth, palp bud-like with two distal points close to outer lobe (Figure 3B). Anterior part of second maxilla with many small protuberances and posterior part with sparse setae. Cuticle of second maxilla covered with pubescence (Figure 2A).

Antennae 15–19 segments (16 in holotype), length 250–350 μ m (320 μ m in holotype), approx. 0.2 of the length of the body. First segment cylindrical, greatest diameter




Figure 3. *Scolopendrellopsis glabrus* sp. n. (Holotype) **A** head and 1st–4th tergites **B** mandible and first maxilla **C** 1st–5th segments of right antenna **D–E** 13th–16th of right antenna **D** dorsal view **E** ventral view **F** first leg **G** 12th leg **H** left sense calicles, dorsal view **I** right sense calicles, dorsal view **J** stylus on base of 11th leg **K** right cercus, dorsal view. Scale bars: 50 µm (**A**), 20 µm (**B–I, K**), 5 µm (**J**).

a little wider than long (20–26 µm: 16–25 µm), with four setae in one whorl, the longest seta (6–11 µm) inserted at the inner side and distinctly longer than outer ones (5–8 µm). Second segment wider (20–30 µm) than long (18–22 µm), with six or seven setae evenly inserted around the segment and inner setae (6–10 µm) a little longer than outer ones (5–7 µm). Chaetotaxy of 3rd segment similar to preceding ones (Figure 3C). Setae on the basal segments 1–3 are slender and on proximal and distal segments rather short. Basal and median parts of the antennae with only primary whorl of setae, in subapical

Table 1. Numbers of setae and sensory organs of antennae (holotype).

Segments	Nos. of primary whorl setae	Nos. of secondary whorl setae	Rod-like organ with setae surrounded	Cavity-shaped organs	Mushroom-shaped organs	Bladder-shaped organs	
						Dorsal	Ventral
1 st	4						
2 nd	6						
3 rd	7						
4 th	8						
5 th	8		1				
6 th	8						
7 th	8						
8 th	8						
9 th	9						
10 th	9	1	1	1	1		
11 th	9	1	1	1	1		
12 th	9	2		1	2	1	
13 th	9	2		1	2	1	
14 th	7	2	1	1	2	2	4
15 th	8	2	1	1	2	1	3
16 th	7	2					

segments one or two minute setae present in secondary whorl (Figure 3E). Four kinds of sensory organs present on antenna: rod-like sensory organs with setae surrounded present on dorsal side of 3rd–17th segments (Figs 1C, 3C, 3D); cavity-shaped organs present on dorsal side of subapical 5–6 segments (Figs 1E, 3D); mushroom-shaped organs present on lateral side of subapical 4–7 segments and bladder-shaped organs on subapical 3–6 segments (Figs 1D, 1E, 3D, 3E). Apical segment subspherical, width 21–22 μm , length 19–20 μm , with 10–12 short setae and wide connection to preceding segment and with two fire-shaped and three baculiform organs present on apex (Figs 1D, 3D, E). All segments covered with short pubescence. Chaetotaxy and sensory organs of antennae are given in Table 1.

Trunk: seventeen dorsal tergites present, with 6th, 9th, 12th, and 15th tergites transversely divided, longer than preceding ones (Figs 2E, 2G, 2H, 2J). Intertergal zones between former and later tergites present, except for 14th and 15th, 16th, and 17th tergites. Tergites 2th–13th and 15th each with one pair of slender chitinous processes, slightly finger-like. Basal distances between processes are approx. the same length as their length from base to tip, which is longer than its basal width. All tergites pubescent and the margins of apical part of processes ornamented with rowed coarse granules. Apical seta on processes slightly anteriorly located and anterolateral setae slightly longer than other setae. No seta between apical and inner basal setae (Figs 2B–2H).

Tergites: 1st tergite reduced to a narrow short plate with a pair of diagonal bands and with six short setae in a row (Figs 2B, 3A). Second tergite complete, broader than long, with two slender posterior processes, 1+1 axial setae and 7+6 lateral setae (asymmetrically lack one lateral seta in holotype, 7+7 lateral setae in all paratypes), with anterolateral setae slightly longer than others, processes approx. 1.5 times as long as

broad, basal distance between processes approx. the same as long as their length (Figs 2B, 3A). Third tergite entire with weak middle indentation, broader and longer than preceding one with the ratios mentioned nearly 1.6 and 0.8 respectively, 2+2 axial setae and 9+9 lateral setae (Figs 2C, 3A). Fourth tergite broader than 3rd tergite, with the ratios approx. 1.2 and 0.9 respectively, 2+2 axial setae and 6+6 lateral setae (Figs 2D, 3A). The shape and chaetotaxy of 5th–7th, 8th–10th, and 11th–13th tergites similar as 2nd–4th tergites. 14th tergite without processes and relevant area replaced by two roundish tubercles with four setae inserted on (Figure 2I). 15th tergite shorter than 3rd, 6th, 9th, and 12th tergites, with smaller processes (Figs 2C, 2E, 2G, 2H, 2J). Chaetotaxy and measurements of tergites are given in Tables 2 and 3.

Legs: all twelve pairs of legs with claws. 1st pair of legs short, 3-segmented, length 35–45 μm , not more than the length of 2th pair of legs, but longer than the tarsus (30–32 μm) of last pair of legs; femur at least 1.2 times wider than long (15–22 μm : 12–15 μm), with two setae at the outer side; tibia approx. 1.4 times wider than long (14–20 μm : 10–14 μm), with dorsal seta (8–10 μm) longer than ventral one (4–6 μm); tarsus longer than wide (12–19 μm : 10–17 μm), with four setae, three dorsal (5–7 μm) and one ventral (6–8 μm); claws simple and the anterior one a little larger and broader than posterior (Figs 1I, 3F). 12th pairs of legs approx. three-fourths as long as the length of the head. Trochanter longer than wide (30–40 μm : 23–31 μm), with 6 subequal setae; femur approx. as long as wide (19–25 μm : 19–25 μm), with three setae transversely, one (10–14 μm) distinctly longer than other two (6–9 μm); tibia longer than wide (19–25 μm : 15–21 μm), with four dorsal setae, of which one (10–14 μm) distinctly longer than others (6–9 μm); tarsus not more than 3 times as long as wide (30–32 μm : 11–15 μm) with 8–9 setae, of which 3 are protruding and 2 depressed, longest setae (12 μm) approx. as long as the greatest width of the joint. Claws rather curved, anterior one a little longer and broader than posterior one (10 μm : 8 μm) (Figure 3G). All legs covered with dense pubescence (Figs 1I, J, K).

Coxal sacs present at bases of 3rd–9th pairs of legs, fully developed, each with 3 setae (Figs 1J, 1K).

Styli present at base of 3rd–12th pairs of legs, reduced into small knobs with tuft of setae, on 9th–12th legs larger than on former legs, especially on 11th legs (5–6 μm), distinctly longer than anterior ones (2–4 μm) (Figs 1G, 1H, 3J).

Sense calicles with smooth margin to pit, length nearly two times longer than outer diameter (25–35 μm : 12–16 μm). Sensory seta inserted in the center of cup, extremely long, length 100–120 μm , at least 8.5 times longer than other two lateral setae (11–14 μm , 7–10 μm respectively) that inserted at the edge of cup (Figs 3H, I).

Cerci subuliform, short, approx. half as long as head, somewhat shorter than 12th pairs of legs, length at least three times as long as its greatest width (80–95 μm : 25–30 μm), sparsely covered with long and straight setae, with the longest one (12.5–17 μm) approx. half of the greatest width of the cerci, terminal area (10–13 μm) short, with length at most half of the greatest width of the cerci and circled by 6–8 layers of curved ridges. Terminal setae length 18–20 μm , distinctly longer than terminal area (Figure 1L, 3K).

Table 2. Chaetotaxy of tergites (holotype).

Tergites	Axial setae	Lateral setae
1 st	3+3	–
2 nd	1+1 ¹	7+6 ⁸
3 rd	2+2 ²	9+9 ⁹
4 th	2+2	6+6
5 th	2+2	5+5
6 th	3+3	9+9 ¹⁰
7 th	2+2 ³	6+6
8 th	2+2 ⁴	5+5
9 th	3+3	9+9 ¹¹
10 th	2+2	6+6
11 th	2+2 ⁴	5+5
12 th	3+3 ⁴	9+9 ¹²
13 th	2+2	6+6 ¹³
14 th	3+3 ⁵	4+4 ¹⁴
15 th	3+3 ⁶	7+7 ¹⁵
16 th	1+1 ⁷	2+2 ¹⁶
17 th		5+5 ¹⁷

Notes on chaetotaxy variations: ¹ with single middle seta (in 5 specimens) or without setae (1); ² asymmetrically lack one seta (3) or lack one pair of seta (1). ³ with 1+1 setae (1); ⁴ asymmetrically present of one additional seta (1); ⁵ with 2+2 setae (5); ⁶ with 1+1 or 2+2 setae (4); ⁷ asymmetrically present of one additional setae (1); ⁸ with 7+7 setae in all paratypes (7); ⁹ asymmetrically lack 1 or 2 setae (2); ¹⁰ with 7+7 setae (1); ¹¹ asymmetrically present of one additional setae (1); ¹² asymmetrically lack one or present of one additional seta (2), or with 10+10 setae (1); ¹³ with 4+4 setae (2); ¹⁴⁻¹⁶ asymmetrically present of one additional seta (2); ¹⁷ asymmetrically lack one or two seta (2).

Table 3. Measurements of tergites and processes (holotype in brackets) (in μm).

No. of tergites	Length	Width	Length of processes	Width of processes	Basal distance between processes
1 st	30–40 (40)	75–100 (80)	–	–	–
2 nd	45–55 (55)	78–100 (100)	25–45 (31)	15–35 (20)	25–35 (30)
3 rd	50–85 (82)	95–130 (112)	30–48 (40)	23–30 (25)	28–35 (33)
4 th	45–61 (61)	100–125 (122)	30–50 (50)	30–40 (40)	35–45 (45)
5 th	41–59 (59)	103–125 (107)	50–55 (50)	25–30 (30)	40–50 (40)
6 th	55–123 (116)	130–150 (140)	40–58 (55)	25–35 (35)	40–55 (55)
7 th	60–71 (71)	140–160 (160)	50–65 (50)	40–50 (50)	40–66 (66)
8 th	55–75 (75)	110–135 (110)	50–65 (55)	25–35 (33)	58–68 (68)
9 th	100–110 (110)	150–170 (160)	60–70 (60)	25–40 (30)	70–80 (80)
10 th	67–71 (71)	150–187 (166)	40–70 (70)	30–50 (50)	45–74 (74)
11 th	60–75 (68)	130–138 (138)	30–65 (50)	33–40 (33)	55–70 (60)
12 th	75–120 (120)	150–165 (160)	54–60 (55)	25–40 (33)	55–65 (64)
13 th	45–69 (69)	110–180 (156)	30–50 (50)	25–50 (50)	50–60 (60)
14 th	55–80 (65)	95–140 (140)	–	–	–
15 th	70–98 (85)	110–150 (140)	38–45 (40)	25–30 (30)	35–50 (42)
16 th	35–43 (38)	95–130 (115)	–	–	–
17 th	58–65 (65)	75–125 (90)	–	–	–

Etymology. The species name *glabrus*, meaning bald, to indicate the lower number of setae on cerci.

Distribution. China (Zhejiang, Jiangsu, Hainan).

Remarks. *Scolopendrellopsis glabrus* sp. n. is similar to *S. hirta* (Scheller, 1971) and *S. spinosa* (Sheller, 1979) in the shape of 3rd tergite which is not divided, shape of processes on tergites, shape of sensory organs on antennae. It differs from the latter two species in the absence of anterior part of central rod (anterior part present but indistinct in *S. hirta*, distinct in *S. spinosa*), chaetotaxy of the 2nd and 3rd tergites (with four and five lateromarginal setae in *S. glabrus* respectively, five and six in the other two species), cerci with lower number of setae (more setae in *S. hirta* and *S. spinosa*), all setae on cerci long and straight (setae on inner side of cerci slightly curved in *S. hirta*, most setae on cerci short and curved in *S. spinosa*). It is also similar to the worldwide species *S. subnuda* in the shape of the first three tergites, number of lateromarginal setae of the 3rd tergite, shape and number of setae of the cerci, but differs in the absence of anterior part of central rod (anterior part present in *S. subnuda*), apical seta on processes slightly anteriorly located (rather close to the apex in *S. subnuda*).

Acknowledgements

We give our great gratitude to Dr. Xin Ke for his fund supports during the collection in Gutian Mountain. We also appreciate Professor José G. Palacios-Vargas (Mexico) for his linguistic corrections of the manuscript as well as his valuable advices. We specially thank Professor Hong Ying Song (China) and Dr. Miguel Domínguez Camacho (Spain) for their careful revisions and valuable comments that greatly improved our manuscript. This study was supported by the National Natural Science Foundation of China (no. 31772509), the Natural Science Foundation of Shanghai (no. 17ZR1418700), the project for construction of Chinese forest biodiversity monitoring network (no. CforBio20120106) and the strategic biological resources service network project of the Chinese Academy of Sciences (no. ZSBZ-003).

References

- Bagnall RS (1913) On the classification of the order Symphyla. *Journal of the Linnean Society London Zoology* 32: 195–199. <https://doi.org/10.1111/j.1096-3642.1913.tb01775.x>
- Bu Y, Jin YL (2018) Progress on the systematic study of Symphyla. *Chinese Bulletin of Life Sciences* 30(5): 500–509. [In Chinese with English summary]
- Domínguez Camacho M, Vandenspiegel D (2012) Scolopendrellidae (Myriapoda, Symphyla) from the Afrotropics with descriptions of seven new species. *European Journal of Taxonomy* 32: 1–28. <https://doi.org/10.5852/ejt.2012.32>
- Edwards CA (1959) Keys to the genera of the Symphyla. *Journal of the Linnean Society Zoology* 44: 164–169. <https://doi.org/10.1111/j.1096-3642.1959.tb01603.x>

- Hansen HJ (1903) The genera and species of the order Symphyla. *Quarterly Journal of Microscopical Science* 47: 1–101.
- Michelbacher AE (1938) The biology of the Garden Centipede, *Scutigereilla immaculate*. *Hilgardia* 11: 55–148. <https://doi.org/10.3733/hilg.v11n03p055>
- Scheller U (1971) Symphyla from Ceylon and Peninsular India. *Entomologica Scandinavica Supplement* 1: 98–187.
- Scheller U (1979) The Pauropoda and Symphyla of the Geneva Museum VI. Symphyla from Rhodesia and South Africa (Myriapoda). *Revue Suisse de Zoologie* 86(4): 947–955. <https://doi.org/10.5962/bhl.part.82352>
- Scheller U (1988) Two new species of Symphyla from the Krakatau Islands and the Ujung Kulon Peninsula (Myriapoda: Symphyla: Scolopendrellidae, Scutigereillidae). *Zoological Expedition to the Krakatau Islands 1984 and 1985. Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* 322(1211): 401–411. <https://doi.org/10.1098/rstb.1988.0134>
- Scheller U, Golovatch SI (1982) Myriapods of the class Symphyla in the USSR. *Zoologicheskii Zhurnal* 61(1): 143–145. [In Russian with English summary]
- Scheller U, Kavianpour MR, Esfandiari M (2011) First record of Symphyla (Myriapoda) from Iran, with description of a new species in *Scolopendrellopsis* (Scolopendrellidae). *Zootaxa* 3041: 66–68.
- Scheller U, Mikhaljova EV (2000) New records of Symphyla (Myriapoda) from the Russian Far East. *Arthropoda Selecta* 9(1): 29–30.
- Szucsich N, Scheller U (2011) Symphyla. In: Minelli A (Ed.) *Treatise on Zoology-Anatomy, Taxonomy, Biology: The Myriapoda* (Vol. 1). Brill, Leiden, 445–466. https://doi.org/10.1163/9789004188266_021
- Zhang CZ, Wang DQ (1992) Symphyla. In: Yin WY, et al. (Eds) *Subtropical Soil animals of China*. Science Press, Beijing, 383–387. [In Chinese]