

# ***Ptychoptera deleta* Novák, 1877 from the Early Miocene of the Czech Republic: redescription of the first fossil attributed to Ptychopteridae (Diptera)**

Wiesław Krzemiński<sup>1</sup>, Jakub Prokop<sup>2</sup>

**1** *Institute of Biology, Pedagogical University of Cracow, ul. Podbrzezie 3, 31-054 Kraków, Poland* **2** *Charles University in Prague, Faculty of Science, Department of Zoology, Viničná 7, CZ-128 44, Praha 2, Czech Republic*

Corresponding author: *Jakub Prokop* (jprokop@natur.cuni.cz)

---

Academic editor: *D. Shcherbakov* | Received 18 April 2011 | Accepted 20 July 2011 | Published 24 September 2011

---

**Citation:** Krzemiński W, Prokop J (2011) *Ptychoptera deleta* Novák, 1877 from the Early Miocene of the Czech Republic: redescription of the first fossil attributed to Ptychopteridae (Diptera). In: Shcherbakov DE, Engel MS, Sharkey MJ (Eds) *Advances in the Systematics of Fossil and Modern Insects: Honouring Alexandr Rasnitsyn*. ZooKeys 130: 299–305. doi: 10.3897/zookeys.130.1401

---

## **Abstract**

The first fossil that was described in Ptychopteridae, *Ptychoptera deleta* Novák, 1877 from the classical Early Miocene locality Mokřina (Krottensee) in western Bohemia is re-examined. The re-description of the holotype including a new line drawing and remarks summarizing the scarce fossil record of this group is provided.


## **Keywords**

Diptera, Ptychopteridae, Ptychoptera, revision, Tertiary, Neogene, Miocene, Ottnangian/Karpatian, Cypris Formation, Cheb Basin, Czech Republic

## **Introduction**

The family Ptychopteridae is a very small group with about 70 Recent species belonging to two subfamilies (Ptychopterinae and Bittacomorphinae), mostly distributed in the Holarctic, Ethiopian and Oriental Regions. In the Neotropics one species is currently known (Hancock et al. 2006). Ptychopteridae are not present in the Australian and Antarctic Regions. The oldest true representative, *Ptychoptera mesozoica* Kalugina, 1989 was described from the Lower Cretaceous (Neocomian) of Baissa in Buryatia (Siberia, Russia). *Ptychoptera deleta* Novák, 1877, known from the Early Miocene of Mokřina (Krot-

tensee) in western Bohemia (Czech Republic), was the first described fossil representative of the family. Handlirsch (1909) supposed that the species did not belong to the genus *Ptychoptera* Meigen, 1803 and created a new genus *Ptychopterula* mainly on the basis of a considerably narrow wing base as present in *Etoptychoptera* Handlirsch, 1909 and a free Sc. However, he emphasized also the common characters present in *Ptychopterula* and the Recent genera *Ptychoptera* and *Bittacomorpha* Westwood, 1835 connection of R and Rs, the position of crossvein r-m, etc. Alexander (1927) also supposed that the species described by Novák did not belong to *Ptychoptera*. Peus (1958), in his monograph on the Ptychopteridae, referred to this species as *Liriope* (“*Ptychopterula*”) *deleta* (Novák, 1877). These opinions were based on mistakenly Novák’s drawing only, without revision of the holotype (see Fig. 2B). This paper presents a re-description of the holotype including a new line-drawing and photograph. Five other compressed fossil ptychopterid species were described till now: *Ptychoptera miocenica* (Cockerell, 1910) from the Oligocene of Florissant (Colorado, USA) originally placed in genus *Bittacomorpha*; *Brodilka mitchelli* Lukashovich, Coram & Jarzembowski, 2001 and *Zhiganka woolgari* Lukashovich, Coram & Jarzembowski, 2001 both from the Lower Cretaceous of Purbeck and Wealden groups (UK); *Zhiganka comitans* Lukashovich, 1995 from the Lower Cretaceous of Yakutia (Russia), and *Probittacomorpha christenseni* Freiwald & Willmann, 1992 from



**Figure 1.** **A** Geographical position of northwestern Bohemia within Europe **B** detailed map of the Cheb and Sokolov basins with indication of position of Mokřina (Krottensee) locality.

the lowermost Eocene of Mo-clay (Denmark) (Freiwald and Willmann 1992; Evenhuis 1994; Lukashovich 1995; Ansoerge and Schröder 1999; Lukashovich et al. 2001). Two genera, *Zhiganka* and *Probittacomorpha* are attributed to the subfamily Bittacomorphinae and others to Ptychopterinae. A single species, *Ptychoptera eocenica* Podenas, 2007 was described from an Eocene Baltic amber inclusion (Podenas 2007).

Lukashovich (2008) proposed a new system of Ptychopteroidea and synonymized the family Eoptychopteridae known from the Upper Triassic to Lower Cretaceous with Ptychopteridae. Eoptychopteridae is a considerably variable group of flies with common occurrence throughout the Jurassic up to the Lower Cretaceous (Kalugina 1989; Krzemiński 1992; Lukashovich 1993; Lukashovich et al. 1998; Ren and Krzemiński 2002; Krzemiński and Krzemińska 2003).

Novák (1877) described a fossil entomofauna from Mokřina (Krottensee) located in Cheb county of western Bohemia (Czech Republic) (Fig. 1). This classical Early Miocene locality in Cheb Basin yielded greenish marls of lake sedimentation belonging to the Cypris Formation (Rojík 2004). Fossiliferous layers previously called "Cypris shales" are well known for abundant occurrence of ostracods e.g. *Cypris angusta* (Reuss, 1852) when Cheb and Sokolov basins were interconnected (Obrhelová and Obrhel 1987). Mokřina (Krottensee) locality belongs to biostratigraphic zone MN4 dated by mammals from nearby locality Dolnice (Fejfar 1974), and to ichthyozone IV defined by Obrhelová and Obrhel (1983). Fossil record from Mokřina is also well known by plants and bird remains (Bůžek et al. 1996; Mlíkovský 1996). Novák's collection from Mokřina housed in National Museum in Prague contains 73 insect specimens classified in 10 families of five insect orders (Prokop et al. 2003).

## Material and methods

The holotype specimen was observed under stereomicroscope Leica MZFLIII & MZ16 in dry state. The line drawing of the venational pattern was drawn directly using a stereomicroscope and finally readjusted to the photograph scale using image-editing software (Adobe Photoshop CS). Photography was made simultaneously using a digital camera Canon PowerShot S80 attached to the stereomicroscope. The type material referred to as 'NM' is housed at the National Museum in Prague, Czech Republic.

## Systematic paleontology

### Family Ptychopteridae Osten-Sacken, 1879

### Genus *Ptychoptera* Meigen, 1803

**Type species.** *Tipula contaminata* Linnaeus, 1758

***Ptychoptera deleta* Novák, 1877**

[http://species-id.net/wiki/Ptychoptera\\_deleta](http://species-id.net/wiki/Ptychoptera_deleta)

Figs 2A–C


*Ptychoptera deleta* Novák, 1877: p. 88, Pl. II, Fig.1.

*Ptychopterula deleta* (Novák, 1877): Handlirsch (1909): p. 264, 269.

*Liriope* (“*Ptychopterula*”) *deleta* (Novák, 1878): Peus (1958):p. 12, fig. 17.

**Diagnosis.** Rs very short; R4+5 five times longer than Rs; R4 nearly as long as R4+5; wing coloration pattern with isolated subapical spot and without spot on R2.

**Redescription.** Wing with original coloration pattern of dark clouds visible in medial and distal part; Sc rather long, ending opposite proximal third part of R3; R1



**Figure 2A-C.** *Ptychoptera deleta* Novák, 1877 **A** photograph of holotype specimen No. NM-P947 **B** original line drawing of wing venation from Novák (1877) **C** line drawing of wing venation (reconstruction) **D** *Ptychoptera agnes* Krzemiński and Zwick, 1993, drawing of wing venation (scale bar represents 1 mm).

long; cross vein r-r (R2) at its about two lengths before tip of R1; Rs very short; R4+5 five times longer than Rs; R4 nearly as long as R4+5; cross vein r-m just before forks of Rs and of M; distal part R5 and most part of medial veins not preserved, cross vein m-cu about its length behind origin of M3+4; A1 rather long, its distalmost section strongly curved to posterior wing margin.

**Dimensions.** Length of wing fragment about 7 mm, probable total length about 7.1 mm; maximum width about 1.8 mm.

**Holotype.** No. NM-P947 (imprint of nearly complete wing with medio-apical part missing, posterior wing margin is partially broken, venation well preserved with traces of original pattern of coloration). Specimen is housed in collection of National Museum in Prague, Czech Republic.

**Age and layer.** Early Miocene (Ottangian/Karpatian), Cypris Formation (grey claystone) *sensu* Rojík (2004).

**Discussion.** We provide a re-description of the holotype specimen with several inaccurate characters in wing venation corrected in comparison to the original drawing of Novák, e.g., ending of R1 to anterior wing margin, indication of coloration pattern, A1 basally running parallel to Cu for long distance and ending in posterior wing margin, several missing or present cross-veins. Moreover, we supplement the description by the first photograph of this specimen (unavailable in 1877).

Unfortunately the single diagnostic wing character (M1+2 fork) of the subfamily Ptychopterinae is not preserved. However, the wing venation pattern with very well visible crossvein r-m positioned just before M forking into M1+2 and M3+4, and original color pattern of *P. deleta* show considerable similarities to recent species of *Ptychoptera*. These are distinct apomorphic characters present only in this genus within the family Ptychopteridae. The wing venation of *P. deleta* is somewhat similar to the recent *P. agnes* Krzemiński & Zwick, 1993 described from Hungary (Figs 2C–D). This fact is not surprising due to the Early Miocene age of our fossil (about 18–16 Ma) and considerably high morphological stability of insects (Nel and Prokop 2009; Hörnschmeyer et al. 2009). Finally, our present re-examination of the holotype confirms the great erudition of the Czech pioneer palaeontologist Dr Ottomar Novák who correctly attributed the fossil species to the otherwise modern genus *Ptychoptera*.

## Acknowledgements

This paper is devoted to the 75th anniversary of Prof. Alexandr Rasnitsyn, leader of Russian paleontological laboratory at Paleontological Institute RAS, Moscow. Authors are grateful to Elena Lukashevich, journal editor Dmitry Shcherbakov (both Palaeontological Institute RAS, Moscow), and anonymous reviewer for their insightful comments and suggestions to the early version of the manuscript. We are grateful to Kamil Zagoršek (National Museum in Praha) for the loan of the holotype specimen and Martin Fikáček (National Museum in Praha) for his kind help with translation from German. Special thank goes to Iwona Kania who kindly helped with scanning of

line drawings to digital form. Second author (JP) acknowledges the research support from Grant Agency of the Czech Republic No. 205/08/0643 and Ministry of Education MSM 0021620828.

## References

- Alexander CP (1927) Diptera family Ptychopteridae. In: Wytzman P (Ed) *Genera Insectorum* 188: 1–12.
- Ansorge J, Schröder F (1999) Bittacomorphinae (Diptera: Ptychopteridae) aus der altpaläozänen Fur-Formation (Moler) von Dänemark. *Studia dipterologica* 6: 305–320.
- Bůžek Č, Holý F, Kvaček Z (1996) Early Miocene flora of the Cypris Shale (western Bohemia). *Acta Musei Nationalis Pragae, Series B, Historia naturalis* 52: 1–72.
- Evenhuis NL (1994) *Catalogue of the Fossil Flies of the World (Insecta: Diptera)*. Backhuys Publishers, Leiden, ii + 600 pp.
- Fejfar O (1974) Die Eomiden und Cricetiden (Rodentia, Mammalia) des Miozäns der Tschechoslowakei. *Palaeontographica A*, 146: 100–180.
- Freiwald A, Willman R (1992) Insekten aus der Fur-Formation von Dänemark (Moler, ob. Paläozän/unt. Eozän?). 7. Ptychopteridae (Diptera). *Meyniana* 44: 179–87.
- Hancock E.G., Marcos-Garcia M.A., Rotheray GE (2006) Ptychopteridae — a family of flies (Diptera) new to the Neotropical Region and description of a new species. *Zootaxa* 1351: 61–68.
- Handlirsch A (1909) Zur Phylogenie und Flügelmorphologie der Ptychopteriden (Dipteren). *Annalen des K.K. Naturhistorischen Hofmuseums Wien* 23: 263–272.
- Hörschemeyer T, Wedmann S, Poinar G (2009) How long can insect species exist? Evidence from extant and fossil *Micromalthus* beetles (Insecta: Coleoptera). *Zoological Journal of the Linnean Society* 158: 300–311.
- Kalugina NS (1989) New Mesozoic psychodomorph dipteran insects from Siberia (Diptera: Eoptychopteridae, Ptychopteridae). *Paleontologicheskii Zhurnal* 1989(1): 65–77. [in Russian]
- Krzemiński W (1992) Triassic and Lower Jurassic stage of Diptera evolution. *Mitteilungen der schweizerischen entomologischen Gesellschaft* 65: 39–59.
- Krzemiński W and Krzemińska E (2003) Triassic Diptera: descriptions, revisions and phylogenetic relations. *Acta zoologica cracoviensia* 46(suppl.-Fossil Insects): 153–184.
- Lukashevich ED (1993) New Eoptychopteridae (Diptera: Psychodomorpha) from the Jurassic of Asia. *Paleontological Journal* 27 (1A): 103–122.
- Lukashevich ED (1995) First pupae of the Eoptychopteridae and Ptychopteridae from the Mesozoic of Siberia. *Paleontological Journal* 29 (4): 164–171.
- Lukashevich ED (2008) Ptychopteridae (Insecta: Diptera): History of its study and limits of the family. *Paleontological Journal* 42: 66–74.
- Lukashevich ED, Ansorge J, Krzemiński W, Krzemińska E (1998) Revision of Eoptychopteridae (Diptera: Eoptychopteridae). *Polskie Pismo Entomologiczne* 67: 311–343.

- Lukashevich ED, Coram RA, Jarzembowski EA (2001) New true flies (Insecta: Diptera) from the Lower Cretaceous of southern England. *Cretaceous Research* 22 (4): 451–460. doi: 10.1006/cres.2001.0265
- Mlíkovský J (1996) Tertiary avian localities of the Czech Republic. *Acta Universitatis Carolinae Geologica* 39(1995): 551–557.
- Nel A, Prokop J (2009) Go to the species in paleoentomology: illusion or necessity. *Mémoires de la Société Entomologique de France* 8 (2009): 49–55.
- Novák O (1877) Fauna der Cyprisschiefer des Egerer Tertiärbeckens. *Sitzungsberichte der Koeniglichen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Classe* 76: 71–96.
- Obrhelová N, Obrhel J (1983) [Biostratigraphy of Miocene strata overlying the main brown-coal seam in the Cheb and Sokolov basins.] *Acta Universitatis Carolinae Geologica* 3: 171–192. [in Czech with English summary]
- Obrhelová N, Obrhel J (1987) Paläoichthyologie und Paläoökologie des kontinentalen Tertiärs und Quartärs der ČSSR. *Zeitschrift für Geologische Wissenschaften, Berlin* 15 (1987): 709–731.
- Peus F (1958) 10b. Familie Liriopidae. In: Lindner E (Ed) *Die Fliegen der paläarktischen Region* 3(1) Lief. 200: 10–44.
- Podenas S (2007) First phantom crane flies (Diptera, Ptychopteridae) from Baltic amber (Eocene). *Proceedings of the Academy of Natural Sciences of Philadelphia* 156: 195–198. doi: 10.1635/0097-3157(2007)156[195:FPCFDP]2.0.CO;2
- Prokop J (2003) Remarks on palaeoenvironmental changes based on reviewed Tertiary insect associations from the Krušné hory piedmont basins and the České středohoří Mts. in northwestern Bohemia (Czech Republic). *Acta Zoologica Cracoviensia* 46 (suppl. – Fossil Insects): 329–344.
- Prokop J, Fleck G, Nel A (2003) New dragonflies from the Lower Miocene (Ottomanian/Karpatian) of the Cypris Shale in western Bohemia (Odonata: Libellulidae). *Neues Jahrbuch für Geologie und Paläontologie Monatshefte* 2003: 561–576.
- Ren D, Krzemiński W (2002) Eoptychopteridae (Diptera) from the Middle Jurassic of China. *Annales Zoologici* 52: 207–210.
- Rojík P (2004) New stratigraphic subdivision of the Tertiary in the Sokolov Basin. *Journal of the Czech Geological Society* 49: 173–185.