

New data on freshwater psammic Gastrotricha from Brazil

André R. S. Garraffoni¹, Thiago Q. Araujo¹, Anete P. Lourenço¹, Maria Balsamo²

1 Universidade Federal dos Vales do Jequitinhonha e Mucuri, Departamento de Ciências Biológicas, Campus II, Rodovia BR-367, CEP: 39.100-000, Diamantina, MG, Brazil **2** Dipartimento di Scienze dell'Uomo, dell'Ambiente e della Natura, Università di Urbino, Campi Scientifici, loc. Crocicchia, Urbino, Italy

Corresponding author: André R. S. Garraffoni (garraffoni@gmail.com)

Academic editor: Antonio Todaro | Received 8 June 2010 | Accepted 22 July 2010 | Published 7 October 2010

Citation: Garraffoni ARS, Araujo TQ, Lourenço AP, Balsamo M (2010) New data on freshwater psammic Gastrotricha from Brazil. ZooKeys 60: 1–12. doi: 10.3897/zookeys.60.495

Abstract

Current knowledge of freshwater gastrotrich fauna from Brazil is underestimated as only two studies are available. The present communication is a taxonomic account of the first-ever survey of freshwater Gastrotricha in Minas Gerais State. Samplings were carried out yielding six species of three Chaetonotidae genera: *Aspidiophorus* cf. *pleustonicus*, *Ichthyidium* cf. *chaetiferum*, *Chaetonotus acanthocephalus*, *C. heideri*, *C. cf. succinctus*, *Chaetonotus* sp., and also an undescribed species belonging to the genus *Redudasys* (*incertae sedis*): this is the first finding of specimens of *Redudasys* outside of original type locality. These preliminary observations suggest that the knowledge of the biodiversity of Gastrotricha in the Minas Gerais State, as well as in the whole Brazil, will certainly increase as further investigations are undertaken, and that freshwater Macrotrichida may be more common than previously thought.

Keywords

freshwater Gastrotricha, Macrotrichida, Chaetonotida, biodiversity, meiofauna

Introduction

Gastrotricha are aquatic free-living microinvertebrates (< 1 mm), with a worldwide distribution in freshwater, estuarine, and marine benthic habitats where they are an important component of the benthos and periphyton (Hochberg and Litvaitis 2000; Balsamo and Todaro 2002; Balsamo et al. 2005, 2008). Although many species are common and occasionally abundant, freshwater gastrotrichs are still insufficiently

known, possibly due to their microscopical size, body fragility, which make their study very difficult (Hochberg and Litvaitis 2000; Balsamo and Todaro 2002; Balsamo et al. 2005, 2008). However, despite the minute body size, they are recognized to have both a complex anatomy and life cycle (Weiss 2001).

The taxon consists of nearly 750 named species grouped into two orders, Macro-dasyida and Chaetonotida (but see Kieneke et al. 2008), which are greatly different in morphology, reproductive biology and ecology (Balsamo and Todaro 2002; Balsamo et al. 2008; Todaro and Hummon 2008). Macro-dasyida comprise about 300 worm-like species, all interstitial in marine and estuarine habitats except for the two freshwater ones recorded only from their type locality (Hummon and Todaro, 2010): *Marinellina flagellata* Ruttner-Kolisko, 1955 (Austrian river Ybbs) and *Redudasys fornerisae* Kisielewski, 1987 (Brazilian dam on the savannah near São Carlos city). The roughly 450 species of Chaetonotida are smaller, tenpin-shaped, and colonize marine, brackish and especially freshwater habitats, where two thirds of the species can be found.

The biodiversity of the Gastrotricha fauna in Brazil is still underestimated (Kisielewski 1991; Marques and Lama 2006; Garraffoni and Araujo in press) because, until now, only few studies have focused on the diversity and distribution of this taxon both in fresh waters (Kisielewski 1987, 1991) and in marine waters (Todaro and Rocha 2004, 2005).

Regarding the freshwater habitat, Kisielewski (1991) reported 14 genera (including three new genera), and 59 species (26 of which new species), from various regions of São Paulo State (cities of São Paulo and São Carlos; Juréia Ecological Reserve), of Mato Grosso do Sul State (city of Corumbá), and Pará State (cities of Belém and Benevides), and from different habitats, such as ponds, reservoirs, rivers, puddles in the tropical forest, mangrove and estuaries. In this study, the author stressed that the diversity of Brazilian fauna of inland-water Gastrotricha appears unusually high, and recommended further faunistic, detailed studies. However, no survey was done later on (Garraffoni and Araujo in press).

Thus, the aim of the present study is to provide the first records of the Gastrotricha fauna from the State of Minas Gerais. This is the first of a series of surveys that will be realized as an effort to increase the taxonomic and biogeographic knowledge of the Brazilian Gastrotricha, with special emphasis on the State of Minas Gerais. Furthermore, with the aim to stimulate new research on this group in Brazil, Garraffoni and Araujo (in press) prepared a taxonomic key for all Brazilian freshwater and marine Gastrotricha, and listed the main morphological characters used to identify species as a glossary with terminologies in Portuguese.

Material and methods

Samples of the upper sediment were taken from 7 distinct stations located along two small watercourses and one river near Diamantina city at an altitude of 1300 m: Soberbo (18°11'38.11"S – 43°34'13.03"W), Água Limpa (18°12'51.95"S – 43°37'01.96"W), and Preto River (18°7.50'23"S – 43°20.15'53"W). Other sampling locations were: an unnamed stream in the Itambé Peak (18°23'50"S – 43°19'44"W), at an altitude of

1680 m, and unnamed stream in Cabral Mountains (17°46'03.7"S – 44°17'09.6"W), at an altitude of 1209 m, and an unnamed stream, near Gouveia City, (18°31'48.5"S – 43°53'55.8"W; 18°32'19.2"S – 43°53'52.8"W), at an altitude of 1174 m. Gastrotrichs were extracted after repeated washing of small amounts of sediment with 2% MgCl₂ aqueous solution. Living individuals were located by examining the supernatant under an Olympus SZ40 stereomicroscope at 40x magnification, and were removed by micropipette to a glass slide. Further observations and photographs were done under a Zeiss Photomicroscope equipped with differential interference contrast optics (DIC) and an Olympus CH30 microscope without DIC.

The morphological study and the identification of gastrotrichs were performed using the terminologies and identification keys presented in Kisielewski (1987), Balsamo and Todaro (2002), Balsamo et al. (2005) and Todaro and Hummon (2008). The descriptions followed the convention of Hummon et al. (1992), whereas the locations of some morphological characters along the length of the body were given in percentage units (U) measured from anterior to posterior. In-group systematization of *Chaetonotus* and *Ichthydium* followed Balsamo et al. (2009).

Descriptions of putative new taxa are beyond the scope of the present study, and their definitive affiliation will be made at the end of the ongoing taxonomical surveys in forthcoming papers. However, we provide a photograph of each taxon and the measurement of the main structures, for the benefit of researchers working in the same area who might find them in the meantime.

All adult formalin–glycerin whole-mounts specimens are kept in the meiofauna collection of the senior author at the Universidade Federal dos Vales do Jequitinhonha e Mucuri.

Taxonomy

Order Chaetonotida Remane, 1925

Family Chaetonotidae Gosse, 1864

Genus *Aspidiophorus* Voigt, 1904


Aspidiophorus cf. *pleustonicus* Kisielewski, 1991

Fig. 1, Table 1

Aspidiophorus pleustonicus – Kisielewski (1991: 79, Figs 85–86, Tab. 36); Balsamo et al. (2009: 08, appendix 1).

Material. Soberbo: 2 specimens, Água limpa: 2 specimens, Preto River: 2 specimens, Gouveia: 2 specimens.

Description. The description is based on a single adult specimen, 212.5 µm in total length. Head with oval edge and body long and wide. Body medium-sized, with



Figures 1–9. Freshwater psammic Gastrotricha from Brazil. **1** *Aspidiophorus* cf. *pleustonicus*: ventral view **2** *Ichthyidium* cf. *chaetiferum* **3** *Chaetonotus acanthocephalus*: dorsal view **4** *Chaetonotus heideri*: dorsal view **5** *C. heideri*: close-up of the dorsal scales **6** *Chaetonotus* sp. ventral view **7** *Chaetonotus* cf. *succinctus*: lateral view **8** *C. cf. succinctus*: dorsal view **9** *Redudasys* sp.: dorso-lateral view.

head and neck weakly defined, but trunk and caudal base clearly distinct. Head with slightly five lobes and two pairs of ciliary tufts. Hypostomion weakly developed as a fine transverse furrow appearing as a thin line. Pharynx 56.25 μm in length from the posterior edge of the mouth to the pharyngo-intestinal junction, that lies at U26. Alternating columns of pedunculated, unkeeled, elongate scales along the body.

Remarks. The genus *Aspidiophorus* counts 30 species in the world with 9 marine species and 21 freshwater (Balsamo et al. 2009) and in Brazil there are five freshwater species (Kisielewski 1991) and three marine (Todaro and Rocha 2004, 2005). Our specimens resemble *A. pleustonicus* Kisielewski, 1991 due to the long body, five-lobed head and shape of pedunculated scales. However, they can be distinguished from *A. pleustonicus*, from the size of the mouth, larger than in the original description (Table 1), and the absence of the cuticular rods internal to the pharynx.

Distribution. Brazil: Diamantina, Gouveia (Minas Gerais State); São Paulo (São Paulo State).

Table 1. Morphometrical features of *Aspidiophorus* cf. *pleustonicus*. N= number of specimens measured.

| Features | Range | N | Literature data from Brazil |
|-------------------------|---------------------------|---|-----------------------------|
| Body length | 212.5–275 μm | 6 | 191–208 μm |
| Length of adhesive tube | 31.25–37.5 μm | 6 | 9.5–11.5 μm |
| Pharynx length | 56.25–93.75 μm | 6 | 22.6–27.2 μm |
| Diameter of mouth ring | 11.25–18.75 μm | 6 | 5 μm |
| Cephalion length | 25.97 μm | 6 | - |
| Cephalion width | 19.48 μm | 1 | 16 μm |

Genus *Ichthydium* Ehrenberg, 1830

Ichthydium cf. *chaetiferum* (Müller, 1786)

Fig. 2, Table 2

Ichthydium chaetiferum – Kisielewski (1991: 81, Figs 87–89, Tab. 37); Hummon (2007: 10); Balsamo et al. (2009: 14, Appendix 1); Kånneby et al. (2009: 35).

Material. Água limpa: 1 specimen.

Description. The description is based on a single adult specimen, 108.49 μm in total length. Head with five lobes and two pair of ciliary tufts, with a pair of large “ocellar” granules. Pharynx 28.30 μm in length from the posterior edge of the mouth to the pharyngo-intestinal junction, that is at U26. 8 Spines present on the ventrolateral body side.

Remarks. The specimens collected in the present study resemble those described in Kisielewski (1991) due to the presence of a bulbous pharynx and “ocellar” granules. However, they can be distinguished from those in Kisielewski (1991) for the absence of the dorsal cephalic sensorial bristle and the number of spines in the ventrolateral body side (8 against 12).

Distribution. Brazil: Diamantina (Minas Gerais State), Juréia Ecological Reserve (São Paulo State).

Genus *Chaetonotus* Ehrenberg, 1830

Subgenus *Primochaetus* Kisielewski, 1997

Chaetonotus (*Primochaetus*) *acanthocephalus* Valkanov, 1937

Fig. 3, Table 3

Chaetonotus acanthocephalus – Kisielewski (1981; 1991:54, Figs 65–68, Tab. 22); Hummon (2007: 5); Balsamo et al. (2009: 12, Appendix 1).

Material. Água Limpa: 2 specimens, Soberbo: 1 specimens, Preto River: 5 specimens.

Table 2. Morphometrical features of *Ichthyidium* cf. *chaetiferum*. N= number of specimens measured.

| Features | Measures | N | Literature data from Brazil |
|-------------------------|----------------------|---|-----------------------------|
| Body length | 108.49 μm | 1 | 107–117 μm |
| Length of adhesive tube | 16.98 μm | 1 | 12.5–14 μm |
| Pharynx length | 28.30 μm | 1 | 25–29 μm |
| Diameter of mouth ring | 4.01 μm | 1 | 3 μm |
| Cephalion length | 4.8 μm | 1 | 8.5–9 μm |
| Cephalion width | 9.6 μm | 1 | 11 μm |

Table 3. Morphometrical features of *Chaetonotus acanthocephalus*. N= number of specimens measured.

| Features | Range | N | Literature data from Brazil | Literature data from Europe |
|---|------------------------|---|-----------------------------|-----------------------------|
| Body length | 169–236 μm | 2 | 123–175 μm | 100–148 μm |
| Length of adhesive tube | 27.5–28 μm | 2 | 11–17 μm | 14–16 μm |
| Pharynx length | 62.5–65 μm | 2 | 34–54 μm | 37–49 μm |
| Diameter of mouth ring | 8–8.75 μm | 2 | 5 μm | 6.5–8 μm |
| Cephalion length | 20 μm | 1 | - | - |
| Length of neck scales | 7 μm | 1 | 6–7 μm | 4–7 μm |
| Length of trunk scales | 10 μm | 1 | 5.5–9.5 μm | 5.5–8 μm |
| Maximum length of the neck spines | 24–27 μm | 2 | 7.5–19.5 μm | 11–15 μm |
| Maximum length of the trunk spines | 30–35 μm | 2 | 12.5–28 μm | 16.5–22 μm |
| Length of terminal spines | 19–31.25 μm | 2 | 8–19.5 μm | - |
| Number of scales in a single longitudinal row | 17 | 2 | 17 | 16–18 |

Description. The description is based on an adult specimen, 236 μm in total length. Head with three lobes and a one pair of ciliary tufts. Five peculiar cephalic scales with long spines present on the head. Two ventral plates at the sides of the hypostomion. Pharynx 65 μm in length from the posterior edge of the mouth to the pharyngo-intestinal junction that is at U27. The general long-spine distribution pattern shows two pairs of conspicuous lateral neck spines. Two pairs of long lateral spines at the furcal base.

Remarks. Kisielewski (1991) reported three distinct morphotypes of *Chaetonotus acanthocephalus* in Brazilian inland waters: two of them were collected in São Carlos city and one in Juréia Reserve. Our specimens appear to be close to one of the morphotypes found in São Carlos due to the presence of two pairs of long spines at the furcal base, and the peculiar transversal row of trunk spines (Kisielewski 1991:54, Figs 65–66). However, the body length, width and the posterior spines of the Diamantina specimens are larger than those observed from São Carlos (Table 3).

Distribution. Brazil: Diamantina (Minas Gerais State); São Carlos (São Paulo State), Juréia Reserve (São Paulo State), Corumbá (Mato Grosso do Sul State); Poland: Lake Piaseczno; Germany; Bulgaria.

***Chaetonotus (Primochaetus) heideri* Brehm, 1917**

Figs 4–5, Table 4

Chaetonotus heideri – Emberton (1981: 95, Figs 1–2); Balsamo (1990: 173); Kisielewski (1991: 17, Tab. 3); Nesteruk (2004: 444, Tab. 1; 2007: 836, Tab. 1); Grilli et al. (2008: 174, Fig. h).

Material. Água Limpa: 1 specimen, Soberbo: 2 specimens, Preto River: 2 specimens.

Description. The description is based on an adult specimen 137.5 µm in total length. Head with three lobes and two pairs of ciliary tufts. Pharynx 41 µm in length from the posterior edge of the mouth to pharyngo-intestinal junction, that is at U29. Anterior scales rounded and posterior ones pentagon-like shaped. Lateral spine denticle located near to the spine end.

Distribution. Brazil: Diamantina (Minas Gerais State), Juréia Ecological Reserve and São Carlos (São Paulo State), Benevides (Pará State); USA: Ohio; Germany; England; Italy; Poland; Romania; Russia; Czech Republic; Switzerland; France: Gironde.

Subgenus *Lepidochaetus* Kisielewski, 1991 [Balsamo et al. 2009, p.11]***Chaetonotus* sp.**

Fig. 6, Table 5

Material. Gouveia: 3 specimens.

Description. The description is based on an adult specimen 236.95 µm in total length. Head with three lobes and one pair of ciliary tufts. Pharynx 64.93 µm in length from the posterior edge of the mouth to the pharyngo-intestinal junction (PhIJ), at U27. Hypostomion as a weak transverse furrow. Three pairs of lateral parafurcal spines, the two posteriormost longer than the adhesive tube. Adhesive tubes very long and thin.

Remarks. The genus *Lepidochaetus* was originally described by Kisielewski (1991) to group some *Chaetonotus* species characterized by numerous, rounded, unkeeled

Table 4. Morphometrical features of *Chaetonotus heideri*. N= number of specimens measured.

| Features | Range | N | Literature data from Brazil | Literature data from Europe |
|-------------------------|-----------------|---|-----------------------------|-----------------------------|
| Body length | 137.5–137.96 µm | 2 | 188–196 µm | 106–220 µm |
| Length of adhesive tube | 25–38.8 µm | 2 | 22–25 µm | 21–32 µm |
| Pharynx length | 41 µm | 2 | 48.5–50 µm | 45–56 µm |
| Diameter of mouth ring | 7.55–10.62 µm | 2 | 10–11.5 µm | 10.5–13 µm |
| Length of trunk spines | 37.5–37.96 µm | 2 | 22–37 µm | 46–68 µm |
| Length of egg | 11 µm | 1 | - | - |

Table 5. Morphometrical features of *Chaetonotus* sp. N= number of specimens measured.

| Features | Range | N |
|-----------------------------------|-----------------------------|---|
| Body length | 150.76–236.95 μm | 3 |
| Length of adhesive tube | 26.15–32.46 μm | 3 |
| Pharynx length | 35.38–63 μm | 3 |
| Diameter of mouth ring | 10–14 μm | 3 |
| Cephalion length | 30.96 μm | 3 |
| Length of the egg | 100 μm | 3 |
| Length of rearmost lateral spines | 38.46–71.42 μm | 3 |

scales, provided with long and thin spines covering both the dorsal and the ventral body surfaces. However, here we follow Balsamo et al. (2009) who considered this taxon as a subgenus of *Chaetonotus*. Our specimens resemble *C. (Lepidochaetus) brasiliensis* (Kisielewski 1991), due a similar scale shape and distribution, and rearmost lateral spines arranged in three pairs, which gradually grow in length in a caudal direction. However, they can be distinguished from the previously described species by the absence of cuticular rods and the neck sensorial bristles.

Distribution. Brazil: Diamantina, Gouveia (Minas Gerais State).

Subgenus *Zonochaeta* Remane, 1927

Chaetonotus (Zonocheta) cf. succinctus Voigt, 1902

Figs 7–8, Table 6

Chaetonotus succinctus - Anderson and Robbins (1980: 226); Kisielewski (1991, 60); Hummon (2007: 6); Weiss (2001: 313); Balsamo et al. (2009: 13, Appendix 1).

Material. Cabral Mountains: 1 specimen; Gouveia: 1 specimen; Preto River: 1 specimen.

Description. The description is based on an adult specimen 201.38 μm in total length. Head with five lobes and two pairs of ciliary tufts. Pharynx 55.48 μm in length from the posterior edge of the mouth to the pharyngo-intestinal junction, lying at U27. On the middle trunk region, a transverse band of five long dorsal spines, all terminally bifurcated, and of equal length (77.6 μm) and thickness. Paired spines at the furca base, not extending beyond the adhesive tube end.

Distribution. Brazil: Diamantina, Cabral Mountains (Minas Gerais State), Belém (Pará State); Poland; Romania; England; Italy; Germany; South Korea.

Remarks. Within the subgenus *Zonochaeta*, four species (*C. bisacer*, *C. cestacanthus*, *C. dracunculus*, *C. succinctus*) are characterized by the presence of a series of long dorsal spines with concave apices (Balsamo 1990, 1999). In Brazil, only *C. succinctus* and *C. bisacer* were previously identified (Kisielewski 1991), and the main difference between the two species is the presence of a pair of long spines at the furca base, which extend beyond the adhesive tube tip.

Table 6. Morphometrical features of *Chaetonotus* cf. *succinctus*. N= number of specimens measured.

| Features | Range | N |
|-------------------------------|-----------------------------|---|
| Body length | 165.27–201.38 μm | 2 |
| Length of adhesive tube | 41.6–43.05 μm | 2 |
| Pharynx length | 54.1–55.55 μm | 2 |
| Diameter of mouth ring | 4.13–6.89 μm | 2 |
| Length of trunk “band” spines | 71.42–72.6 μm | 2 |

Order *Macrodasysida* Remane, 1925

Genus *Redudasys* Kisielewski, 1987

Redudasys sp.

Fig. 9, Table 7

Material. Água limpa: 8 specimens; Cabral Montains: 4 specimens. Video sequence (format .mov) is available at <http://www.megaupload.com/?d=1F7NJ1XI>

Description. The description is based on an adult specimen 461.54 μm in total length. Cephalic cilia occur in one transverse dorsal row as well as in irregularly distributed tufts located at the anterolateral head margin. The mouth opening has a diameter of 10.1 μm . Pharynx 153.85 μm in length from the posterior edge of the mouth to the junction with the intestine. Two elongated caudal lobes, 25.64 μm long and 4.76 μm wide. Median caudal cone absent. Only anterior and caudal adhesive tubes are typically present. One anterior tube per side located laterally in the anterior part of the body. Seven tactile bristles per side along the lateral body and one per side on the caudal end. Two pairs of caudal adhesive tubes. The inner tube (7.14 μm long) is usually 2/5 shorter than the external one (11.9 μm long).

Distribution. Brazil: Diamantina, Cabral Mountains (Minas Gerais State).

Remarks. The specimens found in Minas Gerais State are undoubtedly members of the genus *Redudasys*, an *incertae sedis* Macrodasysida taxon recorded from a freshwater environment (Kisielewski, 1987). However, the data gathered in the present study allow to exclude their affiliation to the single species described in this genus so far, *Redudasys fornerisae* Kisielewski, 1987. Our specimens presented seven of tactile bristles per side along the body, and one per side on the caudal body end, that is a total of eight bristles per side, and in addition they had one anterior tube per side. Many of the great number of specimens found in our samples showed a large egg in the trunk region. The species *Redudasys fornerisae* presents six tactile bristles per side along the body and one per side on caudal body end, that is a total of seven bristles per side, and in addition they had two anterior tube per side. Furthermore, the description of *Redudasys fornerisae* was based on six adult individuals (Kisielewski, 1987). Indeed, the specimen used for the description was considered an adult individual by the presence of mature oocytes and a mature egg. For this reason, we strongly believe that the specimen here

Table 7. Morphometrical features of *Redudasys* sp. N= number of specimens measured.

| Features | Range | N |
|--------------------------------|---------------------------|----|
| Body length | 280–461.54 μm | 14 |
| Pharynx length | 87.5–153.85 μm | 14 |
| Diameter of mouth ring | 10.1–11.0 μm | 14 |
| Length of external caudal tube | 11.9–17.5 μm | 14 |
| Length of inner caudal tube | 7.14–12.5 μm | 14 |

described is an adult and cannot be considered as an early stage of development of *Redudasys fornerisae*. Moreover, even in the smaller specimens from the twelve specimens (0,280 mm; Table 7) there are eight tactile bristles per side.

Discussion

The findings presented here allow us to draw some remarks on the Gastrotricha of the State of Minas Gerais. It is worthwhile noting that a poor sampling effort has allowed us to obtain very interesting faunistic data and to identify seven distinct species, which suggest a high biodiversity of Gastrotricha in this State. Up to now, 22 species of Gastrotricha Chaetonotida had been described from Brazilian rivers with slow water current and quiet habitats (Kisielewski 1991).

The most striking result of this study was the report of *Redudasys* specimens from different streams in Minas Gerais State. Thus, the discovery of *Redudasys* specimens outside of the original record is of great biogeographic interest, as the adaptation of this macrodasyidan taxon to the freshwater habitats could have been followed by considerable radiation, mainly in the neotropical region.

Albeit a high diversity of endemic gastrotrich chaetonotidans has been recorded in the Brazilian fauna (e.g. *Undula*, *Arenotus* - Kisielewski 1987, 1991; Balsamo et al. 2008), most of Gastrotricha species and genera found in Brazil have a cosmopolitan distribution. As pointed out by Kisielewski (1987, 1991) and Hummon (2007), species of the Chaetonotidae genera *Aspidiophorus*, *Chaetonotus*, *Heterolepidoderma*, *Ichthyidium*, *Lepidodermella*, *Polymerurus* have a very wide distribution and are known from Europe, Asia, North and South America (*Chaetonotus* species were also collected in Africa and Oceania, and *Polymerurus* species in Oceania). Furthermore, up to now, 60 Chaetonotida species have been reported in Brazilian inland waters and 34 were also found elsewhere in the world, showing that 57% of the freshwater species have intercontinental or cosmopolitan distribution.

Based on preview studies (Kisielewski 1987, 1991; Todaro and Rocha 2004, 2005) and on our own results, it is important to emphasize that further investigations are needed to increase the knowledge of Brazilian gastrotrich fauna, which likely include a quite higher number of species.

Acknowledgements

We would like to express our gratitude to Fundação de Amparo a Pesquisa do Estado de Minas Gerais (FAPEMIG) for financial support (CRA-APQ-00292-09). To Cecília Amaral, at Universidade Estadual de Campinas, and Gustavo Henrique Bahia de Oliveria, at Universidade Federal dos Vales do Jequitinhonha e Mucuri, for the use of laboratory space and equipment. To Sílvio Nihei, at Universidade de São Paulo, for helping us with literature, and Fabiane Costa, Rafael Pessoa and Leonardo Lessa, at Universidade Federal dos Vales do Jequitinhonha e Mucuri, for collecting samples in Cabral Mountains, Gouveia city, and Preto River. Two anonymous referees are also acknowledged for offering suggestions that greatly improved the paper.

References

- Anderson MT, Robbins CE (1980) A new species of gastrotrich from Illinois. *Transactions of the American Microscopical Society* 99: 225–227.
- Balsamo M (1990) Gastrotrichs from Lakes Bolsena, Chiusi and Montepulciano (central Italy), with the description of four new species. *Bollettino di Zoologia* 57: 165–178.
- Balsamo M, d'Hondt JL, Kisielewski J, Pierboni L (2008) Global diversity of gastrotrichs (*Gastrotricha*) in fresh waters. *Hydrobiologia* 595: 85–91.
- Balsamo M, d'Hondt JL, Pierboni L, Grilli P (2009) Taxonomic and nomenclatural notes on freshwater *Gastrotricha*. *Zootaxa* 2158: 1–19.
- Balsamo M, Todaro MA (2002) *Gastrotricha*. In: Rundle SD, Robertson A, Schmidt-Araya J (Eds) *Freshwater meiofauna: Biology and ecology*. Backhuys Publishers, Leiden, 45–61.
- Balsamo M, Todaro MA, Tongiorgi P (2005) *Gastrotricha*. In: Yule CM, Yong HS (Eds) *Freshwater Invertebrates of the Malaysian region*. Academy of Science Malaysia, Singapore, 127–135.
- Emberton KC Jr. (1981) First record of *Chaetonotus heideri* (*Gastrotricha*: *Chaetonotidae*) in North America. *Ohio Journal of Science* 95: 95–96.
- Garraffoni ARS, Araujo TQ (in press) Chave de identificação de *Gastrotricha* de águas continentais e marinhas do Brasil. *Papéis Avulsos de Zoologia*.
- Grilli P, d'Hondt JL, Balsamo M (2008) Contribution à la connaissance des *Gastrotriches* dulcicoles du département de la Gironde. *Bulletin de la Société Linnéenne de Bordeaux* 36: 171–179.
- Hochberg R, Litvaitis MK (2000) Phylogeny of *Gastrotricha*: a morphology-based framework of gastrotrich relationships. *Biological Bulletin* 198: 299–305.
- Hummon WD (2007) Synopsis of the described *Chaetonotida* of the world. <http://hummonas.biosci.ohiou.edu/2010%20Global%20DB%20MarGast/Syst%20&%20Bibliogr/> [accessed 25.VI.2010]
- Hummon WD, Balsamo M, Todaro MA (1992) Italian marine *Gastrotricha*: I. Six new and one redescribed species of *Chaetonotida*. *Bollettino di Zoologia* 59: 499–516.

- Hummon WD, Todaro MA (2010) Analytic taxonomy and notes on marine, brackish-water and estuarine Gastrotricha. *Zootaxa* 2392: 1–32.
- Kånneby T, Todaro MA, Jondelius U (2009) One new species and records of *Ichthyidium* Ehrenberg, 1830 (Gastrotricha: Chaetonotida) from Sweden with a key to the genus. *Zootaxa* 2278: 26–46.
- Kieneke A, Riemann O, Ahlrichs W (2008) Novel implications for the basal internal relationships of Gastrotricha revealed by an analysis of morphological characters, *Zoologica Scripta* 4: 429–460.
- Kisielewski J (1981) Gastrotricha from raised and transitional peat bogs in Poland. *Monografie Fauny Polski, Państwowe Wydawnictwo Naukowe, Warszawa*, 142 pp.
- Kisielewski J (1987) Two new interesting genera of Gastrotricha (Macrodasysida and Chaetonotida) from the Brazilian freshwater psammon. *Hydrobiologia* 153: 23–30.
- Kisielewski J (1990) Origin and phylogenetic significance of freshwater psammic Gastrotricha. *Stygologia* 5: 87–92.
- Kisielewski J (1991) Inland-water Gastrotricha from Brazil. *Annales Zoologici* 43: 1–168.
- Marques AC, Lamas CJE (2006) Taxonomia zoológica no Brasil: estado da arte, expectativas e sugestões de ações futuras. *Papéis Avulsos de Zoologia* 46: 139–174.
- Nesteruk T (2004) Lake zonality influence on species diversity formation of Gastrotricha. *Acta Agrophysica* 4: 441–447.
- Todaro MA, Dal Zotto M, Maiorova AS, Adrianov AV (2009) A new species of *Aspidiophorus* (Gastrotricha, Chaetonotida) from the Russian Far East with a key to marine species of the genus. *Marine Biological Research* 5: 297–333.
- Todaro MA, Hummon WD (2008) An overview and a dichotomous key to genera of the phylum Gastrotricha. *Meiofauna Marina* 16: 3–20.
- Todaro MA, Rocha CEF (2004) Diversity and distribution of marine Gastrotricha along the northern beaches of the state of São Paulo (Brazil), with description of a new species of *Macrodasys* (Macrodasysida, Macrodasysidae). *Journal of Natural History* 38: 1605–1634.
- Todaro MA, Rocha CEF (2005) Further data on marine gastrotrichs from the State of São Paulo and the first records from the State of Rio de Janeiro (Brazil). *Meiofauna Marina* 14: 27–31.
- Tongiorgi P, Fregni E, Balsamo M (1999) Gastrotricha from Italian brackish environment with description of a new species of *Chaetonotus*. *Journal of the Marine Biological Association of United Kingdom* 79: 585–592.
- Weiss M (2001) Widespread hermaphroditism in freshwater gastrotrichs. *Invertebrate Biology* 120: 308–431.