

Megaloptera of Canada

Xingyue Liu¹

¹ Department of Entomology, China Agricultural University, Beijing 100193, China

Corresponding author: Xingyue Liu (xingyue_liu@yahoo.com)

Academic editor: D. Langor | Received 29 January 2018 | Accepted 2 March 2018 | Published 24 January 2019

<http://zoobank.org/E0BA7FB8-0318-4AC1-8892-C9AE978F90A7>

Citation: Liu X (2019) Megaloptera of Canada. In: Langor DW, Sheffield CS (Eds) The Biota of Canada – A Biodiversity Assessment. Part 1: The Terrestrial Arthropods. ZooKeys 819: 393–396. <https://doi.org/10.3897/zookeys.819.23948>

Abstract

An updated summary on the fauna of Canadian Megaloptera is provided. Currently, 18 species are recorded in Canada, with six species of Corydalidae and 12 species of Sialidae. This is an increase of two species since 1979. An additional seven species are expected to be discovered in Canada. Barcode Index Numbers are available for ten Canadian species.

Keywords

alderflies, biodiversity assessment, Biota of Canada, dobsonflies, fishflies, Megaloptera

The order Megaloptera (dobsonflies, fishflies, and alderflies) is one of the three orders of Neuropterida, and is characterized by the prognathous adult head, the broad anal area of hind wing and the exclusively aquatic larval stages (New and Theischinger 1993). Currently, there are ca. 380 described species of Megaloptera worldwide (Yang and Liu 2010, Oswald 2016). Extant Megaloptera are composed of only two families; Corydalidae, which is divided into Corydalinae (dobsonflies) and Chauliodinae (fishflies), and Sialidae (alderflies). Major species diversity of Megaloptera is confined to the subtropical and warm temperate regions, e.g., the Oriental, Neotropical, and Australasian Regions (Yang and Liu 2010, Liu et al. 2012, 2015).

Southern Canada is probably the northern limit of the distribution range of Nearctic Megaloptera. Recent phylogeographic studies suggest that the Canadian population of a fishfly species (i.e., *Nigronia serricornis* (Say)) was formed very rapidly after the Pleistocene glacial period (Heilveil and Berlocher 2006). Thus, the Canadian Megaloptera may play an important part in understanding the evolutionary history of Nearctic Megaloptera.

All known Canadian species of Megaloptera are also found in the United States of America. Kevan (1979) recorded 16 native species of Megaloptera in Canada, including six species of Corydalidae and 10 species of Sialidae. Since then, no more species of Corydalidae have been found in Canada. Evans (1984) reported two new fishflies, *Orohermes crepusculus* (Chandler) and *Protochauliodes cascadius* Evans, from the northwestern USA where they coexist with *Dysmicohermes disjunctus* (Walker) and *Protochauliodes spenceri* Munroe in Oregon (Evans 1972). As the latter two species are known from southwestern British Columbia, *O. crepusculus* and *P. cascadius* may also be found in similar habitats there. Additionally, *Neohermes concolor* (Davis), which is widespread in the eastern USA (Liu et al. 2016), may also occur in adjacent areas of Canada. It should be also noted that in the Species Catalogue of Lacewing Digital Library (LDL), the most comprehensive and frequently updated database of species of Neuropterida (Oswald 2016), *Chauliodes rastricornis* Rambur is not recorded in Canada, although the Canadian record of this species is reported by van der Weele (1910). Herein the distribution of *C. rastricornis* in Canada is confirmed.

For Canadian alderflies, the most important faunal work after Kevan (1979) is that of Whiting (1991) in which *Sialis infumata* Newman and *S. joppa* Ross were newly recorded in Canada. An additional four species of *Sialis* are estimated to be found in Canada, namely *S. aequalis* Banks, *S. driesbachi* Flint, *S. hasta* Ross, and *S. spangleri* Flint, because these species are distributed around the Great Lakes region of northeastern USA that is in close proximity to Canada.

DNA barcodes are available for all but one species of known Canadian Corydalidae, but for less than half of Sialidae (Table 1). Species with DNA barcodes comprise *Chauliodes pectinicornis* (Linnaeus) (BOLD:AAH3593), *C. rastricornis* (BOLD:AAH3594), *D. disjunctus* (BOLD:ACA3660), *N. serricornis* (BOLD:AAA1274), *P. spenceri* (BOLD:ACP8653), *Sialis concava* Banks (BOLD:AAL6477), *S. hamata* Ross (BOLD:ACA3407), *S. joppa* Ross (BOLD:AAG9766), *S. vagans* Ross (BOLD:AAH7456), and *S. velata* Ross (BOLD:AAG9765). Current barcode information does not indicate the presence of cryptic species.

Thirteen of the 18 species of Canadian megalopterans have their larval stage described, and their life history is known (Davis 1903, Cuyler 1958, Neunzig 1966, Azam and Anderson 1969, Evans 1972, Leischner and Pritchard 1973, Lilly et al. 1978); however, most of the information is based on studies of materials or populations from the United States.

Additional surveys of Megaloptera habitats in southern Alberta and British Columbia, especially southwestern British Columbia, are warranted to fill in gaps in distribution and to ascertain whether other species are present. Fresh material of all megalopterans, especially Sialidae, is needed for obtaining DNA barcodes.

Table 1. Census of Megaloptera in Canada.

Taxon ¹	No. species reported in Kevan (1979)	No. species currently known from Canada	Number of BINs ² available for Canadian species	Est. no. undescribed or unrecorded species in Canada	General distribution by ecozone ³	Information sources
Corydalidae	6	6	5	3	Boreal Shield, Pacific Maritime, Atlantic Maritime, Mixedwood Plains	van der Weele 1910, Evans 1972, Yang and Liu 2010, Oswald 2016, Liu and Winterton 2016
Sialidae	10	12	5	4	Boreal Shield, Boreal Plains, Pacific Maritime, Atlantic Maritime, Montane Cordillera, Mixedwood Plains, Prairies	Whiting 1991, Liu et al. 2015
Total	16	18	10	7		

¹Classification follows that of Yang and Liu (2010). ²Barcode Index Number, as defined in Ratnasingham and Hebert (2013).

³See figure 1 in Langor (2019) for a map of ecozones.

References

- Azam KM, Anderson NH (1969) Life history and habits of *Sialis rotunda* and *S. californica* in Western Oregon. *Annals of the Entomological Society of America* 62: 549–558. <https://doi.org/10.1093/aesa/62.3.549>
- Cuyler RD (1958) The larvae of *Chauliodes* Latreille (Megaloptera: Corydalidae). *Annals of the Entomological Society of America* 51: 582–586. <https://doi.org/10.1093/aesa/51.6.582>
- Davis KC (1903) Sialididae of North and South America. *Bulletin of the New York State Museum* 68: 442–486, 499.
- Evans ED (1972) A study of the Megaloptera of the Pacific coastal region of the United States. PhD thesis, Oregon State University, Corvallis, 210 pp.
- Evans ED (1984) A new genus and a new species of dobsonfly from the far western United States (Megaloptera: Corydalidae). *Pan-Pacific Entomologist* 60: 1–3.
- Heilveil JS, Berlocher SH (2006) Phylogeography of postglacial range expansion in *Nigronia serricornis* Say (Megaloptera: Corydalidae). *Molecular Ecology* 15: 1627–1641. <https://doi.org/10.1111/j.1365-294X.2006.02876.x>
- Kevan DK McE (1979) Megaloptera. In: Danks HV (Ed.) *Canada and its insect fauna*. *Memoirs of the Entomological Society of Canada* No. 108, 351–352.
- Langor DW (2019) The diversity of terrestrial arthropods in Canada. In: Langor DW, Sheffield CS (Eds) *The Biota of Canada – A Biodiversity Assessment. Part 1: The Terrestrial Arthropods*. *ZooKeys* 819: 9–40. <https://doi.org/10.3897/zookeys.819.31947>
- Leischner TG, Pritchard G (1973) The immature stages of the alderfly, *Sialis cornuta* (Megaloptera: Sialidae). *Canadian Entomologist* 105: 411–418. <https://doi.org/10.4039/Ent105411-3>
- Lilly CK, Ashley DL, Tarter DC (1978) Observations on a population of *Sialis itasca* Ross in West Virginia (Megaloptera: Sialidae). *Psyche* 85: 209–217. <https://doi.org/10.1155/1978/96295>

- Liu XY, Hayashi F, Yang D (2015) Phylogeny of the family Sialidae (Insecta: Megaloptera) inferred from morphological data, with implications for generic classification and historical biogeography. *Cladistics* 31: 18–49. <https://doi.org/10.1111/cla.12071>
- Liu XY, Wang YJ, Shih CK, Ren D, Yang D (2012) Early evolution and historical biogeography of fishflies (Megaloptera: Chauliodinae): implications from a phylogeny combining fossil and extant taxa. *PLoS ONE* 7: e40345. <https://doi.org/10.1371/journal.pone.0040345>
- Liu XY, Winterton SL (2016) A new fishfly species (Megaloptera: Corydalidae: *Neohermes* Banks) discovered from North America by a systematic revision, with phylogenetic and biogeographic implications. *PLoS ONE* 11: e0148319. <https://doi.org/10.1371/journal.pone.0148319>
- Neunzig HH (1966) Larvae of the genus *Nigronia* Banks (Neuroptera: Corydalidae). *Proceedings of the Entomological Society of Washington* 68: 11–16.
- New TR, Theischinger G (1993) Megaloptera (Alderflies, Dobsonflies). *Handbuch der Zoologie* (Berlin) 4: 1–97.
- Oswald JD (2016) Neuropterida Species of the World. Version 9.0. <http://lacewing.tamu.edu/Species-Catalogue/> [Accessed on 26 December 2016]
- Ratnasingham S, Hebert PDN (2013) A DNA-based registry for all animal species: the Barcode Index Number (BIN) system. *PLoS ONE* 8: e66213. <https://doi.org/10.1371/journal.pone.0066213>
- van der Weele HW (1910) Megaloptera Monographic Revision. *Collections Zoologiques du Baron Edm. de Selys Longchamps* 5: 1–93.
- Whiting MF (1991) A distributional study of *Sialis* (Megaloptera: Sialidae) in North America. *Entomological News* 102: 50–56.
- Yang D, Liu XY (2010) *Fauna Sinica Insecta Vol.51 Megaloptera*. Science Press, Beijing, 457 pp + 14 pls.