

On a new species of freshwater crab, *Indochinamon khinpyae*, from northern Myanmar (Crustacea, Brachyura, Potamidae)

Peter K. L. Ng¹, Win Mar²

1 Lee Kong Chian Natural History Museum, 2 Conservatory Drive, National University of Singapore, Singapore 117377, Republic of Singapore **2** Department of Zoology, Banmaw University, Banmaw, Kachin State, Republic of the Union of Myanmar

Corresponding author: Peter K. L. Ng (peterng@nus.edu.sg)

Academic editor: S. De Grave | Received 19 August 2018 | Accepted 1 November 2018 | Published 31 December 2018

<http://zoobank.org/C1B05F42-E931-45F8-9339-F691FDD59BB6>

Citation: Ng PKL, Mar W (2018) On a new species of freshwater crab, *Indochinamon khinpyae*, from northern Myanmar (Crustacea, Brachyura, Potamidae). ZooKeys 811: 47–63. <https://doi.org/10.3897/zookeys.811.29187>

Abstract

A new species of freshwater crab of the genus *Indochinamon* Yeo & Ng, 2007 (family Potamidae), is described from highlands north of Myitkyina in Kachin State, Myanmar. *Indochinamon khinpyae* **sp. n.** is distinguished from congeners by its very rugose carapace, broad male pleon and distinctively structured male first gonopod; and is the first potamid species recorded from northern Myanmar.

Keywords

Taxonomy, freshwater crab, Burma, Potaminae, *Indochinamon*, new species, description

Introduction

The freshwater crabs (Brachyura, Potamidae, Gecarcinucidae) of Indochina are very diverse, with the fauna still in a survey and discovery stage. The fauna of Myanmar (= Burma) in particular, is poorly known, with most of the recognised species described in the early 1900s (see Yeo and Ng 1999; Cumberlidge et al. 2012). Recent efforts in recording the crab fauna have only just started (e.g., Ng 1996, 2018; Ng and Kosuge 1997; Ng and Whitten 2017), with many parts of the country still barely explored. The second author recently obtained several lots of freshwater crabs from northern Myanmar, one of which proved to belong to a new species of Potamidae.

Material and methods

The terminology used follows Ng (1988) with recent changes by Davie et al. (2015). The abbreviations G1 and G2 are used for the male first and second gonopods, respectively. Measurements provided, in millimetres, are of the maximum carapace width and length, respectively. The material examined is deposited in the Museo Civico di Storia Naturale “Giacoma Doria” (MGE), Genova, Italy; Naturhistorisches Museum Basel (MBA), Basel, Switzerland; Muséum national d’Histoire naturelle (MNHN), Paris, France; Naturalis [formerly Rijksmuseum van Natuurlijke Histoire, RMNH], Leiden, The Netherlands; Senckenbergischen Naturforschenden Gesellschaft (SMF), Frankfurt am Main, Germany; Zoological Reference Collection (ZRC), Lee Kong Chian Natural History Museum, National University of Singapore; and the Zoological Survey of India (ZSI), Calcutta, India.

Systematics

Family Potamidae Ortmann, 1896 *sensu* Yeo and Ng (2004)

Genus *Indochinamon* Yeo & Ng, 2007

Type species. *Potamon villosum* Yeo & Ng, 1998, by original designation.

Remarks. The genus currently contains 38 species from Thailand, Vietnam, Laos, Myanmar, India and China (Table 1, updated from Ng et al. 2008; Naruse et al. 2018). Established by Yeo and Ng (2007) for Indochinese species previously placed in *Potamon* Savigny, 1816, *s. lato*, *Indochinamon* is defined by a suite of characters: carapace low with a relatively flat dorsal surface; the epigastric cristae are separated from the postorbital cristae by a distinct groove; the postorbital cristae is not confluent with the epibranchial tooth; the exopod of the third maxilliped has a long flagellum; the ambulatory legs are relatively short and stout; the male pleon is narrowly triangular; the sternopleonal cavity reaches an imaginary line joining the median parts of the coxae of the chelipeds; and the G1 terminal segment is relatively short, with the groove for the G2 marginal in position, and the dorsal flap is either absent or only low and broad.

Indochinamon khinpyae sp. n.

<http://zoobank.org/878C53F6-38DA-42C3-B282-A1F6A56C0E92>

Figs 1–4

Material examined. Holotype: male (57.1×43.2 mm) (ZRC 2018.0713), Malikha River, about 3.2 km from confluence point of Ayeyarwady River, north of Myitkyina, Kachin State, Myanmar, coll. Khin Pyae Pyae Thaw Thar, May 2018. Paratypes: 5 males (47.2×36.4 mm, 48.9×37.7 mm, 46.4×35.8 mm, 34.3×26.6 mm, 26.7×21.1 mm), 1 female (39.3×31.1 mm) (ZRC 2018.0714), same data as holotype.

Table I. List of recognised *Indochinamon* species.

<i>Indochinamon abkense</i> Naruse, Chia & Zhou, 2018 [type locality: Guangnan County, Yunnan Province, China]
<i>Indochinamon andersonianum</i> (Wood-Mason, 1871) [type locality: Momein, west Yunnan Province, China]
<i>Indochinamon asperatum</i> (Alcock, 1909) [type locality: Cachar Hills, India]
<i>Indochinamon bavi</i> Naruse, Nguyen & Yeo, 2011 [type locality: Ha Tay Province, northern Vietnam]
<i>Indochinamon beieri</i> (Pretzmann, 1966) [type locality: Dawane Hills, India]
<i>Indochinamon bhumbibol</i> (Naiyanetr, 2001) [type locality: Loei Province, northern Thailand]
<i>Indochinamon boshanense</i> (Dai & Chen, 1985) [type locality: Boshan, Yunnan Province, China]
<i>Indochinamon changpoense</i> (Dai, 1995) [type locality: Jingping, Yunnan Province, China]
<i>Indochinamon chinghungense</i> (Dai, Song, He, Cao, Xu & Zhong, 1975) [type locality: Ching Hung, Yunnan Province, China]
<i>Indochinamon chuahuong</i> Do, Nguyen & Le, 2016 [type locality: Ha Noi province, northern Vietnam]
<i>Indochinamon cua</i> (Yeo & Ng, 1998) [type locality: Vinh Phu Province, northern Vietnam]
<i>Indochinamon dangi</i> Naruse, Nguyen & Yeo, 2011 [type locality: Dien Bien Province, northern Vietnam]
<i>Indochinamon dawuishanense</i> (Dai, 1995) [type locality: Daweishan, Yunnan Province, China]
<i>Indochinamon edwardsii</i> (Wood-Mason, 1871) [type locality: Pensee, Kakhien Hills, Yunnan Province, China]
<i>Indochinamon flexum</i> (Dai, Song, Li & Liang, 1980) [type locality: Napo, Guangxi Province, China]
<i>Indochinamon gengmaense</i> (Dai, 1995) [type locality: Gengma, Yunnan Province, China]
<i>Indochinamon guttum</i> (Yeo & Ng, 1998) [type locality: Muang Saisombun, northern Laos]
<i>Indochinamon hirtum</i> (Alcock, 1909) [type locality: Sheetee Hills, Kakhien Hills, Yunnan Province, China]
<i>Indochinamon hispidum</i> (Wood-Mason, 1871) [type locality: Pensee, Kakhien Hills, Yunnan Province, China]
<i>Indochinamon jianchuanense</i> (Dai & Chen, 1985) [type locality: Hengduan, Yunnan Province, China]
<i>Indochinamon jinpingense</i> (Dai, 1995) [type locality: Yunnan Province, China]
<i>Indochinamon khinpyae</i> sp. n. [type locality: Kachin State, Myanmar]
<i>Indochinamon kimboiense</i> (Dang, 1975) [type locality: Kim Boi Province, northern Vietnam]
<i>Indochinamon lipkei</i> (Ng & Naiyanetr, 1993) [type locality: Chiang Rai Province, northern Thailand]
<i>Indochinamon lui</i> Naruse, Chia & Zhou, 2018 [type locality: Yun County, Yunnan Province, China]
<i>Indochinamon manipurensis</i> (Alcock, 1909) [type locality: Manipur Hills, India]
<i>Indochinamon menglaense</i> (Dai & Cai, 1998) [type locality: Xishuangbana, Yunnan Province, China]
<i>Indochinamon mieni</i> (Dang, 1967) [type locality: Son La Province, northern Vietnam]
<i>Indochinamon orleansi</i> (Rathbun, 1904) [type locality: northern Vietnam]
<i>Indochinamon ou</i> (Yeo & Ng, 1998) [type locality: Phongsali Province, northern Laos]
<i>Indochinamon parpidum</i> Naruse, Chia & Zhou, 2018 [type locality: Shiping County, Yunnan Province, China]
<i>Indochinamon phongnha</i> Naruse, Nguyen & Yeo, 2011 [type locality: Quang Binh Province, central Vietnam]
<i>Indochinamon prolatum</i> (Brandis, 2000) [type locality: Uthai Thani Province, central Thailand]
<i>Indochinamon tammanti</i> (Rathbun, 1904) [type locality: Lao Koi, Yunnan Province, China]
= <i>Potamon hokuoense</i> Dai, Song, He, Cao, Xu & Zhong, 1975 [type locality: Hekou, Yunnan Province, China]
<i>Indochinamon tritum</i> (Alcock, 1909) [type locality: Sheetee Hills, Kakhien Hills, Yunnan Province, China]
<i>Indochinamon tuijense</i> Naruse, Chia & Zhou, 2018 [type locality: Nanhua County, Yunnan Province, China]
<i>Indochinamon villosum</i> (Yeo & Ng, 1998) [type locality: Luang Nam Tha Province, northern Laos]
<i>Indochinamon xinpingense</i> (Dai & Bo, 1994) [type locality: Yuxi, Yunnan Province, China]
= <i>Potamon hispidum xinpingense</i> Bo, He, Huang, Fan, Dai & Chen, 1997 [type locality: Yuxi, Yunnan Province, China]
<i>Indochinamon yunlongense</i> (Dai, 1995) [type locality: Yunlong, Yunnan Province, China]

Comparative material. *Indochinamon abkense* Naruse, Chia & Zhou, 2018 – paratypes: 4 males (largest 38.4×29.7 mm), 2 females (larger 43.1×33.2 mm) (ZRC 2013.0551), Shaping Village, Ahke Town, Guangnan County, Yunnan Province, China, coll. Z.L. Chen, 1 February 2004. *Indochinamon andersonianum* (Wood-Mason, 1871) – syntypes: 2 males (larger 36.4×28.3 mm), 1 female (42.9×33.5 mm) (ZSI 4045/4), Momein, West Yunnan, China, coll. J. Wood-Mason, no date; 1 male (49.5×37.0 mm) (SMF 2805), Mt. Carien, Myanmar, coll. L. Fea, 1885–1889; 1 female (47.9×36.9 mm) (ZSI 6916/3), Yunnan, coll. J. Anderson, no date; 11 juveniles (ZSI 6932/3), West Yunnan and Kahkhyen Hills, coll. J. Anderson, no date; 1 juvenile male, 3 females (largest 33.8×26.8 mm) (ZSI 6906/3), Kahkhyen Hills, Ponsee, Upper Burma, coll. J. Anderson, no date. *Indochinamon asperatum* (Alcock, 1909) – syntypes: 4 juvenile males (largest 18.1×15.3 mm), 1 juvenile female (19.2×15.7 mm) (ZSI 5543/10), Ganjam, Cachar Hills, coll. W. Partridge, no date. *Indochinamon bavi* Naruse, Nguyen & Yeo, 2011 – paratypes: 2 males (46.9×35.8 mm, 47.5×36.3 mm) (ZRC 2010.0167), Ba Vi National Park, Ha Tay Province, Vietnam, coll. V.Q. Nguyen, 19 June 2001. *Indochinamon bhumibol* (Naiyanetr, 2001) – 3 males (ZRC), Ban Nam Tob, Khao Luang, Amphoe Wang Saphung, Loei Province, Thailand, coll. W. Senama, 26 October 1982; 1 male (ZRC), Huai Phai Waterfall, Phu Rua, Amphoe Phu Rua, Loei Province, coll. Wiroon, 24 July 1982; 2 males (ZRC), Ban Na Wa, Amphoe Dan Sai, Loei Province, Thailand, coll. P. Naiyanetr, 11 April 1987. *Indochinamon boshanense* (Dai & Chen, 1985) – 1 male (50.4×37.1 mm) (ZRC 1998.811), Boshan, Yunnan Province, China, coll. A. Dai, 20 October 1981. *Indochinamon chinghungense* (Dai, Song, He, Cao, Xu & Zhong, 1975) – 1 male (50.5×38.2 mm) (ZRC 1997.749), Menghai County, 100 m asl, Yunnan County, China, coll. Y. Cai, 11 May 1994. *Indochinamon cua* (Yeo & Ng, 1998) – holotype: male (46.9×36.6 mm) (ZRC 1998.267), Tam Dao, Vinh Phu Province, northern Vietnam, coll. X.Q. Nguyen, 8 June 1997; paratypes: 2 females (larger 42.6×33.0 mm), 2 juvenile males (ZRC 1998.268–271), same locality and collector as holotype, March 1997. *Indochinamon dangi* Naruse, Nguyen & Yeo, 2011 – 4 males (29.6×23.0 – 48.4×37.6 mm), 2 females (45.2×33.9, 36.7×28.1 mm) (ZRC 2010.0175), upstream and waterfall of Muong Phang stream, Muong Phang, Dien Bien Province, 21°27.000'N, 103°10.548'E, 1070 m asl, coll. D.C.J. Yeo and A.D. Tran, 28 July 2004; 4 males (26.7×20.7 – 41.8×32.2 mm), 1 female (46.7×35.5 mm) (ZRC 2010.0176), Muong Phang stream, Muong Phang, Dien Bien Province, 21°27.159'N, 103°09.921'E, 976m asl, coll. D.C.J. Yeo and A.D. Tran, 26 July 2004. *Indochinamon edwardsii* (Wood-Mason, 1871) – 1 male (40.9×30.3 mm), 1 female (37.5×27.5 mm) (MGE III 228 bis), Mt. Catcin, Birmania (= Myanmar), coll. L. Fea, June–October 1886; 1 male (38.5×28.9 mm), 2 females (larger 33.9×25.5 mm) (MBA 51a), Katein Berge, northern Burma; 1 male (about 34.4 mm carapace width) (ZRC 1984.7036), Mount Katun, Burma (= Myanmar), coll. L. Fea, 1893. *Indochinamon flexum* (Dai, Song, Li & Liang, 1980) – 1 male (47.5×35.0 mm) (ZRC 1997.0750), Guangxi Autonomous Region, China, coll. Y. Song, 16 September 1997. *Indochinamon guttum* (Yeo & Ng, 1998) – holotype: male (62.2×45.7 mm) (ZRC 1998.272), Ban Long Cheng, Muang Saisombun, Saisombun

Special Zone, northern Laos, coll. V. Kittikoon, May 1995; paratypes: 1 male (43.8×34.7 mm) (ZRC 1998.273), same data as holotype; 1 female (33.1×25.3 mm) (ZRC 1998.0274), side of dam, Muang Saisombun, Saisombun Special Zone, northern Laos, coll. V. Kittikoon, May 1995; 8 specimens (largest male 51.3×38.8 mm) (MNHN-B 5316) “Haut Laos, Ban Nong”, coll. Mission Permanente, 10 January 1906. *Indochinamon hirtum* (Alcock, 1909) – holotype: female (32.9×24.9 mm) (ZSI 6961/3), Sheete Hill (= Shitee Doung), Kakhyen Hills, Yunnan, China, coll. J. Anderson, no date; 5 males (largest 37.8×28.0 mm), 2 females (ZSI 6961/3), same data as holotype. *Indochinamon hispidum* (Wood-Mason, 1871) – 1 juvenile female (ZSI 4007/4), Kakhyen Hills, Pongsee, Upper Burma, coll. J. Anderson, no date; 1 female (34.6×26.9 mm), 1 juvenile male (ZSI 7089-90/9), Moug Sal, Mehkok River, coll. Dr. Grey, no date. *Indochinamon jinpingense* (Dai, 1995) – 2 males (larger 64.0×47.0 mm) (ZRC 1998.266), Sin Ho District, Lai Chau Province, northern Vietnam, coll. V.D. Nguyen, November 1997. *Indochinamon kimboiense* (Dang, 1975) – 2 males (71.8×56.6 mm, 71.5×56.8 mm), 2 females (63.0×49.5 mm, 52.8×41.1 mm) (ZRC 2010.0165), Kim Boi area, Hoa Binh Province, Vietnam, purchased from villagers, 14 and 15 April 2007; 1 male (58.8×45.3 mm), 2 females (69.4×53.6 mm, 49.9×37.4 mm) (ZRC 2010.0166), stream in Cuc Phuong National Park, about 6 km from main gate, Ninh Binh Province, northern Vietnam, 20°18'N, 105°38'E, coll. D.C.J. Yeo, H.H. Ng and X.Q. Nguyen, 16 September 1997. *Indochinamon lipkei* (Ng & Naiyanetr, 1993) – holotype: male (56.8×42.8 mm) (RMNH D 42353), Chiang Khong District, Chiang Rai Province, northwestern Thailand, coll. P. Naiyanetr, June 1987. *Indochinamon manipurensis* (Alcock, 1909) – syntypes: 1 male (39.9×31.5 mm), 1 female (40.1×30.4 mm) (ZSI 6923/3), Manipur Hills, India, coll. H. H. Godwin-Austen, no date. *Indochinamon menglaense* (Dai & Cai, 1998) – 1 male (42.9×31.9 mm), 1 female (ZRC), Shangyong, Xishuangbana, Yunnan, China, coll. Y. Cai, 23 April 1994. *Indochinamon mieni* (Dang, 1967) – neotype: male (57.1×43.5 mm) (ZRC 1998.265), Thuan Chau District, Son La Province, northern Vietnam, coll. V.D. Nguyen, 1997; 1 juvenile female (ZRC), same data as neotype. *Indochinamon orleansi* (Rathbun, 1904) – holotype: male (42.4×32.3 mm) (MNHN-B 5262), “Tonkin, rivière Noire” (river Song Da), coll. Prince Henri d’Orleans, no date. *Indochinamon ou* (Yeo & Ng, 1998) – holotype: male (35.6×27.1 mm) (ZRC 1998.275), Nam Ou at confluence with Huay Nam, 21°4'10"N, 102°31'44"E, 3 km ESE of Muang Khoa, Phongsali Province, northern Laos, coll. M. Kottelat, 17 May 1997; 1 male (47.6×36.8 mm) (ZRC), dry evergreen forest mixed with bamboo, Nam Sa River, tributary of Nam Ou, 600 m asl, 22°5'31"N, 102°6'19"E, Phou Dendin, Phongsali, northern Laos, coll. and date not known. *Indochinamon parpidum* Naruse, Chia & Zhou, 2018 – paratypes: 2 males (larger 43.4×32.3 mm), 2 females (larger 35.0×26.2 mm) (ZRC 2013.0558), Niujiu Town, Shiping County, Yunnan Province, China, coll. H.C. Li, 23 February 2004. *Indochinamon phongnha* Naruse, Nguyen & Yeo, 2011 – paratypes: 5 males (25.4×20.2 – 44.2×33.2 mm), 6 females (17.9×14.1 – 43.0×32.6 mm), 1 juvenile (15.4×12.5 mm) (ZRC 2010.0168), Khe Con Khai stream, Cha Noi, Phong Nha, Quang Binh Province, Vietnam, 17°38.196'N, 106°05.928'E, 263 m asl, coll. D.C.J. Yeo and A.D.

Tran, 13 July 2004; 2 males (34.3×26.6 mm, 31.9×24.9 mm), 3 females (38.3×29.9 – 54.8×41.4 mm), 1 juvenile (17.0×13.1 mm) (ZRC 2010.0169), Cha Noi, Phong Nha, Quang Binh Province, Vietnam, Stream under bridge, 17°38.397'N, 106°06.975'E, 261 m asl, coll. D.C.J. Yeo and A.D. Tran, 13 July 2004; 11 males (16.4×13.3 – 64.2×48.0 mm), 3 females (32.6×25.9 – 38.9×30.1 mm) (ZRC 2010.0170), Vuc Tro stream, Phong Nha, Quang Binh Province, 17°38.188'N, 106°12.810'E, coll. D.C.J. Yeo and A.D. Tran, 14 July 2004; 3 females (36.1×28.1 – 42.8×33.0 mm), 2 juveniles (19.6×15.3 mm, 17.3×13.6 mm) (ZRC 2010.0171), stream near Forest Ranger station 37, Phong Nha, Quang Binh Province Vietnam, 17°31.395'N, 106°17.716'E, 86 m asl, coll. D. C. J. Yeo and A. D. Tran, 15 July 2004; 3 males (49.8×37.8 – 53.0×41.5 mm) (ZRC 2010.0172), Chay stream, Quang Binh Province, Vietnam, 17°33.146'N, 106°14.425'E, 94 m asl, coll. D.C.J. Yeo and A.D. Tran, 17 July 2004; 1 male (61.9×47.3 mm) (ZRC 2010.0173), Km 23 + 800 HCM Way, near Hang So Dua, Pong Nha National Park, Quang Binh Province, Vietnam, coll. A.D. Tran, 11 August 2001; 1 male (56.9×44.0 mm), 1 female (54.2×40.8 mm) (ZRC 2010.0174), Thac Xoi waterfall, Phong Nha National Park, Quang Binh Province, Vietnam, coll. Q.K. Hoang and V.K. Dinl, 10 August 2002. *Indochinamon tannanti* (Rathbun, 1904) – holotype: female (35.5×27.7 mm) (MNHN-B 5313), “Tonkin, montagnes du Yunnan (via Lao Koi)”, coll. Tannant, no date; 1 male (56.1×42.6 mm) (ZRC 1998.264), Hekou, Yunnan Province, southern China, coll. A.-Y. Dai, 29 November 1995. *Indochinamon tritum* (Alcock, 1909) – holotype: female (35.8×27.4 mm) (ZSI 4075/4), Sheetee Hill (= Shitee Doung), Kakhien Hills, Yunnan, China, coll. J. Anderson, no date. *Indochinamon tuijense* Naruse, Chia & Zhou, 2018 – paratypes: 2 males (larger 33.1×25.0 mm), 1 female (31.4×23.6 mm) (ZRC 2013.0555), Mang Huai Town, Yun County, Yunnan Province, China, coll. Y.F. Lu, 24 February 2004; 2 males (larger 37.5×28.8 mm), 2 females (larger 38.6×28.6 mm) (ZRC 2013.0557), Mongku Town, Shuangjiang County, Yunnan Province, China, coll. O.C. Li, 26 February 2004. *Indochinamon villosum* (Yeo & Ng, 1998) – holotype: male (44.8×34.3 mm) (ZRC 1998.276), tributary of Nam Tha River about 800 m asl, Luang Nam Tha Province, northern Laos, coll. H. Morioka, 13 November 1997; paratypes: 7 males (largest 55.9×41.4 mm), 4 females, 1 juvenile (ZRC 1998.277–285, 807–809), same data as holotype; 2 females (larger 32.9×24.5 mm) (ZRC 1998.286–287), Nam Luang about 1 km upstream of Ban Nam Luang, Nam Tha watershed, Mekong basin, Luang Nam Tha Province, northern Laos, 21°9'5"N, 101°20'34"E, coll. M. Kottelat, 22 May 1997; 1 female (39.9×30.8 mm) (ZRC 1998.288), tributary of Nam Talan about 3 km S of Ban Nateuy, Nam Tha watershed, Mekong basin, Luang Nam Tha Province, northern Laos, coll. M. Kottelat, 20 May 1997.

Diagnosis. Carapace with dorsal surface prominently rugose in large specimens (ca. 45 mm carapace width), frontal and orbital regions prominently rugose, lateral parts of anterolateral and branchial regions with strong oblique striae; mesogastric, urogastric, cardiac and intestinal regions with distinct rugosities and distinct granules (Fig. 2A, D, F); postorbital cristae distinct, margin uneven, outer edge relatively low,




Figure 1. *Indochinamon khinpyae* sp. n., colour in life, holotype male (57.1×43.2 mm) (ZRC 2018.0713). **A** dorsal view **B** ventral view of cephalothorax.

not well marked (Fig. 2A, D, F); external orbital tooth distinct, separated from anterolateral margin by deep V-shaped cleft; epibranchial tooth prominent (Fig. 2A, F); anterolateral margin lined with sharp granules, appears serrated (Fig. 2A, F); posterior margin of epistome with distinct median triangle (Fig. 2D); outer surface of chela strongly rugose, upper part rugose with granules (Fig. 3H, I); male thoracic sternum, notably sternites 3 and 4, relatively broad, surface with pits and scattered short, stiff setae (Fig. 3A); male pleon triangular; telson triangular, lateral margins gently sinuous; somite 6 transversely rectangular, much wider than long, lateral margin gently convex (Fig. 3A, B); G1 relatively stout; outer margin of subterminal segment with distinct broad cleft on distal part, terminal segment subcylindrical, gently curving outwards, no visible dorsal flap, distal part tapering to rounded tip (Fig. 4A–D).

Description of male holotype. Carapace transversely ovate, distinctly wider than long (width to length ratio 1.32); dorsal surface gently convex from frontal view, regions not prominently inflated; with scattered very short setae, appears glabrous (Fig. 2A, D, F). Frontal and orbital regions prominently rugose; lateral parts of anterolateral and branchial regions covered with strong oblique striae; mesogastric, urogastric, cardiac and intestinal regions covered with rugosities and distinct granules; suborbital region with small granules on lateral parts; pterygostomial, subhepatic and sub-branchial regions rugose to granulose (Fig. 2A, D, F). Epigastric cristae distinct, rugose, not cristate, separated by broad, median Y-shaped furrow; epigastric cristae just anterior of postorbital cristae, separated by short furrow; postorbital cristae distinct, margin uneven, prominently raised, subparallel to frontal margin, outer edge relatively low, not prominent (Fig. 2A, D, F). Cervical grooves deep, not reaching lateral margins, connected to deep H-shaped median gastric groove (Fig. 2A, F). Frontal margin almost straight, appears entire in dorsal view, gently sinuous in frontal view (Fig. 2A, D, F). External orbital tooth distinct, triangular, outer margin more than twice length of inner margin, demarcated from rest of anterolateral margin by deep V-shaped cleft; epibranchial tooth prominent, sharp (Fig. 2A, F). Anterolateral margins convex, cristate, lined with sharp granules, appears serrated (Fig. 2A, F). Posterolateral margin gently sinuous, converging towards convex posterior carapace margin (Fig. 2A, F). Orbits subovate; eye filling orbital space; eye peduncle relatively short, stout; cornea large, round, pigmented (Fig. 2D). Supraorbital margin almost straight (Fig. 2F). Suborbital margin concave, complete, lined with low granules (Fig. 2D). Antennae short, stretching across base of eyes; antennules short, folding transversely in rectangular fossa (Fig. 2D). Posterior margin of epistome with distinct median triangle, lateral margin sinuous (Fig. 2D).

Third maxillipeds covering most of buccal cavity when closed; ischium subrectangular, with distinct median groove, surface with scattered pits and short setae; merus subquadrate, slightly wider than long, surface rugose, margins cristate, anteroexternal angle angular but not produced; exopod slender, reaching to about one-third length of merus, with elongate flagellum that reaches across width of merus (Fig. 3F).

Chelipeds asymmetrical, right larger (Fig. 2A). Anterior margin of basis-ischium lined with small sharp granules; margins of merus lined with low sharp granules, appears weakly serrated. Outer surface of carpus rugose, inner distal angle with large sharp




Figure 2. *Indochinamon khinpyae* sp. n. **A, D, F** holotype male (57.1×43.2 mm) (ZRC 2018.0713) **B, E** paratype male (47.2×36.4 mm) (ZRC 2018.0714) **C, G** paratype male (34.3×26.6 mm) (ZRC 2018.0714). **A–C** overall habitus **D, E** frontal view of cephalothorax **F, G** dorsal view of carapace.

tooth and basal tooth (Fig. 2A). Outer surfaces of chelae strongly rugose, upper part rugose with granules; major chela stouter, shorter than minor chela (Fig. 3H, I). Fingers of major chela short, stout, gently curved, subequal to palm, outer surface lined with 3 rows of pits; cutting edges of both fingers with variously sized sharp teeth and denticles; dorsal margin of dactylus with low tubercles and granules (Figs 2A, 3H). Fingers of minor chela similar to major chela in form but relatively more slender (Figs 2A, 3I).

Ambulatory legs short, segments relatively stout; second pair longest, last pair shortest (Fig. 2A). Merus short, stout, outer surface rugose, dorsal margin uneven, subcristate, without subdistal spine or tooth; carpus rugose, dorsal margin with crista, outer surface with low submedian crista on first to third legs, that on fourth leg




Figure 3. *Indochinamon khinpyae* sp. n. **A, B, E, F, H, I** holotype male (57.1×43.2 mm) (ZRC 2018.0713) **C, G** paratype male (47.2×36.4 mm) (ZRC 2018.0714) **D** paratype male (34.3×26.6 mm) (ZRC 2018.0714). **A, C, D** anterior thoracic sternum and pleon **B** posterior thoracic sternum and pleon **E** sternopleonal cavity **F, G** right third maxilliped **H** outer view of right chela **I** outer view of left chela.

smooth; dorsal margin of propodus with crista, outer surface with low, submedian crista; dactylus relatively short, gently curved, quadrate in cross section, margins with short, sharp pectinate spines (Fig. 2A).

Thoracic sternum, notably sternites 3 and 4, relatively broad, surface with pits and scattered short, stiff setae (Fig. 3A). Sternites 1, 2 completely fused to form broadly triangular plate; separated from sternite 3 by distinct, gently concave suture (towards buccal cavity); sternites 3, 4 completely fused, with shallow incomplete groove demarcating suture (Fig. 3A, B, E). Penis coxal, on condyle of coxa of fourth ambulatory leg. Sternopleonal cavity deep, reaching imaginary line connecting posterior edges of cheliped coxae (Fig. 3A, E). Male pleonal locking tubercle low, round, on posterior third of sternite 5 (Fig. 3E).

Pleon triangular, all somites and telson free; telson triangular, lateral margins gently sinuous; somite 6 transversely rectangular, much wider than long, lateral margin gently convex; somites 3–5 trapezoidal, gradually decreasing in width, increasing in length; somites 1 and 2 subrectangular, very wide, reaching to bases of coxae of fourth ambulatory legs, thoracic sternite 8 not visible when pleon closed (Fig. 3A, B).

G1 relatively stout; subterminal segment gently sinuous, proximal part broad, gradually tapering to median part, outer margin with distinct broad cleft on distal part; clearly separated from terminal segment by prominent dilation; terminal segment subcylindrical, no dorsal flap visible, gently curving outwards, distal part tapering to rounded tip (Fig. 4A–D). G2 elongate, much longer than G1; basal segment longer than distal segment (Fig. 4E).

Variation. The carapace tends to get less broad in smaller specimens and females (width to length ratio 1.26–1.30). The regions in smaller specimens is less sculptured (Fig. 2B, E) with the rugosities restricted mostly to lateral margins (Fig. 2C, G). The third maxilliped ischium is slightly longer in smaller individuals (Fig. 3G). The male pleon is proportionately less broad in smaller individuals with somite 6 more quadrate as they are smaller (Fig. 3C, D). In adult males, the G1 does not vary substantially although the cleft on the outer part of the distal section of the G1 subterminal segment is relatively less distinct (Fig. 4F). Smaller males (ca. 30 mm carapace width), however, not only have the G1 terminal segment relatively shorter and less curved, the cleft on the subterminal segment is also not discernible (Fig. 4H). The adult female has the pleon completely covering the thoracic sternum (Fig. 5A), the vulva is large, raised, ovate and positioned on the anterior half of sternite 6, pushing into the margin with sternite 5 (Fig. 5B).

Etymology. The species is named after Ms Khin Pyae Pyae Thaw Thaw who collected the specimens used for this study. Her name is used here as a noun in apposition.

Colour. In life, the dorsal surfaces of the carapace and outer surfaces of the chelipeds are dark brown; with the ventral surfaces orangish-red; and the ambulatory legs are dark brown to orangish red (Fig. 1).

Habitat. The type locality, Malikha, is a fast-flowing river, the substrate consisting of rocks of various sizes, with the bank sandy. The banks are densely lined with tall trees. This river is a branch of the Ayeyarwady River (= Ayrwarwady River or Myit-sone) and is about 43 km north of Myitkyina, the capital city of Kachin State.

Remarks. Five species of *Indochinamon* have been reported from and near Myanmar: *I. andersonianum*, *I. edwardsii*, *I. hirtum*, *I. hispidum* and *I. tritum* (cf. Alcock 1909, 1910; Bott 1970; Yeo and Ng 2007). All these species were collected by John Anderson from the area east of Bhamo, mostly in the Kakhyen Hills (= Kachin Moun-




Figure 4. *Indochinamon khinpyae* sp. n. **A–E** holotype male (57.1×43.2 mm) (ZRC 2018.0713); **F, G** paratype male (47.2×36.4 mm) (ZRC 2018.0714) **H** paratype male (34.3×26.6 mm) (ZRC 2018.0714). **A** left G1 (ventral view) **B** left G1 (dorsal view) **C** terminal segment of left G1 (ventral view) **D** terminal segment of left G1 (dorsal view) **E** left G2 **F, H** left G1 (ventral view, setae not drawn) **G** left G2 (setae not drawn). Scale bar: 1.0 mm.

tains), in what is today Myanmar and Yunnan (China). One site, Ponsee, which is the type locality of *I. edwardsii* and *I. hispidum* (and where *I. andersonianum* has also been found), does not appear in most modern maps but this village is in the Dehong, Longchuan area in Yunnan, China (ca. 24°25'34.5"N, 97°53'57.3"E) (cf. Anderson 1876). Until the present record of *I. khinpyae*, no species had been reported from the mountains north of Bhamo and Myitkyina in Myanmar.




Figure 5. *Indochinamon khinpyae* sp. n., paratype female (39.3×31.1 mm) (ZRC 2018.0714). **A** anterior thoracic sternum and pleon **B** sternopleonal cavity and vulvae.

Adult male specimens of *I. khinpyae* have a strongly sculptured and very rough carapace (Fig. 2F), the G1 terminal segment is relatively long, gently curved, distally bent and the dorsal margin has no trace of a flap (Fig. 4A–D, F, H). In *I. andersonianum*, even large males (50 mm carapace width) have the gastric regions relatively smooth with the rest of the surfaces also less rugose and granulose, and the male pleon is proportionately more narrow (Wood-Mason 1871: pl. 27 figs 16, 17, 20; Bott 1970: pl. 44 fig. 14; unpublished data). The G1 of *I. andersonianum* is also quite different with the terminal segment straight, slender and tapering towards the tip (Bott 1970: pl. 37 fig. 16). The taxonomy of *I. andersonianum* has been confused and many species previously referred to it have turned out to be other taxa (see Ng and Naiyanetr 1993). The figure of *I. andersonianum* by Alcock (1910: pl. 10 fig. 40) is actually a separate species, *Potamiscus rangoonensis* (Rathbun, 1904) (unpublished data). The G1 of the smaller paratype male of *I. khinpyae* (34.3×26.6 mm, ZRC 2018.0714) superficially resembles that of *I. edwardsii* (the type of which is about the same size) but in *I. edwardsii*, the anterolateral margins are prominently serrated even in smaller specimens (Wood-Mason 1871: pl. 27 figs 11, 12; Alcock 1910: pl. 14 fig. 43; unpublished data) (versus anterolateral margins finely granulated or weakly serrated in *I. khinpyae*; Fig. 2F, G); the upper part of the palm of the chela has many large tubercles (Wood-Mason 1871: pl. 27 figs 11, 14; Alcock 1910: pl. 14 fig. 43; unpublished data) (versus with no large tubercles present in *I. khinpyae*; Figs 2A–C, 3H–I); and the lateral margins of the male telson are concave (Wood-Mason 1871: pl. 27 fig. 14; unpublished data) (versus lateral margins gently sinuous to almost straight in *I. khinpyae*; Fig. 3A, C, D). These differences also apply for *I. hirtum* (cf. Alcock 1919: pl. 10 fig. 42; unpublished data). Compared to *I. tritum*, known only from a 35.8×27.4 mm female from the Shitee Hills in the Kakhyen Hills in Yunnan (just north of Ponsee, cf. Anderson 1876: 420), the lateral margin of the postorbital crista of *I. khinpyae* is less clearly marked (Fig. 2F, G) (versus distinctly formed and clearly demarcated from the lateral branchial region in *I. tritum*; cf. Alcock 1910: pl. 14 fig. 69); and the propodus of the last ambulatory leg is proportionately longer (Fig. 2A–C) (versus much shorter in *I. tritum*; cf. Alcock 1910: pl. 14 fig. 69). Compared to *I. hispidum*, described from a male 43.0×31.0 mm from Ponsee, *I. khinpyae* can easily be distinguished by its more rugose dorsal carapace surface (Fig. 2A, B,

F) (versus smooth dorsal carapace surface in *I. hispidum*; cf. Wood-Mason 1871: pl. 27 figs 1, 2); rugose outer surface of the chela (Fig. 3H, I) (versus smooth in *I. hispidum*; cf. Wood-Mason 1871: pl. 27 fig. 4); and the male pleon is proportionately broader with the telson more broadly triangular (Fig. 3A, C, D) (versus male pleon more narrow with the telson acutely triangular in *I. hispidum*; cf. Wood-Mason 1871: pl. 27 fig. 5).

With regards to the other species of *Indochinamon*, they can be separated into several groups on the basis of their G1s. The type species, *I. villosum*, has a relatively short and stout G1 terminal segment which is gently bent and is conical to subconical in shape without an obvious dorsal flap, a character shared with *I. ahkense*, *I. bavi*, *I. bhumibol*, *I. boshanense*, *I. changpoense*, *I. chinghungense*, *I. dangi*, *I. daweishanense*, *I. flexum*, *I. guttum*, *I. jianchuan*, *I. jinpingense*, *I. kimboiense*, *I. menglaense*, *I. mieni*, *I. orleansi*, *I. ou*, *I. parpidum*, *I. phongnha*, *I. tannanti*, *I. xinpingense*, and *I. yunlongense* (including *I. edwardsii* and *I. hispidum*) (cf. Bott 1970; Yeo and Ng 1998; Dai 1999; Naruse et al. 2011, 2018; unpublished data). The other species have G1 terminal segments which are slender, elongate, and straight or curved; or relatively short and strongly bent (cf. Ng and Naiyanetr 1993; Dai 1999; Do et al. 2016; Naruse et al. 2018; unpublished data). The G1 of *I. khinpyae* closely resembles that of *I. changpoense* and *I. daweishanense* (both from Yunnan) but these two species have only a shallow cleft on the outer margin of the subdistal part of the subterminal segment, and the terminal segment is proportionately shorter and straighter (Dai 1999: figs 85–4, 5; 87–4, 5), even though the types are comparable in size to the holotype of *I. khinpyae*. Similarly, *I. yunlongense* (described from a small male 19.0×16.1 mm from Yunnan) has a superficially similar G1 structure to *I. khinpyae*, except that the terminal segment is much straighter (Dai 1999: fig. 84–4, 5). The strongly sculptured and rugose carapace of large *I. khinpyae* allies it with large species like *I. kimboense* and *I. bavi* (both from Vietnam) but in these species, the cleft on the outer margin with of the G1 subterminal segment is shallow and not distinct (cf. Naruse et al. 2011: fig. 3a, b, d, e), even for specimens larger than the holotype of *I. khinpyae*, which has a prominent broad cleft (Fig. 4A–D). The G1 terminal segment of *I. kimboense* and *I. bavi* (as well as *I. cua*, *I. orleansi*, *I. ou* and *I. tannanti*) are also distinctly tapering towards the tip (Naruse et al. 2011: fig. 3a, b, d, e), unlike the subtruncate condition in *I. khinpyae* (Fig. 4A–D). Compared to *I. phongnha* (from Vietnam), which also has the carapace regions distinct, the surfaces are smoother, notably the median and posterior parts which are smooth, even in large specimens (Naruse et al. 2011: fig. 7) (strongly rugose in large *I. khinpyae*; Figs 1A, 2A, F); and the G1, while it has a strong cleft on the outer margin of the subterminal segment, the terminal segment is sharply tapering (Naruse et al. 2011: fig. 9a, b) (terminal segment subcylindrical in large *I. khinpyae*; Fig. 4A–D). The strong cleft on the outer margin of the G1 subterminal segment of *I. khinpyae* is character also shared with *I. cua* from Thailand, but in this species, the cleft is relatively broader and the terminal segment is tapering distally (Yeo and Ng 1998: fig. 4B, C, E, G); and the carapace regions are proportionately much smoother (Yeo and Ng 1998: fig. 7A). In *I. lipkei* from Thailand, the dorsal carapace surface, even in large specimens, is less well marked with the median parts much less rugose (Ng and Naiyanetr 1993: fig. 12A);

pleonal somite 6 is distinctly trapezoidal in shape (Ng and Naiyanetr 1993: fig. 12C); and the G1 terminal segment is strongly bent at about 60° along the longitudinal axis (Ng and Naiyanetr 1993: fig. 47B–E) (versus the dorsal carapace regions are more rugose, pleonal somite 6 is weakly trapezoidal, and the G1 terminal segment is bent at about 45° in *I. khinpyae*; Figs 2F, 3A, C, D, 4A–D).

Indochinamon khinpyae is not known to be threatened by any developments, and the forests and streams where it has been found are isolated and not easily assessable by man. As such, the species is classified under taxa of Least Concern for the moment (cf. Cumberlidge et al. 2009, 2012).

A note on *I. manipurensis* (Alcock, 1909) is necessary. Takeda et al. (2012: 207) noted that specimens they had of this species did not possess a flagellum on the exopod of the third maxilliped, and as such, the species should be transferred to *Potamiscus* Alcock, 1909. However, the types of this species do have a flagellum (Yeo and Ng 2007; unpublished data), so Takeda et al.'s (2012) specimens will need to be checked to ascertain their identity. As such, for the moment, we retain the species in the genus *Indochinamon*.

Acknowledgements

The second author is most grateful to Khin Pyae Pyae Thaw Thar (Department of Zoology, Myitkyina University, Kachin State) for collecting the specimens and passing them to her for this study. The first author thanks Darren Yeo for sharing his unpublished photographs and data of various *Indochinamon* species; and to Tohru Naruse for his many helpful comments on the manuscript.

References

- Alcock A (1909) Diagnoses of new species and varieties of freshwater crabs. No. 4. Records of the Indian Museum 3(4): 375–381.
- Alcock A (1910) Catalogue of the Indian decapod Crustacea in the collection of the Indian Museum. Part I. Brachyura. Fasciculus II. The Indian fresh-water crabs – Potamonidae. Calcutta: Trustees of the Indian Museum, 1–130.
- Anderson J (1876) Mandalay to Momien: a narrative of the two expeditions to Western China of 1808 and 1875 under Colonel Edward B. Sladen and Colonel Horace Browne. London: Macmillan and Co., 479 pp.
- Bott R (1970) Die Süßwasserkrabben von Europa, Asien, Australien und ihre Stammesgeschichte. Eine Revision der Potamoidea und der Parathelphusoidea (Crustacea, Decapoda). Abhandlungen der Senckenbergischen naturforschenden Gesellschaft 526: 1–338. [figs 1–8, pls 1–58, map 1]
- Cumberlidge N, Ng PKL, Yeo DCJ (2012) Freshwater crabs of the Indo-Burma hotspot: diversity, distribution, and conservation. In: Allen DJ, Smith KG, Darwall WRT (Eds) The status and distribution of Freshwater biodiversity in Indo-Burma, IUCN, Cambridge, 102–113.

- Cumberlidge N, Ng PKL, Yeo DCJ, Magalhães C, Campos MR, Alvarez F, Naruse T, Daniels SR, Esser LJ, Attipoe FYK, Clotilde-Ba F-L, Darwall W, Mcivor A, Collen B, Ram M (2009) Freshwater crabs and the biodiversity crisis: importance, threats, status, and conservation challenges. *Biological Conservation* 142: 1665–1673. <https://doi.org/10.1016/j.biocon.2009.02.038>
- Dai AY (1999) *Fauna Sinica: Arthropoda Crustacea Malacostraca Decapoda Parathelphusidae Potamidae*. Science Press, Beijing, 501 pp. [figs 1–238, pls 1–30]
- Davie PJF, Guinot D, Ng PKL (2015) Anatomy and functional morphology of Brachyura. In: Castro P, Davie PJF, Guinot D, Schram FR, von Vaupel Klein JC (Eds) *Treatise on Zoology – Anatomy, Taxonomy, Biology. The Crustacea. Volume 9C–I. Decapoda: Brachyura (Part 1)*, 11–163. https://doi.org/10.1163/9789004190832_004
- Do VT, Nguyen TC, Le HA (2016) A new species of the genus *Indochinamon* Yeo & Ng, 2007 (Crustacea: Brachyura: Potamoidea: Potamidae) from northern Vietnam. *Raffles Bulletin of Zoology* 64: 187–193.
- Naruse T, Chia JE, Zhou X (2018) Biodiversity surveys reveal eight new species of freshwater crabs (Decapoda: Brachyura: Potamidae) from Yunnan Province, China. *PeerJ* 6: e5497. <https://doi.org/10.7717/peerj.5497>
- Naruse T, Nguyen XQ, Yeo DCJ (2011) Three new species of *Indochinamon* (Crustacea: Decapoda: Potamidae: Potamiscinae) from Vietnam, with a redescription of *Ranguna (Ranguna) kimboiensis* Dang, 1975. *Zootaxa* 2732: 33–48.
- Ng PKL (1988) *The Freshwater Crabs of Peninsular Malaysia and Singapore*. Department of Zoology, National University of Singapore, Shinglee Press, 156 pp. [figs 1–63, 4 colour pls]
- Ng PKL (1996) On a new genus and new species of potamid crab (Brachyura) from Burma. *Crustaceana* 69(8): 1005–1013. <https://doi.org/10.1163/156854096X00439>
- Ng PKL (2018) On two new species of potamid and sesarmid crabs (Decapoda, Brachyura) from caves in Myanmar. *Crustaceana* 91(2): 185–197. <https://doi.org/10.1163/15685403-00003735>
- Ng PKL, Guinot D, Davie PJF (2008) *Systema Brachyurorum: Part I. An annotated checklist of extant brachyuran crabs of the world*. *Raffles Bulletin of Zoology, Supplement* 17: 1–286.
- Ng PKL, Kosuge T (1997) *Lobothelphusa gibbosa*, a new species of potamid crab (Decapoda, Brachyura) from rice fields in Burma. *Crustaceana* 70(7): 813–821. <https://doi.org/10.1163/156854097X00258>
- Ng PKL, Naiyanetr P (1993) New and recently described freshwater crabs (Crustacea: Decapoda: Brachyura: Potamidae, Gecarcinucidae and Parathelphusidae) from Thailand. *Zoologische Verhandlungen* 284: 1–117. [figs 1–68]
- Ng PKL, Whitten AJ (2017) On a new species of *Shanphusa* Yeo & Ng, 2007 (Brachyura, Potamoidea, Potamidae), from a cave in central Myanmar. *Crustaceana* 90(2): 235–245. <https://doi.org/10.1163/15685403-00003646>
- Takeda M, Hiromu S, Singh TS (2012) Some freshwater crabs from northeast India bordered on Myanmar. *Journal of Teikyo Heisei University* 23(1): 199–213.
- Wood-Mason J (1871) *Contribution to Indian Carcinology – On Indian and Malayan Telphusidae, Part I*. *Journal of the Asiatic Society of Bengal*, 40(2): 189–207, 449–454. [pls 11–14, 17]

- Yeo DCJ, Ng PKL (1998) Freshwater crabs of the *Potamon tannanti* species group (Crustacea: Decapoda: Brachyura: Potamidae) from northern Indochina. *Raffles Bulletin of Zoology* 46(2): 627–650.
- Yeo DCJ, Ng PKL (1999) The state of freshwater crab taxonomy in Indochina (Decapoda, Brachyura). In: Schram FR, von Vaupel Klein JC (Eds) *Crustaceana, Special Volume, Crustaceans and the Biodiversity Crisis*, 637–646.
- Yeo DCJ, Ng PKL (2004) Recognition of two subfamilies in the Potamidae Ortmann, 1896 (Brachyura, Potamidae) with a note on the genus *Potamon* Savigny, 1816. *Crustaceana* 76(10): 1219–1235. <https://doi.org/10.1163/156854003773123456>
- Yeo DCJ, Ng PKL (2007) On the genus “*Potamon*” and allies in Indochina (Crustacea: Decapoda: Brachyura: Potamidae). *Raffles Bulletin of Zoology, Supplement* 16: 273–308.