

# New records of leaf-miner flies (Diptera, Agromyzidae) from Morocco

Mourad Doukale Daief<sup>1</sup> , Michael von Tschirnhaus<sup>2</sup> , Kawtar Kettani<sup>1</sup> 

1 Laboratory of Ecology, Systematics and Conservation of Biodiversity (LESCB), URL-CNRST N°18, FS, Abdelmalek Essaadi University, Tetouan, Morocco

2 Faculty of Biology, Biological Collection, University of Bielefeld, P.O. Box 100 131, 33501 Bielefeld, Germany

Corresponding author: Mourad Doukale Daief ([mouraddoukale@gmail.com](mailto:mouraddoukale@gmail.com))

## Abstract

New records are provided of the family Agromyzidae (leaf-miner flies) from Morocco. Thirty agromyzid species are newly recorded, seven of these in Agromyzinae: *Agromyza* (*Agromyza anthracina*, *A. conjuncta*, *A. mobilis*, *A. myosotidis*, *A. nigrescens*), *Hexomyza* (*H. simples*), and *Ophiomyia* (*O. pinguis*). The other 23 new records are in Phytomyzinae: *Amauromyza* (*Cephalomyza*) *karli*, *Calycomyza* *flavomaculata*, *C. solidaginis*, *Cerodontha* (*Butomomyza*) *eucaricis*, *C. (Dizygomyza)* *luctuosa*, *C. (Icteromyza)* *geniculata*, *Chromatomyia* *centaurii*, *Liriomyza* *amoena*, *L. brassicae*, *L. strigata*, *Phytoliriomyza* *arctica*, *P. perpusilla*, *P. pteridii*, *Phytomyza* *anemones*, *Ph. clematidis*, *Ph. crassisetata*, *Ph. notata*, *Ph. plantaginis*, *Ph. rufipes*, *Ph. vitalbae*, *Pseudonapomyza* *confusa*, *Ps. palavae*, and *Ps. vota*. The genus *Calycomyza* Hendel is newly reported from the country. The present study is based on recent entomological surveys and has increased the total number of agromyzid species in Morocco to 92. Information on the distribution and ecology of each species is provided.

**Key words:** Agromyzidae, faunistics, flies, Moroccan fauna



Academic editor: Owen Lonsdale

Received: 15 November 2024

Accepted: 14 January 2025

Published: 5 March 2025

ZooBank: <https://zoobank.org/57CCF242-9E56-4623-A6A1-B9D04706C71D>

Citation: Doukale Daief M, von Tschirnhaus M, Kettani K (2025) New records of leaf-miner flies (Diptera, Agromyzidae) from Morocco. ZooKeys 1230: 81–97. <https://doi.org/10.3897/zookeys.1230.141900>

Copyright: © Mourad Doukale Daief et al.  
This is an open access article distributed under  
terms of the Creative Commons Attribution  
License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Introduction

The family Agromyzidae, or leaf-miner flies (Diptera, Acalyptratae), is widely distributed in the Palaearctic Region, in which it comprises more than 1,303 species (von Tschirnhaus and Groll 2024a, 2024b). Of these, 984 species are recorded from Europe (Černý, pers. comm. to MvT, 24.ix.2024). They are small to medium-sized acalyptratae flies (0.8–6.5 mm), with variable colour, from black to yellow. They are considered as the third largest of Acalyptratae families which possess entirely phytophagous larvae that develop in a wide range of hosts in the plant kingdom (Hering 1957, 1966; Spencer 1972, 1990). They inhabit a wide variety of habitats in forests, high mountains, arctic tundra and all types of lowlands, wetlands, and marine coasts. They play a major role in agricultural entomology and are therefore of economic interest (Spencer 1973), as larvae of many species are known to damage crop plants (Dempewolf 2004; Lonsdale et al. 2023). This family of Diptera is also of significant ecological importance (Spencer 1973). The larvae of most species are monophagous or

oligophagous, but polyphagy is the exception. They attack different parts of plants such as leaves, stems, sometimes the roots, flower heads, seeds and the young xylem of shrubs and trees, erroneously addressed by authors as the cambium. After host plant identification, the form of the leaf-mine can be useful to identify the species that caused the damage (Černý et al. 2018).


In Morocco, the first record of the family Agromyzidae was reported by Becker and Stein (1913) who found *Amauromyza morionella* [at that time included in the genus *Agromyza*] at Tangier. After the 1930s, new records were added by Séguy's study of the Diptera of Morocco. He reported seven species for the country (Séguy 1936, 1953) with a few other records by Hendel (1931–1936) and Kozlowsky and Rungs (1932). Only three species have been described from Morocco after 1936. These are *Aulagromyza cydoniae* [described by Hendel (1936) in combination with *Phytomyza*], *Aulagromyza atlantidis* [described by Spencer (1967) in combination with *Paraphytomyza*], and *Cerodontha (Icteromyza) rozkosnyi* described by Černý (2007). It was not until the 1960s that more records were made with Spencer's (1967, 1973) studies on the Agromyzidae of Morocco and whose contribution had slightly improved the knowledge of these flies by adding 17 species to the Moroccan fauna. After that, few further records have been added by foreign researchers such as Griffiths (1974, 1967), Geipert et al. (1994), Maarouf (2003), and Hanafi and Schnitzler (2004). The most important contribution to the improvement of our knowledge of Moroccan Agromyzidae was made recently by Černý through his many studies (Černý 2007, 2009, 2010, 2012, 2013, 2019; Černý and Merz 2006, 2007; Černý and von Tscharnhaus 2014; Černý et al. 2020). Those articles provided a further 20 species to the country. Beyond that, our knowledge of Moroccan agromyzid fauna is still poor and many gaps remain for this diverse family of Diptera. Many regions of Morocco were still unexplored such as the Rif region in northern Morocco. Indeed, only 62 species of Agromyzidae are listed so far in the catalogue of the Diptera of Morocco compiled by Kettani et al. (2022). Of these, only seven species were recorded from the Rif.

The aim of this study is to present new faunistic records, in particular from some regions which have not previously been explored as an attempt to fill gaps in our knowledge of Moroccan leaf-miners.

The distribution of the species given in this study was mainly gathered from the literature provided by Papp and Černý in their four volumes (Papp and Černý 2015, 2016, 2017, 2020 [as 2019]), Černý et al. 2020; Lonsdale 2021, Lonsdale et al. 2023, and using the file cards and bibliography for world Agromyzidae in von Tscharnhaus and Groll (2024a, 2024b).

## Materials and methods

The specimens examined in the present study were collected by MD and to a large extent by KK and her students. Insects were captured by sweep net and by Malaise trap at different localities throughout Morocco from 2012 to 2023. Of the 140 sites surveyed, 38 were home to the agromyzid species mentioned in this article. The sites are mainly in the Rif and Atlas Mountains region (Fig. 1) where a variety of habitats including forests, grasslands, riverbanks, and wetlands were prospected. Table 1 summarizes the main geographical cartographic data and types of habitats of the prospected sites.



**Figure 1.** Location of the studied sites in Morocco **A** the Rif at the north of Morocco **B** central Morocco with Atlantic Plain and Atlas Mountains.

The Rif region consists of a mountainous domain overlooking the Mediterranean Sea at the north of Morocco. The climate is mainly of the Mediterranean type and is characterized by high levels of precipitation and is considered very rich and diverse in vegetation cover (Benabid 1982; Valdés Castrillón et al. 2002). It integrates five bioclimatic stages: per-humid and humid which are mainly restricted to the high mountain summits, while the subhumid, semi-arid and arid parts are more widespread in the rest of the Rif. The Atlas region consists of mountain massifs located in the centre and further south of Morocco, made up of three large mountain ranges: High Atlas, Middle Atlas and Anti-Atlas with high summits incised by deep valleys. The bioclimates of this area include perhumid, humid, sub-humid, semi-arid and arid zones with more than 800 mm of annual precipitation on the highest peaks (HCEFLCD 2017) which have favoured a great diversity of vegetation such as *Tetraclinis articulata*, *Argania spinosa*, *Quercus suber*, *Quercus faginea*, *Quercus ilex*, *Cedrus atlantica*, and *Juniperus thurifera* (Mokhtari et al. 2014).

Most specimens captured in the field were stored in 70% ethanol until identification. Each specimen was examined in detail before and after its genitalia were mounted. The genitalia, cut off from the rest of the abdomen, were cleared in 10% KOH at 80 °C for 8 or 10 min and then transferred into 10% acetic acid for ~ 5 min, and finally placed in a droplet of glycerine. The genitalia are firmly embedded in glycerine on glass-slides which were sealed by resistant nail varnish to protect them from dust and from drying out. All species have been identified based on the male genitalia. The second author confirmed and validated all the species identifications by checking all photographs.

**Table 1.** Localities with respective biogeographical areas, coordinates and altitudes of the studied sites.

<b>Code of locality</b>	<b>Biogeographic area, locality, collecting site (protected area)</b>	<b>Geographic coordinates</b>	<b>Altitude (m)</b>	<b>Habitat</b>
1	Rif, Talassemiane, Maison forestière (NPTL)	35°8'6.11"N, 5°8'18.06"W	1696	Fir forest
2	Rif, Chefchaouen, Bab Tariouant	35°01'7.68"N, 5°0'35.999"W	1429	Oak forest ( <i>Quercus pyrenaica</i> )
3	Rif, Tarzout, Riba (PNPB)	35°15'755"N, 5°99'691"W	1421	Mixed forest
4	Rif, Talassemiane, Place Espagna (NPTL)	35°05'938"N, 5°08'956"W	1386	Fir forest
5	Rif, El Anasser, Dayat El Anasser	35°00'59.8"N, 4°59'59.77"W	1383	Around pond
6	Rif, Targuist, Beni Bounzar	34°50'36"N, 4°25'23"W	1340	Cedar forest
7	Rif, Afeska, Oued Afeska (NPTL)	35°10'18.4"N, 5°11'10.5"W	1293	Banks of mountain stream
8	Rif, Jbel Kelti, Oued Tanina	35°20'45.50"N, 5°19'01.70"W	1195	Banks of mountain stream
9	Rif, Bouhachem, Amsemlil (PNPB)	35°15'44.424"N, 5°26'0.276"W	1059	Edges of mountain peat bog
10	Rif, Ametrasse, Oued Ametrasse	35°05'01"N, 5°05'03"W	841	Riverbank
11	Rif, Tazia, Dayat Tazia	35°20'52"N, 5°33'12"W	720	Edges of pond
12	Rif, Akchour, Chellal Kbir (NPTL)	35°13'8"N, 5°8'19"W	680	Banks of waterfall
13	Rif, Bni Hassan, Triwa	35°16'50.7"N, 5°22'01.6"W	654	Meadow
14	Rif, Bni Darkoul, Oued Souk Lhad	35°3'26"N, 5°4'10"W	613	Riverbank
15	Rif, Chefchaouen, Bni Ahmed	35°20'45.50"N, 5°19'01.70"W	590	Agricultural field
16	Rif, Tazrout, Adrou (PNPB)	35°13'32.30"N, 5°19'24.35"W	556	Mixed forest
17	Rif, Akchour, Pont de Dieu (NPTL)	35°13'42"N, 5°10'31"W	536	Riverbank
18	Rif, Akchour, Chellal Sghir (NPTL)	35°14'21"N, 5°10'37"W	424	Banks of waterfall
19	Rif, Talembote, Oued Talembote (NPTL)	35°14'20.586"N, 5°10'42.037"W	405	Riverbank
20	Rif, Fahs-Anjra, Rmel	35°38'57.4"N, 5°35'41.4"W	372	Riverbank
21	Rif, El Hamma, Masjid El Hamma	35°23'6"N, 5°30'46"W	330	Matorral dominated by cork oak trees
22	Rif, Tétouan, Oued Raouz	35°26'23.03"N, 5°18'41.70"W	265	Matorral dominated by pine trees
23	Rif, Rmilat, Parc Perdicaris	35°47'26.675"N, 5°51'12.938"W	223	Urban park
24	Rif, Oulad Sidi Mansour, Oued Tassikeste	35°14'05"N, 3°56'37"W	144	Riverbank
25	Rif, Souk El Had, Oued El Koub	35°1'17.88"N, 5°25'19.98"W	124	Riverbank
26	Rif, M'Diq, Koudiat Taifour	35°40'28.706"N, 5°19'1.841"W	100	Pine forest
27	Rif, Ben Karrich, Oued Mhajrate	35°30'39"N, 5°26'27"W	67	Riverbank
28	Rif, Tétouan, Khemis Anjra	35°33'42.453"N, 5°27'34.246"W	63	Riverbank
29	Rif, Tétouan, Oued Khemis	35°39'51.00"N, 5°30'29.00"W	61	Riverbank
30	Rif, Kitane, Oued Zarka	35°32'24.72"N, 5°20'23.58"W	49	Riverbank
31	Eastern Morocco, Taourirt, El Aïoun Sidi Mellouk	34°35'25.336"N, 2°31'4.263"W	589	Agricultural field
32	Atlantic Plain, Sidi Yahya Gharb, forest of Sidi Yahya Gharb	34°18'16"N, 6°18'38"W	20	Eucalyptus forest
33	Middle Atlas, Ifrane, Lac Zarouka (NPIF)	33°32'36.163"N, 5°5'45.277"W	1615	Edges of mountain lake
34	Middle Atlas, Ifrane, Zaouiet Ifrane (NPIF)	33°33'32.884"N, 5°51'0.31"W	1535	Cedar forest
35	High Atlas, Al Haouz, Amizmiz (NPTB)	31°12'54.93"N, 8°13'6.14"W	991	Mixed forest
36	Anti-Atlas, Taroudant, Taliouine	30°31'56"N, 7°55'27"W	928	Agricultural field
37	Anti-Atlas, Errachidia, Aoufous	31°44'31.607"N, 4°11'54.896"W	906	Agricultural field
38	Anti-Atlas, Agadir Idaoutanane, Vallée du Paradis	30°21'6.84"N, 9°18'46.8"W	587	Riverbank

Specimens are housed in the Laboratory of Ecology, Systematics and Conservation of the Biodiversity (**LESCB**) in the Faculty of Sciences of Tetouan in the personal collection of Mourad DOUKALE DAIEF.

The following abbreviations are used for the material examined:

- NPIF** National Park of Ifrane  
**NPTB** National Park of Toubkal  
**NPTL** National Park of Talassemtane  
**PNPB** Project of Natural Park of Bouhachem

## Results

### Subfamily Agromyzinae

#### *Agromyza anthracina* Meigen, 1830

**Material examined.** MOROCCO. • 1 ♂; Rif, M'Diq, Koudiat Taifour; 3 Apr. 2018; K. Kettani leg.; sweep net; LESC-B-R18/21.

**Distribution.** Palaearctic species, widespread in Europe. First record from Morocco.

**Habitat.** Collected in pine forest (*Pinus pinaster*).

#### *Agromyza conjuncta* Spencer, 1966

**Material examined.** MOROCCO. • 2 ♂♂; Rif, Souk El Had, Oued El Koub; 23 Apr. 2018; K. Kettani leg.; sweep net; LESC-B-R18/23. • 1 ♂; Rif, Bni Hassan, Triwa; 8 May 2018; K. Kettani leg.; sweep net; LESC-B-R18/25. • 1 ♂; Rif, Tazrout, Riba (PNPB); 29 Apr.–22 May 2019; K. Kettani leg.; Malaise trap; LESC-B-R19/28.

**Distribution.** Great Britain, Greece (Crete), Hungary, Romania, Serbia, Italy (Sicily), Slovakia and Spain. First record from Morocco.

**Habitat.** This species has a large distribution in the Rif Region across a wide altitudinal range (124–1421 m). It was found on the banks of wetlands bordered by *Erica arborea*, *Pistacia lentiscus*, and *Rubus ulmifolius*, in a meadow dotted by Asteraceae and Poaceae and in a mixed oak forest (*Quercus suber*, *Quercus canariensis*).

#### *Agromyza mobilis* Meigen, 1830

**Material examined.** MOROCCO. • 1 ♂; Rif, Akchour, Chellal Sghir (NPTL); 7 June–14 July 2017; K. Kettani leg.; Malaise trap; LESC-B-R19/29.

**Distribution.** Andorra, Belgium, Czech Republic, Denmark, Estonia, Finland, France (including Corsica), Germany, Great Britain, Greece (Dodecanese, Rhodes), Hungary, Italy, Japan, Latvia, Lithuania, Netherlands, Norway, Poland, Portugal, Russia, Slovakia, Spain, Sweden, Switzerland, Turkey, Ukraine, and Yugoslavia. First record from Morocco.

**Habitat.** Banks of waterfall lined by *Pistacia lentiscus*, *Nerium oleander*, *Ficus carica*, *Mentha* sp., and *Thuya* sp.

***Agromyza myosotidis* Kaltenbach, 1864**

**Material examined.** MOROCCO. • 1 ♂; Rif, El Hamma, Masjid El Hamma; 6–21 June 2016; K. Kettani leg.; Malaise trap; LESC-B-R16/10.

**Distribution.** Belgium, Bulgaria, Czech Republic, Denmark, Finland, Germany, Great Britain, Hungary, Kashmir, Malta, Poland, Spain incl. Canary Islands, Sweden, Switzerland, Turkey and Ukraine. First record from Morocco.

**Habitat.** Collected in maquis composed of *Quercus coccifera*, *Olea europaea*, *Fraxinus angustifolia*, *Phillyrea angustifolia*, *Dittrichia viscosa*, *Pistacia lentiscus*, and *Erica arborea*.

***Agromyza nigrescens* Hendel, 1920**

**Material examined.** MOROCCO. • 1 ♂; Rif, Rmilat, Parc Perdicaris; 16 May 2015; K. Kettani leg.; sweep net; LESC-B-R15/06.

**Distribution.** Austria, Belgium, Croatia, Czech Republic, Denmark, Estonia, Finland, Germany, Great Britain, Hungary, Iraq, Italy, Japan, Kyrgyzstan, Lithuania, Malta, Norway, Poland, Romania, Russia, Serbia, Slovenia, Spain incl. Canary Islands, Sweden, Switzerland, and Turkey. First record from Morocco.

**Habitat.** This species was collected in the forest of Perdicaris Park, at an altitude of 223 m. The habitat is a preserved forest with significant natural and replanted vegetation such as *Olea europaea*, *Cupressus macrocarpa*, *Quercus canariensis*, *Pittosporum undulatum*, *Quercus suber*, *Quercus lusitanica*, and *Acacia longifolia*.

***Hexomyza simplex* (Loew, 1869)**

**Material examined.** MOROCCO. • 1 ♂; Anti-Atlas, Errachidia, Aoufous; 26 Sept. 2022; S. Fekrani leg.; sweep net; LESC-B-AA22/42.

**Distribution.** This Holarctic species is known from Albania, Austria, Great Britain, Denmark, Finland, France, Germany, Greece, Hungary, Israel, Italy, Netherlands, Poland, Turkey, Ukraine and in the USA from California, Connecticut, Hawaii, Iowa, Massachusetts, Michigan, and New Jersey. First record from Morocco.

**Habitat.** This species was found at medium altitude in agricultural fields in arid areas in the south of Morocco.

***Ophiomyia pinguis* (Fallén, 1820)**

**Material examined.** MOROCCO. • 8 ♂♂, 7 ♀♀; Rif, Tazrout, Adrou (PNPB); 14 July–15 Aug. 2013; K. Kettani leg.; Malaise trap; LESC-B-R13/02. • 1 ♂; Rif, Kitane, Oued Zarka; 2 Dec. 2017; K. Kettani leg.; sweep net; LESC-B-R17/19.

**Distribution.** Recorded from Austria, Belgium, Great Britain, Bulgaria, China, Croatia, Czech Republic, Denmark, Egypt, Finland, France, Germany, Greece, Hungary, Israel, Italy, Kashmir, Kyrgyzstan, Lithuania, Netherlands, Norway, Poland, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Tajikistan, Turkey, Ukraine, Uzbekistan, and Yugoslavia. First record from Morocco.

**Habitat.** This species was collected in a mixed forest (*Quercus suber*, *Pinus pinaster*) and on the banks of a lowland stream (Oued Zarka) bordered by *Nerium oleander*.

### Subfamily Phytomyzinae

#### *Amauromyza (Cephalomyza) karli* (Hendel, 1927)

**Material examined.** MOROCCO. • 1 ♂, Rif, Kitane Oued Zarka; 2 Dec. 2017; K. Kettani leg.; sweep net; LESCB-R17/20.

**Distribution.** Recorded from **Canada:** AB, BC\*, MB, NS\*, ON, QC, SK\*. **USA:** MD\*. Europe: Austria, Croatia, Czech Republic, Great Britain, Finland, France, Germany, Greece, Hungary, Korea, Netherlands, Poland, Romania, Russia, Serbia, Slovakia, Spain, Sweden, Switzerland, and Ukraine. China. Mongolia, South Korea. First record from Morocco.

**Habitat.** This species was collected in one site at low altitude (49 m) on the banks of a lowland stream (Oued Zarka) bordered by *Nerium oleander*.

#### *Calycomyza flavomaculata* (Spencer, 1960)

**Material examined.** MOROCCO. • 1 ♂; Rif, Akchour, Chellal Kbir (NPTL); 25 Apr.–6 June 2016; K. Kettani leg.; Malaise trap; LESCB-R16/12.

**Distribution.** Croatia, Greece, and Spain. First record from Morocco.

**Habitat.** Collected on the banks of the Akchour waterfall, lined by *Pistacia lentiscus*, *Nerium oleander*, *Ficus carica*, *Mentha* sp., and *Thuya* sp.

#### *Calycomyza solidaginis* (Kaltenbach, 1869)

**Material examined.** MOROCCO. • 1 ♂; Rif, Jbel Kelti, Oued Tanina; 6 June 2021; L. Zouhair leg.; sweep net; LESCB-R21/38.

**Distribution.** China, Germany, Hungary, Lithuania, Poland, Sweden, Switzerland, Yemen, and **Canada:** AB, NB, NS, ON, QC, YT. **USA:** widespread outside of AK and HI. First record from Morocco.

**Habitat.** Collected on the banks of a mountain river bordered by *Pistacia lentiscus*, *Juniperus oxycedrus*, *Tetraclinis articulata* and crossing a mixed forest dominated by *Quercus suber*, *Pinus pinaster*, and *Pinus halepensis*. A wide variety of shrubs and herbaceous plant species (*Prunus lusitanica*, *Stachys fontqueri*, *Paeonia coriacea*, *Eryngium caespitiferum*, *Eryngium triquetrum*, *Digitalis laciniata*, *Ptilostemon rhiphaeus*, *Ruscus hypophyllum*, *Astragalus armatus*, *Merendera filifolia*, *Viola* sp.) occupy the undergrowth.

#### *Cerodontha (Butomomyza) eucaricis* Nowakowski, 1967

**Material examined.** MOROCCO. • 1 ♂; Anti-Atlas, Taroudant, Taliouine; 2 Dec. 2017; Y. Fekrani leg.; sweep net; LESCB-AA17/14.

**Distribution.** It is known from Belgium, Great Britain, Canada, Czech Republic, Denmark, Finland, France, Germany, Greece, Hungary, Japan, Lithuania, Poland, Slovakia, Sweden, and Switzerland. First record from Morocco.

**Habitat.** It was collected in an agricultural field bordered by *Carex acuta* and *Carex nigra* in an arid area.

#### *Cerodontha (Dizygomyza) luctuosa* (Meigen, 1830)

**Material examined.** MOROCCO. • 1 ♂, 1 ♀; Rif, El Anasser, Dayat El Anasser; 22 Mar. 2019; K. Kettani leg.; sweep net; LESCB-R19/30.

**Distribution.** Species known from Alaska, Albania, Austria, Belarus, Belgium, Canada, Great Britain, Bulgaria, Byelorussia, Czech Republic, Denmark, Finland, France, Germany, Greece, Hungary, Iraq, Ireland, Israel, Italy including Sicily, Latvia, Lithuania, Netherlands, Norway, Poland, Portugal, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Tunisia, Uzbekistan, Yugoslavia, and USA (Oklahoma). First record from Morocco.

**Habitat.** This species was collected in high altitude on the edge of a pond bordered by *Quercus pyrenaica*.

#### *Cerodontha (Icteromyza) geniculata* (Fallén, 1823)

**Material examined.** MOROCCO. • 1 ♂; Middle Atlas, Ifrane, Lac Zarouka (NPIF); 13 Oct. 2023; M. Doukale leg.; sweep net; LESCB-MA23/46.

**Distribution.** A widespread species. It has been recorded from Albania, Austria, Belarus, Great Britain, Bulgaria, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Italy, Latvia, Lithuania, and Macedonia. First record from Morocco.

**Habitat.** This species was collected on the edge of a mountain lake lined by *Eriophorum latifolium*, *Populus alba*, *Carex* sp., *Juncus bufonius*, *Schoenoplectus lacustris*, *Typha latifolia*, and *Salix* sp.

#### *Chromatomyia centaurii* Spencer, 1990

**Material examined.** MOROCCO. • 1 ♂; Rif, Akchour, Pont de Dieu (NPTL); 1 July 2019; M. Nourtli leg.; sweep net; LESCB-R19/32.

**Distribution.** Known from Great Britain, Germany, Greece, Hungary, Ireland, Lithuania, and Poland. First record from Morocco.

**Habitat.** This species was found near running water (Akchour) lined by *Nerium oleander*, *Erica* sp., and *Cistus* sp.

#### *Liriomyza amoena* (Meigen, 1830)

**Material examined.** MOROCCO. • 3 ♂♂, 2 ♀♀; Rif, Khemis Anjra, Oued Khemis; 16 Sep. 2012; K. Kettani leg.; sweep net; LESCB-R12/01.

**Distribution.** The species is recorded from Austria, Belgium, Great Britain, Bulgaria, Czech Republic, Denmark, Finland, France, Germany, Greece, Ireland, Italy, Japan, Latvia, Lithuania, Moldavia, Netherlands, Poland, Portugal (Madeira Island), Romania, Russia, Serbia, Slovakia, Slovenia, Spain, Sweden, Turkey, and in the Oriental Region (India). First record from Morocco.

**Habitat.** Collected on the bank of a lowland river bordered by *Nerium oleander*, *Phragmites australis*, *Tamarix africana*, *Rubus ulmifolius*, and *Juncus acutus*.

#### *Liriomyza brassicae* (Riley, 1884)

**Material examined.** MOROCCO. • 1 ♂; Rif, Souk El Had, Oued El Koub; 23 Apr. 2018; K. Kettani leg.; sweep net; LESCB-R18/24.

**Distribution.** Wide distribution as an agricultural pest, including in the Afro-tropical Region: Cape Verde Islands, Ethiopia, Kenya, Mozambique, Namibia, Oman, Senegal, Spain (Canary Islands), South Africa, Yemen, and Zimbabwe. In the Palaearctic Region including Egypt, France (Corsica), Germany, Iraq, Japan, Malta, Poland, Portugal, Romania, Saudi Arabia, Spain, and Turkey. First record from Morocco.

**Habitat.** The male was captured on the banks of a river bordered by *Erica arborea*, *Pistacia lentiscus* and *Rubus ulmifolius*.

#### *Liriomyza strigata* (Meigen, 1830)

**Material examined.** MOROCCO. • 2 ♂♂; Anti-Atlas, Agadir Idaoutanane, Vallée du Paradis; 4 Dec. 2017; Y. Fekrani leg.; sweep net; LESCB-AA17/15.

**Distribution.** Ubiquitous polyphagous species. It is recorded from Albania, Belarus, Belgium, Bosnia-Herzegovina, Great Britain, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iraq, Ireland, Italy, Kazakhstan, Kyrgyz Republic, Lithuania, Netherlands, Portugal (Madeira Isl.), Norway, Poland, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Turkey, Ukraine, and Uzbekistan. First record from Morocco.

**Habitat.** This species was collected on the sandy riverbank in an arid region.

#### *Phytoliriomyza arctica* (Lundbeck, 1901)

**Material examined.** MOROCCO. • 1 ♂; Rif, Chefchaouen, Bni Ahmed; 10 May 2021; K. Kettani leg.; sweep net; LESCB-R21/39.

**Distribution.** Known from Brazil, Chile, Greenland, North America. Europe. Iran, South Korea, Taiwan, and Sri Lanka. First record from Morocco.

**Habitat.** It was collected in a *Cannabis* cultivation field with *Cannabis sativa* and *Cannabis indica*.

#### *Phytoliriomyza perpusilla* (Meigen, 1830)

**Material examined.** MOROCCO. • 1 ♂; High Atlas, Al Haouz, Amizmiz (NPTB); 26 Mar. 2017; Y. Fekrani leg.; sweep net; LESCB-HA17/16.

**Distribution.** Known from Albania, Austria, Great Britain, Bulgaria, Cape Verde Islands, Czech Republic, Estonia, Finland, France, Germany, Greece, Hungary, Italy including Sardinia, Japan, Lesotho, Lithuania, Malta, Nepal, Netherlands, Norway, Oman, Poland, Portugal incl. Azores Islands, Romania, Russia (Yakutia), Serbia, Spain (Canary Islands), South Africa, Sweden, Switzerland, Taiwan, Tunisia, Turkey, and Yemen. First record from Morocco.

**Habitat.** Collected in a pine forest (*Pinus pinaster*) with herbaceous plants composed of *Trifolium* spp.

### ***Phytoliriomyza pteridii* Spencer, 1973**

**Material examined.** MOROCCO. • 1 ♂; Rif, Fahs-Anjra, Rmel; 27 Feb. 2020; M. Nouri leg.; sweep net; LESC-B-R20/36.

**Distribution.** The species is known from Andorra, Great Britain, Bulgaria, Croatia, France, Germany, Hungary, Ireland, Italy, Montenegro, Poland, Portugal, Scotland, Slovakia, Slovenia, Switzerland, and Yugoslavia. First record from Morocco.

**Habitat.** It was collected at 372 m in a scrubland with rocky soil of a clay-calcareous habitat crossed by a small stream and bordered by vegetation mainly composed of *Pistacia lentiscus*, *Carlina racemosa*, *Conyza canadensis*, *Erica arborea*, and *Cistus monspeliensis*. On a nearby field, fruit trees such as *Olea europaea* and *Ficus carica* were growing.

### ***Phytomyza anemones* Hering, 1925**

**Material examined.** MOROCCO. • 2 ♂♂, 1 ♀; Rif, Chefchaouen, Bab Tariouant; 31 May 2015; K. Kettani leg.; sweep net; LESC-B-R15/07.

**Distribution.** Denmark, Finland, France, Germany, Great Britain, Greece, Hungary, Ireland, Italy, Lithuania, Malta, Netherlands, Poland, Spain (Mallorca), Sweden, and Yugoslavia. First record from Morocco.

**Habitat.** Collected in an oak forest at high altitude (*Quercus pyrenaica*).

### ***Phytomyza clematidis* Kaltenbach, 1859**

**Material examined.** MOROCCO. • 1 ♂; Rif, Tazia, Dayat Tazia; 12 May 2015; K. Kettani leg.; sweep net; LESC-B-R15/05.

**Distribution.** Recorded from Andorra, Austria, Cyprus, Czech Republic, France incl. Corsica, Germany, Great Britain, Greece, Hungary, Israel, Italy, Lithuania, Maltese Islands, Netherlands, Norway, Portugal, Spain incl. Balearic and Canary Islands, Slovakia, Switzerland, and Turkey. First record from Morocco.

**Habitat.** Collected on the edge of a pond surrounded by cork oak forest.

### ***Phytomyza crassiseta* Zetterstedt, 1860**

**Material examined.** MOROCCO. • 2 ♂♂; Rif, Bni Darkoul, Oued Souk Lhad; 30 Apr. 2016; F.Z. Bahid leg.; sweep net; LESC-B-R16/08. • 6 ♂♂, 4 ♀♀;

Rif, Ametrasse, Oued Ametrasse; 30 Apr. 2016; F.Z. Bahid leg.; sweep net; LESC-R16/09. • 1 ♂, 1 ♀; Rif, Tétouan, Oued Raouz; 18 May 2018; A. Adghir leg.; sweep net; LESC-R18/26. • 5 ♂♂, 1 ♀♀; Rif, Tazrout, Riba (PNPB); 29 Apr.–22 May 2019; K. Kettani leg.; Malaise trap; LESC-R19/31. • 4 ♂♂; Rif, Akchour, Pont de Dieu (NPTL); 1 July 2019; M. Nourtí leg.; sweep net; LESC-R19/33. • 1 ♂, 1 ♀; Rif, Oulad Sidi Mansour, Oued Tassikeste; 16 May 2023; K. Kettani leg.; sweep net; LESC-R23/47.

**Distribution.** Known from Albania, Andorra, Austria, Belarus, Bulgaria, Croatia, Czech Republic, Denmark, Estonia, Finland, France, Germany, Great Britain, Greece, Hungary, Ireland, Italy, Latvia, Liechtenstein, Lithuania, Montenegro, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Spain incl. Canary Islands, Sweden, Switzerland. Canada. ON\*, QC. USA: CA, ID, IN\*, MA, MD, ME, NC, NJ\*, NY, PA, VA, WA, WV. Argentina, Chile. Japan, Turkey, and Russia. First record from Morocco.

**Habitat.** This species has a wide distribution in the Rif across a wide altitudinal range (144–1421 m). It was mainly caught at the banks of rivers bordered by *Nerium oleander*, *Rubus ulmifolius*, *Erica* sp., and *Cistus* sp. It was also found in a mixed forest of *Quercus faginea*, *Quercus suber*, and *Pinus pinaster* at 1421 m altitude.

#### ***Phytomyza notata* Meigen, 1830**

**Material examined.** MOROCCO. • 6 ♂♂, 4 ♀♀; Rif, Talassemteane, Maison forestière (NPTL); 7 June–17 July 2014; K. Kettani leg.; Malaise trap; LESC-R14/04.

**Distribution.** This species was reported from Belarus, Bulgaria, Czech Republic, Denmark, Finland, France, Germany, Great Britain, Hungary, Italy, Latvia, Lithuania, Poland, Romania, Russia, Serbia, Slovakia, Spain, Sweden, and Switzerland. First record from Morocco.

**Habitat.** This species was captured in a fir forest (*Abies marocana*) at an altitude of 1696 m in Jbel Talassemteane.

#### ***Phytomyza plantaginis* Robineau-Desvoidy, 1851**

**Material examined.** MOROCCO. • 1 ♂, 2 ♀♀; Rif, Ametrasse, Oued Ametrasse; 30 Apr. 2016; F.Z. Bahid leg.; sweep net; LESC-R16/11. • 6 ♂♂, 3 ♀♀; Rif, Talassemteane, Place Espagna (NPTL); 16 Apr. 2018; K. Kettani leg.; sweep net; LESC-R18/22. • 1 ♂; Rif, Bouhachem, Amsemil (PNPB); 1 July 2019; K. Kettani leg.; sweep net; LESC-R19/34. • 2 ♂♂; Atlantic Plain, Sidi Yahya Gharb, forest of Sidi Yahya Gharb; 4 May 2020; M. Doukale leg.; sweep net; LESC-AP20/35. • 2 ♂♂; Rif, Talembote, Oued Talembote (NPTL); 1 July 2018; K. Kettani leg.; sweep net; LESC-R18/27. • 1 ♂, 1 ♀; Rif, Tétouan, Khemis Anjra; 27 Mar. 2021; L. Zouhair leg.; sweep net; LESC-R21/37. • 1 ♂; Rif, Talassemteane, Maison forestière (NPTL); 10 June 2021; K. Menouar & L. Zouhair leg.; sweep net; LESC-R21/41. • 3 ♂♂; Rif, Afeska, Oued Afeska (NPTL); 10 June 2021; K. Menouar & L. Zouhair leg.; sweep net; LESC-R21/40.

**Distribution.** It is recorded from Canada: BC\*, ON, QC. USA: Widespread, including HI. Bermuda\*. Widespread in Europe, recorded from almost every coun-

try including Azores, Canary Islands, Cyprus, Italy, Portugal including Madeira. In the Palaearctic, recorded also from Algeria, Egypt, Iran, Israel, Japan, Kyrgyzstan, Russia, Tunisia, Turkey, and Uzbekistan. It is recorded also from South Africa, Australia, New Zealand, Taiwan, and Thailand. First record from Morocco.

**Habitat.** This species exhibits a large distribution in the Rif Region of northern Morocco across a wide altitudinal range (11–1683 m) and inhabits a large variety of habitats such as the edge of wetlands (river, peat bog), mixed forest dominated by oak (*Quercus suber*, *Quercus canariensis*), fir (*Abies marocana*), and cedar (*Cedrus atlantica*) at high altitudes and some other vegetation such *Tetraclinis articulata*, *Eucalyptus globulus*, *Scolymus maculatus*, and *Carlina corymbosa*.

### ***Phytomyza rufipes* Meigen, 1830**

**Material examined.** MOROCCO. • 2 ♂♂, 1 ♀; Rif, Ben Karrich, Oued Mhajrate; 20 July–13 Sep. 2016; K. Kettani leg.; Malaise trap; LESC-B-R16/13.

**Distribution.** It is recorded from Albania, Austria, Belgium, Canada, Croatia (Dalmatia), Czech Republic, Denmark, Egypt, Estonia, Finland, France, Germany, Great Britain, Greece, Iceland, Iraq, Ireland, Italy, Latvia, Lithuania, Netherlands, Norway, Poland, Portugal (Madeira), Russia, Slovakia, Spain (mainland, Canary Islands), Sweden, Switzerland, Turkey, and United Kingdom (Yugoslavia). In the Nearctic Region: Canada (New Brunswick, Newfoundland), USA (California). In the Neotropical Region: Argentina, Colombia. First record from Morocco.

**Habitat.** Collected on a riverbank bordered by *Armoracia rusticana*, *Alliaria petiolata*, *Brassica oleracea*, *Diplotaxis* sp., and *Myagrum perfoliatum* and ground vegetation in a pine forest.

### ***Phytomyza vitalbae* Kaltenbach, 1872**

**Material examined.** MOROCCO. • 1 ♂; Rif, Targuist, Bni Bounzar; 1–30 Apr. 2022; K. Kettani leg.; Malaise trap; LESC-B-R22/44.

**Distribution.** The species is very widely distributed, being known from Andorra, Austria, Belgium, Bulgaria, Canada, China, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France incl. Corsica, Germany, Great Britain, Greece, Hungary, Ireland, Italy, Montenegro, Nepal, Netherlands, New Zealand, Poland, Romania, Russia (Yakutia), Serbia, South Africa, Spain incl. Canary Islands, Sweden, Switzerland. First record from Morocco.

**Habitat.** Collected in cedar forest (*Cedrus atlantica*).

### ***Pseudonapomyza confusa* Zlobin, 1993**

**Material examined.** MOROCCO. • 2 ♂♂; Middle Atlas, Ifrane, Zaouiet Ifrane (PNIF); 2 Apr. 2017; Y. Fekrani leg.; sweep net; LESC-B-MA17/17. • 1 ♂; Anti-Atlas, Errachidia, Aoufous; 26 Sep. 2022; S. Fekrani leg.; sweep net; LESC-B-AA22/43. • 1 ♂; Eastern Morocco, Taourirt, El Aioun Sidi Mellouk; 9 Jul. 2023; M. Doukale leg.; sweep net; LESC-B-EM23/45.

**Distribution.** It is recorded from Cape Verde Islands, Greece (Crete), Madagascar, and Namibia. First record from Morocco.

**Habitat.** This species was found at medium altitude in agricultural fields in arid areas in Eastern Morocco as well as in the Anti-Atlas. It was also captured at high altitude in cedar forest (*Cedrus atlantica*) in the Middle Atlas.

### ***Pseudonapomyza palavae* Černý, 1998**

**Material examined.** MOROCCO. • 1 ♂; Eastern Morocco, Taourirt, El Aïoun Sidi Mellouk; 9 July 2023; M. Doukale leg.; sweep net; LESCB-EM24/48.

**Distribution.** It is recorded from Czech Republic and Hungary. First record from Morocco.

**Habitat.** This species was found at medium altitude in agricultural fields in arid areas in the Eastern Morocco.

### ***Pseudonapomyza vota* Spencer, 1973**

**Material examined.** MOROCCO. • 1 ♂; Rif, Tazrout, Adrou (PNPB); 14 Jul.–15 Aug. 2013; K. Kettani leg.; Malaise trap; LESCB-R13/03. • 2 ♂♂; Rif, Akhour, Chellal Sghir (NPTL); 7 June–14 July 2017; K. Kettani leg.; Malaise trap; LESCB-R17/18.

**Distribution.** This species is known from Croatia, Cyprus, France, Greece, Hungary, Italy, Malta, Portugal, Serbia, Spain, Switzerland, and Turkey. First record from Morocco.

**Habitat.** Collected in an oak-dominated forest and on the banks of a waterfall lined by *Pistacia lentiscus*, *Nerium oleander*, *Ficus carica*, *Mentha* sp., and *Thuya* sp.

## **Discussion**

This article presents additions to the Moroccan fauna represented by 30 new records of leaf-miner flies, bringing the total number of Agromyzidae currently known for Morocco to 92 species. This faunal assemblage is subdivided into 11 genera, three genera in Agromyzinae, in which *Agromyza* (*A. anthracina*, *A. conjuncta*, *A. mobilis*, *A. myosotidis*, *A. nigrescens*) dominates in the Rif region and was found in various habitats. The second genus, *Hexomyza* (*H. simplex*) is quite rare, with only one species found in the Atlas region in an agricultural field. The third genus is *Ophiomyia* with the single species *O. pinguis* and 16 individuals present only in the Rif region in various habitats (forest and riverbank). The eight genera in Phytomyzinae are widely distributed across Morocco. *Calycomyza* is newly reported from the country, represented by two species (*C. flavomaculata* and *C. solidaginis*) found in the Rif region on riverbanks at a significant altitude of 1463 m. *Phytomyza* was the prevalent genus in the Rif region and the most diversified with seven species (*Ph. anemones*, *Ph. clematidis*, *Ph. crassisetosa*, *Ph. notata*, *Ph. plantaginis*, *Ph. rufipes*, *Ph. vitalbae*). Some other genera were less diversified such as *Cerodontha* (*C. (Butomomyza) eucaricis*, *C. (Dizygomyza) luctuosa*, *C. (Icteromyza) geniculata*), *Liriomyza* (*L. amoena*, *L. brassicae*, *L. strigata*), *Phytoliriomyza* (*P. arctica*,

*P. perpusilla*, *P. pteridii*), and *Pseudonapomyza* (*Ps. confusa*, *Ps. palavae*, and *Ps. vota*). The other two genera were less diversified and were found in forest habitats such as *Amauromyza* (*A. (Cephalomyza) karli*) and *Chromatomyia* (*C. centaurii*). Most obtained species belonged into *Phytomyza* (23,4%) with *Phytomyza crassiseta* being the most frequent species (38.8%). *Agromyza nigrescens* has a wide distribution in Europe.

These numbers, however, cannot be considered final because further collecting in unexplored areas will undoubtedly bring new additions to the Moroccan fauna

## Acknowledgements

We are grateful to the Water and Forests National Agency of Morocco for providing permits for the entomological surveys.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Funding

No funding was reported.

### Author contributions

Writing - original draft: MD. Writing - review and editing: MT, KK.

### Author ORCIDs

Mourad Doukale Daief  <https://orcid.org/0009-0000-6946-2244>

Michael von Tscharnhaus  <https://orcid.org/0000-0002-1903-4767>

Kawtar Kettani  <https://orcid.org/0000-0003-2361-3996>

### Data availability

All of the data that support the findings of this study are available in the main text.

## References

- Becker T, Stein P (1913) Dipteren aus Marokko. Annuaire du Musée zoologique de l'Académie impériale de Sciences de St.-Petersbourg 18: 62–95.
- Benabid A (1982) Étude phytoécologique, biogéographique et dynamique des associations et séries sylvatiques du Rif occidental (Maroc). Thèse de Doctorat Es. Sc. Nat. Faculté de St. Jérôme Marseille III, France, 165 pp.
- Černý M (2007) New faunistic records of Agromyzidae (Diptera) from Andorra including descriptions of three new species. Boletín de la Sociedad entomológica aragonesa, Zaragoza 41: 43–51.
- Černý M (2009) New faunistic data on the Agromyzidae (Diptera) from the West Palaearctic Region. Klapalekiana, Praha 45(1–2): 9–21.

- Černý M (2010) Agromyzidae: 185 + 233 - In: Borges PAV, Costa A, Cunha R, Gabriel R, Gonçalves V, Martins AF, Melo I, Parente M, Raposeiro P, Rodrigues P, Santos RS, Silva L, Vieira P, Vieira V (Eds): Listagem dos organismos terrestres e marinhos dos Açores. / A list of the terrestrial and marine biota from the Azores. Princípia, Cascais [ISBN 978-989-8131-75-1], Parede, Portugal, 429 pp.
- Černý M (2012) The fauna of Agromyzidae (Diptera) in the Gemer region (Central Slovakia), with descriptions of three new species from Slovakia. Časopis slezského zemského Muzea Opava (A), Opava 61: 49–76.
- Černý M (2013) Additional records of Agromyzidae (Diptera) from the West Palaearctic Region. Časopis slezského zemského Muzea Opava (A), Opava 62: 281–288.
- Černý M (2019) Additional new records of Agromyzidae (Diptera) from the Palaearctic Region. Acta Musei Silesiae, Scientiae naturales 67(2018): 117–137. <https://doi.org/10.2478/cszma-2018-0010>
- Černý M, Merz B (2006) New records of Agromyzidae (Diptera) from Palaearctic Region. Mitteilungen der schweizerischen entomologischen Gesellschaft, Zürich 79(1): 77–106. <http://doi.org/10.2478/cszma-2018-0010>
- Černý M, Merz B (2007) New records of Agromyzidae (Diptera) from the West Palaearctic Region, with an updated checklist for Switzerland. Mitteilungen der schweizerischen entomologischen Gesellschaft, Zürich 80(1/2): 107–121.
- Černý M, von Tschirnhaus M (2014) New records of Agromyzidae (Diptera) from the Afrotropical Region, with a checklist. Acta Musei Silesiae, Scientiae naturales, Opava 63(2): 159–176. <https://doi.org/10.2478/cszma-2014-0017>
- Černý M, Andrade R, Gonçalves AR, von Tschirnhaus M (2018) New records of Agromyzidae (Diptera) from Portugal, with an updated checklist. Acta Musei Silesiae, Scientiae naturales 67: 7–57. <https://doi.org/10.2478/cszma-2018-0002>
- Černý M, von Tschirnhaus M, Winqvist K (2020) First records of Palaearctic Agromyzidae (Diptera) from 40 countries and major islands. Acta Musei Silesiae, Scientiae naturales, Opava 69(3): 193–229. <https://doi.org/10.2478/cszma-2020-0017>
- Dempewolf M (2004) Arthropods of Economic Importance – Agromyzidae of the World (CD-ROM). ETI. University of Amsterdam, Amsterdam. <https://nlbif.eti.uva.nl/bis/agromyzidae.p>
- Geipert S, Kroschel J, Sauerborn J (1994) Incidence of *Phytomyza orobanchia* Kalt. (Diptera: Agromyzidae) in northern Morocco and perspectives for a biological control of Orobanche spp. Results of a field survey. In: Fifth Arab Congress of Plant Protection, 27<sup>th</sup> November – 2<sup>nd</sup> December 1994, Fès, Morocco, 235 pp
- Griffiths GCD (1967) Revision of the *Phytomyza syngenesiae* group (Diptera, Agromyzidae), including species hitherto known as "*Phytomyza atricornis* Meigen". Stuttgarter Beiträge zur Naturkunde 177: 1–28.
- Griffiths GCD (1974) Studies on boreal Agromyzidae (Diptera). V. On the genus *Chromatomyia* Hardy, with revision of Caprifoliaceae-mining species. Quaestiones entomologicae, Edmonton 10(1): 35–69. <https://biostor.org/reference/194955>
- Hanafi A, Schnitzler WH (2004) Integrated production and protection in greenhouse tomato in Morocco. Acta Horticulturae, ISHS (International Society for Horticultural Science), Leuven, Belgium 659: 323–330. <https://doi.org/10.17660/ActaHortic.2004.659.38>
- HCEFLCD [Haut-Commissariat aux Eaux et Forêts et à la Lutte Contre la Desertification] (2017) Plan d'Aménagement et de Gestion du Parc National de Toubkal. Mission I – Diagnostic: analyse du périmètre d'intervention et du territoire environnant. Mai 2017, 305 pp.

- Hendel F (1936) Fam. Agromyzidae. In: Frey R, von O. Duda B, Hendel Fr, Lackschewitz P, Schmitz H, Storå R, Villeneuve J (Eds) Die Dipterenfauna der Kanarischen Inseln und ihre Probleme. Mit - Commentationes biologicae, Helsingfors und Berlin 6(1): 120–124.
- Hendel FG (1931–36) 59. Agromyzidae. In: Lindner E (Ed.) Die Fliegen der palaearktischen Region 6(2): xi + 570 pp. [pls i–xvi + 16 pp]
- Hering EM (1957) Minierfliegen in Lebermoosen (Dipt. Agromyzidae) = (Leaf miner flies in liverworts (Dipt. Agromyzidae)). Mitteilungen der deutschen entomologischen Gesellschaft, Berlin 16: 48–51. <https://doi.org/10.1002/mmnd.4820160311>
- Hering EM (1966) Minierfliegen in Lebermoosen II (Dipt. Agromyzidae) = (Leaf miner flies in liverworts II (Dipt. Agromyzidae)). Deutsche entomologische Zeitschrift (neue Folge), Berlin 13(I/III): 231–235.
- Kettani K, Ebejer MJ, Ackland DM, Bächli G, Barraclough D, Barták M, Carles-Tolrá M, Černý M, Cerretti P, Chandler P, Dakki M, Daugeron C, De Jong H, Dils J, Disney H, Droz B, Evenhuis N, Gatt P, Graciolli G, Grichanov IY, Haenni JP, Hauser M, Himmi O, Macgowan I, Mathieu B, Mouna M, Munari L, Nartshuk EP, Negrobov OP, Oosterbroek P, Pape T, Pont AC, Popov GV, Rognes K, Skuhravá M, Skuhravý V, Speight M, Tomasovic G, Trari B, Tschorasnig HP, Vala JC, von Tscharnhaus M, Wagner R, Whitmore D, Woźnicka AJ, Zatwarnicki T, Zwick P: Catalogue of the Diptera (Insecta) of Morocco - an annotated checklist, with distributions and a bibliography. ZooKeys 1094: 279–283. [466 pp] <https://doi.org/10.3897/zookeys.1094.62644>
- Kozlowsky S, Rungs C (1932) Sur quelques insectes ennemis des plantes maraîchères au Maroc. Bulletin de la Société des Sciences naturelles du Maroc, Rabat 12(1–3): 66–68.
- Lonsdale O (2021) Manual of North American Agromyzidae (Diptera, Schizophora), with revision of the fauna of the “Delmarva” states. ZooKeys 1051: 1–481. <https://doi.org/10.3897/zookeys.1051.64603>
- Lonsdale O, Murphy ST, Scheffer SJ (2023) Agromyzidae (Diptera) Plant Pests. Pensoft Publishers, Sofia, i + 110 pp. <https://doi.org/10.3897/ab.e108410>
- Maarouf A (2003) Diptères Cyclorrhaphes et Brachycères nouveaux pour le Maroc. L'entomologiste 59(1–2): 41–47.
- Mokhtari N, Mrabet R, Lebailly P, Bock L (2014) Spatialisation des bioclimats, de l'aridité et des étages de végétation du Maroc. Revue marocaine des Sciences agronomiques et vétérinaires 2(1): 50–66.
- Papp L, Černý M (2015) Agromyzidae (Diptera) of Hungary. Volume 1. Agromyzinae - Pars Ltd, Nagykovácsi, Hungary, 416 pp. <https://doi.org/10.18655/Agromyzidae.Vol.1>
- Papp L, Černý M (2016) Agromyzidae (Diptera) of Hungary. Volume 2. Phytomyzinae I - Pars Ltd, Nagykovácsi, Hungary, 385 pp. <https://doi.org/10.18655/Agromyzidae.Vol.2>
- Papp L, Černý M (2017) Agromyzidae (Diptera) of Hungary. Volume 3. Phytomyzinae II - Pars Ltd, Nagykovácsi, Hungary, 427 pp. <https://doi.org/10.18655/Agromyzidae.Vol.3>
- Papp L, Černý M (2020) [as 2019]: Agromyzidae (Diptera) of Hungary. Volume 4. Phytomyzinae III - Pars Ltd, Nagykováksi, Hungary, 708 pp.
- Séguay E (1936) Un nouveau Muscide nuisible à l'orge au Maroc. Liste des Diptères vivant sur les Hordeum. Bulletin de la Société des sciences naturelles du Maroc 16: 3–5.
- Séguay E (1953) Diptères du Maroc. Encyclopédie entomologique. Série B II. Mémoires et notes. (1947–1953) 11: 77–92.
- Spencer KA (1967) Some Agromyzidae (Diptera) from Morocco. Entomologist's Monthly Magazine 103: 126–130.

- Spencer KA (1972) Diptera, Family Agromyzidae - Handbooks for the Identification of British Insects, Royal Entomological Society of London, London 10(5g): 1–136.
- Spencer KA (1973) Agromyzidae (Diptera) of economic importance. Series entomologica, Volume 9. W. Junk B.V., The Hague, xi + 418 pp. <https://doi.org/10.1007/978-94-017-0683-4>
- Spencer KA (1990) Host Specialization in the World Agromyzidae (Diptera). In: Spencer KA (Ed.): Series entomologica 45: - Kluwer Academic Publishers, continued as Springer Science + Business Media Netherlands, Dordrecht, XII + 444. <https://doi.org/10.1007/978-94-009-1874-0>
- Valdés Castrillón B, Rejdali M, Achhal el Kadmiri A, Jury JL, Montserrat JM (2002) Catalogue des plantes vasculaires du nord du Maroc, incluant des clés d'identification. Vol. 1. CSIC, Centre Mixt del Consejo superior de Investigaciones Científicas, Madrid, Biblioteca de Ciencias, Vol. 39479, X + 491 pp.
- von Tscharnhaus M, Groll E (2024a) World Agromyzidae Online. <https://sdei.senckenberg.de/tscharnhaus-agromyzidae>
- von Tscharnhaus M, Groll E (2024b) World Bibliography of Agromyzidae and Chloropidae Online. <https://sdei.senckenberg.de/tscharnhaus-bibliography>