

Two new species of the genus *Viscosia* de Man, 1890 (Nematoda, Enoplea, Enoplida, Oncholaimidae) from the intertidal zone of the Yellow Sea, China

Lingyun Sun¹, Huimin Gu¹, Yong Huang¹

¹ College of Agriculture and Biology, Liaocheng University, Liaocheng 252059, China
Corresponding author: Yong Huang (huangy@lcu.edu.cn)

Abstract

Two new species of *Viscosia* from the intertidal zone along the Yellow Sea are described and illustrated. *Viscosia media* **sp. nov.** is characterized by a heavily cuticularized and relatively shallow buccal cavity with stubby teeth; cephalic setae 7–8 μm long; amphidial fovea invisible; slender spicules almost straight, cephalated proximally and conical distally; and tail conical, straight in males and slightly bent ventrally in females. *Viscosia sinica* **sp. nov.** is characterized by a relatively large amphidial fovea, conico-cylindrical tail with swollen horseshoe-shaped tip, spicules slightly curved ventrally, and 10–12 setae surrounding the cloaca, each 3–4 μm long. The basic morphological data of males of 26 valid species in *Viscosia* with a body length of 1–2.9 mm are presented.

Key words: Biodiversity, free-living marine nematodes, Huangdao coast, Qingdao, Rizhao coast, Shandong Province, taxonomy



Academic editor: Alexei Tchesunov
Received: 18 November 2024
Accepted: 8 February 2025
Published: 11 March 2025

ZooBank: <https://zoobank.org/A65EAA36-BA3E-4B4B-9EF2-93041E5F621A>

Citation: Sun L, Gu H, Huang Y (2025) Two new species of the genus *Viscosia* de Man, 1890 (Nematoda, Enoplea, Enoplida, Oncholaimidae) from the intertidal zone of the Yellow Sea, China. ZooKeys 1231: 99–117. <https://doi.org/10.3897/zookeys.1231.142078>

Copyright: © Lingyun Sun et al.
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

Free-living marine nematodes are the most widespread and abundant meiofaunal group in marine benthic habitats (Heip et al. 1982). They play a significant role in marine environment and are widely used as pollution indicators in biological monitoring (Lamshead 2004). However, many species of nematodes are still unknown. At present, less than 600 species of free-living marine nematodes have been identified in the Chinese sea area, which is estimated to be less than half of the total species (Sun et al. 2024).

The Yellow Sea is located on the edge of the western Pacific Ocean, between China and the Korean Peninsula. It is a semi-enclosed inland shallow sea basin. Biodiversity surveys and taxonomical studies on nematodes in the Yellow Sea have been carried out in recent years (Huang and Sui 2024; Huang and Zhai 2024; Li et al. 2024). More than 390 species of nematodes have been identified, of which 117 species were new to science (Sun et al. 2024). The new species accounted for 30% of the total known species. However, the total number of nematodes in this sea area is unknown, and new species are routinely found. It is, therefore, important to continue investigating the taxonomy of nematodes in the region. Here, the present paper

describes two new species of the genus *Viscosia* de Man, 1890 that were discovered in the intertidal zone of Qingdao and Rizhao, Yellow Sea.

Viscosia is a large genus with many species. It occurs in marine, brackish, and freshwater sediments (Smol et al. 2014; Leduc and Zhao 2023). The genus was established by de Man in 1890 as a subgenus of *Oncholaimus* Dujardin, 1845. Later, it was raised to genus by Filipjev in 1918. Wieser and Hopper (1967), Gerlach and Riemann (1974), Filipjev (1918), Smol and Sharma (1984), Smol et al. (2014), and Leduc and Zhao (2023) successively reviewed the genus. Wieser and Hopper (1967) stressed the importance of certain characteristics in species identification, such as the size of amphids, the shape of the buccal cavity and teeth, the arrangement of the male circumanal organs like papillae and bursa, and the shape of the tail. Gagarin (2020) provided the basic morphological features of males of 24 valid species with body length of 1–2 mm within the genus *Viscosia*. Currently, the World Database of Nematodes (Nemys eds. 2024) makes a list of 92 valid species including four freshwater species (*Viscosia nicaraguensis* Gerlach, 1957, *V. orientalis* Gagarin, 2020, *V. timmi* Gagarin & Nguyen Thi Thu, 2008 and *V. uipii* Coomans, Vincx & Decraemer, 1985). But in fact, many species were described only based on juveniles or females, and lack the characters of males. In the Chinese sea area, only seven species within *Viscosia* were recorded. They are *Viscosia bandaensis* Kreis, 1932, *V. elegans* (Kreis, 1924) Lorenzen, 1981, *V. franzii* Boucher, 1977, *V. glabra* (Bastian, 1865) de Man, 1890, *V. heterolaima* Smol & Sharma, 1984, *V. longicaudatoides* Nguyen Vu Thanh & Gagarin, 2013 and *V. nuda* Kreis, 1932.

Material and methods

The meiofauna samples were collected from the top sediment layer (0–8 cm deep) using a 2.9 cm diameter sawn-off syringe. The samples were fixed with an equal amount of 10% formalin solution. In the laboratory, samples were stained with 0.1% Rose Bengal for 24 hours. The stained samples were poured through two sieves (500 and 42 µm mesh sizes), and washed with tap water to remove silt and separate macrofauna from meiofauna. The intercepted material in the 42 µm mesh was centrifuged in Ludox-TM (50% colloidal silica, suspension in water; product of Sigma Aldrich Co., USA) with a specific gravity of 1.15 g/ml (de Jonge and Bouwman 1977) to separate meiofauna from the heavier sediment particles. The supernatant obtained after three rounds of centrifugation was poured into the 42 µm mesh to filter Ludox-TM. The sample in the 42 µm mesh was washed into a Petri dish with distilled water and meiofauna was sorted out under a stereoscopic microscope.

Nematodes were transferred into a cavity block containing a solution of 5% glycerol, 5% pure ethanol, and 90% distilled water in volume (McIntyre and Warwick 1984). After ethanol was slowly evaporated, the specimens were mounted in glycerin on permanent slides. The descriptions were made using a differential interference contrast microscope (Leica DM 2500). The photos were taken with a Leica DMC 5400 digital camera. Line drawings were made with the aid of a camera lucida. All measurements were taken using Leica software of LAS X version 3.3.3, and all curved structures were measured along the curved median line. All measurements are in µm. Type specimens were deposited in the Marine Biological Museum of the Chinese Academy of Sciences, Qingdao.

Results and discussion

Taxonomy

Class Enoplea Inglis, 1983

Order Enoplida Filipjev, 1929

Family Oncholaimidae Filipjev, 1916

Genus *Viscosia* de Man, 1890

Diagnosis (according to Smol et al. 2014). Buccal cavity large, barrel-shaped with three unequal immovable teeth, of which the right subventral tooth largest. Females didelphic-amphidelphic with reflexed ovaries. Spicules short, straight or slightly curved. Gubernaculum absent. Demanian system present and simple, consisting of prolongation of the ovary at the reflexed point that connects it with the intestine through the osmosium.

List of valid species (64 species)

- V. abyssorum* (Allgén, 1933) Wieser, 1953.
syn. *Oncholaimus abyssorum* Allgén, 1933.
- V. angustata* (Cobb, 1890) Kreis, 1934.
syn. *Oncholaimus angustatus* Cobb, 1890.
- V. antarctica* Allgén, 1959.
- V. bandaensis* Lorenzen, 1981.
syn. *Mononcholaimus bandaensis* Kreis, 1932.
- V. bandaensis* Kreis, 1932.
syn. *Mononcholaimus bandaensis* Kreis, 1932.
- V. bayensis* Keppner, 1987.
- V. brachylaima* Filipjev, 1927.
- V. brevicaudata* Mawson, 1958.
- V. brevilaima* Allgén, 1959.
- V. cobbi* Filipjev, 1918.
- V. coomansi* Smol & Sharma, 1984.
- V. dossena* Leduc & Zhao, 2023.
- V. elegans* (Kreis, 1924) Lorenzen, 1981.
syn. *Mononcholaimus elegans* Kreis, 1924.
- V. elongata* Filipjev, 1922.
- V. epapillosa* Platonova, 1971.
- V. erasmi* Furstenberg & Vincx, 1989.
- V. falklandiae* Allgén, 1959.
- V. filiformis* (Kreis, 1932) Lorenzen, 1981.
syn. *Mononcholaimus filiformis* Kreis, 1932.
- V. floridana* Keppner, 1987.
- V. franzii* Boucher, 1977.
- V. glabra* (Bastian, 1865) de Man, 1890.
syn. *Oncholaimus glaber* Bastian, 1865.
syn. *Viscosia micoletzkyi* Chitwood, 1951.
- V. grahami* Allgén, 1959.
- V. hanstroemi* Wieser, 1953.

- V. heterolaima* Smol & Sharma, 1984.
V. keiensilis Lorenzen, 1981.
syn. *Mononcholaimus keiensis* Kreis, 1932.
V. keiensis Kreis, 1932.
syn. *Mononcholaimus keiensis* Kreis, 1932.
V. langrunensis (de Man, 1890) Filipjev, 1918.
syn. *Oncholaimus langrunensis* de Man, 1890.
V. longicaudatoides Nguyen Vu Thanh & Gagarin, 2013.
V. macramphida Chitwood, 1951.
V. macrobursata Keppner, 1987.
V. megalaima (Ditlevsen, 1928) Hope & Murphy, 1972.
syn. *Steineria megalaima* Ditlevsen, 1928.
V. meridionalis Kreis, 1932.
V. microseta Wieser, 1953.
V. nicaraguensis Gerlach, 1957.
syn. *Viscosia papillata nicaraguensis* Gerlach, 1957.
V. nona Filipjev, 1946.
V. norvegica (Allgén, 1946) Lorenzen, 1981.
syn. *Mononcholaimus norvegicus* Allgén, 1946.
V. nuda Kreis, 1932.
V. oncholaimelloides Wieser & Hopper, 1967.
V. orientalis Gagarin, 2020.
V. papillata Chitwood, 1951.
V. papillatoides Chitwood, 1960.
V. papillatula Lorenzen, 1981.
syn. *Mononcholaimus papillatus* Kreis, 1932.
V. parafalklandiae Allgén, 1959.
V. parapellucida (Allgén, 1959) Gerlach & Riemann, 1974.
syn. *Viscosia pellucida* Allgén, 1959.
V. parasetosa (Kreis, 1932) Lorenzen, 1981.
syn. *Mononcholaimus parasetosus* Kreis, 1932.
V. pedroensis Allgén, 1947.
V. profunda (Vitiello, 1970) Lorenzen, 1981.
syn. *Mononcholaimus profundus* Vitiello, 1970.
V. propinqua Allgén, 1959.
V. pseudoglabra Kreis, 1932.
V. pygmaea Nguyen Vu Thanh & Gagarin, 2013.
V. rectangulata Wieser, 1953.
V. rustica (Kreis, 1929) Lorenzen, 1974.
syn. *Mononcholaimus rusticus* Kreis, 1929.
V. sedata Gagarin & Nguyen Vu Thanh, 2007.
V. separabilis (Wieser, 1953) Lorenzen, 1981.
syn. *Mononcholaimus separabilis* Wieser, 1953.
V. setosa (Kreis, 1932) Lorenzen, 1981.
syn. *Mononcholaimus setosus* Kreis, 1932.
V. similis Allgén, 1959.
V. stenolaima Filipjev, 1927.
V. tenuissima Allgén, 1959.
V. timmi Gagarin & Nguyen Thi Thu, 2008.

- V. tumidula* Wieser, 1959.
- V. uipii* Coomans, Vincx & Decraemer, 1985.
- V. viscosa* (Bastian, 1865) de Man, 1890.
 - syn. *Mononcholaimus viscosus* Allgén, 1930.
 - syn. *Oncholaimus viscosus* Bastian, 1865.
- V. viscosula* Lorenzen, 1981.
 - syn. *Mononcholaimus viscosus* Allgén, 1930.
- V. wieseri* Mawson, 1958.

Species *inquirendae* (38 species)

- V. brachydonta* Allgén, 1959.
- V. brachylaimoides* Chitwood, 1937.
- V. brevidentata* (Vitiello, 1967) Lorenzen, 1981.
 - syn. *Mononcholaimus brevidentatus* Vitiello, 1967.
- V. conicaudata* (Kreis, 1932) Lorenzen, 1981.
 - syn. *Mononcholaimus conicaudatus* Kreis, 1932.
- V. crassa* Kreis, 1932.
- V. cryptodentata* Allgén, 1959.
- V. diodon* (Wieser, 1951) Lorenzen, 1981.
 - syn. *Mononcholaimus diodon* (Wieser, 1951) Wieser, 1953.
 - syn. *Oncholaimellus diodon* Wieser, 1951.
- V. dubiosa* Kreis, 1932.
- V. fatigans* Filipjev, 1946.
- V. filipjevi* Paramonov, 1929.
- V. gabriolae* (Allgén, 1951) Lorenzen, 1981.
 - syn. *Mononcholaimus gabriolae* Allgén, 1951.
- V. glaberoides* (Allgén, 1932) Lorenzen, 1981.
 - syn. *Mononcholaimus glaberoides* Allgén, 1932.
- V. isotonchula* Kreis, 1932.
- V. klatti* (Allgén, 1941) Lorenzen, 1981.
 - syn. *Mononcholaimus klatti* Allgén, 1941.
- V. leptolaima* Kreis, 1932.
- V. linstowi* (de Man, 1904) Filipjev, 1918.
 - syn. *Oncholaimus linstowi* de Man, 1904.
- V. longicaudata* (Kreis, 1932) Lorenzen, 1981.
 - syn. *Meroviscosia longicaudata* Kreis, 1932.
- V. longidentata* (Schuurmans Stekhoven, 1931) Lorenzen, 1981.
 - syn. *Mononcholaimus longidentatus* (Schuurmans Stekhoven & Adam in Schuurmans Stekhoven, 1931) Kreis, 1934.
 - syn. *Oncholaimus longidentatus* Schuurmans Stekhoven & Adam in Schuurmans Stekhoven, 1931.
- V. longissima* Filipjev, 1946.
- V. macrorhopalocerca* Kreis, 1932.
- V. minor* Filipjev, 1918.
- V. minudonta* Vitiello, 1970.
- V. nijhoffi* Allgén, 1935.
- V. nordgaardi* Allgén, 1940.
- V. palmae* Schuurmans Stekhoven, 1942.

- V. papillosa* (Eberth, 1863) Kreis, 1934.
syn. *Oncholaimus papillosus* Eberth, 1863.
- V. paralinstowi* Chitwood, 1937.
- V. parapedroensis* Allgén, 1947.
- V. paridentata* Kreis, 1932.
- V. parva* Kreis, 1929.
- V. pellucida* (Cobb, 1898) Filipjev, 1918.
syn. *Oncholaimus pellucidus* Cobb, 1898.
- V. poseidonica* Belogurov & Belogurova, 1977.
- V. pseudosegmentata* Allgén, 1947.
- V. stenostoma* Platonova, 1971.
- V. strandi* Allgén, 1935.
- V. subantarctica* Allgén, 1959.
- V. tasmaniensis* (Allgén, 1927) Lorenzen, 1981.
syn. *Mononcholaimus tasmaniensis* Allgén, 1927.
- V. tenuilaima* Allgén, 1959.

Invalid species

- V. aegyptica* (Steiner, 1921) Schuurmans Stekhoven 1943 = *Oncholaimus aegypticus* Steiner, 1921.
- V. carnleyensis* (Ditlevsen, 1921) Kreis, 1932 = *Viscosia glabra* (Bastian, 1865) de Man, 1890.
- V. carnleyensis tropica* Kreis, 1932 = *V. carnleyensis* (Ditlevsen, 1921) Kreis, 1932.
- V. donsi* Allgén, 1947 = *Oncholaimus donsi* (Allgén, 1947) Wieser, 1953.
- V. micoletzkyi* Chitwood, 1951 = *Viscosia glabra* (Bastian, 1865) de Man, 1890.
- V. pacifica* Allgén, 1951 = *Oncholaimus rapax* Kreis, 1932.
- V. papillata nicaraguensis* Gerlach, 1957 = *Viscosia nicaraguensis* Gerlach, 1957.
- V. paralangrunensis* Allgén, 1947 = *Oncholaimus paralangrunensis* (Allgén, 1947) Allgén, 1959.
- V. pellucida* Allgén, 1959 = *V. parapellucida* (Allgén, 1959) Gerlach & Riemann, 1974.
- V. scanica* Allgén, 1935 = *Oncholaimus scanicus* (Allgén, 1935) Wieser, 1953.
- V. taboguillensis* Allgén, 1947 = *Adoncholaimus taboguillensis* (Allgén, 1947) Wieser, 1953.

Remarks. According to the relevant literature, there are 27 species described only from juveniles or females (five species, *V. brachydonta*, *V. conicaudata*, *V. leptolaima*, *V. macrorhopalocerca* and *V. palmae*, were described only from juveniles; 18 species, *V. brevidentata*, *V. cryptodentata*, *V. fatigans*, *V. filipjevi*, *V. isotonchula*, *V. linstowi*, *V. longidentata*, *V. minor*, *V. papillosa*, *V. paralinstowi*, *V. paridentata*, *V. parva*, *V. pellucida*, *V. poseidonica*, *V. stenostoma*, *V. subantarctica*, *V. tasmaniensis* and *V. tenuilaima*, were described only from females; four species, *V. crassa*, *V. diodon*, *V. longissimi* and *V. minudonta*, were described from females and juveniles). These species should be considered as *inquirendae*. According to the original description of *V. brachylaimoides* by Chitwood (1937) and the opinion of Leduc and Zhao (2023), this species has a large left ventrosublateral tooth instead of large right ventrosublateral tooth, indicating that the species does not belong to *Viscosia*. Here, *V. brachylaimoides* is treated as species *inquirendum*. Based on the

observation and analysis by Smol and Sharma (1984), the largest ventro-sublateral tooth of *V. franzii* is situated on the right side instead of the left as described by Boucher (1977). The species is consistent with the diagnosis of *Viscosia*, and it is a valid species of the genus. Meanwhile, Smol and Sharma considered *V. carnleyensis* as a synonym of *V. glabra*.

***Viscosia media* sp. nov.**

<https://zoobank.org/EE875370-C9F8-4025-A4D0-519431DC0893>

Figs 1–3, Table 1

Diagnosis. Body slender, medium size in the genus. Heavily cuticularized and relatively shallow buccal cavity with three stubby teeth, and a right ventro-sublateral tooth massive. Cephalic setae 7–8 µm long. Amphidial fovea invisible. Spicules slender, almost straight, cephalated proximally and conical distally. Tail conical, straight in males, slightly bent ventrally in females.

Holotype and paratype material. Four males and two females were measured. **Holotype** male 1 on slide RZ080312-9. **Paratype** 1 (male 2) on slide RZ080310-3, both **paratype** 2 (male 3) and **paratype** 3 (male 4) on slide RZ0803123-4, **paratype** 4 (female 1) on slide RZ080312-9, and **paratype** 5 (female 2) on slide RZ0803123-4.

Type locality and habitat. Holotype and paratypes were all collected from the surface layer of fine sand sediment on an intertidal beach along the Rizhao coast of the Yellow Sea, China (35°34'21"N, 119°39'29"E).

Etymology. Species epithet *media* refers to the medium body size of this species within the genus.

Measurements. All measurement data are given in Table 1.

Descriptions. Males. Body medium in size, and slender, tapering slightly towards both extremities. Cuticle smooth. Cervical setae 4–6 µm long, sparse, present only in the anterior portion of pharyngeal region. Cephalic region truncated, continuing with body contour. Six lips, each bearing a single inner labial papilla. Six outer labial setae and four slightly longer cephalic setae in a single circle. Outer labial setae 6–7 µm long, cephalic setae 7–8 µm long. Amphidial fovea invisible. Buccal cavity heavily cuticularized, relatively shallow, 15–18 µm deep and 7–9 µm wide, with three stubby teeth. Right ventro-sublateral tooth massive and larger than left ventro-sublateral tooth and dorsal tooth. The tip of right ventro-sublateral tooth at the same level with outer labial and cephalic setae. The height of left ventro-sublateral tooth and dorsal tooth are almost equal. Pharynx cylindrical, widening slightly towards posterior extremity. Cardia conical, surrounded by intestinal tissue. Nerve ring located pre-mid of length of pharynx. Secretory-excretory system present; excretory pore located slightly posterior to nerve ring, 160–170 µm from anterior end. Ventral gland located at anterior part of the intestine.

Reproductive system with two opposed and outstretched testes located to the right of intestine. Sperm cells oval or irregularly square, 12–16 µm long and 8–10 µm wide. Spicules slender, almost straight, cephalated proximally and conical distally. Gubernaculum absent. Tail conical, narrowing abruptly immediately posterior to cloaca and directed slightly dorsally, without caudal setae. Three caudal glands extending anteriorly to tail region. Spinneret present.




Figure 1. Drawings of *Viscosia media* sp. nov. **A** anterior end of holotype, showing outer labial seta (arrow 1), outer labial and cephalic setae (arrow 2), buccal cavity with three teeth, and cervical setae (arrow 3) **B** anterior end of female, showing large right subventral tooth, small dorsal tooth, outer labial and cephalic setae (arrow 1) and cervical setae (arrow 2) **C** posterior end of holotype, showing spicule and tail **D** spicule **E** posterior end of female, showing anus (arrow) and tail **F** entire view of female (arrow 1, vulva; 2, anus; 3, excretory pore; 4, caudal gland; 5, ventral gland; 6, ovary) **G** posterior part of female (arrow 1, Demanian system; 2, ovary; 3, egg; 4, vulva; 5, anus) **H** entire view of male (arrow 1, ventral gland; arrow 2, testis). Scale bars: 20 μm (**A–E**); 50 μm (**F–H**).




Figure 2. Micrographs of *Viscosia media* sp. nov. **A** anterior end of holotype, showing outer labial and cephalic setae (arrow); buccal cavity with three teeth **B** anterior end of female, showing right subventral tooth and cervical setae (arrow) **C** posterior end of holotype, showing spicule and tail **D** cloacal region of male, showing spicule. Scale bars: 10 μ m.




Figure 3. Micrographs of *Viscosia media* sp. nov. **A** middle region of female body, showing vulva (arrow), ovary and eggs **B** showing gland cells (arrow) at the reflexed point of ovary **C** showing sac-like structure and osmosium of Demanian system (arrows).

Table 1. Individual measurements of *Viscosia media* sp. nov. and *Viscosia sinica* sp. nov. (in μm except a, b, c, c' and V%) a, ratio of body length to maximum body diameter; b, ratio of body length to pharynx length; c, ratio of body length to tail length; c', ratio of tail length to cloacal or anus body diameter; V%, position of vulva from anterior end expressed as a percentage of total body length; -, no data.

Characters	<i>Viscosia media</i> sp. nov.						<i>Viscosia sinica</i> sp. nov.					
	♂1	♂2	♂3	♂4	♀1	♀2	♂1	♂2	♂3	♂4	♀1	♀2
Total body length	1620	1534	1615	1522	1539	1458	1733	1885	1840	1821	1457	1653
Maximum body diameter	32	33	33	26	41	35	28	26	26	25	31	28
Head diameter	17	16	17	16	18	17	13	14	14	14	13	15
Length of outer labial setae	6	7	7	6	6	6	5	7	7	5	5	6
Length of cephalic setae	7	8	7	7	7	6	5	7	7	5	5	6
Depth of buccal cavity	18	15	16	17	16	17	29	22	21	26	16	16
Width of buccal cavity	9	8	8	7	9	9	10	8	8	8	8	9
Height of amphidial fovea	-	-	-	-	-	-	5	5	6	5	4	5
Width of amphidial fovea	-	-	-	-	-	-	10	11	11	9	8	8
Length of pharynx	317	310	320	307	305	312	278	298	287	304	260	304
Body diameter at pharyngeal base	26	27	27	24	32	31	24	23	24	24	31	26
Spicules length along arc	29	24	30	32	-	-	20	21	22	19	-	-
Vulva from anterior end	-	-	-	-	732	690	-	-	-	-	720	820
V%	-	-	-	-	47.6	47.3	-	-	-	-	49.4	49.6
Body diameter at cloaca or anus	17	16	17	15	16	15	16	16	16	16	19	15
Tail length	86	75	85	80	75	83	81	104	106	98	88	97
a	50.6	46.5	48.9	58.5	37.5	41.7	61.9	72.5	70.8	72.8	47.0	59.0
b	5.1	4.9	5.1	5.0	5.0	4.7	6.2	6.3	6.4	6.1	5.6	5.4
c	18.8	20.4	19.0	19.0	20.5	17.6	21.4	18.1	17.4	18.6	16.6	17.0
c'	5.1	4.7	5.0	5.3	4.7	5.5	5.1	6.5	6.6	6.1	4.4	6.5

Females. Similar to males in most morphological characteristics but there are less cervical setae, and tail slightly bent ventrally. Reproductive system with two opposed and reflexed ovaries both located to the right of intestine. Eggs very long, can be up to 220 μm long. Demanian system simple, consisting of a sac-like prolongation of the ovary at the reflexed point of the ovary and a duct-shaped osmosium. Vulva located at mid-body.

Differential diagnosis and discussion. *Viscosia media* sp. nov. is characterized by heavily cuticularized and relatively shallow buccal cavity with three stubby teeth, right ventrosublateral tooth massive, cephalic setae 7–8 μm long, amphidial fovea invisible, spicules slender, almost straight, cephalated proximally and conical distally, tail conical, straight in males, slightly bent ventrally in females. The new species is similar to *V. epapillosa* Platonova, 1971 in the deep of buccal cavity and length of cephalic setae, but differs by the wider head diameter (16–18 μm versus 11–13 μm), buccal cavity with heavily cuticularized parallel walls (versus buccal cavity wider in the middle and narrower at the ends), having cervical setae (versus absent in the latter species), spicules enlarged proximally (versus not enlarged), and a conical tail almost straight in the males (versus conico-cylindrical tail bent in the latter species). The new species is also similar to *Viscosia pygmaea* Nguyen Vu Thanh & Gagarin, 2013 in the tail shape, but differs from *V. pygmaea* by the longer body length (1458–1620 μm versus 838–992 μm), longer cephalic setae (7–8 μm versus 3–3.5 μm), shallower buccal cavity with stubby teeth (1 head diameter versus 1.5 head diameters in depth with slender teeth), amphidial fovea invisible (versus obvious and located close to the base of buccal cavity), and the different spicule shape (slender with cephalated proximal end and tapering distally end versus conical and tapering off). *Viscosia media* sp. nov. can be differentiated from all other species of the genus by its heavily cuticularized and relatively shallow buccal cavity with stubby teeth, 7–8 μm cephalic setae, and a conical tail. The basic morphological characteristics of the valid species with similar size to the new species in the genus are compared in Table 2.

***Viscosia sinica* sp. nov.**

<https://zoobank.org/1EB9C0EF-BDC5-484C-8932-4CFBA8F1D88D>

Figs 4, 5, Table 1

Dianosis. Relatively large amphidial fovea, about 80% corresponding body diameter wider in males and 53–62% in females. Cephalic setae 5–7 μm long. Tail conico-cylindrical with swollen horseshoe-shaped tip. Spicules slender, slightly curved ventrally, not cephalate proximally. 10–12 setae surrounding the cloaca, each 3–4 μm long.

Holotype and paratype material. Four males and two females were measured. **Holotype** male 1 on slide RZ080123-4. **Paratype** 1 (male 2) on slide YST24251-2, paratype 2 (male 3) on slide YST24381-11, paratype 3 (male 4) on slide YST24253-1, paratype 4 (female 1) on slide RZ0803123-4, paratype 5 (female 2) on slide YST2418-12.

Type locality and habitat. Holotype and paratype 4 (female 1) were collected from the surface layer of fine sand sediment on an intertidal beach along the Rizhao coast of the Yellow Sea (35°34'21"N, 119°39'29"E). The other paratypes

were all collected from the surface layer of silt sediment on an intertidal beach of Huangdao along the Yellow Sea (31°44'53"N, 121°55'43"E).

Etymology. The species epithet refers to the country of origin, China.

Measurements. All measurement data are given in Table 1.

Descriptions. Males. Body relatively slender. Cuticle smooth, without somatic setae. Labial region spherical, demarcated by slight constriction. Six lips, each bearing a single internal labial papilla. Six outer labial setae and four cephalic




Figure 4. Drawings of *Viscosia sinica* sp. nov. **A** anterior end of holotype, showing outer labial and cephalic setae, buccal cavity with three teeth **B** anterior end of male, showing amphidial fovea (arrow) **C** anterior end of female **D** spicule and setae surrounding the cloaca **E** posterior end of female, showing anus (arrow) and tail **F** posterior end of holotype, showing cloaca (arrow), spicule and tail **G** entire view of male (arrow showing testis) **H** entire view of female (arrow 1, ovary; 2, vulva; 3, anus). Scale bars: 10 μ m (**A–D, F**); 20 μ m (**E**); 50 μ m (**G, H**).




Figure 5. Micrographs of *Viscosia sinica* sp. nov. **A** anterior end of holotype, showing buccal cavity with large right subventral and dorsal teeth **B** anterior end of male 1, showing outer labial and cephalic setae, amphidial fovea (arrow) **C** cloacal region of male 2, showing spicule and setae surrounding cloaca (arrows) **D** posterior end of holotype, showing spicule and tail. Scale bars: 10 μ m.

setae in a single circle, equal in length. Amphidial fovea relatively large, pocket-shaped, 9–11 μm wide, about 80% corresponding body diameter, and 5–6 μm deep, located at level of the middle of buccal cavity. Buccal cavity 16–29 μm deep and 8–10 μm wide, with three teeth. Right ventrosublateral tooth larger than left ventrosublateral tooth and dorsal tooth. The tip of right ventrosublateral tooth at the same level with outer labial and cephalic setae. The height of left ventrosublateral tooth and dorsal tooth are almost equal. Pharynx cylindrical, widening slightly towards posterior extremity. Cardia conical, about 15 μm long, surrounded by intestinal tissue. Nerve ring located approximately halfway down length of pharynx. Secretory-excretory system and excretory pore not observed.

Reproductive system with two opposed and outstretched testes located to the right of intestine. Spicules slender, slightly curved ventrally, not cephalate proximally. Gubernaculum absent. 10–12 setae surrounding the cloaca, each 3–4 μm long. Tail conico-cylindrical, with swollen horseshoe-shaped tip. Caudal setae absent. Three caudal glands extending anteriorly to tail region. Spinneret present.

Females. Similar to males in the most morphological characteristics but amphidial fovea slightly smaller, 53–62% corresponding body diameter wider, and buccal cavity slightly shallower (21–29 μm versus 16 μm deep). Reproductive system with two opposed and reflexed ovaries both located to the right of intestine. Demanian system indistinct. Vulva located at mid-body.

Table 2. Basic morphological features of males of the valid species in *Viscosia* with body length of 1000–2900 μm (according to Gagarin 2020) (in μm except a, b, c and c') L, body length; a, ratio of body length to maximum body diameter; b, ratio of body length to pharynx length; c, ratio of body length to tail length; c', ratio of tail length to cloacal or anus body diameter.

Species	L	a	b	c	c'	Cephalic sensilla length	Buccal cavity depth	Spicules length
<i>V. bandaensis</i>	1389–1715	41–48	4.8–5.1	12.8–13.8	6.0–6.5	papilla	23	18–20
<i>V. bayensis</i>	1630–1820	41–48	5.2	17.3	5.6	9–11	25–29	43–50
<i>V. dossena</i>	1759–2048	23–25	5–6	19–20	3.0	4–6	31–35	27–33
<i>V. epapillosa</i>	1750	51	5.3	20.7	4.9	6	16	29
<i>V. erasmi</i>	1905	60	5.7	15.9	5.0	5	23	32
<i>V. glabra</i>	1600–2100	42–70	5.3–6.3	7.5–11.0	10–13.0	papilla	23–25	27
<i>V. longicaudatoides</i>	1348–1432	53–54	4.8–5.4	8.3–9.0	10.1–10.8	2.3–3	17–18	27
<i>V. macramphida</i>	1400	39	5.6	7.9	9.6	papilla	15	18–19
<i>V. macrobursata</i>	1680–2200	50–53	5.7	13.2	8.6	10–12	23–29	20
<i>V. meridionalis</i>	1415–1743	30–43	5.3–6.9	8.2–9.6	7.3–9.8	papilla	18–25	32–40
<i>V. media</i> sp. nov.	1522–1620	46.5–58.5	4.9–5.1	18.8–20.4	4.7–5.5	7–8	16–18	24–30
<i>V. microseta</i>	1850	44	6.1	16.5	4.3	1.5–2.0 μm	20–22	33
<i>V. nuda</i>	1625–1931	45–54	5.2–6.0	7.3–8.0	13.0–13.5	papilla	18–22	18
<i>V. oncholaimelloides</i>	1950	89	6.5	13.9	9.1	papilla	13	17
<i>V. orientalis</i>	991–1122	17–20	4.7–5.1	15.7–19.9	2.8–3.0	2.0 μm	18–20	34–36
<i>V. papillatoides</i>	1520–2200	40–51	5.5–5.6	19.1–23.0	3.4–4.6	papilla	24–32	26–29
<i>V. papillate</i>	1520	31	5.0	11.0	6.0	papilla	18	24
<i>V. parasetosa</i>	1721–1748	52–55	5.7–5.9	11.6–12.0	8.2–8.3	2.7 μm	18–20	24
<i>V. profunda</i>	1311–1501	47–50	5.0–5.6	14.8–15.4	4.5–5.2	2.0–2.5 μm	17.5–18	17.5
<i>V. separabilis</i>	1320–2200	32–78	4.8–7.8	12.9–16.6	5.0–8.0	7.5 μm	20–26	19–26
<i>V. setosus</i>	1784–1857	51–57	6.0–6.1	11.9–12.1	8.0–9.0	5.0 μm	20–21	22
<i>V. sinica</i> sp. nov.	1733–1885	62–73	6.1–6.4	17.4–21.4	5.1–6.6	5–7 μm	26–29	19–22
<i>V. stenolaimus</i>	1870	37	4.5	12.6	4.6	papilla	25–27	37
<i>V. timmi</i>	1016–1509	22–37	4.1–5.1	16.0–25.6	2.0–3.5	2.5–3.0 μm	21–28	24–28
<i>V. viscosa</i>	1700–2600	52–60	5.0–6.5	12.0–16.0	6.0–7.5	4.0 μm	22–30	27–30
<i>V. weiseri</i>	1600–2900	29–46	4.3–5.2	13.3–17.7	4.5	5.0 μm	28–30	42

Differential diagnosis and discussion. *Viscosia sinica* sp. nov. is characterized by relatively large amphidial fovea, conico-cylindrical tail with horse-shoe-shaped tip, spicules slender, slightly curved ventrally, not cephalate proximally, 10–12 setae surrounding the cloaca, each 3–4 μm long. The new species resembles *V. brachylaima* Filipjev, 1927 and *V. filipjevi* Paramonov, 1929 in tail shape, but differs from *V. brachylaima* by larger amphidial fovea (80% versus 30% corresponding body diameter) and higher value of de Man ratio *a* in males (61.9–72.8 versus 42.5). The new species differs from *V. filipjevi* which description is based only in females by larger amphidial fovea (wider than width of buccal cavity versus narrower than width of buccal cavity), and smaller body size (1.46–1.65 mm in body length and 28–31 μm maximum body diameter in females versus 2.17–2.20 mm in length and 40.5–48.6 μm maximum body diameter in *V. filipjevi*). The new species is also similar to *V. elegans* (Kreis, 1924) Lorenzen, 1981 in body size and shape, but differs by larger amphidial fovea (80% versus 33% corresponding body diameter) and horseshoe-shaped tail tip. The new species differs from *V. media* sp. nov. in having deeper buccal cavity with slender teeth in males (21–29 μm deep versus 15–18 μm with stubby teeth), larger amphidial fovea (versus invisible), and different tail shape (conico-cylindrical with swollen horseshoe-shaped end versus conical). *Viscosia sinica* sp. nov. is distinguished from all other known species of the genus by its relatively large amphidial fovea, 5–7 μm cephalic setae, and conico-cylindrical tail with swollen horseshoe-shaped end.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (No. 41176107). Authors are very thankful to Prof. Dr Virag Venekey (the famous nematologist from Brazil) for constructive suggestions and for improving English writing of this article. We are sincerely grateful to anonymous referees for reviewing and improving the manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research was funded by the National Natural Science Foundation of China (grant number 41676146) and the Natural Science Foundation of Shandong Province (grant number ZR2022QC218).

Author contributions

Project management, work program and taxonomy: Y.H.; methodology, experiment, and data collection: L.Y.S. and H.M.G; writing and editing of paper: Y.H. and L.Y.S. All authors have read and agreed to the submitted version of the manuscript.

Author ORCIDs

Lingyun Sun  <https://orcid.org/0009-0005-3553-3645>

Huimin Gu  <https://orcid.org/0009-0008-5820-4678>

Yong Huang  <https://orcid.org/0000-0002-1846-8088>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Allgén CA (1927) Freilebende marine Nematoden von der Küste Tasmaniens. *Zoologischer Anzeiger* 9(10): 197–217.
- Allgén CA (1930) Über einige neue oder wenig bekannte Brachwasser-Nematoden von der Litoralzone des Öresunds. *Zoologischer Anzeiger* 85: 58–72.
- Allgén CA (1932) Weitere Beiträge zur Kenntnis der marinen nematodenfauna der Campbellinsel. *Nyt Magazin for Naturvidenskaberne* 70: 97–198.
- Allgén CA (1933) Freilebende Nematoden aus dem Trondhjemsfjord. *Capita Zoologica* 4(2): 1–162.
- Allgén CA (1935) Die freilebenden Nematoden des Öresunds. *Capita Zoologica* 6(3): 1–192.
- Allgén CA (1940) Über einige neue freilebende Nematoden von der Nordwest- und Nordküste Norwegens. *Folia Zoologica et Hydrobiologica* 10: 443–449.
- Allgén CA (1941) Über einige freilebende Nematoden aus Bandirma an der Nordküste Kleinasiens (Marmarameer, Türkei). *Zoologischer Anzeiger* 135: 281–294.
- Allgén CA (1946) Zur Kenntnis norwegischer Nematoden VI. Neue freilebende marine Nematoden von der Insel Frøya. *Kongelige Norske Videnskabers Selskab Forhandlingar* 18(38): 160–162.
- Allgén CA (1947) Die Nematoden-Familie Tripyloididae, ihre Arten und Verwandtschaft. *Arkiv for Zoologi* 39: 1–35.
- Allgén CA (1951) Pacific Freelifving Marine Nematodes. (Papers from Dr. Th. Mortensen's Pacific Expedition 1914–1916. LXXVI). *Videnskabelige Meddelelser fra Dansk naturhistorisk Forening* 113: 263–411.
- Allgén CA (1959) Freelifving marine nematodes. Further zoological results of the Swedish Antarctic expedition, 1901–1903 under the direction of Dr. Otto Nordenskjöld. 5(2) P.A. Norstedt & Söner, Stockholm, 293 pp.
- Bastian HC (1865) Monograph of the Anguillulidae, or Free Nematoids, Marine, Land, and Freshwater; with Descriptions of 100 New Species. *The Transactions of the Linnean Society of London* 25(2): 73–184. <https://doi.org/10.1111/j.1096-3642.1865.tb00179.x>
- Belogurov OI, Belogurova LS (1977) The morphology of *Viscosia poseidonica* n. sp. and *V. stenostoma* Platonova, 1971 (Nematoda, Oncholaimidae) free-living nematodes from the neighbourhood of the island of Furugel'ma Islands (Translated from Russian). *Nauchnye Doklady Vyssei Shkoly, Biologicheskie Nauki* 6: 44–47.
- Boucher G (1977) Nématodes des sables fins infralittoraux de la Pierre Noire (Manche occidentale). IV. Enoplida. *Bulletin du Muséum National d'Histoire Naturelle* 468: 733–752. <https://doi.org/10.5962/p.282054>
- Chitwood BG (1937) A new genus and ten new species of marine nematodes from North Carolina. *Proceedings of the Helminthological Society of Washington* 4: 54–59.
- Chitwood BG (1951) North American nematodes. *The Texas Journal of Science* 4: 617–672.

- Chitwood BG (1960) A preliminary contribution on the marine nemas (Adenophorea) of Northern California. *Transactions of the American Microscopical Society* 79: 347–384. <https://doi.org/10.2307/3224119>
- Cobb NA (1890) Arabian Nematodes. *Proceedings of the Linnean Society of New South Wales* 5: 449–468. <https://doi.org/10.5962/bhl.part.18647>
- Cobb NA (1898) Australian free-living marine nematodes. *Proceedings of the Linnean Society of New South Wales* 23(3): 383–407.
- Coomans AV, Vincx M, Decraemer W (1985) Nematodes from a fresh-water pool on a coral island in the Solomon Islands. *Hydrobiologia* 123(3): 265–281. <https://doi.org/10.1007/BF00034389>
- de Jonge VN, Bouwman LA (1977) A simple density separation technique for quantitative isolation of meiobenthos using the colloidal silica Ludox-TM. *Marine Biology* 42: 143–148. <https://doi.org/10.1007/BF00391564>
- de Man JG (1890) Quatrième note sur les nématodes libres de la mer du Nord et de la Manche. *Mémoires de la Société Zoologique de France* 3: 169–194.
- de Man, JG (1904) Nématodes libres (Expédition. Antarctique Belge). *Résultats du Voyage S. Y. Belgica*, 1–51.
- Ditlevsen H (1921) Papers from Dr. Th. Mortensens Pacific Expedition 1914–1916. III Marine free-living Nematodes from the Auckland and Campbell Islands. *Videnskabelige Meddelelser fra Dansk naturhistorisk Forening i Kjøbenhavn* 73: 1–39.
- Ditlevsen H (1928) Free-living marine nematodes from Greenland waters. *Meddelelser om Grønland* 23: 199–250.
- Dujardin F (1845) *Histoire naturelle des helminthes ou vers intestinaux*. Librairie Encyclopedique de Roret. Paris, 654 pp. <https://doi.org/10.5962/bhl.title.10123>
- Eberth CJ (1863) *Untersuchungen über Nematoden*. Leipzig, Verlag von Wilhelm Engelmann, 77 pp.
- Filipjev IN (1916) Les nematodes libres contenus dans les collections du musee Zoologique de l'Academie Imperiale des Sciences de Petrograd. *Annales du Museum Zoologique de Academie des Sciences Petrograd* 21: 59–116.
- Filipjev IN (1918) Free-living marine Nematodes from the vicinity of Sevastopol. Part 1. *Trudy Osoboï Zoologicheskoi Laboratorii Sebastopolskoi Stantsii Rossiïkoi, Series 2* 4: 1–614.
- Filipjev IN (1922) New data on free nematodes of the Black Sea (Novye Dannye o Svobodnykh Nematodakh Chernogo Moria.). *Trudy Stavropolskogo Selskokhoziaistvennogo Instituta* 1: 13–184.
- Filipjev IN (1927) Les Nématodes libres des mers septentrionales appartenant a la famille des Enoplidae. *Archiv für Naturgeschichte* 91: 1–216.
- Filipjev IN (1929) Classification of freelifing Nematoda and relations to parasitic forms. *Journal of Parasitology* 15: 281–282.
- Filipjev IN (1946) Nématodes libres du bassin polaire. *Trudy Dreifuiushchiae Ekspeditsiia Glavsevmorputi na Ledokol'nom Parokhode G. Sedov 1937–1940* 3: 158–184.
- Furstenberg JP, Vincx M (1989) Two oncholaimid species from a South African estuary (Nematoda, Oncholaimidae). *Hydrobiologia* 184: 43–50. <https://doi.org/10.1007/BF00014300>
- Gagarin VG, Nguyen Vu Thanh (2007) *Ingenia communis* sp. n., *Adoncholaimus longispiculatus* sp. n. and *Viscosia sedata* sp. n. (Nematoda: Enoplida) from Mekong River Delta, Vietnam. *International Journal of Nematology* 17(2): 205–212.
- Gagarin VG (2020) *Halalaimus borealis* sp. nov. and *Viscosia orientalis* sp. nov. (Nematoda, Enoplida) from the mouth of the Cam River in Vietnam. *Amurian Zoological Journal* 12: 26–42. <https://doi.org/10.33910/2686-9519-2020-12-1-26-42>

- Gagarin VG, Nguyen TT (2008). Free-living nematodes from the Red River delta, Vietnam. *Inland Water Biology* 1(4): 320–325. <https://doi.org/10.1134/S1995082908040020>
- Gerlach SA (1957) Die Nematodenfauna des Sandstrandes an der Küste von Mittelbrasilien (Brasilianische Meerse-Nematoden IV). *Mitteilungen aus dem Zoologischen Museum Berlin* 33: 411–459. <https://doi.org/10.1002/mmnz.19570330206>
- Gerlach SA, Riemann F (1974) The Bremerhaven Checklist of Aquatic Nematodes. A catalogue of Nematoda Adenophorea, excluding the Doryaimida. Part 2. Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven Supplement 4(2): 404–735.
- Heip C, Smol N, Vranken G (1982) The systematics and ecology of free-living marine nematodes. *Helminthological Abstracts Series B* 51(1): 1–31.
- Hope WD, Murphy DG (1972) A taxonomic hierarchy and checklist of the genera and higher taxa of marine nematodes. *Smithsonian Contributions to Zoology* 137: 1–101. <https://doi.org/10.5479/si.00810282.137>
- Huang M, Sui XX (2024) Description of *Amphimonhystrella sinica* sp. nov. and *Cobbia zhangii* sp. nov. of the family Xyalidae (Nematoda: Monhysterida) from the Yellow Sea, China *Zootaxa* 5471(3): 343–354. <https://doi.org/10.11646/zootaxa.5471.3.4>
- Huang M, Zhai HX (2024) Two new species of the genus *Halichoanolaimus* (Nematoda, Selachinematidae) from the intertidal zone of the Yellow Sea, China. *Zookeys* 1208: 259–274. <https://doi.org/10.3897/zookeys.1208.124047>
- Inglis WG (1983) An outline classification of the phylum Nematoda. *Australian Journal of Zoology* 31: 243–255. <https://doi.org/10.1071/ZO9830243>
- Keppner EJ (1987) Five new and one known species of free-living marine nematodes of the family Oncholaimidae (Nematoda: Enoplida) from Northwest Florida, U.S.A. *Transactions of the American Microscopical Society* 106(3): 214–231. <https://doi.org/10.2307/3226251>
- Kreis HA (1924) Zur Kenntnis der freilebenden marinen Nematoden. *Schriften für Süßwasser und Meereskunde, Büsum* 2: 157–170.
- Kreis HA (1929) Freilebende marine Nematoden von der Nordwest-Küste Frankreichs (Trébeurden Côtes du Nord). *Capita Zoologica* II 7: 1–98.
- Kreis HA (1932) Freilebende marine Nematoden von den Sunda-Inseln II. Oncholaiminae. *Videnskabelige Meddelelser fra Dansk naturhistorisk Forening* 93: 23–69.
- Kreis HA (1934) Oncholaiminae Filipjev 1916. Eine monographische Studie. *Capita Zoologica* 4(5): 1–271.
- Lambshhead PJD (2004) Marine nematode biodiversity. In: Chen ZX, Chen SY, Dickson DW (Eds) *Nematology: Advances and Perspectives*, Vol. 1. Nematode Morphology, Physiology and Ecology. CABI, Wallingford, 438–468. <https://doi.org/10.1079/9780851996455.0438>
- Leduc D, Zhao ZQ (2023) The Marine Biota of Aotearoa New Zealand. Ngā toke o Parumoana: Common free-living Nematoda of Pāuatahanui Inlet, Te-Awarua-o-Porirua Harbour, Wellington. *NIWA Biodiversity Memoir*, 135, 212 pp. <https://niwa.co.nz/oceans/niwa-biodiversity-memoirs>
- Li T, Huang Y, Huang M (2024) Two new species of the order Monhysterida (Nematoda) from the sea of China, *Zookeys* 1193: 63–79. <https://doi.org/10.3897/zookeys.1193.110188>
- Lorenzen S (1974) Die Nematodenfauna der sublitoralen Region der Deutschen Bucht, insbesondere im Titan-Abwassergebiet bei Helgoland. *Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven* 14: 305–327.

- Lorenzen S (1981) Entwurf eines phylogenetischen System der freilebenden Nematoden. Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven 7: 1–472.
- Mawson PM (1958) Free-living nematodes. Section 3: Enoploidea from subantarctic stations. B.A.N.Z. Antarctic Research Expedition Reports, Series B 6(14): 307–358.
- McIntyre AD, Warwick RM (1984) Meiofauna techniques. In: Holme NA, McIntyre AD (Eds) Methods for the study of marine benthos. Blackwell Scientific Publications, Oxford, 217–244.
- Nemys eds (2024) Nemys: World Database of Nematodes. [accessed 1 December 2024] <https://doi.org/10.14284/366>
- Nguyen Vu Thanh, Gagarin VG (2013) Three new species of nematodes (Nematoda: Enoplida) from coastal waters of Vietnam. Russian Journal of Marine Biology 39(6): 420–428. <https://doi.org/10.1134/S1063074013060060>
- Paramonov AA (1929) Die freilebenden Nematoden der Kinburnsenge und der angrenzenden Gewässer. Trudy Gosudarstvennoi Ikhtologicheskoi Opytnoi Stantsii 4(1): 59–129.
- Platonova TA (1971) Exploration of the fauna of the seas VIII (XVI). Fauna and flora of the Possjet Bay of the Sea of Japan. Zoological Institute Academy of Sciences of the USSR 8(41): 72–108.
- Schuermans Stekhoven Jr JH (1931) Ökologische und morphologische Notizen über Zuiderseenematoden. I. Die westliche Hälfte der Zuidersee. Zeitschrift für Morphologie Ökologie der Tiere 20 (4): 613–678. <https://doi.org/10.1007/BF00407681>
- Schuermans Stekhoven Jr JH (1942) The free living nematodes of the Mediterranean. III. The Balearic Islands. Zoologische Mededeelingen 23: 229–262.
- Schuermans Stekhoven Jr JH (1943) Freilebende marine Nematoden des Mittelmeeres. IV. Freilebende marine Nematoden der Fischereigründe bei Alexandrien. Zoologische Jahrbücher (Systematik) 76(4): 267–396.
- Smol N, Sharma J (1984) Two new and three redescribed species of *Viscosia* (Nematoda, Oncholaimidea). Hydrobiologia 114: 123–147. <https://doi.org/10.1007/BF00018110>
- Smol N, Muthumbi A, Sharma J (2014) Order Enoplida In: Schmidt-Rhaesa A (Ed.) Handbook of Zoology: Gastrotricha, Cycloneuralia and Gnathifera, Vol. 2: Nematoda. De Gruyter, Berlin, 193–249. <https://doi.org/10.1515/9783110274257.193>
- Steiner G (1921) Beiträge zur Kenntnis mariner Nematoden. Zoologische Jahrbücher (Systematik) 44: 1–68.
- Sun X Y, Bai RB, Zhai HX, Huang Y (2024) Species list of free-living marine nematodes in the Yellow Sea, China. Journal of Liaocheng University (Natural Science Edition) 37(6): 34–52. <https://doi.org/10.19728/j.issn1672-6634.2024050010>
- Vitiello P (1967) Nématodes libres marins de Roscoff. I. Description de cinq espèces nouvelles. Cahiers de Biologie Marine 8: 403–416.
- Vitiello P (1970) Nématodes libres marins des vases profondes du Golfe du Lion. I. Enoplida. Téthys 2: 139–210.
- Wieser W (1951) Untersuchungen über die algenbewohnende Mikrofauna mariner Hartböden. I. Zur Ökologie und Systematik der Nematodenfauna von Plymouth. Österreichische Zoologische Zeitschrift 3: 425–480.
- Wieser W (1953) Reports of the Lund University Chile expedition 1948–1949: 1. Enoploidea. Lunds Universitets Årsskrift, N. F. 49(6): 1–155.
- Wieser W (1959) Free-living nematodes and other small invertebrates of Puget Sound beaches. University of Washington Publications in Biology 19: 1–179.
- Wieser W, Hopper B (1967) Marine nematodes of the east coast of North America. I. Florida. Bulletin of the Museum of Comparative Zoology 135: 239–344.