



# A new species of *Monstrilla* (Copepoda, Monstrilloida) in coastal waters of northern South China Sea

Zhiqian Zhou<sup>1,2</sup>, Xiping Lian<sup>1</sup>, Yehui Tan<sup>1,2</sup>

<sup>1</sup> Marine Biodiversity Collections of South China Sea, Key Laboratory of Tropical Marine Bio-resources and Ecology, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, 510301, China

<sup>2</sup> University of Chinese Academy of Sciences, Beijing, 100049, China

Corresponding author: Xiping Lian ([xplian@scsio.ac.cn](mailto:xplian@scsio.ac.cn))

## Abstract

A new monstrilloid copepod species, *Monstrilla pseudograndis* **sp. nov.**, from coastal waters in the northern South China Sea is described and illustrated. The diagnostic characters of this new species include the presence of small setae and sensilla on the forehead and four pairs of nipple-like scars on the ventral surface of the cephalothorax. Although the new species closely resembles *Monstrilla grandis* Giesbrecht, 1891 in overall morphology, it can be distinguished by the shorter antennule, the thumb-like process on the fifth leg, and the distinctive shape of the oral papilla. This represents the eighth record of this genus from China seas.

**Key words:** copepods, *Monstrilla*, zooplankton, taxonomy, China seas



Academic editor: Maria Cristina Bruno

Received: 19 December 2024

Accepted: 26 February 2025

Published: 18 March 2025

ZooBank: <https://zoobank.org/4FA13ABA-0D63-4648-B4D4-8C0522F7BBDE>

Citation: Zhou Z, Lian X, Tan Y (2025) A new species of *Monstrilla* (Copepoda, Monstrilloida) in coastal waters of northern South China Sea. ZooKeys 1232: 237–248. <https://doi.org/10.3897/zookeys.1232.144830>

Copyright: © Zhiqian Zhou et al.  
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Introduction

The order Monstrilloida Sars, 1901 is renowned for its unique and enigmatic copepods that exhibit a fascinating life cycle transitioning from endoparasitic juveniles to free-living, non-feeding adults (Huys and Boxshall 1991; Suárez-Morales 2011, 2018). Monstrilloids are known to parasitize various marine benthic invertebrates, including polychaetes and molluscs, during their preadult stages (Huys et al. 2007; Suárez-Morales 2011; Suárez-Morales et al. 2014). Adults are characterized by their non-feeding, free-swimming nature, and lack of mouthparts; they are typically encountered in plankton samples from coastal and estuarine environments (Sale et al. 1976; Suárez-Morales 2011; Lee et al. 2016).

In recent decades, the records of monstrilloids have increased rapidly, with seven valid genera, about 183 accepted species, recognized within the single family Monstrillidae Dana, 1849 (Grygier and Ohtsuka 2008; Suárez-Morales 2011, 2018, 2019; Suárez-Morales and Mckinnon 2014; Jeon et al. 2018; Walter and Boxshall 2024). Currently, research on the taxonomy and diversity of Monstrilloida continues to expand globally, with studies covering marine regions of Australia (Suárez-Morales and Mckinnon 2014), Korea (Lee and Chang 2016; Lee et al. 2016; Jeon et al. 2018), China (Lian and Tan 2019), the Philippines (Suárez-Morales 2021), Brazil (Da Cruz Lopes da Rosa et al. 2021), Canada

(Delaforge et al. 2017), Mexico (Suárez-Morales 2024; Suárez-Morales and Velazquez-Ornelas 2024), and the Mediterranean Sea (Suárez-Morales et al. 2017). In addition to diversity studies, research has focused on the biology and ecology of monstrilloids, particularly their parasitic behavior, host interactions, and effects on host health, especially on mussels (Suárez-Morales et al. 2010, 2014; Carneiro-Schaefer et al. 2017).

*Monstrilla* Dana, 1849–1852, is one of the most diverse genera within Monstrillidae (Suárez-Morales 2011, 2018; Jeon et al. 2024), and the proportion of antennule and the shape of fifth leg are important identification criteria in the morphological identification of females of this genus (Lee and Chang 2016; Suárez-Morales 2024). *Monstrilla* has a global distribution comprising 53 species to date and includes about 29% of all recorded monstrilloids (Walter and Boxshall 2024). Despite its extensive contribution to marine biodiversity, taxonomic studies on monstrilloid copepods in China are limited (Suárez-Morales 2018), with only seven species of *Monstrilla* having been recorded in the seas of China (Chen and Li 2008; Chen and Huang 2012; Walter and Boxshall 2024). The absence of data on Chinese species may lead to an underappreciation of the diversity of monstrilloids and hinder adequate risk assessments for mussels and other benthic marine invertebrate aquaculture industries.

During a recent re-examination of zooplankton deposited in the South China Sea marine biodiversity collections of the Chinese Academy of Sciences (SCSMBC), an adult female specimen of *Monstrilla* was discovered. This specimen, after taxonomic analysis and comparison with known congeners, was identified as a new species, *M. pseudograndis* sp. nov. This discovery not only contributes to the growing knowledge of monstrilloid diversity but also expands our understanding of the distribution of the genus *Monstrilla*.

## Material and methods

Zooplankton was collected in the coast near Fangchenggang (21°22'53"N, 108°19'38"E), Guangxi Province, China on 7 June 2023 by vertical tow net (0.505 mm mesh, 0.8 m diameter at towing speed of 0.5 m/s) from the surface to a depth of 15 m (Fig. 1). The material was immediately preserved in 5% formaldehyde. Observation and measurements were carried out under a microscope (SMZ18, Nikon, Japan), and drawings were made with the aid of a camera lucida (Leica MC 190HD). Morphologic terminology follows Huys and Boxshall (1991). The nomenclature for the female monstrilloid antennular armature proposed by Grygier and Ohtsuka (1995) is adopted. The type specimen is deposited in the South China Sea marine biodiversity collections, Chinese Academy of Sciences (SCSMBC).


## Taxonomy

**Subclass Copepoda Milne Edwards, 1840**

**Order Monstrilloida Sars, 1901**

**Family Monstrillidae Dana, 1849**

**Genus *Monstrilla* Dana, 1849–1852**



**Figure 1.** Sampling site of *Monstrilla pseudograndis* sp. nov. near Fangchenggang, Guangxi Province, China.

***Monstrilla pseudograndis* sp. nov.**

<https://zoobank.org/D8D11F64-1FCB-482D-BE80-7DF0CF40076B>

Figs 2–4, Tables 1, 2

**Type material. Holotype:** • adult female (SCSMBC 240185); Zhiqian Zhou leg.; 7 June 2023; partially dissected, formaldehyde preserved.

**Type locality.** CHINA • Guangxi Province; coast near Fangchenggang. 21°22'53"N, 108°19'38"E; salinity 29.42, temperature 31 °C; depth 17 m.

**Etymology.** The species name is derived from the Greek word *pseudo*, meaning “false”, and the name of the closely similar *M. grandis* Giesbrecht, 1891.

**Diagnosis.** Female *Monstrilla* with smooth cuticle on cephalothorax; forehead medially concave, bearing a pair of short sensilla and small setae bilaterally near antennule bases. Cephalothorax ventrally marked by four pairs of small, nipple-like scars, arranged symmetrically anterior to oral papilla. Oral papilla located at approximately midlength of cephalothorax, ventrally posteriorly-bent. Antennule two-segmented, segments fused distally, reaching 37.8% of total body length. Legs 1–4 with relatively short outer exopodal spines. Fifth legs bilobed, outer lobe elongate, with three plumose setae; inner lobe shorter, bearing two plumose setae and a basal protuberance on inner margin. Caudal rami 2.1 times as long as wide, divergent posteriorly, each armed with six well-developed caudal setae.





Figure 2. *Monstrilla pseudograndis* sp. nov., female holotype **A** habitus, dorsal **B** habitus, lateral **C** habitus, ventral. **A–C** share the same scale bar.

**Description of adult female holotype.** Body moderately elongate (Fig. 2A–C), about 1.68 mm, measured from anterior end of cephalothorax to posterior margin of caudal rami, excluding antennules and caudal setae. Cephalothorax rather large and relatively long, accounting for about 57.7% of total body length, transparent, dorsal surface smooth; anterior 2/5 slightly swollen laterally and ventrally. Nauplius eye present, weakly developed, elliptical, ocelli unpigmented with separate oval hyaline bodies, separated by  $1\frac{1}{2}$  eye diameter (Fig. 2A). Anteriormost part of cephalothorax with ventral, rounded convex protuberance with irregular margin in lateral view (Fig. 2B); cuticular ornamentation observed on the surface of cephalothorax in lateral and ventral view. Forehead slightly concave medially between antennular bases in dorsal view, without rostral protrusion, bearing a pair of short, slender sensilla in the middle and a pair of setae near first antennule (Fig. 3B at arrow); weak, fine, longitudinal and transverse



**Figure 3.** *Monstrilla pseudograndis* sp. nov., female holotype **A** cephalosome view, ventral (arrow indicates nipple-like processes) **B** anterior part of cephalothorax, ventral (s1 = sensilla; s2 = setae) **C** postgenital urosomites and caudal rami, ventral **D–G** leg 1–4. **D–G** share the same scale bar.

wrinkles running behind antennular bases on each side of lateroventral surfaces, flanked by four pairs of small nipple-like scars ahead of oral papilla (Fig. 3A at arrow), without sensory pore. Oral papilla situated slightly posterior to mid-length of cephalothorax, accounting for about 52.9%, protruding ventrally, with distal half posteriorly-bent (Fig. 3A).

Antennule long (Fig. 4A, B), about 37.8% of total body length, about 65.5% of the cephalothorax; antennule two-segmented, only first segment distinctly separate, remaining segments fused, with constrictions along antennular body representing places of intersegmental divisions (purported 2–5), length ratio of antennule segments, from basal to distal one: 14.0:86.0 (= 100). In terms of the pattern described by Grygier and Ohtsuka (1995) for female monstrilloid antennular armature, setae (Roman numerals) and spines (Arabic numerals), short, slender element 1 present on first segment; purported segment two with elements 2d<sub>1</sub>, 2d<sub>2</sub>, 2v<sub>1</sub>, 2v<sub>2</sub>, 2v<sub>3</sub>, IId; purported segment three with elements 3, IIIId and IIIv; purported segment four with elements 4v<sub>1-2</sub>, 4d<sub>1</sub>, IVd and IVv as well as 4aes (aesthetasc); purported segment five with elements 5, Vd, Vv and Vm; setae b<sub>1-3</sub>, b<sub>5</sub> all dichotomously branched from proximal half or third, setae b<sub>4</sub> simple, without b<sub>6</sub>. Apical elements 6<sub>1</sub>, 6<sub>2</sub>, and 6aes present, but 6aes absent on right antennule in dorsal view.

Legs 1–4 (Fig. 3D–G) all with both endopod and exopod three-segmented. Coxa without setae and lacking marginal rows of setae or spines. Basis not fully divided medially from coxa; all outer basal setae on legs 1–4 slender, naked; seta on leg 3 much longer. First and second exopodal segments of legs 1–4 slightly swollen, third exopodal segment undulate along outer distal margin; outer margins of all endopodal segments swollen and smooth. Outer distal spines on first and third exopodal segments of legs 1–4 feeble, much shorter than segments bearing them. Seta/spine armature of swimming legs 1–4 as in Table 1.


**Table 1.** Armature of legs 1–4 including basis, exopods, and endopods in *Monstrilla pseudograndis* sp. nov. Roman numerals indicate numbers of spines, and Arabic numerals indicate numbers of setae.

	Coxa	Basis	Endopod	Exopod
Leg 1	0-0	1-0	0-0; 0-1; 1,2,2	I-1;0-1;I,1,3
Leg 2–4	0-0	1-0	0-1; 0-1; 1,2,2	I-1;0-1;I,2,3

Leg 5 bilobed, both lobes confluent basally and divided distally. Outer lobe elongate, armed with three long, plumose setae apically or subapically, of subequal lengths. Inner lobe relatively short, with a basal protuberance on inner margin (arrowed in Fig. 3C), its tip exceeding the half of the outer lobe, armed with two plumose setae apically or subapically.

Urosome consisting of four urosomites: fifth pedigerous somite, genital double-somite, free postgenital somite and anal somite, accounting for 17.9% of total body length, excluding caudal setae; ratio of lengths 30.3:44.0:15.4:10.3 (=100). Genital double-somite representing almost half-length of urosome (44.0%), somewhat swollen laterally, partial suture visible; about 1.7 times longer than the combined length of the next two segments. Genital double-somite bearing pair of long ovigerous spines, these being inserted on middle of ventral surface, basally separated, with pointed tips extending far beyond tips of caudal setae, in total equal to about 38.2% of total body length. Anal somite trapezoidal; lateral margin nearly smooth in dorsal but with apparent notch in ventral; lacking wrinkles or striae both on dorsal and ventral surfaces.

Caudal rami long (Fig. 3C), about 2.1 times as long as wide; divergent outward; with small cuticular protuberance at basal part of outer face and slightly swollen at



**Figure 4.** *Monstrilla pseudograndis* sp. nov., female holotype **A** left antennule, in dorsal view but tilted little from the medial side **B** right antennule, lateral. **A, B** share the same scale bar.

distal part of inner face; each ramus armed with six well-developed caudal setae, consisting of two distal, two lateral, one inner distal, and one dorsomedial setae.

**Remarks.** The new species is assigned to the genus *Monstrilla* based on the presence of one free postgenital somite and anal somite in the female, six caudal setae, and the oral papilla located ventrally at nearly midlength of the cephalothorax (Isaac 1975; Huys and Boxshall 1991; Suárez-Morales 2011). Among the females of *Monstrilla*, there are two main types of fifth leg, one of which is formed by a single lobe, such as *M. mariaeugeniae* Suarez-Morales & Islas-Landeros, 1993 (Suárez-Morales and Islas-Landeros 1993). The type II fifth leg is bilobed. Six species, *M. annulata* Suárez-Morales, 2024, *M. cymbula* A. Scott, 1909, *M. gibbosa* Suárez-Morales & Palomares-García, 1995, *M. grandis*, *M. grygieri* Suárez-Morales, 2000, and *M. investigatoris* Sewell,

1949, are similar to the new species in having the type II fifth leg and the antennule that exceeds  $\frac{1}{2}$  the length of the cephalothorax (Table 2) (Giesbrecht 1891; Scott 1909; Sewell 1949; Suárez-Morales and Palomares-García 1995; Suárez-Morales 2000b, 2000a, 2024; Chang 2014). Notably, only two species, *M. cymbula* and *M. grandis*, share similar features with *M. pseudograndis* sp. nov., including six well-developed caudal setae on the caudal rami and five setae on the bilobed fifth leg. These three species exhibit the same setation pattern in the fifth leg, a plesiomorphic character state with three exopodal and two endopodal setae, largest number of setae found on the fifth leg in monstrilloids (Huys and Boxshall 1991; Suárez-Morales 2024).

The forehead between the antennular bases exhibits different cuticular dorsal ornamentations in various species. This area may appear concave with a pair of small setae (Lee and Chang 2016), they may form a protrusion (Suárez-Morales et al. 2013, 2017; Suárez-Morales and Castellanos-Osorio 2019; Suárez-Morales 2024), or they present a flat surface lacking both sensilla and setae (Suárez-Morales and Gasca 2003; Suárez-Morales et al. 2020; Suárez-Morales 2021). Notably, setae and sensilla are either present singly or absent in other species. The new species possesses two distinctive features that render it readily distinguishable among its congeners and support its status as a new member of *Monstrilla*. First, *M. pseudograndis* sp. nov. bears a pair of bilateral setae on the forehead and a pair of medial sensilla (Fig. 3B), marking a unique combination that has not been previously reported in *Monstrilla*. Second, the new species has four pairs of small, nipple-like scars on the cephalothorax, flanked in ventral view, which are commonly recorded as 1–3 pairs in *Monstrilla* (Suárez-Morales 2011, 2018; Jeon et al. 2024).

**Table 2.** Comparison of main characters of females in seven similar *Monstrilla* species (elongate cephalothorax and fifth leg with bilobed). Sequence from A to G: *Monstrilla*, *Monstrilla*, *Monstrilla* sp. nov.

Item	<i>M. annulata</i>	<i>M. cymbula</i>	<i>M. gibbosa</i>	<i>M. grandis</i>	<i>M. grygieri</i>	<i>M. investigatoris</i>	<i>M. pseudograndis</i> sp. nov.
Antennule length $>\frac{1}{2}$ cephalothorax	Yes	Yes	–	Yes	Yes	Yes	Yes
Number of segments of antennule	4	2	3	3	2	2	2
Nauplius eye with pigmented	No	–	No	No	No	Yes	No
Nauplius eye elliptical	–	–	Yes	No	Yes	–	Yes
Oral papilla with posteriorly-bent distal half	Yes	Yes	Yes	No	Yes	–	Yes
Oral papilla near midlength of cephalothorax	No	Yes	No	Yes	Yes	–	Yes
Sensory pores on ventral cephalothorax	No	No	No	Yes	No	No	No
Fifth leg bilobed	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Basal protuberance on inner margin of inner lobe	No	No	No	No	No	No	Yes
Distal thumb-like process on inner margin of outer lobe	No	No	No	Yes	No	No	No
Number of setae on fifth leg	2	5	4	5	4	3	5
Number of setae on caudal rami	5	6	5	6	6	6	6



Among all known species of *Monstrilla*, the new species is most closely related to *M. grandis*, which was reported by Giesbrecht (1891) in the southeastern Atlantic Ocean from southern Patagonia. As a widespread species, *M. grandis* has been extensively redescribed by researchers worldwide (Huys and Boxshall 1991; Suárez-Morales 2000b; Chang 2014). *Monstrilla grandis* and the new species share similar body proportions, including total body length, the position of the oral papilla, and the relative lengths of the cephalothorax and ovigerous spines. In addition to the two distinctive features above mentioned, several significant differences also exist: 1) the relative length of the antennule and the number of segments, which is 37.8% and two in *M. pseudograndis* sp. nov., compared to 48% of the total body length and three in *M. grandis*; 2) the ventral sensory pores of cephalothorax are absent in *M. pseudograndis* sp. nov., whereas they are present in *M. grandis*; 3) *M. pseudograndis* sp. nov. bears a protruding ventral oral papilla, with the distal half posteriorly-bent, in contrast to the slightly protruding, unbent midventral papilla in *M. grandis*.; 4) the setae on the first segment of endopod of leg 2 present in *M. pseudograndis* sp. nov., whereas absent in *M. grandis*; 5) a thumb-like process is absent on the distal part of the inner margin of the outer lobe in *M. pseudograndis* sp. nov., and it is present in *M. grandis*; 6) the inner margin of the inner lobe bears a basal protuberance in *M. pseudograndis* sp. nov., which is absent in *M. grandis*.

## Acknowledgements

We thank the two anonymous reviewers for their valuable taxonomic insights. Special gratitude is extended to the editors for their meticulous editorial review and constructive feedback during the revision process. We acknowledge Jun Wang for his skilled assistance in specimen illustration and Haochen Huang for his critical advice on manuscript submission.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Funding

This work was supported by Funding Biological Resources Program, Chinese Academy of Sciences (CAS-TAX-24-042); National Animal Collection Resource Center, China; Tropical Ocean Environment in Western Coastal Waters Observation and Research Station of Guangdong Province (2024B1212040008); the Program for Capacity Building for Strategic Biological Resources, Chinese Academy of Sciences (KFJ-BRP-017-48).

### Author contributions

Conceptualization: ZZ, XL. Funding acquisition: XL. Investigation: ZZ. Methodology: ZZ, XL. Visualization: ZZ. Supervision: YT, XL. Project administration: YT, XL. Writing – original draft: ZZ. Writing – review & editing: ZZ, XL.

## Author ORCIDs

Zhiqian Zhou  <https://orcid.org/0009-0006-8587-586X>

Xiping Lian  <https://orcid.org/0009-0002-3301-1999>

Yehui Tan  <https://orcid.org/0000-0002-6378-2156>

## Data availability

All of the data that support the findings of this study are available in the main text.

## References

- Carneiro-Schaefer AL, Sühnel S, Dias CDO, Melo CMRD, Magalhães ARM (2017) Ocorrência de *Monstrilla* sp. em Perna perna cultivado no Brasil. Boletim do Instituto de Pesca 43: 437–445. <https://doi.org/10.20950/1678-2305.2017v43n3p437>
- Chang CY (2014) Two new records of monstrilloid copepods (Crustacea) from Korea. Animal Systematics, Evolution and Diversity 30: 206–214. <https://doi.org/10.5635/ASED.2014.30.3.206>
- Chen QC, Huang ZG (2012) Family Monstrillidae Dana, 1849. In: Huang ZG, Lin M (Eds) The Living Species in China's Seas. China Ocean Press, Beijing, 722 pp.
- Chen QC, Li KZ (2008) Family Monstrillidae Dana, 1849. In: Liu RY (Ed.) Checklist of Marine Biota of Chinese Seas. Science Press, Beijing, 632 pp.
- Da Cruz Lopes da Rosa J, Dias CDO, Suárez-Morales E, Weber LI, Fischer LG (2021) Record of *Caromiobenella* (Copepoda, Monstrilloida) in Brazil and Discovery of the male of *C. brasiliensis*: morphological and molecular evidence. Diversity 13: 241. <https://doi.org/10.3390/d13060241>
- Delaforge A, Suárez-Morales E, Walkusz W, Campbell K, Mundy CJ (2017) A new species of *Monstrillopsis* (Crustacea, Copepoda, Monstrilloida) from the lower Northwest Passage of the Canadian Arctic. ZooKeys 709: 1–16. <https://doi.org/10.3897/zookeys.709.20181>
- Giesbrecht W (1891) Elenco dei copepodi: pescati dalla R. Corvetta "Vettor Pisani", secondo la loro distribuzione geografica. Atti della Accademia Nazionale dei Lincei, Classe di Scienze Fisiche Matematiche e Naturali Rendiconti (4)7 sem. 2: 63–68, 276–282.
- Grygier MJ, Ohtsuka S (1995) Sem observation of the nauplius of *Monstrilla hamatapex*, new species, from Japan and an example of upgraded descriptive standards for monstrilloid copepods. Journal of Crustacean Biology 15: 703. <https://doi.org/10.2307/1548820>
- Grygier MJ, Ohtsuka S (2008) A new genus of monstrilloid copepods (Crustacea) with anteriorly pointing ovigerous spines and related adaptations for subthoracic brooding. Zoological Journal of the Linnean Society 152: 459–506. <https://doi.org/10.1111/j.1096-3642.2007.00381.x>
- Huys R, Boxshall GA (1991) Copepod Evolution. The Ray Society, London, 468 pp.
- Huys R, Llewellyn-Hughes J, Conroy-Dalton S, Olson PD, Spinks JN, Johnston DA (2007) Extraordinary host switching in siphonostomatoid copepods and the demise of the Monstrilloida: integrating molecular data, ontogeny and antennular morphology. Molecular Phylogenetics and Evolution 43: 368–378. <https://doi.org/10.1016/j.ympev.2007.02.004>
- Isaac MJ (1975) Copepoda, sub-order: Monstrilloida. Fiches d'Identification du Zooplancton 144/145: 1–10.

- Jeon D, Lee W, Soh HY (2018) A new genus and two new species of monstrilloid copepods (Copepoda: Monstrillidae): integrating morphological, molecular phylogenetic, and ecological evidence. *Journal of Crustacean Biology* 38: 45–65. <https://doi.org/10.1093/jcbiol/rux095>
- Jeon D, Song C, Jeong HG, Ohtsuka S, Lee W, Soh HY, Eyun S (2024) An integrated phylogenomic approach for potential host-associated evolution of monstrilloid copepods. In: Todd PA, Russell BD (Eds) *Oceanography and Marine Biology, an Annual Review*, Volume 62. Taylor & Francis, 350–375. <https://doi.org/10.1201/9781003477518-9>
- Lee J, Chang CY (2016) A new species of *Monstrilla* Dana, 1849 (Copepoda: Monstrilloida: Monstrillidae) from Korea, including a key to species from the north-west Pacific. *Zootaxa* 4174: 396–409. <https://doi.org/10.11646/zootaxa.4174.1.24>
- Lee J, Kim D, Chang CY (2016) Two new species of the genus *Monstrillopsis* Sars, 1921 (Copepoda: Monstrilloida: Monstrillidae) from South Korea. *Zootaxa* 4174: 410–423. <https://doi.org/10.11646/zootaxa.4174.1.25>
- Lian X, Tan Y (2019) A new species of *Cymbasoma* Thompson, 1888 (Copepoda: Monstrilloida) from the Fujian coast, China. *Journal of Oceanology and Limnology* 37: 1708–1713. <https://doi.org/10.1007/s00343-019-8219-5>
- Sale PF, McWilliam PS, Anderson DT (1976) Composition of the near-reef zooplankton at Heron Reef, Great Barrier Reef. *Marine Biology* 34: 59–66. <https://doi.org/10.1007/BF00390788>
- Scott A (1909) The Copepoda of the Siboga Expedition, Part I. Free swimming, littoral and semi-parasitic Copepoda. *Siboga Expeditie* 29a: 1–323. <https://doi.org/10.5962/bhl.title.58653>
- Sewell RBS (1949) The littoral and semi-parasitic Cyclopoida, Monstrilloida and Notodelphyoida. *John Murray Expedition 1933–34 Scientific Reports* 9: 17–199.
- Suárez-Morales E (2000a) A new species and new geographic records of *Monstrilla* (Copepoda: Monstrilloida) from the Philippines. *Journal of Crustacean Biology* 20: 680–686. <https://doi.org/10.1163/20021975-99990092>
- Suárez-Morales E (2000b) Taxonomic report on some monstrilloids (Copepoda, Monstrilloida) from Toulon Bay, France. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Biologie* 70: 107–118.
- Suárez-Morales E (2011) Diversity of the Monstrilloida (Crustacea: Copepoda). *PLoS ONE* 6: e22915. <https://doi.org/10.1371/journal.pone.0022915>
- Suárez-Morales E (2018) Monstrilloid copepods: the best of three worlds. *Bulletin, Southern California Academy of Sciences* 117: 92–103. <https://doi.org/10.3160/3646.1>
- Suárez-Morales E (2019) A new genus of the Monstrilloida (Copepoda) with large rostral process and metasomal spines, and redescription of *Monstrilla spinosa* Park, 1967. *Crustaceana* 92: 1099–1112. <https://doi.org/10.1163/15685403-00003925>
- Suárez-Morales E (2021) A new species of *Monstrilla* (Crustacea: Copepoda: Monstrilloida: Monstrillidae) collected by the R/V albatross (1908) from the Sulu Sea, Philippines. *Journal of Natural History* 55: 3041–3052. <https://doi.org/10.1080/00222933.2022.2032444>
- Suárez-Morales E (2024) Two new species of *Monstrilla* (Copepoda: Monstrilloida: Monstrillidae) from a protected reef system of the Mexican Caribbean. *European Journal of Taxonomy* 917: 152–169. <https://doi.org/10.5852/ejt.2024.917.2395>
- Suárez-Morales E, Castellanos-Osorio IA (2019) A new species of *Monstrilla* (Copepoda, Monstrilloida) from the plankton of a large coastal system of the northwestern Caribbean with a key to species. *ZooKeys* 876: 111–123. <https://doi.org/10.3897/zookeys.876.38400>

- Suárez-Morales E, Gasca R (2003) On the invalidity of *Strilloma* Isaac (Copepoda: Monstrilloida): observations from the type species. *Zoological Studies* 43: 292–299.
- Suárez-Morales E, Islas-Landeros ME (1993) A new species of *Monstrilla* (Copepoda: Monstrilloida) from a reef lagoon off the Mexican coast of the Caribbean Sea. *Hydrobiologia* 271: 45–48. <https://doi.org/10.1007/BF00005694>
- Suárez-Morales E, McKinnon AD (2014) The Australian Monstrilloida (Crustacea: Copepoda) I. *Monstrillopsis* Sars, *Maemonstrilla* Grygier & Ohtsuka, and *Australomonstrillopsis* gen. nov. *Zootaxa* 3779: 301–340. <https://doi.org/10.11646/zootaxa.3779.3.1>
- Suárez-Morales E, Palomares-García R (1995) A new species of *Monstrilla* (Copepoda: Monstrilloida) from a coastal system of the Baja California Peninsula, Mexico. *Journal of Plankton Research* 17: 745–752. <https://doi.org/10.1093/plankt/17.4.745>
- Suárez-Morales E, Velazquez-Ornelas K (2024) A new species of the genus *Monstrilla* (Copepoda: Monstrilloida: Monstrillidae) from the Gulf of California, Mexico. *Revista Mexicana de Biodiversidad* 95: e955361. <https://doi.org/10.22201/ib.20078706e.2024.95.5361>
- Suárez-Morales E, Scardua MP, Da Silva PM (2010) Occurrence and histopathological effects of *Monstrilla* sp. (Copepoda: Monstrilloida) and other parasites in the brown mussel *Perna perna* from Brazil. *Journal of the Marine Biological Association of the United Kingdom* 90: 953–958. <https://doi.org/10.1017/S0025315409991391>
- Suárez-Morales E, Carrillo A, Morales-Ramírez A (2013) Report on some monstrilloids (Crustacea: Copepoda) from a reef area off the Caribbean coast of Costa Rica, Central America with description of two new species. *Journal of Natural History* 47: 619–638. <https://doi.org/10.1080/00222933.2012.742933>
- Suárez-Morales E, Harris LH, Ferrari FD, Gasca R (2014) Late postnaupliar development of *Monstrilla* sp. (Copepoda: Monstrilloida), a protelean endoparasite of benthic polychaetes. *Invertebrate Reproduction & Development* 58: 60–73. <https://doi.org/10.1080/07924259.2013.816787>
- Suárez-Morales E, Goruppi A, De Olazabal A, Tirelli V (2017) Monstrilloids (Crustacea: Copepoda) from the Mediterranean Sea (northern Adriatic Sea), with a description of six new species. *Journal of Natural History* 51: 1795–1834. <https://doi.org/10.1080/00222933.2017.1359698>
- Suárez-Morales E, Vásquez-Yeomans L, Santoya L (2020) A new species of the *Cymbasoma longispinosum* species-group (Copepoda: Monstrilloida) from Belize, western Caribbean. *Revista Mexicana de Biodiversidad* 91: 913176. <https://doi.org/10.22201/ib.20078706e.2020.91.3176>
- Walter TC, Boxshall G (2024) World of Copepods Database. *Monstrilla* Dana, 1849–1852. <https://www.marinespecies.org/aphia.php?p=taxdetails&id=119780> [Accessed on 2024-10-29]