

The first two complete mitochondrial genomes for the genus *Anagyrus* (Hymenoptera, Encyrtidae) and their phylogenetic implications

Cheng-Hui Zhang¹, Hai-Yang Wang¹, Yan Wang¹, Zhi-Hao Chi¹, Yue-Shuo Liu¹, Guo-Hao Zu¹

¹ College of Horticulture and Landscape, Tianjin Agricultural University, Tianjin 300392, China

Corresponding author: Guo-Hao Zu (zuguohao@tjau.edu.cn)

Abstract

Anagyrus, a genus of Encyrtidae (Hymenoptera, Chalcidoidea), represents a successful group of parasitoid insects that attack various mealybug pests of agricultural and forestry plants. Until now, only 20 complete mitochondrial genomes have been sequenced, including those in this study. To enrich the diversity of mitochondrial genomes in Encyrtidae and to gain insights into their phylogenetic relationships, the mitochondrial genomes of two species of *Anagyrus* were sequenced, and the mitochondrial genomes of these species were compared and analyzed. Encyrtid mitochondrial genomes exhibit similarities in nucleotide composition, gene organization, and control region patterns. Comparative analysis of protein-coding genes revealed varying molecular evolutionary rates among different genes, with six genes (*ATP8*, *ND2*, *ND4L*, *ND6*, *ND4* and *ND5*) showing higher rates than others. A phylogenetic analysis based on mitochondrial genome sequences supports the monophyly of Encyrtidae; however, the two subfamilies, Encyrtinae and Tetracneminae, are non-monophyletic. This study provides valuable insights into the phylogenetic relationships within the Encyrtidae and underscores the utility of mitochondrial genomes in the systematics of this family.

Key words: Encyrtid, genome structure, mitogenome, protein-coding genes, phylogenetic analyses, Tetracneminae



Academic editor: Zachary Lahey

Received: 29 February 2024

Accepted: 12 June 2024

Published: 5 July 2024

ZooBank: <https://zoobank.org/C6737E11-5992-4B82-B0C1-48C6C32DF550>

Citation: Zhang C-H, Wang H-Y, Wang Y, Chi Z-H, Liu Y-S, Zu G-H (2024) The first two complete mitochondrial genomes for the genus *Anagyrus* (Hymenoptera, Encyrtidae) and their phylogenetic implications. ZooKeys 1206: 81–98. <https://doi.org/10.3897/zookeys.1206.121923>

Copyright: © Cheng-Hui Zhang et al.
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

Encyrtidae is a large hymenopteran family in the superfamily Chalcidoidea, comprising 518 known genera, of which 495 are recognized as valid (totaling more than 4830 species), along with 23 fossil genera (26 species) worldwide (Simutnik et al. 2022; Simutnik et al. 2023; Simutnik and Perkovsky 2023; Wang et al. 2023). The genus *Anagyrus* Howard, 1986 is one of the largest genera in Encyrtidae, comprising 289 valid species (Noyes 2019). This genus was established by Howard and Ashmead (1896) based on the type species, *Anagyrus greeni* Howard, 1896. Diagnostics for the genus include a broadened, flattened scape (normally 2–3× as long as broad), funicle segments longer than broad, occipital margin normally quite sharp but often rounded, postmarginal vein normally not longer than the stigma vein, and ovipositor at least half the length of

the mid tibia (Noyes 1980; Noyes and Hayat 1994). *Anagyrus* species are primary parasitoids of Pseudococcidae; for example, *Anagyrus galinae* has been utilized in classical biocontrol and integrated pest management of *Trionymus copiosus* (Japoshvili and Hansen 2015; Noyes 2019).

Insect mitochondrial genomes are usually small, circular molecules containing 37 genes: 13 protein-coding genes (PCGs), two ribosomal RNA genes (rRNAs), and 22 transfer RNA genes (tRNAs), as well as a large non-coding element known as the A+T-rich or control region (CR), which regulates transcription and replication (Wolstenholme 1992a, 1992b; Boore 1999; Cameron 2014). Due to their distinct characteristics, including gene-content conservation, maternal inheritance, and rapid evolutionary rate, mitogenome sequences serve as valuable molecular markers for various evolutionary studies (Boore 1999; Krzywinski et al. 2006). Although the mitochondrial genome of Chalcidoidea exhibits structural resemblance to other insects, significant rearrangements characterize it, along with a relatively high A+T content in its sequence composition, deviating from the presumed ancestral pattern (Brown et al. 1979; Cameron and Whiting 2008).

The exploration of hymenopteran mitochondrial genomes commenced with the sequencing of *CYTB* and *ATP8* genes of *Apis mellifera*, and it was not until 1993 that the first complete mitochondrial genome was deciphered (Crozier and Crozier 1992, 1993). The first comprehensive phylogenetic analysis of Chalcidoidea based on molecular data was conducted using 18S and 28S rDNA (Munro et al. 2011). Subsequently, Heraty et al. (2013) conducted an in-depth exploration of the phylogenetic relationships within Chalcidoidea based on both morphological and molecular data. Zhang et al. (2020a) further reconstructed the phylogenetic relationships within Chalcidoidea using transcriptome data, providing valuable insights for achieving more accurate phylogenetic relationships. Recently, Cruaud et al. (2024) conducted a comprehensive phylogenetic study using data from PCGs and ultra-conserved elements (UCEs), while Zhu et al. (2023) conducted a comprehensive phylogenetic study using 139 mitochondrial genomes from the main clades of Chalcidoidea. These studies have significantly advanced our understanding of the phylogenetic relationships within Chalcidoidea. However, to obtain a more accurate reconstruction of evolutionary relationships, it is necessary to expand the sampling range to include more understudied species. This approach will help construct a more comprehensive and precise phylogenetic tree, revealing deeper levels of phylogenetic relationships. Additionally, integrating different types of data, such as rDNA genes, mitochondrial genomes, and UCEs, is crucial. By comprehensively utilizing morphological, biological, and molecular data and conducting multidimensional phylogenetic analyses, we can improve the accuracy of classification and phylogenetic research. Such integrative approaches will provide a more robust framework for understanding the evolutionary relationships within Hymenoptera.

At present, there are only morphology-based classification systems for Encyrtidae (Noyes and Hayat 1984, 1994; Trjapitzin 1989), lacking auxiliary verification from molecular data, particularly from the mitochondrial genome (mitogenome). Consequently, the monophyly and phylogenetic relationships of Encyrtidae have been controversial for a long time. Problems that are difficult to distinguish in taxonomy indicate the requirement for using various molecular data to understand the systematic position and the monophyly of Encyrtidae in

Chalcidoidea. Mitogenome data seem sufficient to solve these problems (Wei et al. 2010; Li et al. 2016; Liu et al. 2023). There are currently only 1291 complete mitochondrial genomes of Hymenoptera on GenBank, and the number of encyrtid genomes is small (Sayers et al. 2024). This limited data negatively impacts our ability to resolve potential systematic ambiguity within Encyrtidae.

In this study, we conducted the sequencing and annotation of the mitogenomes of *Anagyrs galinae* (accession number: [OR652687](#)) and *Anagyrs jenniferae* (accession number: [OR790122](#)), analyzing their respective characteristics. In addition, we reconstructed the molecular phylogenetic relationships of these two new mitochondrial genomes and other species of Encyrtidae. The molecular data presented in this study will contribute to a better understanding of the characteristics of the Encyrtidae mitogenome. Further, a phylogenetic analysis was performed, including 19 uploaded mitogenomes together with our newly acquired data, which represented Encyrtidae. The goal of our study was to place two new species of *Anagyrs* within the context of the known mitogenome diversity of Encyrtidae by performing mitogenomic and phylogenetic analyses.

Materials and methods

Sample collection, DNA extraction and sequencing

The specimens, *A. galinae* and *A. jenniferae*, were collected from Tianjin Agricultural University (39°5'21"N, 117°5'38"E), Xiqing District, Tianjin City, China, in September 2022. Freshly collected specimens were promptly immersed in 100% ethanol for initial preservation and subsequently stored at -40 °C in the Insect Herbarium of Tianjin Agricultural University. Following morphological identification, total DNA from each specimen was extracted from the body, excluding the abdomen, using the DNeasy Blood & Tissue Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. The purity and concentration of the extracted total DNA were assessed through 1% agarose gel electrophoresis and optical density value detection. The total DNA of two encyrtids underwent sequencing using the Illumina NovaSeq 6000 platform with a 350 bp insert size and a paired-end 150 bp sequencing strategy. Sequencing was conducted by Novogene Co., Ltd. (Beijing, China).

Mitogenome assembly, annotation and analysis

After initial data acquisition, with adapter sequences removed, additional filtering was carried out using fastp 0.23.4 (Chen et al. 2018) to filter low-quality reads (quality value <30), ensuring that each sample retained clean data of no less than 4 Gb. The software MitoZ v. 3.6 (Meng et al. 2019) and GetOrganelle v. 1.7.7.0 (Jin et al. 2020) were used for the de novo assembly of mitogenomes. Homologous sequences of other Encyrtidae species from GenBank were used for comparison, and the mitogenomes were annotated using the Mitos WebServer (Donath et al. 2019). The secondary structures of tRNAs were predicted using Mitos WebServer and further visualized using VARNA v. 3.9 (Darty et al. 2009). The structures of the mitochondrial genome were mapped using the online tool CGview Server. The nucleotide composition and relative synonymous codon usage (RSCU) of protein-coding genes were calculated and analyzed by

MEGA v. 11.0.13 (Tamura et al. 2021). The skew analysis of nucleotide composition was calculated using the formulas: AT-skew = $(A-T)/(A+T)$ and GC-skew = $(G-C)/(G+C)$, where A, T, G and C were the base contents of the same chain (Perna and Kocher 1995; Hassanin et al. 2005). The nonsynonymous mutation rate (Ka) and synonymous mutation rate (Ks) of protein coding genes were calculated using DnaSP 6.12.03 (Rozas et al. 2017). Tandem repeats in the CR were identified by Tandem Repeats Finder (Benson 1999).

Molecular phylogenetic analyses

A total of 21 mitogenomes from two families of Chalcidoidea, including 20 Encyrtidae species and a Aphelinidae species as outgroup, were used for the phylogenetic analysis (Table 1). The phylogenetic trees were reconstructed using both maximum-likelihood (ML) and Bayesian-inference (BI) methods. For this, each PCG was individually aligned using the MAFFT 7 online service with the L-INS-i strategy, followed by optimization using MACSE (Ranwez et al. 2018; Katoh et al. 2019). The individual PCG alignments were trimmed using GBlocks and concatenated into a PCG dataset using PhyloSuite v. 1.2.3 (Talavera and Castresana 2007; Zhang et al. 2020b). The best nucleotide substitution model was obtained using ModelFinder v. 2.2.0 with Bayesian Information Criterion (BIC) (Kalyanamorthy et al. 2017). BI analysis was performed using MrBayes v. 3.2.7a with four chains (Ronquist et al. 2012). Two independent runs of

Table 1. GenBank accession numbers of species used in phylogenetic reconstruction and their original publications.

Superfamily	Family	Species	Accession Number	References
Chalcidoidea	Aphelinidae	<i>Encarsia formosa</i>	MG813797	Zhu et al. 2018
	Encyrtidae	<i>Aenasius arizonensis</i>	NC_045852	Ma et al. 2019
		<i>Anagyrus galinae</i>	OR652687	This study
		<i>Anagyrus jenniferae</i>	OR790122	This study
		<i>Blastothrix speciosa</i>	NC_082111	Unpublished
		<i>Cheiloneurus chinensis</i>	NC_084192	Unpublished
		<i>Cheiloneurus elegans</i>	NC_071192	Unpublished
		<i>Diaphorencyrtus aligarhensis</i>	NC_046058	Du et al. 2019
		<i>Encyrtus aurantii</i>	OR120384	Unpublished
		<i>Encyrtus eulecaniumiae</i>	NC_051459	Rudoy et al. 2022
		<i>Encyrtus infelix</i>	NC_041176	Xiong et al. 2019
		<i>Encyrtus rhodococcusiae</i>	NC_051460	Rudoy et al. 2022
		<i>Encyrtus sasakii</i>	NC_051458	Rudoy et al. 2022
		<i>Exoristobia philippinensis</i>	NC_084171	Unpublished
		<i>Lamennaisia ambigua</i>	NC_082113	Unpublished
		<i>Lamennaisia nobilis</i>	NC_061411	Unpublished
		<i>Leptomastidea bifasciata</i>	OR790123	Unpublished
		<i>Metaphycus eriococci</i>	NC_056349	Zhou et al. 2021
		<i>Ooencyrtus plautus</i>	NC_068223	Xing et al. 2022
		<i>Psyllaephagus</i> sp.	OP787025	Unpublished
		<i>Tassonia gloriae</i>	NC_082112	Unpublished

2,000,000 generations were carried out with sampling every 1,000 generations. The first 25% of trees were discarded as burn-in. After the average standard deviation of split frequencies fell below 0.01 and the potential scale reduction factor (PSRF) approached 1.0, stationarity was assumed. ML analysis was performed using IQ-TREE v. 2.2.0 (Nguyen et al. 2015) under the standard bootstrap approximation approach with 1,000 replicates.

Results

Mitogenome organization and nucleotide composition

The assembled mitochondrial genome of *A. galinae* was a 15,364 bp, and the *A. jenniferae* mitochondrial genome was 15,396 bp, which both had the same gene organization, including 13 PCGs, 22 tRNAs, two rRNAs and a control region located between *trnM* and *trnI* (Fig. 1). For the mitogenomes of two species, the majority strand (J-strand) encodes 10 PCGs (*ND3*, *CO3*, *ATP6*, *ATP8*, *CO2*, *CO1*, *ND5*, *ND4*, *ND4L*, *ND1*), 15 tRNAs (*trnI*, *trnY*, *trnS1*, *trnC*, *trnR*, *trnG*, *trnD*, *trnL2*, *trnF*, *trnH*, *trnP*, *trnL1*, *trnA*, *trnV*, *trnM*) and 2 rRNAs (*IrRNA*, *srRNA*), while the remaining three PCGs (*ND2*, *ND6*, *CYTB*) and seven tRNAs (*trnW*, *trnN*, *trnK*, *trnE*, *trnT*, *trnS2*, *trnQ*) are located on the minority strand (Table 2). Two mitogenomes both obtained 13 overlapping nucleotides, and up to 53 bp ranging from 1 to 16 bp. The longest overlap was located between *CO1* and *trnE* in *A. jenniferae*. There were 17 and 16 intergenic spacers each from *A. galinae* and *A. jenniferae*, totaling 171 bp and 115 bp, ranging 1 to 77 bp and 1 to 27 bp, respectively.

The nucleotide composition of the mitogenome from *A. galinae* was biased toward A and T, with 83.12% of A+T content (A = 45.12%, T = 38.00%, C = 10.82%, G = 6.05%), A+T content was 82.94%, 87.20% in PCGs and




Figure 1. Circular map of the mitochondrial genome of *Anagyrus galinae* and *Anagyrus jenniferae*.

Table 2. Gene organization of the mitochondrial genomes of *Anagyryrus galinae* and *Anagyryrus jenniferae*.

Gene	Direction		Anticodon	<i>Anagyryrus galinae</i>				<i>Anagyryrus jenniferae</i>				
				Position	Length	Start codon	Stop codon	Intergenic Nucleotides	Position	Length	Start codon	Stop codon
<i>trnI</i>	-	GAU	1-70	70				1-67	67			
<i>ND2</i>	+		98-1087	990	ATT	TAA	27	74-1081	1008	ATT	TAA	6
<i>trnW</i>	+	UCA	1087-1149	63			-1	1080-1146	67			-2
<i>trnY</i>	-	GUA	1155-1221	67			5	1148-1212	65			1
<i>trnS1</i>	-	UCU	1222-1280	59			0	1216-1275	60			3
<i>trnC</i>	-	GCA	1283-1348	66			2	1293-1361	69			17
<i>trnN</i>	+	GUU	1369-1434	66			20	1368-1431	64			6
<i>trnR</i>	-	UCG	1433-1497	65			-2	1439-1504	66			7
<i>ND3</i>	-		1498-1842	345	ATT	TAA	0	1505-1858	354	ATA	TAA	0
<i>trnG</i>	-	UCC	1843-1906	64			0	1856-1919	64			-3
<i>CO3</i>	-		1911-2714	804	ATG	TAA	4	1925-2710	786	ATG	TAA	5
<i>ATP6</i>	-		2715-3387	673	ATG	T	0	2710-3383	674	ATG	TA	-1
<i>ATP8</i>	-		3381-3542	162	ATT	TAA	-7	3377-3538	162	ATC	TAA	-7
<i>trnD</i>	-	GUC	3543-3608	66			0	3539-3602	64			0
<i>trnK</i>	+	UUU	3612-3683	72			3	3606-3676	71			3
<i>CO2</i>	-		3688-4365	678	ATT	TAG	4	3678-4355	678	ATT	TAA	1
<i>trnL2</i>	-	UAA	4369-4434	66			3	4365-4428	64			9
<i>CO1</i>	-		4440-5987	1548	ATT	TAA	5	4431-5969	1539	ATG	TAA	2
<i>trnE</i>	+	UUC	5972-6036	65			-16	5972-6034	63			2
<i>trnF</i>	-	GAA	6036-6102	67			-1	6034-6099	66			-1
<i>ND5</i>	-		6102-7769	1668	ATA	TAA	-1	6099-7763	1665	ATT	TAG	-1
<i>trnH</i>	-	GUG	7767-7833	67			-3	7764-7829	66			0
<i>ND4</i>	-		7844-9169	1326	ATG	TAG	10	7829-9156	1328	ATG	TA	-1
<i>ND4L</i>	-		9163-9450	288	ATT	TAA	-7	9150-9437	288	ATT	TAA	-7
<i>trnT</i>	+	UGU	9453-9518	66			2	9440-9505	66			2
<i>trnP</i>	-	UGG	9520-9582	63			1	9506-9574	69			0
<i>ND6</i>	+		9584-10151	568	ATG	T	1	9575-10143	569	ATG	TA	0
<i>CYTB</i>	+		10152-11300	1149	ATG	TAA	0	10143-11285	1143	ATG	TAA	-1
<i>trnS2</i>	+	UGA	11300-11365	66			-1	11290-11354	65			4
<i>ND1</i>	-		11356-12291	936	ATT	TAG	-10	11345-12283	939	ATA	TAG	-10
<i>trnL1</i>	-	UAG	12292-12358	67			0	12284-12348	65			0
<i>lrRNA</i>	-		12364-13674	1311			5	12353-13654	1302			4
<i>trnA</i>	-	UGC	13682-13744	63			7	13651-13719	69			-4
<i>trnQ</i>	+	UUG	13761-13831	71			16	13797-13864	68			77
<i>srRNA</i>	-		13831-14602	772			-1	13891-14646	756			26
<i>trnV</i>	-	UAC	14602-14669	68			-1	14646-14710	65			-1
<i>trnM</i>	-	CAU	14668-14735	68			-2	14709-14770	62			-2
CR			14736-15364	629			0	14771-15396	626			0

rRNAs, respectively. The nucleotide composition of the mitogenome from *A. jenniferae* was biased toward A and T, with 82.64% of A+T content (A = 46.41%, T = 36.23%, C = 11.33%, G = 6.02%), A+T content was 82.32%, 85.20% in PCGs and rRNAs, respectively. The values of AT-skew and GC-

Table 3. Nucleotide features of the mitochondrial genome across *Anagyryus galinae* and *Anagyryus jenniferae*.

Feature	Length (bp)	T%	C%	A%	G%	A+T%	AT-Skew	GC-Skew
Whole genome	15364/15396	38.00/36.23	10.82/11.33	45.12/46.41	6.05/6.02	83.12/82.64	0.086/0.123	-0.283/-0.306
<i>ATP6</i>	673/674	46.66/47.63	7.43/8.01	34.92/34.27	11.00/10.09	81.58/81.90	-0.144/-0.163	0.194/0.115
<i>ATP8</i>	162/162	48.77/48.77	4.32/4.94	43.83/36.42	3.09/9.88	92.59/85.19	-0.053/-0.145	-0.167/0.333
<i>CO1</i>	1524/1539	45.41/46.39	10.37/10.98	29.86/27.23	14.37/15.40	75.26/73.62	-0.207/-0.260	0.162/0.167
<i>CO2</i>	678/678	45.58/45.72	8.55/8.41	33.04/33.19	12.83/12.68	78.61/78.91	-0.159/-0.159	0.200/0.203
<i>CO3</i>	804/786	46.64/49.75	7.84/8.52	32.21/29.90	13.31/11.83	78.86/79.64	-0.183/-0.249	0.259/0.163
<i>CYTB</i>	1149/1143	43.69/41.91	14.36/14.7	32.64/34.82	9.31/8.57	76.33/76.73	-0.145/-0.092	-0.213/-0.263
<i>ND1</i>	936/939	46.47/48.35	7.05/6.71	32.37/31.31	14.1/13.63	78.85/79.66	-0.179/-0.214	0.333/0.340
<i>ND2</i>	990/1008	50.10/47.52	9.19/9.62	37.58/39.19	14.10/13.63	87.68/86.71	-0.143/-0.096	-0.492/-0.448
<i>ND3</i>	345/351	51.01/52.99	5.22/5.41	33.91/31.34	9.86/10.26	84.93/84.33	-0.201/-0.257	0.308/0.309
<i>ND4</i>	1326/1328	50.08/52.41	4.98/5.20	34.01/30.20	10.94/12.20	84.09/82.61	-0.191/-0.269	0.374/0.403
<i>ND4L</i>	288/288	53.13/53.82	2.78/2.08	34.03/36.46	10.07/7.64	87.15/90.28	-0.219/-0.192	0.568/0.571
<i>ND5</i>	1665/1665	50.81/51.11	5.77/5.77	33.09/32.61	10.33/10.51	83.90/83.72	-0.211/-0.221	0.284/0.292
<i>ND6</i>	568/569	46.13/45.34	8.45/10.54	42.25/41.48	3.17/2.64	88.38/86.82	-0.044/-0.045	-0.455/-0.600
<i>srRNA</i>	772/756	44.30/44.84	4.15/4.10	43.52/40.08	8.03/10.98	87.82/84.92	-0.009/-0.056	0.319/0.456
<i>lrRNA</i>	1311/1302	44.55/46.08	4.27/4.15	42.03/39.40	9.15/10.37	86.58/85.48	-0.029/-0.078	0.364/0.429
CR	629/626	42.61/40.57	7.15/8.47	46.26/48.72	3.98/2.24	88.87/89.29	0.041/0.091	-0.285/-0.582

skew were often used to indicate the nucleotide composition of the mitochondrial genome. In this study, the nucleotide features of two new mitogenomes were investigated by calculating the percentages of AT-skew and GC-skew (Table 3). The skew analysis showing the mitogenome of *A. galinae* had a positive AT-skew (0.086) and a negative GC-skew (-0.283), and the mitogenome of *A. jenniferae* had a positive AT-skew (0.123) and a negative GC-skew (-0.306).

Protein-coding genes and codon usage

By comparing the known mitochondrial genome structure of Encyrtidae, we found that the sequence of 13 PCGs was consistent, except for *ND3* rearranged in *Diaphorencyrtus aligarhensis* and *Leptomastidea bifasciata*. The sequence of PCGs in these mitochondrial genomes were the same (Fig. 2). Additionally, this arrangement is consistent with the mitochondrial gene order in other Encyrtidae, which is also consistent with inferred ancestry.

The total lengths of 13 PCGs are 11,108 bp in *A. galinae*, 11,130 bp in *A. jenniferae*. In these mitochondrial genomes, the length of each PCG ranges from 162 bp (*ATP8*) to 1665 bp (*ND5*). Two mitogenomes of *Anagyryus* exhibited similar start and stop codons. All the initiation codons of PCGs were ATN (ATA, ATG and ATT). Three kinds of stop codons existed on the new mitogenomic sequences: TAA, TAG and truncated termination codons (TA existed on *ATP6*, *ND4*, *ND6* in *A. jenniferae*, T existed on *ATP6*, *ND6* in *A. galinae*), TAA were the most frequently used. Truncated termination codons are commonly used in metazoan mitogenomes, which could be completed by post-transcriptional poly-adenylation (Ojala et al. 1981).

<i>Aenasius arizonensis</i>	J	ND2	W	N	Y	SI	C	R	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	F	E	ND5	H	ND4	ND4L	P	T	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M		
<i>Anagyrus galinae</i>	J	ND2	W	I	SI	C	N	R	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Anagyrus jenniferae</i>	J	ND2	W	I	SI	C	N	R	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Blastothrix speciosa</i>	ND2	W	SI	N	R	I	Y	C	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Cheiloneurus elegans</i>	F	M	CR	Y	I	ND2	G	R	N	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR
<i>Cheiloneurus chinensis</i>	CR	J	ND2	R	Q	ND3	SI	N	Y	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Diaphorencyrtus aligarhensis</i>	ND2	W	SI	Y	C	Q	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	J			
<i>Encyrtus aurantii</i>	N	SI	Y	C	W	I	ND2	R	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Encyrtus eulecaniumiae</i>	N	SI	Y	C	W	I	ND2	R	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Encyrtus infelix</i>	N	SI	Y	C	W	I	ND2	R	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Encyrtus rhodococcidae</i>	N	SI	Y	C	W	I	ND2	R	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Encyrtus sasakii</i>	N	SI	Y	C	W	I	ND2	R	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Exoristobia philippinensis</i>	J	ND2	C	N	R	SI	Q	Y	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Lamennaisia ambigua</i>	ND2	I	Y	R	SI	ND3	C	Q	G	N	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Lamennaisia nobilis</i>	Q	SI	R	C	W	ND2	J	Y	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Leptomastidea bifasciata</i>	CR	J	ND2	W	Y	C	Q	SI	R	A	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Metaphycus eriococi</i>	J	ND2	W	Y	SI	N	C	R	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Ooencyrtus plautus</i>	ND2	R	Q	N	W	C	SI	Y	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Psyllapbagus</i> sp.	J	ND2	W	N	SI	Y	R	C	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	
<i>Tassonia glorieae</i>	R	N	I	SI	W	ND2	Q	C	ND3	G	CO3	ATP6	ATP8	D	K	CO2	L2	COI	E	F	ND5	H	ND4	ND4L	T	P	ND6	CYTB	S2	ND1	LI	rRNA	A	Q	rRNA	V	M	CR	

Figure 2. Gene order of mitochondrial genomes of different Encyrtidae species.




Figure 3. Relative synonymous codon usage in mitochondrial genomes of *Anagyrus galinae* and *Anagyrus jenniferae*.

The codon UUA (Leu2) was the most commonly used in both mitogenomes. Mitochondrial protein coding genes have obvious bias towards A and T, and for mitochondrial protein-coding gene of *A. galinae* the three most frequently used codons were UUA (Leu2) 469 times, AUU (Ile) 440 times and UUU (Phe) 432 times. For *A. jenniferae*, the three most used codons were UUA (Leu2) 463 times, UUU (Phe) 431 times and AUU (Ile) 393 times. Mitochondrial protein-coding genes in Encyrtidae prefer A and U in the third codon, which is like some hymenopteran insects (Fan et al. 2017; Peng et al. 2017). The RSCU values of *A. galinae* and *A. jenniferae* are shown in Fig. 3.




Figure 4. Evolutionary rates of protein-coding genes in the mitochondrial genomes of Encyrtidae.

In this study, based on 20 mitochondrial genomes of Encyrtidae, DnaSP was used to calculate the non-synonymous substitution rate, synonymous substitution, and Ka/Ks ratio of 13 PCGs in the mitochondrial genome and then to compare the evolution rate between genes (Fig. 4). The results showed that among the 13 protein-coding genes in the mitochondrial genome of encyrtids, *CYTB* had the highest Ks, whereas *ATP8* had the highest Ka and Ka/Ks value, and *ATP8* had the largest variation and *COI* had the slowest evolution rate. The evolution rate of 13 genes was in the order of *ATP8* > *ND2* > *ND4L* > *ND6* > *ND4* > *ND5* > *ND3* > *ATP6* > *ND1* > *CO3* > *CO2* > *CYTB* > *CO1*.

Ka/Ks values of 12 PCGs (all PCGs except *ATP8*) were far lower than 1.0, indicating that they were subject to purifying selection, a phenomenon first discovered in Chalcidoidea. In addition, the Ka/Ks value of *ATP8* is higher than 1.0, higher value of *ATP8* was also found in other species (Ma et al. 2019; Jia et al. 2020; Xu et al. 2021). The reason for this phenomenon may be that the evolution speed of a gene is related to its function (Wang et al. 2011).

Transfer RNA genes, ribosomal RNA genes, and control region

The mitochondrial genomes of the two species both included 22 tRNA genes, and the total lengths of the tRNAs of *A. galinae* and *A. jenniferae* are 1455 bp and 1445 bp, respectively. The length of tRNA genes in two *Anagyrus* species ranged from 59 to 72 bp. The secondary structures of the 22 tRNAs of the two species are shown in Suppl. materials 1, 2. The 22 tRNA genes in the mitochondrial genome are identical with the anticodon of tRNA corresponding to the mitochondrial genome of other Hymenoptera, except that *trnL* and *trnS* have two tRNA structures, and the others only have one corresponding tRNA structure. Most tRNAs could be folded into a typical clover-leaf structure, except for *trnS1* which lost a dihydrouridine (DHU) arm and became a simple loop. A lack of the DHU arm in *trnS1* was found in the mitochondrial genomes of most insects (Downton et al. 2002). Changes in the length of the DHU and TΨC arms led to

differences in the size of the tRNA sequence (Shao et al. 2001). In addition, the anticodon of *trnS1* became UCU instead of the more common GCU. In addition to typical Watson-Crick pairings (A-U and G-C), G-U pairings also exist, which are called atypical pairings or wobble base pairs. A total of 30 mismatched base pairs were found in the arm structures of the tRNAs.

Hymenopteran mitochondria have a high rearrangement rate, which mainly occurs in A+T-rich regions, *ND2*, *ND2-CO2*, *CO2-ATP8*, and *ND3-ND5* regions (Wei 2009). The gene arrangement of the suborder Symphyta was conserved and less rearranged than that of suborder Apocrita. However, there are a large number of rearrangements in the Apocrita, including displacement, inversion in situ, and ectopic inversion (Song 2015; Zhao et al. 2021). The rearrangement of mitochondrial genomes in Encyrtidae species was compared (Fig. 2), and the rearrangement was mainly found in tRNA genes. The rearrangement of tRNA occurred at many sites, and the pattern was complicated. Except that *trnD-trnK* (*trnK-trnD* in *Ooencyrtus plautus*), *trnL2*, *trnE-trnF* (*trnF-trnE* in *Aenasius arizonensis* and *Diaphorencyrtus aligarhensis*), *trnH*, *trnT-trnP* (*trnP-trnT* in *Aenasius arizonensis* and *Lamennaisia ambigua*), *trnS2* and *trnL1* are stable between *ATP8* to *lrRNA*, there was no exclusion, and the other tRNA genes had been rearranged.

As for the rRNAs of two *Anagyrs* species, both *lrRNA* and *srRNA* genes are encoded on the N-strand and have a heavy AT nucleotide bias. The lengths of *lrRNA* and *srRNA* in *A. galinae* are 1311 bp and 772 bp, with the different A+T contents of 86.58% and 87.82%, and in *A. jenniferae* are 1302 bp and 756 bp, with the different A+T contents of 85.48% and 84.92%.

In the mitogenome, the largest non-coding region is normally the A+T-rich region, also known as the control region, which regulates the replication and transcription of mitochondrial DNA (Boore 1999; Cameron 2014). In the mitogenomes of the two *Anagyrs* species sequenced in this study, the CR is located between *trnM* and *trnI* (Fig. 5). The length of the CR is 629 bp in *A. galinae* and 626 bp in *A. jenniferae*. The A+T content is 88.87% and 89.29% in the CR of *A. galinae* and *A. jenniferae*. Analysis of AT-skew and CG-skew indicates that both *Anagyrs* species exhibit A and C usage bias. Three structural elements were found in each CR of two *Anagyrs* species: (1) a leading sequence adjacent to *trnM*; (2) four tandem repeats (TPs); (3) the remaining area of the control region.




Figure 5. Control region structure of two *Anagyrs* species. TR, tandem repeat.

Phylogenetic relationships

The phylogenetic analysis of the concatenated dataset was conducted using BI and ML, which were shown in Fig. 6. With *Encarsia formosa* as an outgroup, the phylogenetic trees of Encyrtidae were constructed based on 13 protein-coding gene sequences of the 21 mitochondrial genomes, including NCBI data and the two newly sequenced *Anagyrus* genomes reported in this study.

The result of maximum-likelihood and Bayesian analysis both indicate that the taxonomic relationship of each genus of Encyrtidae is (*Metaphycus* + *Aenasius*) + (((*Anagyrus* + *Leptomastidea*) + *Encyrtus*) + ((*Blastothrix* + *Psyllaephagus*) + (((*Cheiloneurus* + *Tassonia*) + *Diaphorencyrtus*) + (*Ooencyrtus* + (*Exoristobia* + *Lamennaisia*)))))).

Overall, the phylogenetic trees reconstructed by both methods indicate that species belonging to the same tribe are clustered into one or adjacent clades, while species belonging to the same genus are clustered into the same clade, consistent with the morphological classification system. At the subfamily level, according to the morphological classification system, Encyrtidae is divided into two subfamilies: Tetracneminae and Encyrtinae. *Aenasius*, *Anagyrus*, and *Leptomastidea* all belong to Tetracneminae, while the remaining genera belong to Encyrtinae. However, in the phylogenetic trees reconstructed in this study, the results of both methods show that, except for *Encyrtus* and *Metaphycus*, Encyrtidae is divided into two main parts, which essentially conforms to the morphological classification system. *Metaphycus* and *Aenasius* form a monophyletic clade as sister groups, which is consistent with the previous phylogenetic results (Zhao et al. 2021; Xing et al. 2022).

While the *Anagyrus* species were not clustered on one branch with *Aenasius arizonensis* but clustered with *Encyrtus*, this may be due to different dietary habits. The five genera *Metaphycus*, *Aenasius*, *Anagyrus*, *Leptomastidea*, and *Encyrtus* exclusively parasitize scale insects within Hemiptera. In contrast, other species of Encyrtinae have a broader host range, including species from Lepidoptera, Diptera, Coleoptera, Hymenoptera, and more families within Hemiptera (Noyes 2019). Specifically, *Anagyrus jenniferae* parasitizes *Phenacoccus indicus*, *Anagyrus galinae* parasitizes *Trionymus copiosus*, and *Leptomastidea bifasciata*




Figure 6. Phylogenetic tree of Encyrtidae based on nucleotide sequence of PCGs. Numbers at the nodes are Bayesian posterior probabilities (left) and ML bootstrap values (right). Each color block represents the corresponding family and tribe.

parasitizes *Phenacoccus aceris* and *Planococcus vovae* (Noyes and Hayat 1994; Japoshvili and Hansen 2015; Trjapitzin 1989; Zhang and Xu 2009). These Anagyrini species, which exclusively parasitize the Pseudococcidae, form a distinct clade in both phylogenetic trees. The hosts of *Encyrtus sasakii* include *Takahashia japonica* and *Eulecanium kuwanai*; *Encyrtus eulecaniumiae* parasitizes *Eulecanium kuwanai* and *Eulecanium giganteum*; *Encyrtus rhodococcusiae* targets *Rhodococcus sariuoni*; and *Encyrtus infelix* parasitizes *Ceroplastes destructor*, *Saissetia coffeae*, and *Saissetia oleae* (Trjapitzin 1989; Öncüer 1991; Noyes and Hayat 1994; Zhang and Huang 2001; Gupta and Poorani 2009; Wang et al. 2016), which were exclusively parasitize the Coccidae. Additionally, *Encyrtus aurentii* can parasitize members of the Coccidae (*Saissetia coffeae*), Eriococcidae (*Eriococcus buxi*), and Pseudococcidae (*Planococcus citri*) (Hayat et al. 2003). Consequently, in the phylogenetic trees, the clustering of Anagyrini and Encyrtini species together in the phylogenetic analysis might be attributed to the close genetic relationship between Coccidae and Pseudococcidae (Cook et al. 2002). This phenomenon also indicates the need for further mitochondrial genome sequencing of Encyrtidae species to obtain a more accurate classification status.

Discussion

In this study, we determined two newly sequenced mitogenomes, which are from *A. galinae* and *A. jenniferae*, then found them consistent with previously reported mitogenomes of Encyrtidae. Two new mitogenomes exhibited quite similar features in the genome size, base content, AT nucleotide bias, AT-skew, GC-skew, codon usage of protein genes, secondary structure of tRNAs and gene rearrangement. The BI and ML phylogenetic analysis among the major lineages based on the concatenated datasets yielded well-resolved topologies with moderate to high support for most branches. These results provide a relatively holistic framework and valuable data toward the future resolution of phylogenetic relationships in Encyrtidae. This study provided insights into the phylogenetic relationships of certain taxa within Encyrtidae, the limited sample size and scarcity of molecular evidence remain challenges. Therefore, future studies should aim to augment the number of sampled species and expand the dataset of mitochondrial genomes, utilizing a broader range of data for robust phylogenetic analysis and a comprehensive assessment of the taxonomic status within Encyrtidae.

Acknowledgments

We extend our heartfelt gratitude to Tao Wang from the College of Life Sciences, Nankai University, China, for providing valuable assistance in software analysis. Special thanks also go to Miss Zi-Yan Wang from the University of Sheffield, UK, as well as Mr. Shuai Zhang and Mr. Mark Sharples from the University of Manchester, UK, for their dedicated efforts in reviewing and revising the article. Their contributions have significantly enriched the quality and clarity of our work.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

No funding was received for conducting this study.

Author contributions

Conceptualization: GHZ, CHZ. Data curation: YW, CHZ. Formal analysis: CHZ. Investigation: CHZ, HYW. Methodology: CHZ, YW. Project administration: YW, CHZ, HYW. Resources: GHZ. Software: CHZ, YSL, ZHC. Supervision: GHZ, CHZ. Validation: CHZ, HYW. Visualization: CHZ. Writing – original draft: YW, CHZ, HYW. Writing – review and editing: CHZ, HYW.

Author ORCIDs

Cheng-Hui Zhang  <https://orcid.org/0000-0001-8234-0903>

Hai-Yang Wang  <https://orcid.org/0009-0007-5665-2111>

Yan Wang  <https://orcid.org/0000-0002-5001-975X>

Zhi-Hao Chi  <https://orcid.org/0009-0006-0447-6948>

Guo-Hao Zu  <https://orcid.org/0000-0002-9892-2171>

Data availability

Data presented in this study are openly available in the NCBI repository with accession numbers: [OR652687](https://ncbi.nlm.nih.gov/nucl/OR652687), [OR790122](https://ncbi.nlm.nih.gov/nucl/OR790122).

References

- Benson G (1999) Tandem repeats finder: A program to analyze DNA sequences. *Nucleic Acids Research* 27(2): 573–580. <https://doi.org/10.1093/nar/27.2.573>
- Boore JL (1999) Animal mitochondrial genomes. *Nucleic Acids Research* 27(8): 1767–1780. <https://doi.org/10.1093/nar/27.8.1767>
- Brown WM, George Jr M, Wilson AC (1979) Rapid evolution of animal mitochondrial DNA. *Proceedings of the National Academy of Sciences of the United States of America* 76(4): 1967–1971. <https://doi.org/10.1073/pnas.76.4.1967>
- Cameron SL (2014) Insect mitochondrial genomics: Implications for evolution and phylogeny. *Annual Review of Entomology* 59(1): 95–117. <https://doi.org/10.1146/annurev-ento-011613-162007>
- Cameron SL, Whiting MF (2008) The complete mitochondrial genome of the tobacco hornworm, *Manduca sexta*, (Insecta: Lepidoptera: Sphingidae), and an examination of mitochondrial gene variability within butterflies and moths. *Gene* 408(1–2): 112–123. <https://doi.org/10.1016/j.gene.2007.10.023>
- Chen SF, Zhou YQ, Chen YR, Gu J (2018) fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics (Oxford, England)* 34(17): i884–i890. <https://doi.org/10.1093/bioinformatics/bty560>
- Cook LG, Gullan PJ, Trueman HE (2002) A preliminary phylogeny of the scale insects (Hemiptera: Sternorrhyncha: Coccoidea) based on nuclear small-subunit ribosomal DNA. *Molecular Phylogenetics and Evolution* 25(1): 43–52. [https://doi.org/10.1016/S1055-7903\(02\)00248-8](https://doi.org/10.1016/S1055-7903(02)00248-8)
- Crozier RH, Crozier YC (1992) The cytochrome *b* and ATPase genes of honeybee mitochondrial DNA. *Molecular Biology and Evolution* 9(3): 474–482. <https://doi.org/10.1093/oxfordjournals.molbev.a040729>

- Crozier RH, Crozier YC (1993) The mitochondrial genome of the honeybee *Apis mellifera*: Complete sequence and genome organization. *Genetics* 133(1): 97–117. <https://doi.org/10.1093/genetics/133.1.97>
- Craud A, Rasplus JY, Zhang JX, Burks R, Delvare G, Fusu L, Gumovsky A, Huber JT, Janšta P, Mitroiu MD, Noyes JS, van Noort S, Baker A, Böhmová J, Baur H, Blaimer BB, Brady SG, Bubeníková K, Chartois M, Copeland RS, Dale-Skey Papilloud N, Dal Molin A, Dominguez C, Gebiola M, Guerrieri E, Kresslein RL, Krogmann L, Lemmon E, Murray EA, Nidelet S, Nieves-Aldrey JL, Perry RK, Peters RS, Polaszek A, Sauné L, Torrén J, Triapitsyn S, Tselikh EV, Yoder M, Lemmon AR, Woolley JB, Heraty JM (2024) The Chalcidoidea bush of life: Evolutionary history of a massive radiation of minute wasps. *Cladistics* 40(1): 34–63. <https://doi.org/10.1111/cla.12561>
- Darty K, Denise A, Ponty Y (2009) VARNA: Interactive drawing and editing of the RNA secondary structure. *Bioinformatics (Oxford, England)* 25(15): 1974–1975. <https://doi.org/10.1093/bioinformatics/btp250>
- Donath A, Jühling F, Al-Arab M, Bernhart SH, Reinhardt F, Stadler PF, Middendorf M, Bernhart M (2019) Improved annotation of protein-coding genes boundaries in metazoan mitochondrial genomes. *Nucleic Acids Research* 47(20): 10543–10552. <https://doi.org/10.1093/nar/gkz833>
- Downton M, Castro LR, Austin AD (2002) Mitochondrial gene rearrangements as phylogenetic characters in the invertebrates: The examination of genome ‘morphology’. *Invertebrate Systematics* 16(3): 345–356. <https://doi.org/10.1071/IS02003>
- Du YM, Song X, Liu XJ, Zhong BL (2019) Mitochondrial genome of *Diaphorencyrtus aligarhensis* (Hymenoptera: Chalcidoidea: Encyrtidae) and phylogenetic analysis. *Mitochondrial DNA. Part B, Resources* 4(2): 3190–3191. <https://doi.org/10.1080/23802359.2019.1667913>
- Fan XL, Gong YJ, Chen PY, Tan QQ, Tan JL, Wei SJ (2017) Next-generation sequencing of the mitochondrial genome of *Dolichovespula panda* (Hymenoptera: Vespidae) with a phylogenetic analysis of Vespidae. *Journal of Asia-Pacific Entomology* 20(3): 971–976. <https://doi.org/10.1016/j.aspen.2017.07.009>
- Gupta A, Poorani J (2009) Taxonomic studies on a collection of Chalcidoidea (Hymenoptera) from India with new distribution records. *Journal of Threatened Taxa* 1(5): 300–304. <https://doi.org/10.11609/JoTT.o1861.300-4>
- Hassanin A, Léger N, Deutsch J (2005) Evidence for multiple reversals of asymmetric mutational constraints during the evolution of the mitochondrial genome of Metazoa, and consequences for phylogenetic inferences. *Systematic Biology* 54(2): 277–298. <https://doi.org/10.1080/10635150590947843>
- Hayat M, Narendran TC, Remadevi OK, Manikandan S (2003) Parasitoids (Hymenoptera: Chalcidoidea; Ceraphronoidea) reared mainly from Coccoidea (Homoptera) attacking Sandalwood, *Santalum album* L. *Oriental Insects* 37(1): 309–334. <https://doi.org/10.1080/00305316.2003.10417352>
- Heraty JM, Burks RA, Craud A, Gibson GAP, Liljebblad J, Munro J, Rasplus JY, Delvare G, Janšta P, Gumovsky A, Huber J, Woolley JB, Krogmann L, Heydon S, Polaszek A, Schmidt S, Darling DC, Gates MW, Mottern J, Murray E, Dal Molin A, Triapitsyn S, Baur H, Pinto JD, van Noort S, George J, Yoder M (2013) A phylogenetic analysis of the megadiverse Chalcidoidea (Hymenoptera). *Cladistics* 29: 466–542. <https://doi.org/10.1111/cla.12006>
- Howard LO, Ashmead WH (1896) On some reared parasitic hymenopterous insects from Ceylon. *Proceedings of the United States National Museum* 18(1092): 633–648. <https://doi.org/10.5479/si.00963801.18-1092.633>

- Hurst LD (2002) The Ka/Ks ratio: diagnosing the form of sequence evolution. *Trends in Genetics* 18(9): 486–487. [https://doi.org/10.1016/S0168-9525\(02\)02722-1](https://doi.org/10.1016/S0168-9525(02)02722-1)
- Japoshvili O, Hansen LO (2015) New records of Encyrtidae (Hymenoptera, Chalcidoidea) from Norway VI. *Norwegian Journal of Entomology* 62(2): 174–179.
- Jia CH, Zhang XM, Xu SY, Yang TY, Yanagimoto T, Gao TX (2020) Comparative analysis of the complete mitochondrial genomes of three rockfishes (Scorpaeniformes, Sebastiscus) and insights into the phylogenetic relationships of Sebastidae. *Bioscience Reports* 40(12): BSR20203379. <https://doi.org/10.1042/BSR20203379>
- Jin JJ, Yu WB, Yang JB, Song Y, dePamphilis CW, Yi TS, Li DZ (2020) GetOrganelle: A fast and versatile toolkit for accurate de novo assembly of organelle genomes. *Genome Biology* 21(1): 241. <https://doi.org/10.1186/s13059-020-02154-5>
- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods* 14(6): 587–589. <https://doi.org/10.1038/nmeth.4285>
- Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20(44): 1160–1166. <https://doi.org/10.1093/bib/bbx108>
- Krzywinski J, Grushko OG, Besansky NJ (2006) Analysis of the complete mitochondrial DNA from *Anopheles funestus*: An improved dipteran mitochondrial genome annotation and a temporal dimension of mosquito evolution. *Molecular Phylogenetics and Evolution* 39(2): 417–423. <https://doi.org/10.1016/j.ympev.2006.01.006>
- Li Q, Wei SJ, Tang P, Wu Q, Shi M, Sharkey MJ, Chen XX (2016) Multiple Lines of Evidence from Mitochondrial Genomes Resolve Phylogenetic Relationships of Parasitic Wasps in Braconidae. *Genome Biology and Evolution* 8(9): 2651–2662. <https://doi.org/10.1093/gbe/evw184>
- Liu MD, Luo YF, Jallow BJJ, Meng FM (2023) Characterization of Complete Mitochondrial Genome and Phylogenetic Analysis of a Nocturnal Wasps – *Provespa barthelemyi* (Hymenoptera: Vespidae). *Current Issues in Molecular Biology* 45(12): 9368–9377. <https://doi.org/10.3390/cimb45120587>
- Ma Y, Zheng BY, Zhu JC, Tang P, Chen XX (2019) The mitochondrial genome of *Aenasius arizonensis* (Hymenoptera: Encyrtidae) with novel gene order. *Mitochondrial DNA. Part B, Resources* 4(1): 2023–2024. <https://doi.org/10.1080/23802359.2019.1617052>
- Meng GL, Li YY, Yang CT, Liu SL (2019) MitoZ: A toolkit for animal mitochondrial genome assembly, annotation and visualization. *Nucleic Acids Research* 47(11): e63. <https://doi.org/10.1093/nar/gkz173>
- Munro JB, Heraty JM, Burks RA, Hawks D, Mottern J, Cruaud A, Rasplus JY, Janšta P (2011) A Molecular Phylogeny of the Chalcidoidea (Hymenoptera). *PLOS ONE* 6(11): e27023. <https://doi.org/10.1371/journal.pone.0027023>
- Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* 32(1): 268–274. <https://doi.org/10.1093/molbev/msu300>
- Noyes JS (1980) A review of the genera of Neotropical Encyrtidae (Hymenoptera: Chalcidoidea). *Bulletin of the British Museum of Natural History* 41: 107–253.
- Noyes JS (2019) Universal Chalcidoidea Database. World Wide Web electronic publication. <http://www.nhm.ac.uk/chalcidoids>. [Accessed on 27 Jul 2023]
- Noyes JS, Hayat M (1984) A review of the genera of Indo-Pacific Encyrtidae (Hymenoptera: Chalcidoidea). *Bulletin of the British Museum of Natural History* 48: 131–395.
- Noyes JS, Hayat M (1994) Oriental mealybug parasitoids of the Anagyrini (Hymenoptera: Encyrtidae). CAB International, Wallingford, 554 pp.

- Ojala D, Montoya J, Attardi G (1981) tRNA punctuation model of RNA processing in human mitochondria. *Nature* 290(5806): 470–474. <https://doi.org/10.1038/290470a0>
- Öncüer C (1991) A catalogue of the parasites and predators of insect pests of Turkey. Ege University Agricultural Faculty Press No. 505, Izmir, 204 pp.
- Peng Y, Chen B, Li TJ (2017) Sequencing and analysis of the complete mitochondrial genome of *Parapolybia crocea* (Hymenoptera: Vespidae). *Acta Entomologica Sinica* 60(4): 464–474. <https://doi.org/10.16380/j.kcxb.2017.04.011>
- Perna NT, Kocher TD (1995) Patterns of nucleotide composition at fourfold degenerate sites of animal mitochondrial genomes. *Journal of Molecular Evolution* 41(3): 353–358. <https://doi.org/10.1007/BF01215182>
- Ranwez V, Douzery EJP, Cambon C, Chantret N, Delsuc F (2018) MACSE v2: Toolkit for the Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. *Molecular Biology and Evolution* 35(10): 2582–2584. <https://doi.org/10.1093/molbev/msy159>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE, Sánchez-Gracia A (2017) DnaSP 6: DNA sequence polymorphism analysis of large data sets. *Molecular Biology and Evolution* 34(12): 3299–3302. <https://doi.org/10.1093/molbev/msx248>
- Rudoy A, Zhu CD, Ferrari RR, Zhang YZ (2022) Integrative taxonomy based on morphometric and molecular data supports recognition of the three cryptic species within the *Encyrtus sasakii* complex (Hymenoptera, Encyrtidae). *Journal of Hymenoptera Research* 90: 129–152. <https://doi.org/10.3897/jhr.90.75807>
- Sayers EW, Cavanaugh M, Clark K, Pruitt KD, Sherry ST, Yankie L, Karsch-Mizrachi I (2024) GenBank 2024 Update. *Nucleic Acids Research* 52(D1): D134–D137. <https://doi.org/10.1093/nar/gkad903>
- Shao RF, Campbell NJH, Barker SC (2001) Numerous gene rearrangements in the mitochondrial genome of the Wallaby Louse, *Heterodoxus macropus* (Phthiraptera). *Molecular Biology and Evolution* 18(5): 858–865. <https://doi.org/10.1093/oxford-journals.molbev.a003867>
- Simutnik SA, Perkovsky EE (2023) Description of a new genus and species of Encyrtidae (Hymenoptera: Chalcidoidea) from Danish amber, based on a male specimen featuring an antenna with a distinct anellus. *Zootaxa* 5369(3): 437–445. <https://doi.org/10.11646/zootaxa.5369.3.7>
- Simutnik SA, Perkovsky EE, Vasilenko DV (2022) *Electronoyesella antiqua* Simutnik, gen. et sp. nov. (Chalcidoidea, Encyrtidae) from Rovno amber. *Journal of Hymenoptera Research* 94: 105–120. <https://doi.org/10.3897/jhr.94.94773>
- Simutnik SA, Perkovsky EE, Pankowski MV (2023) A new genus and species of Encyrtidae (Hymenoptera: Chalcidoidea) with a four-segmented funicle from late Eocene Baltic amber. *Zootaxa* 5389(1): 119–127. <https://doi.org/10.11646/zootaxa.5389.1.6>
- Song SN (2015) Comparative mitogenomics and phylogeny of the Microsatrinae (Hymenoptera: Braconidae). PhD Thesis, Zhejiang University, Hangzhou, China.
- Talavera G, Castresana J (2007) Improvement of Phylogenies after Removing Divergent and Ambiguously Aligned Blocks from Protein Sequence Alignments. *Systematic Biology* 56(4): 564–577. <https://doi.org/10.1080/10635150701472164>

- Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular Biology and Evolution* 38(7): 3022–3027. <https://doi.org/10.1093/molbev/msab120>
- Trjapitzin VA (1989) Parasitic Hymenoptera of the Fam. Encyrtidae of Palaearctics. Nauka, Leningrad, 489 pp.
- Wang DP, Fei L, Wang L, Huang S, Yu J (2011) Nonsynonymous substitution rate (Ka) is a relatively consistent parameter for defining fast-evolving and slow-evolving protein-coding genes. *Biology Direct* 6(1): 13. <https://doi.org/10.1186/1745-6150-6-13>
- Wang Y, Zhou QS, Qiao HJ, Zhang AB, Yu F, Wang XB, Zhu CD, Zhang YZ (2016) Formal nomenclature and description of cryptic species of the *Encyrtus sasakii* complex (Hymenoptera: Encyrtidae). *Scientific Reports* 6(1): 34372. <https://doi.org/10.1038/srep34372>
- Wang HY, Zhang CH, Xue H, Xi CX, Yang ZN, Zu GH (2023) Encyrtidae (Hymenoptera: Chalcidoidea) from China, with the description of two new species. *Zootaxa* 5361(2): 237–251. <https://doi.org/10.11646/zootaxa.5361.2.5>
- Wei SJ (2009) Characterization and evolution of hymenopteran mitochondrial genomes and their phylogenetic utility. PhD Thesis, Zhejiang University, Hangzhou, China.
- Wei SJ, Shi M, Sharkey MJ, van Achterberg C, Chen XX (2010) Comparative mitogenomics of Braconidae (Insecta: Hymenoptera) and the phylogenetic utility of mitochondrial genomes with special reference to Holometabolous insects. *BMC Genomics* 11(1): 371. <https://doi.org/10.1186/1471-2164-11-371>
- Wolstenholme DR (1992a) Genetic novelties in mitochondrial genomes of multicellular animals. *Current Opinion in Genetics & Development* 2(6): 918–925. [https://doi.org/10.1016/S0959-437X\(05\)80116-9](https://doi.org/10.1016/S0959-437X(05)80116-9)
- Wolstenholme DR (1992b) Animal mitochondrial DNA: Structure and evolution. *International Review of Cytology* 141: 173–216. [https://doi.org/10.1016/S0074-7696\(08\)62066-5](https://doi.org/10.1016/S0074-7696(08)62066-5)
- Xing ZP, Liang X, Wang X, Hu HY, Huang YX (2022) Novel gene rearrangement pattern in mitochondrial genome of *Ooencyrtus plautus* Huang & Noyes, 1994: New gene order in Encyrtidae (Hymenoptera, Chalcidoidea). *ZooKeys* 1124: 1–21. <https://doi.org/10.3897/zookeys.1124.83811>
- Xiong M, Zhou QS, Zhang YZ (2019) The complete mitochondrial genome of *Encyrtus infelix* (Hymenoptera: Encyrtidae). *Mitochondrial DNA. Part B, Resources* 4(1): 114–115. <https://doi.org/10.1080/23802359.2018.1537727>
- Xu W, Lin SP, Liu HY (2021) Mitochondrial genomes of five *Hyphessobrycon* tetras and their phylogenetic implications. *Ecology and Evolution* 11(18): 12754. <https://doi.org/10.1002/ece3.8019>
- Zhang YZ, Huang DW (2001) Two new Encyrtid parasites (Hymenoptera: Chalcidoidea) from China. *Oriental Insects* 35(1): 311–319. <https://doi.org/10.1080/00305316.2001.10417310>
- Zhang YZ, Xu ZH (2009) A review of Chinese species of *Leptomastidea* Mercet (Hymenoptera: Encyrtidae). *Acta Entomologica Sinica* 52(4): 420–423.
- Zhang JX, Lindsey ARI, Peters RS, Heraty JM, Hopper KR, Werren JH, Martinson EO, Woolley JB, Yoder MJ, Krogmann L (2020a) Conflicting signal in transcriptomic markers leads to a poorly resolved backbone phylogeny of chalcidoid wasps. *Systematic Entomology* 45(4): 783–802. <https://doi.org/10.1111/syen.12427>
- Zhang D, Gao FL, Jakovlić I, Zou H, Zhang J, Li WX, Wang GT (2020b) PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data

- management and evolutionary phylogenetics studies. *Molecular Ecology Resources* 20(1): 348–355. <https://doi.org/10.1111/1755-0998.13096>
- Zhao HF, Chen Y, Wang ZT, Chen HF, Qin YG (2021) Two complete mitogenomes of Chalcididae (Hymenoptera: Chalcidoidea): genome description and phylogenetic implications. *Insects* 12(12): 1049. <https://doi.org/10.3390/insects12121049>
- Zhou QS, Xiong M, Luo AR, Zhang YZ, Zhu CD (2021) The complete mitochondrial genome of *Metaphycus eriococci* (Timberlake) (Hymenoptera: Encyrtidae). *Mitochondrial DNA. Part B, Resources* 6(2): 550–552. <https://doi.org/10.1080/23802359.2021.1872450>
- Zhu JC, Tang P, Zhou BY, Wu Q, Wei SJ, Chen XX (2018) The first two mitochondrial genomes of the family Aphelinidae with novel gene orders and phylogenetic implications. *International Journal of Biological Macromolecules* 118(A): 386–396. <https://doi.org/10.1016/j.ijbiomac.2018.06.087>
- Zhu JC, Xiao H, Tang P, Li XF, Li XK, Zhu CD, Wu Q, Xiao JH, van Achterberg C, Huang DW, Chen XX (2023) Evolutionary timescale of chalcidoid wasps inferred from over one hundred mitochondrial genomes. *Zoological Research* 44(3): 467–482. <https://doi.org/10.24272/j.issn.2095-8137.2022.379>

Supplementary material 1

Secondary structures of 22 tRNA genes of *Anagyrus galinae*

Authors: Cheng-Hui Zhang, Hai-Yang Wang, Yan Wang, Zhi-Hao Chi, Yue-Shuo Liu, Guo-Hao Zu

Data type: jpg

Explanation note: Blue gene names indicate that in the major strand, and red names indicate that in the minor strand.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1206.121923.suppl1>

Supplementary material 2

Secondary structures of 22 tRNA genes of *Anagyrus jenniferae*

Authors: Cheng-Hui Zhang, Hai-Yang Wang, Yan Wang, Zhi-Hao Chi, Yue-Shuo Liu, Guo-Hao Zu

Data type: jpg

Explanation note: Blue gene names indicate that in the major strand, and red names indicate that in the minor strand.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1206.121923.suppl2>