

Species diversity of freshwater glass eel (*Anguilliformes*, *Anguillidae*) of Yilan, Taiwan, with remark on two new records

Yen-Ting Lin¹, Yu-San Han¹

¹ Institute of Fisheries Science, College of Life Science, National Taiwan University, Taipei 10617, Taiwan
Corresponding author: Yu-San Han (yshan@ntu.edu.tw)

Abstract

Yilan, Taiwan is the first place in East Asia where freshwater glass eels, the juvenile stage of *Anguilla* species, arrive by ocean currents. We collected glass eels by fyke net in Lanyang River estuary twice a month from July 2010 to November 2023. By morphological examination and sequencing of the mitochondrial cytochrome b gene, we identified seven species of *Anguilla*. Most of the glass eels captured in Yilan belonged to the species *A. japonica*, *A. marmorata*, and *A. bicolor pacifica*. Only a few were *A. luzonensis*, and two *A. celebesensis* were recorded. In addition, two species were recorded for the first time from Taiwan; *A. interioris* and *A. borneensis* were confirmed by cytochrome b sequencing. Thus, we increase the number of *Anguilla* species in Taiwan from five to seven.

Key words: *Anguilla borneensis*, *Anguilla interioris*, glass eel, new records



Academic editor: Yusuke Hibino
Received: 17 April 2024
Accepted: 20 June 2024
Published: 9 December 2024

ZooBank: <https://zoobank.org/584FAC26-31F2-45D5-85A7-FB1288454E92>

Citation: Lin Y-T, Han Y-S (2024) Species diversity of freshwater glass eel (*Anguilliformes*, *Anguillidae*) of Yilan, Taiwan, with remark on two new records. In: Ho H-C, Russell B, Hibino Y, Lee M-Y (Eds) Biodiversity and taxonomy of fishes in Taiwan and adjacent waters. ZooKeys 1220: 5–14. <https://doi.org/10.3897/zookeys.1220.125590>

Copyright: © Yen-Ting Lin & Yu-San Han.
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

The freshwater eel (*Anguilla* spp.) comprises 16 species and three subspecies (Arai 2016a). All *Anguilla* species are catadromous fish, meaning they migrate to the ocean to spawn (Arai and Chino 2012). The leaf-like larvae of *Anguilla* species, known as leptocephali, are carried by ocean currents and undergo metamorphosis into eel-like juveniles, which are known as glass eels (Tsukamoto et al. 2002; Hatakeyama et al. 2022). Glass-eel fishing is crucial for the eel aquaculture industry, as there are no artificial reproduction techniques for commercial purposes (Okamura et al. 2014). In Taiwan, the dispersal of glass eels is primarily influenced by the Kuroshio Current (Hsiung et al. 2022b). Yilan, Taiwan, is renowned as the largest glass-eel fishing ground in Taiwan due to its proximity to the Kuroshio. Notably, Yilan holds the distinction of being the first location in East Asia where glass eels arrive, establishing it as a significant hub for this crucial stage in the eel life cycle (Han et al. 2016a).

To date, five *Anguilla* species have been identified and recorded in Taiwan (Leander et al. 2012; Han et al. 2016b). Among these, *A. japonica*, *A. marmorata*, and *A. bicolor pacifica* are the most prevalent species (Han 2001; Hsu et al. 2019), while *A. luzonensis* and *A. celebesensis* are notably very rare and primarily observed as glass eel in Taiwan (Teng et al. 2009; Han et al. 2016b). Previous studies suggest that *A. japonica*, *A. marmorata*, and *A. bicolor pacifica* share a

common spawning area near the southern West Mariana Ridge (Kuroki et al. 2009; Arai 2016b), whereas other tropical eel species (*A. celebesensis*, *A. borneensis*, *A. luzonensis*, and *A. interioris*) have been identified near southern Mindanao Island as their spawning grounds (Aoyama et al. 2003; Wouthuyzen et al. 2009; Arai 2014, 2016b). Due to the morphological challenges in distinguishing tropical eel glass eels (Minegishi et al. 2005), DNA barcoding techniques, as highlighted by Wibowo et al. (2021), provide a precise method for the identification of species. Previous research also indicates that mitochondrial cytochrome b gene fragments are suitable for the identification of freshwater eels (Han et al. 2008). This study aims to analyse glass-eel samples captured in Yilan from July 2010 to November 2023. Through DNA sequencing, the goal is to confirm the number of freshwater glass-eel species transported to Taiwan during this period.

Materials and methods

Sample collection

Glass eels were collected twice a month at night using a fyke net positioned in the estuary of the Yilan River (24.7162°N, 121.8352°E) from July 2010 to November 2023. Following the capture, all the samples were immersed in a 95% ethanol solution for measurement and preservation. All freshwater glass-eel specimens were deposited in the Institute of Fisheries, National Taiwan University (NTUIFS). Recent research adhered to ethical regulations set forth by the Institutional Animal Care and Use Committee (IACUC) under approval number NTU-110-EL-00152.

Morphological measurement

The method for morphological identification of anguillid glass eels was adapted from Han et al. (2012), and the description of the pigmentation stage followed Fukuda et al. (2013). Four morphological parameters were measured using digital callipers with an accuracy of 0.1 mm: total length (TL), head length (HL), pre-dorsal length (PDL), and pre-anal length (PAL). The fin-difference ratio was then calculated using the formula shown below. Glass eels with fin differences exceeding 13% in Yilan were consistently identified as *A. marmorata* (Han et al. 2012). Therefore, specimens displaying black pigment on the tail and fin differences <13% were chosen for mitochondrial cytochrome b gene sequencing.

$$\text{Fin Difference Ratio (\%)} = \frac{\text{PAL (mm)} - \text{PDL (mm)}}{\text{TL (mm)}} \times 100$$

Mitochondrial cytochrome b gene sequencing

Freshwater glass-eel specimens with a fin-difference ratio <13% were DNA sequenced for precise identification; these amounted to 281 samples. Genomic DNA was extracted from the dorsal-fin tissue of the glass eels using the Favor-Prep Tissue Genomic DNA Extraction Mini Kit (Favorgen, Taiwan). Polymerase chain reaction (PCR) was carried out to amplify a segment of mitochondrial cytochrome b using forward primer: cytb-F (5'-GAT GCC CTA GTG GAT CTA CC-3') and reverse primer: cytb-R (5'-TAT GGG TGT TCT ACT GGT AT-3'), which was adapted from Han et al. (2008). The resulting PCR product (approximately

1000 bp) was sequenced using the primers cytb-F or cytb-R (by Genomic Biotech Inc., Taiwan), following protocols from Han et al. (2008). Sequencing results were submitted to the National Center for Biotechnology Information (NCBI) GenBank (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) to confirm species.

Results

Diversity of freshwater glass eel

A total of 29,442 freshwater glass eels were collected between July 2010 and November 2023. The composition of freshwater glass-eel species is shown in Table 1. *Anguilla japonica* and *A. marmorata* were the most prevalent species, comprising 95.4% of our captures (Table 1). Although *A. bicolor pacifica* and *A. luzonensis* were infrequently captured, they still represented 3.9% and 0.7% of all specimens, respectively, and two *A. celebesensis* were also recorded (Table 1). Additionally, two species were found in Taiwan for the first time: *A. interioris* and *A. borneensis* (Table 1).

New *Anguilla* records from Taiwan

The sequencing results of the two new records, total two specimens (NTUIFS IL13'0812-76 and NTUIFS IL21'0715-207), and the best matched BLAST results are shown in Table 2.

Table 1. Number of species (*n*) and percentage contributions of freshwater glass eels collected in Yilan.

Species	<i>n</i>	Percentage contributions
<i>A. japonica</i>	14217	48.3
<i>A. marmorata</i>	13864	47.1
<i>A. bicolor pacifica</i>	1152	3.9
<i>A. luzonensis</i>	205	0.7
<i>A. celebesensis</i>	2	<0.01
<i>A. interioris</i>	1	<0.01
<i>A. borneensis</i>	1	<0.01

Table 2. Sequencing BLAST results of three new records freshwater glass eel.

Specimen	Species	Percent identity (%)	NCBI accession
NTUIFS IL13'0812-76	<i>Anguilla interioris</i>	99.4	HG965574.1
NTUIFS IL21'0715-207	<i>Anguilla borneensis</i>	99.4	NC_006536.1 *

*[NC_006536.1](https://ncbi.nlm.nih.gov/nucl/NC_006536.1) was identified as *A. malgumora*, which is a junior synonym of *A. borneensis* according to Minegishi et al. (2005).

Family Anguillidae

Anguilla interioris Whitley, 1938

Figs 1, 2, Table 3

Material examined. NTUIFS IL13'0812-76, 46 mm TL, off the estuary of the Yilan River, Yilan, northeastern Taiwan (24.7162°N, 121.8352°E), 12 August 2013, fyke net, collected by Yu-San Han.

Short description. PDL 29.3% in TL; PAL 39.1% in TL; fin-difference ratio 9.78%. Body elongate, head length 13.1% TL. The specimen was in fresh condition, with black pigment distributed on the caudal fin and slightly on the caudal peduncle; pigmentation stages V_{B2} (Fig. 2).

Distribution. New Guinea (Aoyama et al. 2000); Philippines (glass eel only, Wibowo et al. 2021); Indonesia (leptocephalus only, Kuroki et al. 2006; all stages, Zan et al. 2022); Taiwan (glass eel only, present study).

Remarks. The distribution of *A. interioris* has been primarily known from only New Guinea (Aoyama et al. 2000). However, a study by Kuroki et al. (2006) documented the leptocephalus of *A. interioris* in the Indonesian Archipelago, marking the first expansion of the species beyond its then-known range. Additionally, records of *A. interioris* have been identified using DNA sequencing from Indonesia and southern Mindanao, Philippines (Wibowo et al. 2021; Zan et al. 2022). Herein, we present the first record of *A. interioris* glass eel from Taiwan.



Figure 1. *Anguilla interioris* (NTUIFS IL13'0812-76), 46 mm TL. Preserved in 95% alcohol. Scale bar: 10 mm.



Figure 2. Pigmentation on the tail tip of *Anguilla interioris* (NTUIFS IL13'0812-76).

Table 3. The morphological parameters of seven freshwater glass eel collected in Yilan.

Species	TL (mm)	PDL (mm)	PAL (mm)	Fin-difference ratio (%)
<i>A. japonica</i>	61.1±2.5	15.1±0.9	20.1±0.7	9.2±1.3
<i>A. marmorata</i>	51.4±2.7	11.8±0.8	19.4±1.1	15.5±0.8
<i>A. bicolor pacifica</i>	49.2±2.3	18.3±1.6	18.5±1.6	0.5±0.5
<i>A. luzonensis</i>	52.9±2.7	13.7±0.7	19.4±1.0	11.4±1.1
<i>A. celebesensis</i>	45.3	12.8	17.5	10.4
<i>A. interioris</i>	46.0	13.5	18.0	10.1
<i>A. borneensis</i>	49.5	13.0	18.0	9.8

***Anguilla borneensis* Popta, 1924**

Figs 3, 4, Table 3

Material examined. NTUIFS IL21'0715-207, 49.5 mm TL, off the estuary of the Yilan River, Yilan, northeastern Taiwan (24.7162°N, 121.8352°E), 15 July 2021, fyke net, collected by Yen-Ting Lin.

Short description. PDL 26.3% in TL; PAL 36.4% in TL; fin-difference ratio 10.1%. Body extremely elongate, head length 10.1% TL. The specimen was in fresh condition, with black pigment distributed on the caudal peduncle and caudal fin; pigmentation stages V_A (Fig. 4).

Distribution. Indonesia (Watanabe et al. 2014); Taiwan (glass eel only, present study).

Remarks. The best-matched GenBank accession number for NTUIFS IL21'0715-207 was found to be [NC_006536.1](#), which corresponds to *A. malgumora* submitted by Minegishi et al. (2005). However, it is noteworthy that *A. malgumora* was identified as a junior synonym of *A. borneensis* by Minegishi et al. (2005). Based on the comprehensive examination by Minegishi et al. (2005) and the detailed de-



Figure 3. *Anguilla borneensis* (NTUIFS IL21'0715-207), 49.5mm TL. Preserved in 95% alcohol. Scale bar: 10 mm.



Figure 4. Pigmentation on the tail tip of *Anguilla borneensis* (NTUIFS IL21'0715-207).

scription provided in [NC_006536.1](#), we can confidently affirm that our specimen NTUIFS IL21'0715-207 is *A. borneensis*. Herein, we report the first record of *A. borneensis* outside of the Indonesia (Watanabe et al. 2014).

Discussion and conclusion

Leptocephali and glass eels primarily rely on ocean currents for transport (Kuroki et al. 2016). In Taiwan, the main current responsible for transporting glass eels is the North Equatorial Current (NEC), followed by the Kuroshio, which is known to carry the most abundant anguillid species (*A. japonica* and *A. marmorata*) to the region (Hsiung et al. 2022a). Additionally, other tropical eels (*A. bicolor pacifica*, *A. luzonensis*, and *A. celebesensis*) may reach Taiwan via the bifurcation region of the NEC near the Philippine coast, which could potentially transport glass eels from southern Mindanao Island to the Kuroshio (Aoyama et al. 2015; Rudnick et al. 2015). The two species identified in our study align with previous research on the diversity of tropical glass eels (*A. celebesensis*, *A. interioris*, and *A. borneensis*) in southern Mindanao (Shirotori et al. 2016).

Previous research based on differences in Sr:Ca ratios in the leptocephalus otoliths has shown the presence of two populations of *A. interioris*, with one population in the Indian Ocean and another in the Pacific Ocean (Kuroki et al. 2006). Furthermore, leptocephali of the Pacific Ocean population of *A. interioris* potentially are transported to Taiwan via the Mindanao Current which ultimately forms a connection with the Kuroshio and the Mindanao Eddy (Kuroki et al. 2006).

The distribution of leptocephali and glass eels of the Indonesian *A. borneensis*, which is considered the most basal *Anguilla* species, remains unclear (Aoyama et al. 2001). The spawning area of *A. borneensis* may overlap with other basal tropical eels (*A. celebesensis*, *A. interioris*, *A. marmorata*, and *A. bicolor bicolor*) in Indonesia in the western Pacific Ocean (Arai and Abdul Kadir 2017); this suggests the possibility that a similar pathway to Taiwan is followed, as by *A. interioris* and *A. celebesensis* (Han et al. 2016b).

Alternatively, it is possible that if *A. borneensis* and *A. interioris* establish a new population in the western Pacific Ocean, their larvae could be carried to Taiwan via the North Equatorial Current (NEC) and the Kuroshio. Additionally, some alien freshwater eel species have escaped from aquaculture ponds and have been reported to have similar migration behaviour of native eel in East Asia (Okamura et al. 2002). Examples include *A. rostrata*, which has been discovered in Taiwanese waters (Han et al. 2002), and the European eel, *A. anguilla*, which was captured in the East China Sea and Japanese waters (Aoyama 2000; Okamura et al. 2002). Therefore, the possibility of alien eel species establishing new populations in the West Pacific Ocean cannot be discounted, whether caused by human activities (*A. rostrata* and *A. anguilla*) or by natural phenomenon (*A. interioris* and *A. borneensis*) (Aoyama 2000; Han et al. 2002).

Although there are seven species of freshwater glass eel recorded in Taiwan, only elvers of *A. japonica*, *A. marmorata*, *A. luzonensis*, and *A. bicolor pacifica* had been found in streams (Tzeng and Tabeta 1983; Watanabe et al. 2013; Hsu et al. 2019). The existence of the elvers and adults of *A. celebesensis*, *A. interioris*, and *A. borneensis* still need confirmation in the field.

In conclusion, the present study increases the number of freshwater glass-eel species in Taiwan from five to seven (Leander et al. 2012; Han et al. 2016b), with the addition two new species records in this paper.

Acknowledgements

We thank Zhen-Hui Chen (local fisherman) for assisting us in setting the fyke net and providing various sampling support; Chien-Hsiang Lin and Hsin-Wei Liu (Biodiversity Research Center, Academia Sinica) for photographing specimens and offering curatorial insights for this research.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

The authors extend our gratitude to the National Science and Technology Council, Executive Yuan, Taiwan (MOST 111-2313-B-002-016-MY3) for funding this project.

Author contributions

The experiment and sample collecting were performed by Yu-San Han and Yen-Ting Lin. Yen-Ting Lin write the manuscript. Yu-San Han designed and supervised the experiments. All authors participated in manuscript writing and interpretation of results. All authors read and approved the final manuscript.

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Aoyama J (2000) The European eel, *Anguilla anguilla* (L.), in Japanese waters. *Dana* 12: 1–5.
- Aoyama J, Watanabe S, Ishikawa S, Nishida M, Tsukamoto K (2000) Are morphological characters distinctive enough to discriminate between two species of freshwater eels, *Anguilla celebesensis* and *A. interioris*? *Ichthyological Research* 47(2): 157–161. <https://doi.org/10.1007/BF02684236>
- Aoyama J, Nishida M, Tsukamoto K (2001) Molecular phylogeny and evolution of the freshwater eel, genus *Anguilla*. *Molecular Phylogenetics and Evolution* 20(3): 450–459. <https://doi.org/10.1006/mpev.2001.0959>
- Aoyama J, Wouthuyzen S, Miller MJ, Inagaki T, Tsukamoto K (2003) Short-distance spawning migration of tropical freshwater eels. *The Biological Bulletin* 204(1): 104–108. <https://doi.org/10.2307/1543500>
- Aoyama J, Yoshinaga T, Shinoda A, Shirotori F, Yambot AV, Han YS (2015) Seasonal changes in species composition of glass eels of the genus *Anguilla* (Teleostei:

- Anguillidae) recruiting to the Cagayan River, Luzon Island, the Philippines. *Pacific Science* 69(2): 263–270. <https://doi.org/10.2984/69.2.8>
- Arai T (2014) Evidence of local short-distance spawning migration of tropical freshwater eels, and implications for the evolution of freshwater eel migration. *Ecology and Evolution* 4(19): 3812–3819. <https://doi.org/10.1002/ece3.1245>
- Arai T (2016a) Taxonomy and distribution. In: Arai T (Ed.) *Biology and Ecology of Anguillid Eels*. CRC Press, Boca Raton, 1–20. <https://doi.org/10.1201/b19925>
- Arai T (2016b) Spawning ground of tropical eels. In: Arai T (Ed.) *Biology and Ecology of Anguillid Eels*. CRC Press, Boca Raton, 89–107. <https://doi.org/10.1201/b19925>
- Arai T, Abdul Kadir SR (2017) Opportunistic spawning of tropical anguillid eels *Anguilla bicolor bicolor* and *A. bengalensis bengalensis*. *Scientific Reports* 7(1): 41649. <https://doi.org/10.1038/srep41649>
- Arai T, Chino N (2012) Diverse migration strategy between freshwater and seawater habitats in the freshwater eel genus *Anguilla*. *Journal of Fish Biology* 81(2): 442–455. <https://doi.org/10.1111/j.1095-8649.2012.03353.x>
- Fukuda N, Miller MJ, Aoyama J, Shinoda A, Tsukamoto K (2013) Evaluation of the pigmentation stages and body proportions from the glass eel to yellow eel in *Anguilla japonica*. *Fisheries Science* 79(3): 425–438. <https://doi.org/10.1007/s12562-013-0621-x>
- Han YS (2001) Validation of the occurrence of short-finned eel *Anguilla bicolor pacifica* in natural waters of Taiwan. *Acta Zoologica Taiwanica* 12(1): 9–19.
- Han YS, Yu CH, Yu HT, Chang CW, Liao IC, Tzeng WN (2002) The exotic American eel in Taiwan: Ecological implications. *Journal of Fish Biology* 60(6): 1608–1612. <https://doi.org/10.1111/j.1095-8649.2002.tb02454.x>
- Han YS, Sun YL, Liao YF, Liao IC, Shen KN, Tzeng WN (2008) Temporal analysis of population genetic composition in the overexploited Japanese eel *Anguilla japonica*. *Marine Biology* 155(6): 613–621. <https://doi.org/10.1007/s00227-008-1057-1>
- Han YS, Yambot AV, Zhang H, Hung CL (2012) Sympatric spawning but allopatric distribution of *Anguilla japonica* and *Anguilla marmorata*: Temperature-and oceanic current-dependent sieving. *PLOS ONE* 7(6): e37484. <https://doi.org/10.1371/journal.pone.0037484>
- Han YS, Wu CR, Iizuka Y (2016a) Batch-like arrival waves of glass eels of *Anguilla japonica* in offshore waters of Taiwan. *Zoological Studies (Taipei, Taiwan)* 55: 36. <https://doi.org/10.6620/ZS.2016.55-36>
- Han YS, Lin YF, Wu CR, Iizuka Y, Castillo TR, Yambot IU, Mamalangkap MD, Yambot AV (2016b) Biogeographic distribution of the eel *Anguilla luzonensis*: Dependence upon larval duration and oceanic currents. *Marine Ecology Progress Series* 551: 227–238. <https://doi.org/10.3354/meps11728>
- Hatakeyama R, Sudo R, Yatabe T, Yamano K, Nomura K (2022) Developmental features of Japanese eels, *Anguilla japonica*, from the late leptocephalus to the yellow eel stages: An early metamorphosis to the eel-like form and a prolonged transition to the juvenile. *Journal of Fish Biology* 100(2): 454–473. <https://doi.org/10.1111/jfb.14956>
- Hsiung KM, Lin YT, Han YS (2022a) Current dependent dispersal characteristics of Japanese glass eel around Taiwan. *Journal of Marine Science and Engineering* 10(1): 98. <https://doi.org/10.3390/jmse10010098>
- Hsiung KM, Kuo YC, Lin YT, Tseng YH, Han YS (2022b) North Equatorial Current and Kuroshio velocity variations affect body length and distribution of the Japanese eel *Anguilla japonica* in Taiwan and Japan. *Scientific Reports* 12(1): 2888. <https://doi.org/10.1038/s41598-022-06669-8>

- Hsu HY, Chen HW, Han YS (2019) Habitat partitioning and its possible genetic background between two sympatrically distributed eel species in Taiwan. *Zoological Studies* (Taipei, Taiwan) 58: 27. <https://doi.org/10.6620/ZS.2019.58-27>
- Kuroki M, Aoyama J, Wouthuyzen S, Sumardhiharga KO, Miller MJ, Minagawa G, Tsukamoto K (2006) Age and growth of *Anguilla interioris* leptocephali collected in Indonesian waters. *Coastal Marine Science* 30(2): 464–468.
- Kuroki M, Aoyama J, Miller MJ, Yoshinaga T, Shinoda A, Hagihara S, Tsukamoto K (2009) Sympatric spawning of *Anguilla marmorata* and *Anguilla japonica* in the western North Pacific Ocean. *Journal of Fish Biology* 74(9): 1853–1865. <https://doi.org/10.1111/j.1095-8649.2009.02299.x>
- Kuroki M, Okamura A, Takeuchi A, Tsukamoto K (2016) Effect of water current on the body size and occurrence of deformities in reared Japanese eel leptocephali and glass eels. *Fisheries Science* 82(6): 941–951. <https://doi.org/10.1007/s12562-016-1015-7>
- Leander NJ, Shen KN, Chen RT, Tzeng WN (2012) Species composition and seasonal occurrence of recruiting glass eels (*Anguilla* spp.) in the Hsiukuluan River, eastern Taiwan. *Zoological Studies* (Taipei, Taiwan) 51(1): 59–71.
- Minegishi Y, Aoyama J, Inoue JG, Miya M, Nishida M, Tsukamoto K (2005) Molecular phylogeny and evolution of the freshwater eels genus *Anguilla* based on the whole mitochondrial genome sequences. *Molecular Phylogenetics and Evolution* 34(1): 134–146. <https://doi.org/10.1016/j.ympev.2004.09.003>
- Okamura A, Yamada Y, Mikawa N, Tanaka S, Oka HP (2002) Exotic silver eels *Anguilla anguilla* in Japanese waters: Seaward migration and environmental factors. *Aquatic Living Resources* 15(6): 335–341. [https://doi.org/10.1016/S0990-7440\(02\)01190-7](https://doi.org/10.1016/S0990-7440(02)01190-7)
- Okamura A, Horie N, Mikawa N, Yamada Y, Tsukamoto K (2014) Recent advances in artificial production of glass eels for conservation of anguillid eel populations. *Ecology Freshwater Fish* 23(1): 95–110. <https://doi.org/10.1111/eff.12086>
- Rudnick DL, Jan S, Lee CM (2015) A new look at circulation in the western North Pacific: Introduction to the special issue. *Oceanography* (Washington, D.C.) 28(4): 16–23. <https://doi.org/10.5670/oceanog.2015.77>
- Shirotori F, Ishikawa T, Tanaka C, Aoyama J, Shinoda A, Yambot AV, Yoshinaga T (2016) Species composition of anguillid glass eels recruited at southern Mindanao Island, the Philippines. *Fisheries Science* 82(6): 915–922. <https://doi.org/10.1007/s12562-016-1030-8>
- Teng HY, Lin YS, Tzeng CS (2009) A new *Anguilla* species and a reanalysis of the phylogeny of freshwater eels. *Zoological Studies* (Taipei, Taiwan) 48(6): 808–822.
- Tsukamoto K, Aoyama J, Miller MJ (2002) Migration, speciation, and the evolution of diadromy in anguillid eels. *Canadian Journal of Fisheries and Aquatic Sciences* 59(12): 1989–1998. <https://doi.org/10.1139/f02-165>
- Tzeng WN, Tabeta O (1983) First record of the short-finned eel *Anguilla bicolor pacifica* elvers from Taiwan. *Nippon Suisan Gakkaishi* 49(1): 27–32. <https://doi.org/10.2331/suisan.49.27>
- Watanabe S, Aoyama J, Hagihara S, Ai B, Azanza RV, Tsukamoto K (2013) *Anguilla huangi* Teng, Lin, and Tzeng, 2009, is a junior synonym of *Anguilla luzonensis* Watanabe, Aoyama, and Tsukamoto, 2009. *Fisheries Science* 79(3): 375–383. <https://doi.org/10.1007/s12562-013-0620-y>
- Watanabe S, Aoyama J, Tsukamoto K (2014) On the Identities of *Anguilla borneensis*, *A. malgumora*, and *Muraena malgumora*. *Copeia* 2014(3): 568–576. <https://doi.org/10.1643/Ci-14-059>

- Wibowo A, Hubert N, Dahrudin H, Steinke D, Suhaimi RA, Samuel, Atminarso D, Anggraeni DP, Trismawanti I, Baumgartner LJ, Ning N (2021) Assessing temporal patterns and species composition of glass eel (*Anguilla* spp.) cohorts in Sumatra and Java using DNA Barcodes. *Diversity* 13(5): 193. <https://doi.org/10.3390/d13050193>
- Wouthuyzen S, Aoyama J, Sugeha HY, Miller MJ, Kuroki M, Minegishi Y, Suharti SR, Tsukamoto K (2009) Seasonality of spawning by tropical anguillid eels around Sulawesi Island, Indonesia. *Naturwissenschaften* 96(1): 153–158. <https://doi.org/10.1007/s00114-008-0457-x>
- Zan ND, Taha H, Akbar N, Achmad MJ, Arai T (2022) Molecular confirmation of the occurrence of *Anguilla interioris* (Actinopterygii: Anguilliformes) in North Maluku of Indonesia and mitochondrial DNA haplotype diversity among existing specimens. *Zoologia* 39: e22008. <https://doi.org/10.1590/s1984-4689.v39.e22008>

Supplementary material 1

NTUIFS IL13'0812-76 sequencing data

Authors: Yen-Ting Lin, Yu-San Han

Data type: txt

Explanation note: The sequencing results of NTUIFS IL13'0812-76.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1220.125590.suppl1>

Supplementary material 2

NTUIFS IL21'0715-207 sequencing data

Authors: Yen-Ting Lin, Yu-San Han

Data type: txt

Explanation note: The sequencing results of NTUIFS IL21'0715-207.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1220.125590.suppl2>