

A new genus and two new species of Southeast Asian *Bidessini* as well as new synonyms for Oceanian species (Coleoptera, Dytiscidae)

Michael Balke¹, Johannes Bergsten², Liang-Jong Wang³, Lars Hendrich¹

1 SNSB-Zoologische Staatssammlung München, Münchhausenstrasse 21, München, Germany **2** Department of Zoology, Swedish Museum of Natural History, Box 50007, SE-10405 Stockholm, Sweden **3** Division of Forest Protection, Taiwan Forestry Research Institute, Taipei, Taiwan

Corresponding author: Michael Balke (kaefer@zsm.mwn.de)

Academic editor: M. Michat | Received 17 November 2016 | Accepted 14 January 2017 | Published 30 January 2017

<http://zoobank.org/3ED32CCA-1B02-4152-8141-6900E40D9445>

Citation: Balke M, Bergsten J, Wang L-J, Hendrich L (2017) A new genus and two new species of Southeast Asian *Bidessini* as well as new synonyms for Oceanian species (Coleoptera, Dytiscidae). ZooKeys 647: 137–151. <https://doi.org/10.3897/zookeys.647.11231>

Abstract

Rompindessus jenisi Balke, Bergsten & Hendrich, **gen. n.** et **sp. n.** is described from near Rompin village in West Malaysia. The new genus is characterized by the presence of an occipital line and basal pronotal striae, the presence of a thick anterior bead on clypeus and two-segmented parameres as well as by the absence of basal elytral striae, the absence of sutural line on elytron, the absence of basal epipleural transverse carina, and the absence of longitudinal elytral carina. Moreover, male pro- and mesotarsus appear stout, and distinctly dilated laterally; the pronotum is comparably long and parallel-sided and the colour of beetle conspicuous dark orange. *Leiodytes kualalipis* Balke, Wang, Bergsten & Hendrich, **sp. n.** is described from West Malaysia (Pahang) and South Vietnam (Cat Tien). It is well characterized by its large size, elongate body and the form of the median lobe. *Limbodessus fjiensis* (J. Balfour-Browne, 1944), **comb. n.** described from Fiji is a new synonym of *Limbodessus curvuplicatus* (Zimmermann, 1927) described from Samoa.

Keywords

Bidessini, Dytiscidae, new combination, new genus, new species, new synonymy, Oceania, Southeast Asia

Introduction

There are currently 48 genera in the Bidessini (Nilsson 2016; Miller 2016; Miller and Bergsten 2016). With a body length of typically around 1–3 mm, they constitute most of the smaller species of Dytiscidae. Bidessini genera have to date been justified mainly on a diagnostic combination of structural features (Biström 1988; Balke et al. 2002; Miller and Short 2015; Miller and Bergsten 2016) rather than apomorphies and this had to lead to recognition of genera that render other paraphyletic (Balke and Ribera 2004). Some of these features such as presence / absence of elytral striae (plicae) or occipital line have been shown to vary within clades of closely related species (Balke et al. 2015) or even within one species (Balke unpublished). In this context, the use of phylogenetic reconstructions based on DNA sequence data offers a source of information that helps to delineate monophyletic entities (Hendrich and Balke 2009; Balke et al. 2013). Here, we report the discovery of two new species of Bidessini from Peninsula Malaysia. Both species are known from older collections only and therefore we use morphological characters and a pragmatic approach to make tentative generic assignments that lead us to suggest one new genus here.

Material and methods

Drawings of the male genitalia were made based on digital images. The beetles were studied with a Leica MZ 12.5 stereomicroscope at 10–100x and a Scanning Electron Microscope (JSM-5600, JOEL, Tokyo, Japan) at 90–1000X. The terminology to denote the orientation of the genitalia follows Miller and Nilsson (2003). Label data of the type material are cited in quotation marks. The following abbreviations were used in the text: **TL** (total length), **TL-H** (total length without head), and **MW** (maximum width). Google Earth (<http://earth.google.com>) was used to locate localities and the coordinates are given in decimal degree format. Specimens mentioned in this work are deposited in several collections, which are abbreviated in the text as follows:

BMNH	Natural History Museum [former British Museum (Natural History)], London, England
CSR	Collection Saverio Rocchi, Firenze, Italy
CWT	Collection L. J. Wang, Taipei, Taiwan
MNHN	Muséum National d'Histoire Naturelle, Paris, France
NHMW	Naturhistorisches Museum Wien, Austria
TFRI	Taiwan Forestry Research Institute, Taipei, Taiwan
ZSM	Zoologische Staatssammlung München, Germany

Taxonomy

***Rompindessus* Balke, Bergsten & Hendrich, gen. n.**

<http://zoobank.org/2562EA40-B2CA-4909-89B4-8332C6367AE0>

Type species. *Rompindessus jenisi* sp. n. by present designation.

Diagnosis. Of the set of structural features generally used to classify Bidessini, the following combination is present in this taxon: 1) occipital line present; 2) basal pronotal striae present; 3) basal elytral striae absent; 4) sutural line on elytron absent; 5) basal epipleural transverse carina absent; 6) clypeus with thick anterior bead (or margin); 7) longitudinal elytral carina on disc absent; and 8) parameres two-segmented. Moreover, the male pro- and mesotarsus appear stout, and distinctly dilated laterally; the pronotum is comparably long and parallel-sided and the colour of beetle is conspicuously dark orange. This differentiates *Rompindessus* gen. n. from all other Bidessini. In the key to Bidessini genera by Miller and Bergsten (2016) *Rompindessus* keys out to *Platydytes* Biström, 1988, which only occurs in sub-Saharan Africa. Apart from geography, *Rompindessus* can be distinguished from *Platydytes* by the thickly bordered clypeal margin (finely bordered to indistinct in *Platydytes*), the enlarged, laterally expanded male pro- and mesotarsomeres I-III, the dark orange colouration with dark speckles on elytra, and the more discontinuous body outline between pronotum and elytra.

Etymology. Named after the collecting locality, Rompin village, and the ending –dessus as used for many genera in Bidessini.

***Rompindessus jenisi* Balke, Bergsten & Hendrich, sp. n.**

<http://zoobank.org/73B2A0D7-62AC-4891-9510-3504A50D96EC>

Holotype. Male (NHMW). “Malaysia, Pahang, 40km W Rompin, Selendang, 29.4.–6.6., leg. I. Jenis, 1993”.

Type locality. Situated around Malaysia, Pahang, Selendang, between these points 2.622516°N 103.334934°E / 2.609909°N 103.382443°E / 2.643574°N 103.409337°E, it is not known where exactly this particular beetle was discovered. The altitude is around 50 to 250m (Jeniš pers. comm.). According to satellite images in Google Earth, the lower elevations are now heavily transformed into oil palm plantations.

Description. Habitus elongate oval. Measurements: TL = 2.45 mm, TL-H = 2.15 mm; MW = 1.1 mm; elytra 3.5× longer than pronotum.

Colouration. Head, pronotum and elytron dark orange, the latter with few darker speckles (Fig. 1). Ventral side and appendages testaceous.

Surface sculpture. Head with distinct microreticulation posterior of occipital line; finer microreticulation along eyes and clypeus, frons without microreticulation and thus shiny / polished but with few setiferous punctures. Pronotum and elytron with distinct microreticulation and sparse setiferous punctation. Ventral surfaces polished, with faint and sparse setiferous punctation.




Figure 1. *Rompindessus jenisi* gen. et sp. n., holotype, habitus (A), head, pronotum and base of elytra (B). White arrows point to the thick anterior bead of clypeus.




Figure 2. *Rompindessus jeniš* gen. et sp. n., holotype, median lobe of aedeagus, ventral view (A), lateral view (B), paramere, lateral view (C).

Structures. Head with distinct occipital line and broadly beaded clypeus (Fig. 1A). Pronotum with faint lateral bead and distinct basal striae, the latter are curved inwards (Fig. 1). Elytron without basal striae and without sutural line. Basal epipleural transverse carina absent. Metathoracic wings apparently fully developed (not dissected but seen from opening caused by previous removal of the last three ventrites). Pro- and mesotarsus appearing stout as they are distinctly dilated laterally.

Male. Median lobe of aedeagus simply curved (Fig. 2A, B), lateral lobes (parameres) of two parts and of general Bidessini type (e.g. as figured for *Uvarus lacustris* (Say, 1823), *Leiodytes evanescens* (Boheman, 1848), *Platydytes incognitus* Biström, 1988 and *Bidessus unistriatus* (Goeze, 1777) in Biström (1988)) (Fig. 2C).

Female. Unknown.

Etymology. Named after Ivo Jeniš, discoverer of this species. The specific epithet is a substantive in the genitive case.

Distribution. Only known from the type locality.

Habitat. Unknown.

Leiodytes Guignot, 1936

Type species. *Hydroporus evanescens* Boheman, 1848

Diagnosis. Very small, yellowish diving beetles (1.4–2.2 mm for the known species, 2.7 mm with the new one described below) with black markings on elytra. Shape of body globular to elongate, widest in middle, not flattened. 1) Occipital line present; 2) basal pronotal striae present; 3) basal elytral striae present; 4) sutural line on elytron absent; 5) basal epipleural transverse carina absent; 6) clypeus with fore margin narrowly and finely bordered, sometimes unmodified; 7) longitudinal elytral carina on disc absent; and 8) parameres two-segmented.

Includes 27 species (Nilsson 2016), widely distributed in the Afrotropical, Oriental and Palearctic regions.

Leiodytes kualalipis Balke, Wang, Bergsten & Hendrich, sp. n.

<http://zoobank.org/3F824AE1-26C9-4E15-AB50-5974F10ED08A>

Holotype. Male (ZSM). “Malaysia, Pahang, Kuala Lipis, old rubber plantation, iv.1997, Balke & Hendrich”. **Paratypes**, 2 males (CSR, ZSM), 1 female (CSR) “S Vietnam (Cat Tien), 120 km NNE Ho Chi Minh, Cat Tien NP, 3.–15.07.1995, A. Napolov leg.”; 2 males (CWT) “Vietnam, Nam Cat Tien, 200 m, 17-25-VI-1995, leg. Malicky”; 3 males, 2 females (CWT, NHMW, TFRI). “Vietnam, Dong NAI, Nam Cat Tien NP, 120 m, 18-IX-1998, leg. L. J. Wang”.

Type locality. The type locality was around 4.200104°N 102.061570°E, altitude 100m. The entire region is now heavily transformed into oil palm plantations. The paratype locality was in Nam Cat Tien National Park, as a rough reference we obtained a position at the forest border 11.422096°N 107.427578°E.

Description. Habitus elongate oval. Measurements: TL = 2.7 mm; TL-H = 2.4 mm; MW = 1.3 mm; elytra 4.2× longer than pronotum.

Colouration. Beetle dark yellow with few darker basal pronotal markings and darker pattern on elytra (Fig. 3).

Surface sculpture. Head with distinct microreticulation posterior of occipital line; faint to extremely faint microreticulation along eyes and clypeus, frons without microreticulation and thus shiny / polished but with few setiferous punctures. Pronotum and elytron without microreticulation but with dense, coarse setiferous punctation. Ventral surfaces mostly polished, metaventrite, metacoxa and ventrites 1–3 with large punctures bearing short setae, distance between punctures approximately the diameter of a single puncture (similar to *Leiodytes vietnamensis* Wang, Satô & Yang, 1998, p. 165, Fig. 6A in their work).

Structures. Head with faint occipital line and depressed before clypeus leading to the impression that the clypeus is broadly beaded (Figs 3B, 4). Pronotum with faint lateral bead and distinct basal striae, the latter are strongly directed inwards (Figs 3B, 4). Elytron with basal striae but without sutural lines. Basal epipleural transverse carina




Figure 3. *Leiodytes kualalipis* sp. n., holotype, habitus (A), head, pronotum and base of elytra (B).




Figure 4. *Leiodytes kualalipis* sp. n.: male paratype, SEM of head, pronotum and base of elytra.




Figure 5. *Leiodytes kualalipis* sp. n.: female paratype, SEM of microreticulation on metasternum.




Figure 6. *Leiodytes kualalipis* sp. n.: female paratype, SEM of last ventrite with deep suture apically.

absent. Flight wings apparently fully developed (not dissected but seen from opening caused by previous removal of the last three ventrites). Pro- and mesotarsus not conspicuously dilated laterally.

Male. Median lobe of aedeagus in lateral view thin and pointed apically, apical portion spatulate in ventral view (Figs 7A, B); lateral lobes (parameres) of two parts and distally broad, with broad “nose” or broadly twisted (Fig. 7C).

Female. Similar to male, but surface dull due to presence of strong microreticulation dorsally and ventrally (Figs 5, 6).

Diagnosis. This is the largest species of *Leiodytes* and well characterised by its size as the other species in the region are distinctly smaller (below 2.2 mm or even below 2 mm long, see e.g. Régimbart 1899; Wang et al. 1998). Besides the larger size, this new species has a unique feature: the last ventrite has a deep suture apically (Fig. 6). This might be an autapomorphy for the new species.

This is, to our knowledge, the second species reported from the Malayan Peninsula, *L. nicobaricus* (Redtenbacher, 1867) being the other one (Balke et al. 2004; Hendrich et al. 2004).

Etymology. Named after the type locality, Kuala Lipis Town. The species name is a noun in apposition.

Distribution. A species with a wide geographic range, known from the type locality in West Malaysia as well as from Southern Vietnam. The distance between these two localities is roughly 1,000 km measured as a straight line across the Gulf of Thailand.




Figure 7. *Leiodytes kualalipis* sp. n., median lobe of aedeagus, ventral view (A), lateral view (B), paramere, lateral view (C).

Habitat. The holotype was collected in shallow water, among dense layers of rotten leaves, in a shaded forest pool situated in an old rubber plantation overgrown by secondary forest (Fig. 8). The species was associated with the Dytiscidae: *Copelatus* sp., *Hydaticus pacificus pacificus* Aubé, 1838 and *Hyphydrus birmanicus* Régimbart, 1888.

In Vietnam, the species was collected in shallow water made by jeep on a path. Syn-otopically occurring dytiscid species included *Leiodytes nicobaricus* (Redtenbacher, 1867), *Hydroglyphus orientalis* (Clark, 1863) and *Sandracottus maculatus* (Wehncke, 1876).

Note

Nilsson (2016) in the updated World Catalogue of Dytiscidae lists two Bidessini species of unclear generic placement. One of them is *Hydroporus aberrans* Clark, 1863: 426, described from Indonesia (“Java”). The holotype cannot be located in the Natural History Museum London, which houses Clark’s collections. Clark (1863) wrote that he only received one imperfect specimen from Mr. Bowring from Java. It was about 1.6 mm long, with parallel-sided “abdomen” but laterally rounded thorax (narrow anteriorly and posteriorly) and each elytron had eight rows of coarse punctures (Clark referred to them as “striae”). The dorsal colouration was testaceous, with the area between the 1st–6th rows being irregularly rufo-testaceous. This might refer to a species of *Leiodytes*, although we have not seen any species of that genus with rows of coarse serial punctures on the elytron.



Figure 8. Habitat of *Leiodytes kualalipis* sp. n. near Kuala Lipis, West Malaysia. Drawing of beetle by D. Paramonov (Riga).

We can however rule out that any of the new species proposed herein is conspecific with *Hydroporus aberrans*.

The other species of unclear generic placement was *Hydroporus fairmairei* Branden, 1885, which we transfer to *Limbodessus* below.

New combinations and synonymies

Limbodessus curvuplicatus (Zimmermann, 1927)

Fig. 9A, B

Bidessus curvuplicatus Zimmermann, 1927:16 (Samoa).

Limbodessus curvuplicatus (Zimmermann, 1927): Balke and Ribera 2004: 125.

= *Hydroporus dorsoplagiatus* Fairmaire, 1881: 249 (Fiji); preoccupied by Fairmaire (1880: 247), **syn. n.**

= *Hydroporus fairmairei* Branden, 1885: 53 (a replacement name for *H. dorsoplagiatus* Fairmaire, 1881); this replacement name is in turn preoccupied by *Hydroporus fairmairei* Leprieur (1876: 142, currently in *Deronectes*), objective synonym of *Hydroporus dorsoplagiatus* Fairmaire, 1881.

= *Bidessus fijiensis* J. Balfour-Browne, 1944: 99 (Fiji), **syn. n.**

Liodessus fijiensis (J. Balfour-Browne, 1944): Biström 1988: 19.

Limbodessus fijiensis (J. Balfour-Browne, 1944), **comb. n.**

Type material. *Hydroporus dorsoplagiatus*: not located. A loan request was sent to MNHN with request number 66649 on 11.08.2016, and Antoine Mantilleri as well as later Dr. Arnaud Faille searched the collection and did not find potential type material. This might still be stored in the Oberthür collection, however, and might be found at some stage.

Bidessus curvuplicatus: Holotype, male (BMNH) and paratype, male (ZSM), Samoan Is. Mulifanua Upolu, F. Burton and G.H. Hopkins, Type.

Bidessus fijiensis: Holotype, female (BMNH) Namaka, C Fiji 1545, 8.12.43, R.A. Lever, Pres. By Imp. Inst. Ent. B.M. 1945.9, *Bidessus fijiensis* Type! J. Balfour-Browne det.

Additional material. 8 exx (ZSM) Fiji: Viti Levu, Rakiraki, Navara, 50m, 10.xi.2003, 17.416024°S 178.147712°E, Wewalka & Balke (FI 11).

Diagnosis. A stout, yellow to yellow orange *Limbodessus* with slightly enlarged male antenna and distinct angle between base of elytron and base of pronotum (Fig. 9). Female with strongly expanded antennomeres 3–6. Fijian specimens are slightly smaller than the Samoan types: 3.5 mm (Samoa) versus 2.8–3.2 mm (Fiji).

Notes on classification. We have not seen type material of *Hydroporus dorsoplagiatus* but have collected specimens in Fiji that match the description by Fairmaire (1881). At the same time, we found that our Fiji specimens agree with the types of *Limbodessus curvuplicatus* (Zimmermann, 1927) from Samoa. Moreover, *Bidessus fijiensis* J. Balfour-Browne, 1944, later moved to *Liodessus* by Biström (1988), and which we here move to *Limbodessus*, agrees well with our newly collected Fijian specimens and thus establishes another junior synonym to *Hydroporus fairmairei*, the replacement name of *Hydroporus dorsoplagiatus*. However, since the replacement name *Hydroporus fairmairei* suggested by Branden (1885) is also a junior homonym (see above), the oldest valid synonym, *Limbodessus curvuplicatus* (Zimmermann, 1927), becomes the valid name.

Distribution. Samoa, Fiji.




Figure 9. *Limbodesus curviplicatus*: male paratype of *Bidessus curviplicatus* Zimmermann, 1927 from Samoa (**A**) and male specimen from Fiji (**B**).

Acknowledgements

We thank Ivo Jeniš (Prague) for valuable comments on his collecting sites in West Malaysia, Christine Taylor (BMNH, London), Manfred Jäch (NHMW, Vienna) and Saverio Rocchi (Firenze, Italy) for loan of specimens, and Antoine Mantilleri (MNHN Paris) and Arnaud Faille (MNHN Paris) for their help searching for type material. Jiří Hájek (Prague) and Olof Biström (Helsinki) kindly commented on earlier versions of this work. Ignacio Ribera and Helena Shaverdo as well as editor Mariano Michat reviewed this paper and helped with many useful comments. We also very warmly thank D. Paramonov (Riga) for the drawing of the new *Leiodytes* used in Figure 8, and Katja Neven (Munich) for the digital inking of the line drawings used here.

References

- Balfour-Browne J (1944) New and interesting Dytiscidae (Coleoptera) from Fiji. Proceedings of the Royal Entomological Society of London (B) 13: 97–100. <https://doi.org/10.1111/j.1365-3113.1944.tb00796.x>
- Balke M, Ribera I (2004) Jumping across Wallace's line: *Allodessus* Guignot and *Limbodessus* Guignot revisited (Coleoptera: Dytiscidae, Bidessini) based on molecular phylogenetic and morphological data. Australian Journal of Entomology 43: 114–128. <https://doi.org/10.1111/j.1440-6055.2004.00415.x>
- Balke M, Jäch MA, Hendrich L (2004) Coleoptera. In: Yule CM, Yong HS (Eds) Freshwater Invertebrates of the Malaysian Region. Akademi Sains Malaysia, Kuala Lumpur, 555–609.
- Balke M, Hendrich L, Mazzoldi P, Biström O (2002) *Borneodessus zetteli*, new genus and new species, notes on biogeography and a checklist of Dytiscidae from Borneo (Insecta: Coleoptera: Dytiscidae). Journal of Natural History 36: 963–978. <https://doi.org/10.1080/00222930110062345>
- Balke M, Warikar E, Toussaint EFA, Hendrich L (2013) *Papoadessus baueri* spec. nov. from Biak Island, Papua (Coleoptera, Dytiscidae, Hydroporinae). Spixiana 36(2): 283–288.
- Balke M, Ruthensteiner B, Warikar EL, Neven K, Hendrich L (2015) Two new species of *Limbodessus* diving beetles from New Guinea - short verbal descriptions flanked by online content (digital photography, μ CT scans, drawings and DNA sequence data). Biodiversity Data Journal 3: e7096. <https://doi.org/10.3897/bdj.3.e7096>
- Biström O (1988) Generic review of the Bidessini (Coleoptera, Dytiscidae). Acta Zoologica Fennica 184: 1–41.
- Branden C van den (1885) Catalogue des coléoptères carnassiers aquatiques (Halipilidae, Amphizoidae, Pelobiidae et Dytiscidae). Annales de la Société Entomologique de Belgique 29: 5–116.
- Clark H (1863) Descriptions of new East-Asiatic species of Halipilidae and Hydroporidae. The Transactions of the Royal Entomological Society of London (3) 1: 417–428.
- Fairmaire L (1881) Essai sur les coléoptères des Îles Viti (Fidji). Annales de la Société Entomologique de France (6) 1: 243–318.
- Hendrich L, Balke M (2009) *Kakadudessus tomweiri*, a new genus and species of diving beetle from tropical northern Australia, based on molecular phylogenetic and morphological data (Coleoptera, Dytiscidae, Bidessini). Zootaxa 2134: 49–59.
- Hendrich L, Balke M, Yang CM (2004) Aquatic Coleoptera of Singapore – Species Richness, Ecology and Conservation. The Raffles Bulletin of Zoology 52: 97–141.
- Leprieur C-E (1876) Synopsis des *Hydroporus* du groupe de l'*opatrinus*. Bulletin Bimensuel des Séances de la Société Entomologique de France 1876 No. 79 (Séance du 12 Juillet 1876): 141–143.
- Miller KB (2016) *Novadessus viracocha*, a new genus and species of Bidessini Sharp from Peru (Coleoptera, Adephaga, Dytiscidae, Hydroporinae). ZooKeys 623: 125–130. <https://doi.org/10.3897/zookeys.623.10018>
- Miller KB, Bergsten J (2016) Diving Beetles of the World. Systematics and Biology of the Dytiscidae. Johns Hopkins University Press, Baltimore, 320 pp.

- Miller KB, Nilsson AN (2003) Homology and terminology: Communicating information about rotated structures in water beetles. *Latissimus* 17: 1–4.
- Miller KB, Short AEZ (2015) *Belladessus* Miller and Short (Coleoptera: Dytiscidae: Hydrophorinae: Bidessini), new genus for two new species from northern South America: Parthenogenetic diving beetles? *The Coleopterists Bulletin* 69: 498–503. <https://doi.org/10.1649/0010-065X-69.3.498>
- Nilsson AN (2016) A World Catalogue of the Family Dytiscidae, or the Diving Beetles (Coleoptera, Adephaga). Version 1, 300 pp. <http://www.norrent.se> and www.waterbeetles.eu
- Régimbart M (1899) Révision des Dytiscidae de la région Indo-Sino-Malaise. *Annales de la Société Entomologique de France* 68: 186–367.
- Wang LJ, Sato M, Yang PS (1998) Note on the genus *Leiodytes* Guignot (Coleoptera, Dytiscidae) with three new species from Taiwan, Thailand and Vietnam. *Japanese Journal of Systematic Entomology* 4: 161–171.
- Zimmermann A (1927) Dytiscidae. *Insects of Samoa* 4: 15–19.