

# *Liliocerus groehni* sp. n.: the first authentic species of Criocerinae (Coleoptera, Chrysomelidae) from Baltic amber

Andris Bukejs<sup>1</sup>, Michael Schmitt<sup>2</sup>

<sup>1</sup> Institute of Life Sciences and Technologies, Daugavpils University, Vienības Str. 13, Daugavpils, LV-5401, Latvia <sup>2</sup> Ernst-Moritz-Arndt-Universität, Allgemeine & Systematische Zoologie, Soldmannstr. 14, 17489 Greifswald, Germany

Corresponding author: Michael Schmitt ([michael.schmitt@uni-greifswald.de](mailto:michael.schmitt@uni-greifswald.de))

---

Academic editor: J. Santiago-Blay | Received 4 August 2016 | Accepted 9 September 2016 | Published 19 September 2016

---

<http://zoobank.org/8648005D-9B75-4905-AGA5-027830B5E95E>

---

**Citation:** Bukejs A, Schmitt M (2016) *Liliocerus groehni* sp. n.: the first authentic species of Criocerinae (Coleoptera, Chrysomelidae) from Baltic amber. ZooKeys 618: 67–77. doi: 10.3897/zookeys.618.10085

---

## Abstract

Based on a single well-preserved specimen from Eocene Baltic amber, *Liliocerus groehni* sp. n. is described and illustrated using phase-contrast X-ray microtomography. It is the first described species of Criocerinae (Coleoptera: Chrysomelidae) from Baltic amber. A check-list of fossil Criocerinae is provided. Placement of *Criocerus pristiana* (Germar, 1813) is discussed, this species is removed from Criocerinae and placed in Coleoptera *incertae sedis*.

## Keywords

Taxonomy, palaeontology, shining leaf beetles, new species, *Criocerus pristiana*, fossil resin, Tertiary, Eocene

## Introduction

The subfamily Criocerinae (shining leaf beetles) contains ca. 1500 extant species (Schmitt 1996) in 20 genera (Seeno and Wilcox 1982), of which 211 species in 6 genera are recorded from the Palaearctic region (Schmitt 2010). The genus *Liliocerus* Reitter, 1913 comprises ca. 170 extant species distributed over the temperate, subtropical and tropical regions of the Palaearctic, Orientalis, Aethiopsis including Madagascar, and the Australis, and was introduced to North America by man (Monrós 1960, plus records from the Zoological Record up to present as taken from the Index to Organism Names).

Shining leaf beetles are rarely represented in fossil material and especially in Baltic amber (Table 1). Two fossil species from Baltic amber were mistakenly described within Criocerinae. *Electrolema baltica* Schaufuss, 1892 was described as member of Criocerinae but later transferred to Hispinae (Korschefsky 1939). According to modern classification (Staines 2012), it is placed in Gonophorini Chapuis, 1875 within Cassidinae. Another species, *Crioceris pristina* (Germar, 1813) originally described as *Criocerina* (Germar 1813) was mentioned within Criocerinae (e.g. Giebel 1856a, 1856b; Spahr 1981; Santiago-Blay 1994). In our opinion it is not a member of the Criocerinae (see Discussion).

*Crioceris* sp. and *Lema* sp. were mentioned from Eocene Baltic amber without detailed species descriptions (Hope 1836; Giebel 1856; Menge 1856; Scudder 1885, 1886, 1891; Handlirsch 1908; Klebs 1910; Bachofen-Echt 1949; Larsson 1978; Spahr 1981; Santiago-Blay 1994; Poinar 1999). In the current paper, the first extinct species of Criocerinae from Baltic amber is described, figured, and compared with extant species using phase-contrast X-ray microtomography.

## Material and methods

The specimen is included in an amber piece that was polished by hand and faceted on their sides, allowing improved views of the included specimens. The material examined is deposited in the collection of the Geological-Palaeontological Institute of the University of Hamburg, Germany [GPIH], as part of the collection of Carsten Gröhn.

Observations were made using a Nikon SMZ 745T stereomicroscope. Photographs were taken using a Canon EOS 70D with a 100 mm macro lens, and a Canon EOS 5D with the Canon MP E 65 mm macro lens in a visionary digital bk plus lab system by Dun Inc. The microCT-images were produced by means of an Xradia Micro XCT-200 (Carl Zeiss X-ray Microscopy Inc.), using the 4x object lens units, at 30kV and 4W, with a pixel size of 5.36 µm. Tomography projections were reconstructed using the reconstruction software provided by XRadia. Volume rendering of image stacks was performed by using Amira 5.6.0 (FEI Visualization Science Group, Burlington, USA) using the “Volren” or “Voltex” function.

## Systematic Palaeontology

### Chrysomelidae Latreille, 1802

### Criocerinae Latreille, 1804

### Criocerini Latreille, 1804

### *Lilioceris* Reitter, 1913

The specimen considered here was assigned to the family Chrysomelidae based on the pseudoteramerous tarsi and the lack of a rostrum and of antennae not inserted on

pronounced tubercles, to the subfamily Criocerinae because the prothorax does not bear side borders and the frons has distinct diverging grooves behind the antennal insertions, and to the genus *Liliocerus* based on (1) free tarsal claws and (2) divided vertex separated from the neck by a dorsal constriction.

***Liliocerus groehni* sp. n.**

<http://zoobank.org/FD228756-DF55-47E3-BDE4-F0D3937A8A1E>

Figs 1–5

**Type material.** Holotype: Nr. “C 8130” [GPIH]; female. A rather complete beetle (missing apical antennomere of left antenna, and tarsomeres 4 and 5 of left meso- and metatarsi) is included in a small, transparent yellow amber piece (length about 20 mm, width 12 mm, and maximum thickness 5 mm). Syninclusions: one specimen of Nematocera (Diptera), and few stellate Fagaceae trichomes (Figs 1 and 2).

**Type strata.** Baltic amber, mid-Eocene to Upper Eocene.


**Type locality.** Yantarny settlement (formerly Palmnicken), Sambian (Samland) Peninsula, the Kaliningrad region, Russia.

**Differential diagnosis.** Head, body, and elytra of *Liliocerus groehni* sp. n. appear unicolorous black and thus similar to the extant species *L. hitam* Mohamedsaid, 1990 from Borneo, which differs from the new species in (1) the shape of the pronotum (distinctly longer than wide with its constriction at the middle), (2) metaventrite glabrous in the middle, (3) pubescent scutellum, (4) impunctate elytra (with few moderately large punctures at base only), (5) vertex with sparse pubescence, (6) a distinct conical neck between head and pronotum, and (7) a larger body (10 mm).

Additionally, the extant species *L. lilii* Scopoli, 1763 and *L. merdigera* Linnaeus, 1758 from Baltic region differ from *L. groehni* sp. n. in having (1) a pronotum with a longitudinal row of punctures medially, (2) metaventrite, metepisternum and ventrites of abdomen almost glabrous or with very sparse pubescence, and (3) pronotum and elytra rufous to red.

**Description.** Holotype. Body length 7.1 mm, maximum width 4.1 mm; elongate, subparallel, moderately convex dorsally and ventrally, unicolorous black, glabrous dorsally.

Head hypognathous, transverse, widest across eyes, together with eyes nearly as wide as pronotum, strongly constricted behind the eyes forming a neck (Fig. 3); shiny, hairless and without distinct punctures dorsally. Compound eyes large, strongly convex, deeply and acutely notched at antennal insertions; distance between eyes nearly as wide as transverse diameter of one eye. Frontal grooves deep, crossed forming X. Vertex convex, hairless, with median longitudinal groove. Genae large, with sparse pubescence. Antennae poorly visible because of a beetle location in amber piece. Antennae robust, covered with fine pubescence, moderately long, extending nearly to basal one-fourth of elytra, slightly widened apically; antennomere 2 shortest, about 0.4 times as long as antennomere 3, antennomere 4 slightly longer than antennomere




**Figures 1–2.** *Liliocerus groehni* sp. n., holotype: **1** habitus, dorsal view **2** habitus, lateral view. Scale bars: 1 mm.

3, antennomeres 5–7 subequal in length, antennomere 5 about 1.2 times as long as antennomere 4.

Pronotum nearly as long as wide, deeply constricted medially, distinctly narrower than elytra, widest in anterior one-third; impunctate, shiny; disc flattened, with an arcuate transverse depression subbasally (Figs 3 and 4). Anterior margin straight me-





**Figures 3–5.** *Liliocercis groehni* sp. n., holotype, microCT images: **3** habitus, fronto-lateral view, showing the pronounced arcuate constriction behind the disk of the pronotum **4** elytra, dorsal view **5** details of head and prothorax, dorsal view. Not reproduced to the same scale.

dially; posterior margin convex; lateral margins rounded anteriorly and strongly constricted just behind middle; all margins not bordered. Anterior and posterior angles obtusely rounded.

Scutellum large, triangular; apparently hairless and impunctate. Elytra subparallel, widest in the middle, about 1.5 times as long as wide; humeri prominent. Elytral punctures small and dense (in basal one-third deeper), arranged in rows; scutellar row present, short; intervals flat, only at apices weakly convex.

Metaventrite with sparse, fine pubescence; metepisternum and metepimeron densely covered with fine, short, semierect pubescence. Abdomen with sparse, fine pubescence.

Legs moderately long, covered with fine pubescence. Femora spindle-shaped; tibiae slightly curved, dilated apically. Tarsi long, about 0.7 times as long as tibia; metatarsomeres 1–2 subequal in length, distinctly dilated apically, metatarsomere 3 deeply bilobed, metatarsomere 4 subcylindrical, narrow. Tarsal claws free, not fused at base.

The interior of the abdomen does not contain any identifiable structure, as revealed by the microCT-analysis. No traces of an aedeagus could be found, and none of the smaller particles – all covered with homogeneous material – could be addressed as the spermatheca.

**Derivatio nominis.** This new species is named after Carsten Gröhn (Glinde, Germany) – he enabled us to study this specimen.

## Discussion

The specimen of *Liliocerus groehni* sp. n. appears externally complete. However, the fact that we found no traces of internal structures in the abdomen, especially of an aedeagus, does most probably mean that (1) the specimen was a female, and (2) that it remained openly accessible for scavengers and/or detritivores before it was covered by resin. This could also provide a possible reason for its black appearance as the dead individual might have been exposed to humic acids before being fossilised. If this should be the case, the live animal had most probably a habitus similar to the extant lily beetles. Actually, *L. groehni* sp. n. is hardly distinguishable from extant *Liliocerus*-species. In this respect, the new species is quite normal. Hennig (1966) wrote that it is „a long known fact” [„eine altbekannte Tatsache”] that the morphological differences between fossils from Baltic amber and their extant relatives are only minute”.

Santiago-Blay (1994) mistakenly listed two fossil species as members of Criocerinae: *Lema pervetusta* Cockerell, 1921 and *Lema pulchella* Förster, 1891. *Lema pervetusta* was described from Bridgerian lacustrine shale (Eocene, 50.3–48.6 Ma) of the Green River Formation of Colorado, USA (Cockerell 1921), but according to Linsley (1942) this species belongs to the longhorn beetle genus *Clytus* Laicharting, 1784 (Cerambycidae). *Lema pulchella* was described from Oligocene lacustrine (33.9–28.4 Ma) of Riedisheim, Mulhouse, France (Förster 1891), but according to Théobald (1937) it belongs to the weevil genus *Phyllobius* Germar, 1824 (Curculionidae).

Few Quaternary sub-fossil records contain specimens of the extant species: *Lema cyanella* (Linnaeus, 1758) from La Taphanel, Massif Central, France (Ponel and Coope

Table 1. Check-list of records of fossil and sub-fossil Criocerinae.

Taxon	References	Fossil Type	Locality	Age
Criocerinae	Bachofen-Echt 1949; Handlirsch 1925; Spahr 1981	Baltic amber	Kaliningrad region (Russia)	37.2–33.9 Ma
Criocerinae	Hayashi et al. 2002	poorly lithified peat	Mizozone Formation, Yoshimatsucho, Kagoshima Prefecture (Japan)	0.1–0.0 Ma
<i>Crioceridea dubia</i>	Wickham 1912, 1913, 1914a, 1920; Santiago-Blay 1994	lacustrine shale	Florissant, Colorado (USA)	37.2–33.9 Ma
<i>Criocaris margarum</i>	Oustalet 1874; Handlirsch 1908; Théobald 1937; Santiago-Blay 1994	lacustrine shale	Aix-en-Provence (France)	28.4–23.0 Ma
<i>Criocaris vetusta</i>	Heer 1865 ( <i>Lema</i> ); Handlirsch 1908 ( <i>Lema</i> ); Cockerell 1921	lacustrine shale	Oeningen (Germany)	12.7–11.6 Ma
<i>Criocaris</i> sp.	Hope 1836; Giebel 1856a; Menge 1856; Scudder 1885, 1886, 1891; Handlirsch 1908; Klebs 1910; Bachofen-Echt 1949; Spahr 1891	Baltic amber	Kaliningrad region (Russia)	37.2–33.9 Ma
<i>Lema evanescens</i>	Wickham 1910, 1913, 1914a, 1920	lacustrine shale	Florissant, Colorado (USA)	37.2–33.9 Ma
<i>Lema fortior</i>	Wickham 1914a, 1920	lacustrine shale	Florissant, Colorado (USA)	37.2–33.9 Ma
<i>Lema lesquerexi</i>	Wickham 1914b, 1920	lacustrine shale	Florissant, Colorado (USA)	37.2–33.9 Ma
<i>Lema tumulata</i>	Heyden and Heyden 1865; Handlirsch 1908	terrestrial siliclastic	Salzhausem (Germany)	15.9–11.6 Ma
<i>Lema</i> sp.	Scudder 1885, 1886, 1891; Helm 1896; Handlirsch 1908; Larsson 1978; Spahr 1981; Poinar 1999	Baltic amber	Kaliningrad region (Russia)	37.2–33.9 Ma
<i>Lema</i> sp.	Pearson 1962	lacustrine shale	West Cumberland (England)	0.1–0.0 Ma
(?) <i>Lema</i> sp.	Kiselev and Nazarov 2009	unlithified siliclastic sediments	Achchagyat-Allaikha Yana-Indigirka Lowland, nord-east Siberia (Russia)	0.1–0.0 Ma
<i>Liliocerus groehni</i>	present paper	Baltic amber	Kaliningrad region (Russia)	37.2–33.9 Ma
<i>Coleoptera incertae sedis</i>				
<i>Criocaris pristina</i>	Germar 1813 ( <i>Criocarina</i> ); Giebel 1856a, 1856b; Schleichendal ( <i>Criocarina</i> ); Handlirsch 1908; Spahr 1981; Santiago-Blay 1994; Poinar 1999	Baltic amber	Kaliningrad region (Russia)	37.2–33.9 Ma

1990); *Lema trilinea* White, 1981 from late Quaternary Kaetan Cave, Colorado Plateau, Colorado, USA (Elias and Van Devender 1992); and *Oulema obscura* (Stephens, 1831) from the Holocene of Belarus (Nazarov 1984). These records are not mentioned in the current list (Table 1). None of the records of “Criocerinae”, “*Crioceris* sp.”, or “*Lema* sp.” from Baltic amber listed in Table 1 can be assigned to a certain species of shining leaf beetles.

Germar (1813) described *Criocerina pristina* from Baltic amber. Later this species was mentioned as *Crioceris pristina* (Germar, 1813) within Criocerinae (e.g. Giebel 1856a, 1856b; Spahr 1981; Santiago-Blay 1994). The correct subfamily and family placement of *Crioceris pristina* is doubtful in our opinion. According to the original description (Germar 1813: 14), this fossil species has antennae with a club (similar as in members of the genus *Anobium*) [“... Die Fühler von etwas mehr als halber Körperlänge, roth, and der Spitze dunkler, das erste Glied kurz und dick, vor den Augen auf der Stirn eingesetzt, das folgende Glied klein, kugelförmig, die nun folgenden sechs Glieder sehr klein und dicht zusammengedrängt, dass sie als blosse Ringe erscheinen, die drey letzten Glieder lang und dicker, fast wie bey *Anobium* gebaut, sie machen zusammen zwei Drittheil der ganzen Fühlerlänge aus ... ”]. In addition, Germar gave the length of this specimen as  $1\frac{1}{8}$  lines = 2.54 mm. This would be an extremely low value for a species of *Lilioceris*. All extant species are described as being longer than 5 mm. Germar mentioned that his *Criocerina pristina* resembled „*Crioceris testacea* Fabr.”, of which he said it were six times larger – i.e. ca. 1.5 cm. The species Fabricius described as *Crioceris testacea* (Fabricius 1787: 87/88) is currently listed under *Aulacophora indica* (Gmelin, 1790), Galerucinae (Mohamedsaid 2009), and its lectotype is depicted in Lee and Beenen (2015, Figs 42 & 43). We conclude that „*Crioceris pristina*” is actually not a criocerine beetle nor a member of the family Chrysomelidae. Instead, we suggest that it should be better placed as Coleoptera *incertae sedis*. This conclusion leaves *Lilioceris groehni* sp. n. as the first beetle species from Baltic amber that we can classify with certainty as a member of the Coleoptera Chrysomelidae Criocerinae.

## Acknowledgements

The authors are grateful to Carsten Gröhn (Glinde, Germany) for the loan of interesting fossil material, to Vitalii I. Alekseev (Kaliningrad, Russia) for valuable suggestions and constructive advice, to Peter Michalik (Greifswald, Germany) for invaluable help with macro-imaging and the microCT-analysis, and the German Science Foundation (DFG) for financial support (INST 292/119-1 FUGG, INST 292/120-1 FUGG).

## References

Bachofen-Echt A (1949) Der Bernstein und seine Einschlüsse. Springer, Wien, 204 pp. doi: 10.1007/978-3-7091-2303-4



- Cockerell TDA (1921) Eocene Insects from the Rocky Mountains. Proceedings of the United States National Museum 57: 233–260. doi: 10.5479/si.00963801.57-2313.233
- Elias SA, Van Devender TR (1992) Insect fossil evidence of Late Quaternary environments in the northern Chihuahuan desert of Texas and New Mexico: comparisons with the paleobotanical record. The southwestern Naturalist 37(2): 101–116. doi: 10.2307/3671658
- Fabricius JC (1787) Mantissa insectorum sistens eorum species nuper detectas adiectis characteribus genericis, differentiis specificis, emendationibus, observationibus, vol. 2. C.G. Proft, Copenhagen, 382 pp. doi: 10.5962/bhl.title.36471
- Förster B (1891) Die Insekten der “Plattigen Steinmergels” von Brunstatt. Abhandlungen zur Geologischen Specialkarte von Elsass-Lothringen 3: 335–593.
- Germar EF (1813) Insekten in Bernstein eingeschlossen, beschrieben aus dem akademischen Mineralien—Cabinet zu Halle. Magazin der Entomologie 1: 11–18.
- Giebel CGA (1856a) Fauna der Vorwelt mit steter Berücksichtigung der lebenden Thiere. Gliederthiere. Zweite Band: Insecten und Spinnen. F.A. Brockhaus, Leipzig, XVII + 511 pp.
- Giebel CGA (1856b) Geologische Übersicht der vorweltlichen Insecten. Zeitschrift für die gesammten Naturwissenschaften 8: 174–188.
- Handlirsch A (1908) Die fossilen Insekten und die Phylogenie der rezenten Formen. Ein Handbuch für Paläontologen und Zoologen. Wilhelm Engelmann, Leipzig, IX + 1430 pp.
- Handlirsch A (1925) Paläontologie. In: Schröder CWM (Ed.) Handbuch der Entomologie, Volume 3. Gustav Fischer, Jena, 117–306.
- Hayashi M, Yahiro K, Kitabayashi E (2002) Late Pleistocene insects from the Mizozono Formation in Yoshimatsu-cho, Kagoshima Prefecture, Japan. Bulletin of the Mizunami Fossil Museum 29: 161–168.
- Heer O (1865) Die Urwelt der Schweiz. Zürich, Friedrich Schultheß, 622 pp.
- Helm O (1896) Beiträge zur Kenntniss der Insecten des Bernsteins. Schriften der Naturforschenden Gesellschaft in Danzig (Anlage C) N. S. 9: 220–231.
- Hennig W (1966) *Fannia scalaris* Fabricius, eine rezente Art im Baltischen Bernstein? (Diptera: Muscidae). Stuttgarter Beiträge zur Naturkunde Serie A (Biologie) 150: 1–12.
- Heyden C, Heyden L (1865) Fossile Insekten aus der Braunkohle von Salzhausen. Palaeontographica 14(1): 31–35.
- Hope FW (1836) Observations on succinic insects. The Transactions of the Royal Entomological Society of London 1: 133–147. doi: 10.1111/j.1365-2311.1838.tb00157.x
- Index to Organism Names: <http://www.organismnames.com/> [accessed August 1, 2016]
- Kiselev SV, Nazarov VI (2009) Late Cenozoic insects of northern Eurasia. Paleontological Journal 43(7): 732–850. doi: 10.1134/S0031030109070016
- Klebs R (1910) Über Bernsteineinschlüsse in allgemeinen und die Coleopteren meiner Bernsteinsammlung. Schriften der Physikalisch-ökonomischen Gesellschaft zu Königsberg 51: 217–242.
- Korschefsky R (1939) Abbildungen und Bemerkungen zu vier Schaufuß’schen Coleopteren aus dem deutschen Bernstein. Arbeiten über morphologische und taxonomische Entomologie aus Berlin-Dahlem 6(1): 11–12.
- Larsson SG (1978) Baltic Amber – a Palaeobiological Study. Volume 1. Scandinavian Science Press Ltd., Klampenborg, 192 pp.

- Lee C-F, Beenen R (2015) Revision of the genus *Aulacophora* from Taiwan (Coleoptera: Chrysomelidae: Galerucinae). *Zootaxa* 3949(2): 151–190. doi: 10.11646/zootaxa.3949.2.1
- Linsley EG (1942) A review of the fossil Cerambycidae of North America. *Proceedings of the New England Zoölogical Club* 21: 17–42.
- Menge A (1856) Lebenszeichen vorweltlicher, im Bernstein eingeschlossener Thiere. Programm der öffentlichen Prüfung der Schüler der Petrischule. A.W. Kafemann, Danzig, 32 pp.
- Mohamedsaid MS (1990) A New Species of *Lilioceria* from Sabah, Malaysia (Chrysomelidae: Criocerinae). *Entomological Review of Japan* 45(2): 93–95.
- Monrós F (1960) Los Géneros de Chrysomelidae, *Opera Lilloana* III (1959), Tucumán, 337 pp.
- Nazarov VI [Назаров ВИ] (1984) Реконструкция ландшафтов Белоруссии по палеоэнтомологическим данным (антропоген) [Reconstruction of the landscapes of Byelorussia on the basis of palaeoentomological data]. *Труды Палеонтологического института Академии наук СССР* [Trudy paleontologicheskogo instituta Akademii Nauk SSSR] 205: 1–104. [In Russian]
- Oustalet ME (1874) Recherches sur les insectes fossiles des terrains Tertiaires de la France. Deuxième partie. Insectes fossiles d'Aix en Provence. *Annales des Sciences Géologiques* 5(2): 1–347.
- Pearson RG (1962) The Coleoptera from a late-glacial deposit at St. Bees, West Cumberland. *Journal of Animal Ecology* 31: 129–150. doi: 10.2307/2335
- Poinar GO Jr. (1999) Chrysomelidae in fossilized resin: behavioural inferences. In: Cox ML (Ed.) *Advances in Chrysomelidae Biology*. Backhuys Publishers, Leiden, the Netherlands, 1–16.
- Ponel P, Coepe GR (1990) Late glacial and Early Flandrian Coleoptera from La Taphanel, Massif Central, France: climatic and ecological interpretations. *Journal of Quaternary Science* 5: 235–249. doi: 10.1002/jqs.3390050306
- Santiago-Blay JA (1994) Paleontology of leaf beetles. In: Jolivet PHA, Cox ML, Petitpierre E (Eds) *Novel Aspects of the Biology of Chrysomelidae*. Kluwer Academic Publishers, the Netherlands, 1–68. doi: 10.1007/978-94-011-1781-4\_1
- Schaufuss C (1892) Preussens Bernstein-Käfer. Neue Formen aus der Helm'schen Sammlung im Danziger Provinzialmuseum. *Berliner Entomologische Zeitschrift* 36(1): 53–64. doi: 10.1002/mmnd.18910360111
- Schlechtendal D (1888) Mittheilungen über die in der Sammlung aufbewahrten Originale zu Germar's: "Insekten in Bernstein eingeschlossen" mit Rücksicht auf Giebels "Fauna der Vorwelt". *Zeitschrift für Naturwissenschaften* 61: 473–491.
- Schmitt M (1996) The phylogenetic system of the Chrysomelidae - history of ideas and present state of knowledge. In: Jolivet PHA, Cox ML (Eds) *Chrysomelidae Biology, Volume 1: The Classification, Phylogeny and Genetics*. SPB Publishing, the Netherlands, 57–96.
- Schmitt M (2010) Criocerinae. In: Löbl I, Smetana A (Eds) *Catalogue of Palearctic Coleoptera, Volume 6*. Apollo Books, Stenstrup, 359–368.
- Scudder SH (1885) Insecta. Insekten. In: Zittel KA (Ed.) *Handbuch der Palaeontologie, I Abtheilung. Palaeozoologie, II Band. Mollusca und Arthropoda*. Oldenbourg, München & Leipzig, 747–831.

- Scudder SH (1886) Systematic review of our present knowledge of fossil insects, including Myriapods and Arachnids. *Bulletin of the United States geological Survey* 31: 1–128.
- Scudder SH (1891) Index to the known fossil insects of the world including Myriapods and Arachnids. *Bulletin of the United States geological Survey* 71: 1–744.
- Seeno TN, Wilcox JA (1982) Leaf beetle genera (Coleoptera: Chrysomelidae). *Entomography* 1: 1–221.
- Spahr U (1981) Systematischer Katalog der Bernstein und Kopalkäfer (Coleoptera). *Stuttgarter Beiträge zur Naturkunde, Serie B (Geologie und Paläontologie)* 80: 1–107.
- Staines CL (2012) Catalog of the hispines of the world (Coleoptera: Chrysomelidae: Cassidinae). [http://entomology.si.edu/Collections\\_Coleoptera-Hispines.html](http://entomology.si.edu/Collections_Coleoptera-Hispines.html)
- Théobald N (1937) Les insectes fossiles des terrains oligocènes de France. PhD thesis, Imprimerie George Thomas, University of Nancy, France.
- Wickham HF (1910) New fossil Coleoptera from Florissant, with notes on some already described. *American Journal of Science* 29: 47–51. doi: 10.2475/ajs.s4-29.169.47
- Wickham HF (1912) A report of some recent collections of fossil Coleoptera from the Miocene Shale of Florissant. *Bulletins from the Laboratory of Natural History of the State University of Iowa* 6(3): 1–38.
- Wickham HF (1913) Fossil Coleoptera from the Wilson Ranch near Florissant, Colorado. *Bulletins from the Laboratory of Natural History of the State University of Iowa* 6(4): 3–29.
- Wickham HF (1914a) New Miocene Coleoptera from Florissant. *Bulletin of the Museum of Comparative Zoology, Harvard College* 53: 423–494.
- Wickham HF (1914b) Twenty new Coleoptera from the Florissant Shales. *Transactions of the American Entomological Society* 40: 257–270.
- Wickham HF (1920) Catalogue of the North American Coleoptera described as fossils. In: Leng CW (Ed.) *Catalogue of the Coleoptera of America North of Mexico*. John D. Sherman Jr., Mount Vernon, N.Y., 349–365.