

Complete mitochondrial genomes of the slugs *Deroceras laeve* (Agriolimacidae) and *Ambigolimax valentianus* (Limacidae) provide insights into the phylogeny of Stylommatophora (Mollusca, Gastropoda)

Te Zhao¹, Nan Song¹, Xingyu Lin¹, Yang Zhang¹

¹ College of Plant Protection, Henan Agricultural University, Zhengzhou 450002, China
Corresponding author: Nan Song (songnan@henau.edu.cn)



Academic editor: Martin Haase

Received: 1 March 2023

Accepted: 13 July 2023

Published: 31 July 2023

ZooBank: <https://zoobank.org/767DB487-06EF-4066-A504-D4292C2FAAED>

Citation: Zhao T, Song N, Lin X, Zhang Y (2023) Complete mitochondrial genomes of the slugs *Deroceras laeve* (Agriolimacidae) and *Ambigolimax valentianus* (Limacidae) provide insights into the phylogeny of Stylommatophora (Mollusca, Gastropoda). ZooKeys 1173: 43–59. <https://doi.org/10.3897/zookeys.1173.102786>

Copyright: © Te Zhao et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Abstract

In this study, we sequenced two complete mitogenomes from *Deroceras laeve* and *Ambigolimax valentianus*. The mitogenome of *Ambigolimax valentianus* represented the first such data from the family Limacidae. The lengths of the mitogenomes of *Deroceras laeve* and *Ambigolimax valentianus* were 14,773 bp and 15,195 bp, respectively. The entire set of 37 mitochondrial genes were identified for both mitogenomes. Compared with the mitogenome of *Achatina fulica*, the *trnP-trnA* tRNA cluster was rearranged in both *Deroceras laeve* and *Ambigolimax valentianus*. The secondary structures of tRNA and rRNA genes for the two species were predicted. Phylogenetic analyses based on amino acid sequences supported (1) monophyly of Stylommatophora, (2) division of Stylommatophora into the ‘achatinoïd’ clade (i.e., the suborder Achatinina) and the ‘non-achatinoïd’ clade (i.e., the suborder Helicina), (3) placement of the Orthurethra in the ‘non-achatinoïd’ clade, and (4) monophyly of each of the superfamilies Helicoidea, Urocoptoidea, Succineoidea, Arionoidea, Pupilloidea and Limacoidea. The exemplars of Helicidae, Philomycidae and Achatinellidae displayed many more mitochondrial gene rearrangements than other species of Stylommatophora.

Key words: Gene rearrangement, Limacoidea, mitogenome, next generation sequencing, phylogeny

Introduction

Typically, the metazoan mitochondrial genome (mitogenome) is a closed-circular and small (15–20 kb) genome encoding 13 protein-coding genes (PCGs), 22 transfer RNAs (tRNAs) and two ribosomal RNAs (rRNAs) (Boore 1999). Mitogenomes have been widely used to resolve phylogenetic relationships within molluscs (e.g., He et al. 2016; Minton et al. 2016; Xie et al. 2019; Guzmán et al. 2021). With the development of new sequencing technologies and the significantly decreased cost of next generation sequencing, the numbers of available mitogenomes has increased rapidly. Molluscs are the second largest

phylum next to Arthropoda, with about 52,500 extant species (Ponder et al. 2020). However, relatively few mitogenomes are available for this group. Within Mollusca, Stylommatophora includes the vast majority of terrestrial snails and slugs. As of December 2022, only 58 stylommatophoran mitogenomes were available in GenBank.

The Stylommatophora is the largest group within the pulmonate gastropods, containing $20,000 \pm 1500$ species (Rosenberg et al. 2022). Based on the structure of the excretory system Pilsbry (1900) divided the Stylommatophora in three infraorders: Orthurethra, Heterurethra and Sigmurethra. Baker (1955) adapted this system by recognizing a fourth major group, viz. Mesurethra. Of these four taxa, only Orthurethra is still widely accepted as a natural group. Deep-level relationships within Stylommatophora have been controversial. Nordsieck (1992) suggested dividing Stylommatophora into two subclades, Orthurethra and Sigmurethra. The first comprehensive molecular study of stylommatophoran relationships was undertaken by Wade et al. (2001), who recognized an 'achatinoïd' clade and a 'non-achatinoïd' clade. This hypothesis was subsequently supported by Wade et al. (2006). Bouchet et al. (2005) divided the Stylommatophora into three clades, Elasmognatha, Orthurethra and Sigmurethra, based a combined analysis of morphological and molecular data. Elasmognatha contains Succineoidea and Athoracophoroidea. Orthurethra contains Partuloidea, Achatinelloidea, Cochlicopoidea, Pupilloidea and Enoidea. Sigmurethra was suggested to be an informal group, which contains Clausilioidea, Orthalicoidea, Achatinoidea, Aillyoidea, Testacelloidea, Papillodermatoidea, Streptaxoidea, Rhytidoidea, Acavoidea, Plectopyloidea, Punctoidea and Sagdoidea. Bouchet et al. (2017) recognized Stylommatophora as an order and divided it into three suborders, Achatinina, Helicina and Scolodontina. Achatinina comprises Achatinoidea and Streptaxoidea. Helicina includes Coelociontoidea, Papillodermatoidea, Plectopyloidea, Punctoidea, Testacelloidea and Urocoptoidea. Scolodontina contains the single family Scolodontidae. Saa-di and Wade (2019) further refined this hypothesis by recognizing Scolodontidae as sister to all other stylommatophoran groups comprising the 'achatinoïd' and 'non-achatinoïd' clades. Scolodontidae corresponded to the suborder Scolodontina proposed by Bouchet et al. (2017), while the 'achatinoïd' clade corresponded to the suborder Achatinina and the 'non-achatinoïd' clade corresponded to the suborder Helicina.

Limacoidea is a superfamily of Stylommatophora that is subdivided into four families: Agriolimacidae, Limacidae, Boettgerillidae and Vitrinidae (Hausdorf 1998; Bouchet et al. 2017). The slug *Ambigolimax valentianus* is an invasive species in North and South America, Africa and Asia (Robinson 1999). Several studies have proved the usefulness of mitogenome data to resolve phylogenetic relationships in Stylommatophora (Minton et al. 2016; Xie et al. 2019; Guzmán et al. 2021). In this study, we applied next generation sequencing to obtain the complete mitogenomes of *Deroceras laeve* (O. F. Müller, 1774) and *Ambigolimax valentianus* (A. Férussac, 1821). The mitogenome of *Deroceras laeve* represented the second for Agriolimacidae and that of *Ambigolimax valentianus* was the first for Limacidae. This contribution aims at characterizing these two new mitogenomes and using them for a phylogenetic analysis of Stylommatophora.

Material and methods

Specimens and DNA extraction

Specimens of *Deroceras laeve* and *Ambigolimax valentianus* were collected from Zunyi, Guizhou Province, China, in July, 2020. They were identified by checking their adult morphological characters and blasting the mitochondrial *cox1* gene sequences in the BOLD system. The voucher specimens were deposited at the Henan Agricultural University, Zhengzhou, China, under the accession numbers MT-Zy20200701 and MT-Zy20200702. The specimens were preserved in absolute ethanol, and stored at -80 °C until DNA extraction. Total genomic DNA of the individual specimen was extracted with the TIANamp Genomic DNA Kit (TIANGEN BIOTECH CO., LTD), following the manufacturer's protocol.

Genome sequencing, assembly and annotation

Genome sequencing was performed on an Illumina HiSeq2500 platform, with a strategy of 150 paired-end sequencing. Library generation for the Illumina HiSeq sequencing was carried out using the Illumina TruSeq™ DNA Sample Prep Kit (Illumina, San Diego, CA, USA), with 350 bp insert size. NGS QC Toolkit v.2.3.3 (Patel and Jain 2012) was used to check the quality of the data. Adapters, ploy-N, and low-quality reads were removed from raw data. About 3 Gb clean data obtained by NGS for each species were used to assemble the mitochondrial scaffold.

GetOrganelle v.1.7.5.2 (Jin et al. 2020) was used for mitogenome assembly. The GetOrganelle animal database (-F animal_mt) was applied to identify, filter, and assemble target-associated reads. The new mitogenomes were annotated with the MITOS webserver (Bernt et al. 2013) (<http://mitos2.bioinf.uni-leipzig.de/index.py>). The gene boundaries of protein-coding genes were refined by alignment against mitochondrial gene sequences of closely related species. tRNA genes were identified using MITOS (Bernt et al. 2013) and ARWEN (Laslett and Canbäck 2008), and the secondary structures were redrawn in Adobe Illustrator CC 2019. The secondary structures of rRNA genes were predicted with reference to *Omalonyx unguis* (Guzmán et al. 2021). The mitogenome structure images were generated using mtviz (<http://pacosy.informatik.uni-leipzig.de/mtviz>). The annotated mitogenome sequences were submitted to GenBank under the accession numbers of [OQ198714](https://www.ncbi.nlm.nih.gov/nuccore/OQ198714) (*Deroceras laeve*) and [OQ198715](https://www.ncbi.nlm.nih.gov/nuccore/OQ198715) (*Ambigolimax valentianus*).

Characterization of the new mitogenomes

Pairwise comparisons of gene order with the gene order of *Achatina fulica* (He et al. 2016; Yang et al. 2016; Xie et al. 2019) and assessment of rearrangement events were performed using CREx (<http://pacosy.informatik.uni-leipzig.de/crex/form>) (Bernt et al. 2007). The nucleotide compositions of the mitogenome sequences were calculated with MEGA 11 (Kumar et al. 2018). AT and GC-skew values were obtained using the following formulas: AT-skew = $(A-T)/(A+T)$ and GC-skew = $(G-C)/(G+C)$ (Perna and Kocher 1995).

Sequence alignment

Protein-coding genes were aligned individually using MUSCLE as implemented in MEGA 11 (with default settings) (Kumar et al. 2018). Protein-coding genes were translated into amino acid sequences using the invertebrate mitochondrial genetic code. The alignments of genes were concatenated with FASconCAT-G_v.1.04 (Kück and Longo 2014) to create the amino acid dataset PCG_aa.

Phylogenetic analysis

A total of 68 mollusk mitogenome sequences were used in the phylogenetic analyses, of which 59 species were included as ingroup to represent Stylommatophora. Four species from Systellommatophora, two species from Ellobiida and three species from Hygrophila were selected as outgroups (Suppl. material 1). Phylogenetic analyses were performed based on the amino acid dataset mentioned above, under maximum likelihood (ML) and Bayesian inference (BI) criteria. ML analysis was performed with IQ-TREE v.1.6.10 (Nguyen et al. 2015). The data was partitioned by gene types. The best-fitting substitution models for partitions were chosen using ModelFinder (Kalyaanamoorthy et al. 2017) implemented in IQ-TREE. Branch support (BS) values were calculated using ultrafast bootstrap with 10,000 replicates. BI analysis was conducted using MrBayes v.3.2.7 (Ronquist et al. 2012). Two runs with four chains each were performed. The initial number of generations for each run was set to 10 million. Sampling was done every 1000 generations. The Average Standard Deviation of Split Frequencies (ASDSF) were monitored using Tracer v.1.7 (Rambaut et al. 2018). After reaching convergence (ASDSF < 0.01), the tree and branch length information were summarized using the *sumt* command, discarding the first 25% as burn-in. The consensus tree was yielded, and posterior probability (PP) values were used to assess branch support.

Results

Characteristics of the new mitogenomes

The entire mitochondrial genomes of *Deroceras laeve* and *Ambigolimax valentianus* were 14,773 base pairs (bp) and 15,195 bp long, respectively. They contained the entire set of 37 genes usually present in the animal mitogenomes and had identical gene orders (Fig. 1). Compared with the gene order of *Achatina fulica* (He et al. 2016; Yang et al. 2016; Xie et al. 2019), the *trnP_trnA* cluster was rearranged in both *Deroceras laeve* and *Ambigolimax valentianus*. The CREx analysis showed that the rearranged gene order of *Deroceras laeve* and *Ambigolimax valentianus* has evolved as the result of transposition (Fig. 2). The nucleotide compositions of both mitogenomes were heavily biased towards A and T. The overall A+T content of *Deroceras laeve* was 73.0%, while the A+T content of *Ambigolimax valentianus* was 71.4%. Both mitogenomes had the negative AT-skew values (-0.115 for *Deroceras laeve* and -0.096 for *Ambigolimax valentianus*) and the positive GC-skew values (0.181 for *Deroceras laeve* and 0.113 for *Ambigolimax valentianus*) in the major strand. This indicated the occurrence of more T than A and more G than C.




Figure 1. Gene order and gene content of the mitogenome of *Deroceras laeve*. The abbreviation of genes follows MITOS. Starting from *trnI* and ending at *cox3*, count clockwise. The outer numbers indicate the positions of each section.

family diagram for *Achatina fulica*



family diagram for *Deroceras laeve*



scenario:



Figure 2. The rearrangement event assessed from the CREx analysis for the mitogenome of *Deroceras laeve*.

For the protein-coding genes, ATG (five for *Deroceras laeve* and four for *Ambigolimax valentianus*), ATT (six for *Deroceras laeve* and three for *Ambigolimax valentianus*) and ATA (one for *Ambigolimax valentianus*) were used as the start codons. For the *cox2* gene and *atp8* gene of *Ambigolimax valentianus*, GTG was the start codon. For the *cox1* gene and *cob* gene of both *Ambigolimax valentianus* and *Deroceras laeve*, TTG was the start codon. All protein-coding genes terminated with the stop codon TAA or TAG, except for *nad3*

and *nad4L* of *Ambigolimax valentianus*, which had the incomplete stop codon T. For both species, Leu, Ile, Phe and Val were the most frequently used amino acids. Relative synonymous codon usage (RSCU) for 13 protein-coding genes of *Deroceras laeve* is shown in Fig. 3. *Ambigolimax valentianus* (Suppl. material 2) had similar RSCU values to *Deroceras laeve*.




Figure 3. Codon usage of the 13 mitochondrial protein-coding genes of *Deroceras laeve*. RSCU: relative synonymous codon usage.

All 22 tRNA genes were identified by both MITOS and ARWEN, and their lengths ranged from 61 to 68 bp. Most tRNA genes of both species can be folded into the classic clover-leaf structure (Fig. 4 and Suppl. material 3). The *trnT* gene of *Deroceras laeve* had an unusual TΨC loop. The *trnK* gene and *trnS1* gene of *Ambigolimax valentianus* had an incomplete DHU arm. The position of the *rnrL* gene was located between *trnV* and *trnL1*, while the *rnrS* gene was found between *trnE* and *trnM*. In *Deroceras laeve*, *rnrL* had a length of 1052 bp with an A + T content of 74.4%, whereas *rnrS* had a length of 697 bp with an A + T content of 71.5%. In *Ambigolimax valentianus*, *rnrL* was 1091 bp long with an A + T content of 75.9%, while *rnrS* was 711 bp long with an A + T content of 72.7%. The secondary structures for *rnrL* and *rnrS* of *Deroceras laeve* and *Ambigolimax valentianus* are presented in Figs 5, 6 and Suppl. materials 4, 5, respectively. The *rnrL* molecule of both species contained six domains (labeled I–VI) comprising 43 helices. The *rnrS* molecule consisted of three domains (labeled I–III) and 28 helices.

Phylogenetic inference

ML and BI produced similar tree topologies. The monophyly of Stylommatophora was supported under both analyses (Figs 7, 8). Achatinoidea represented by *Achatina fulica* was consistently resolved as the sister group of all other Stylommatophora (BS = 100, PP = 1.0). This lineage corresponded to the suborder Achatinina. The remaining stylommatophorans formed the ‘non-achatinoidea’ clade which corresponded to the suborder Helicina. In Helicina, all superfamilies with more than two representatives were well supported, with the exception of Punctoidea. All families with multiple representatives were supported as monophyletic. The newly sequenced *Deroceras laeve* was sister to *Deroceras reticulatum*, and both together formed the




Figure 4. The secondary structures of tRNA genes inferred for the mitogenome of *Deroceras laeve*.

sister group of *Ambigolimax valentianus*. The three species form a monophyletic Limacoidea clade.

The major differences between the ML and BI analyses were the positions of Xanthonychidae and the clade Orthalicoidea + Punctoidea. The ML analysis placed Xanthonychidae as sister to all other Helicoidea. Whereas, the BI analysis recovered Xanthonychidae as the sister group of a clade Geomitridae + Helicidae. The ML analysis placed Orthalicoidea + Punctoidea between a clade comprising (Clausilioidea + Pupilloidea) and Arionoidea comprising (Philomycidae + Arionidae). But this arrangement received no statistical support. The BI analysis recovered Orthalicoidea + Punctoidea between the clade (Clausilioidea + Pupilloidea) and a clade including Limacoidea, Helicarionoidea, Trochomorpoidea and Punctidae.




Figure 5. The secondary structure of rrmL inferred for *Deroceras laeve*.




Figure 6. The secondary structure of *rrmS* inferred for *Deroceras laeve*.




Figure 7. ML phylogenetic tree inferred with IQ-TREE using amino acid sequences of 13 PCGs. Numbers at the nodes are ultrafast bootstrap values (BS > 70). Blue indicates the newly sequenced species. The branch of *Cepaea nemoralis* is depicted as half of its original branch length. Scale bar represents substitutions/site.

Discussion

Mitochondrial gene rearrangement

Large-scale changes in genomes are considered to be rare events (Rokas and Holland 2000). The gene set remains constant across bilaterian animals, mitochondrial gene rearrangements appear to be unusual, and gene order is selectively neutral (Boore et al. 1995; Dowton et al. 2002; Dowton et al. 2009; Cameron 2014). In a certain animal group, for example the hexapods, most species share an identical mitogenome organization (Dowton et al. 2002; Cameron 2014). However, Stylommatophora mitogenomes have experienced many more mitochondrial gene rearrangements than other groups (Minton et al. 2016; Xie et al. 2019; Guzmán et al. 2021). In this study, two new mitogenomes from *Deroceras laeve* and *Ambigolimax valentianus* had a tRNA gene rearrangement associated with the *trnP_trnA* tRNA cluster. We compared the gene order for all included mitogenomes and found that all the Helicina species have gene rearrangements compared to *Achatina fulica*. The exemplars of Helicidae, Philomycidae and Achatinellidae displayed many more mitochondrial gene rearrangements than others (Fig. 8).

Phylogeny of Stylommatophora

Within Stylommatophora, the division of the order into Achatinina and Helicina is well accepted (Wade et al. 2001; Wade et al. 2006). Recently, some authors added Scolodontidae to the phylogenetic analysis (Ramírez et al. 2012; Bouchet et al. 2017; Saadi and Wade 2019). Currently, no mitogenomes of Scolodontidae have been published, so the phylogenetic position of Scolodontidae could not be assessed in this analysis. Our analyses consistently supported the division of Stylommatophora into two principal clades, Achatinina and Helicina. This result is consistent with the previous mitogenome analyses (Xie et al. 2019; Guzmán et al. 2021).

The present mitogenome data recovered Clausilioidea as a sister group of the orthurethran clade. This result contrasted with Wade et al. (2006), who recovered Orthurethra to be close to a clade comprising Arionoidea and Lima-coidea. In a prior mitogenome analysis, a sister group relationship between Succineoidea and Arionoidea was supported (Xie et al. 2019). In this study, this pattern was supported by the BI analysis (PP = 0.96) based on expanded taxon sampling of mitogenomes. In our analyses, only one species of *Haplotrema minimum* (Ancey, 1888) representing Haplotrematoidea was included due to mitogenome data availability. Both the ML and BI analyses placed Haplotrematoidea as sister to Helicoidea. However, this relationship had low support values. In future researches, larger taxon samples are needed to identify the phylogenetic placement of Haplotrematoidea.

The superfamily Punctoidea included the families Punctidae, Charopidae, Cystopeltidae, Discidae ("Endodontidae"), Helicodiscidae, Oopeltidae and Oreohelicidae (Bouchet et al. 2017). In this study, we included three species of Punctoidea in the phylogenetic analyses, which respectively represented Discidae [*Discus perspectivus* (Megerle von Mühlfeld, 1816)], Oreohelicidae [*Oreohelix idahoensis* (Newcomb, 1866)] and Punctidae [*Punctum randolphii* (Dall, 1895)]. *Discus perspectivus* and *O. idahoensis* were significantly supported to




Figure 8. Bayesian phylogenetic tree inferred in MrBayes using amino acid sequences of 13 PGCs (left), and gene order comparisons for the stylommatophoran mitogenomes (right). Numbers at the nodes in the tree are Bayesian posterior probabilities (PP > 0.9). Blue indicates the newly sequenced species. The branch of *Cepaea nemoralis* is depicted as half of its original branch length. Scale bar represents substitutions/site. In the mitogenome structure maps, gene rearrangements are highlighted by red.

be a sister group. *Punctum randolphii* was placed separately, and clustered with a clade including Trochomorpoidea, Helicarionoidea and Limacoidea. Taxon sampling of Punctoidea was very limited in our analysis. The monophyly of Punctoidea needs to be further tested by additional species in future studies.

Previous analyses based on the multiple gene fragments have demonstrated the monophyly of Helicoidea, which comprised the families Helicidae, Bradybaenidae, Xanthonychidae, Hygromiidae, Camaenidae, Polygyridae and Sagdidae (Wade et al. 2006). Our results strongly supported a monophyletic Helicoidea (BS = 100, PP = 1.0). In addition, the superfamilies Urocoptoidea, Succineoidea, Arionoidea, Pupilloidea and Limacoidea were supported to be monophyletic groups. These patterns are congruent with the current taxonomy of land snails (Bouchet et al. 2017).

For the taxon sampling of outgroups, we included four exemplars from Systellommatophora and two from Ellobiida. The monophyly of Systellommatophora and Ellobiida were consistently supported in both ML and BI analyses. Moreover, Systellommatophora and Ellobiida formed a sister-group relationship (BS = 100, PP = 1). Amphipulmonata Schrödl, 2014 was established as a clade containing Systellommatophora and Ellobiida (Bouchet et al. 2017). Amphipulmonata (comprising Ellobioidea and Systellommatophora) was supported in the previous molecular phylogenetic analysis (Dayrat et al. 2011). Our results confirmed the hypothesis of Amphipulmonata.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was funded by the National Natural Science Foundation of China (U1904104) and Zunyi Branch of Guizhou Tobacco Company Key Research Project (2020XM01). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

N.S. and T.Z. designed the research; N.S., X.L., Y.Z. and T.Z. performed the research and analyzed the data; N.S., T.Z. and X.L. wrote the paper. All authors have read and agreed to the published version of the manuscript.

Data availability

The mitogenome sequences newly generated in this study were deposited in GenBank, with the accession numbers of [OQ198714–OQ198715](#). The dataset used for phylogenetic analyses can be obtained from the corresponding author upon request.

References

Baker H (1955) Heterurethrous and aulacopod. *The Nautilus* 68: 109–112.

- Bernt M, Merkle D, Ramsch K, Fritzsich G, Perseke M, Bernhard D, Schlegel M, Stadler PF, Middendorf M (2007) CREx: Inferring genomic rearrangements based on common intervals. *Bioinformatics* (Oxford, England) 23(21): 2957–2958. <https://doi.org/10.1093/bioinformatics/btm468>
- Bernt M, Bleidorn C, Braband A, Dambach J, Donath A, Fritzsich G, Golombek A, Hadrys H, Jühling F, Meusemann K, Middendorf M, Misof B, Perseke M, Podsiadlowski L, von Reumont B, Schierwater B, Schlegel M, Schrödl M, Simon S, Stadler PF, Stöger I, Struck TH (2013) A comprehensive analysis of bilaterian mitochondrial genomes and phylogeny. *Molecular Phylogenetics and Evolution* 69(2): 352–364. <https://doi.org/10.1016/j.ympev.2013.05.002>
- Boore JL (1999) Animal mitochondrial genomes. *Nucleic Acids Research* 27(8): 1767–1780. <https://doi.org/10.1093/nar/27.8.1767>
- Boore JL, Collins TM, Stanton D, Daehler LL, Brown WM (1995) Deducing the pattern of arthropod phylogeny from mitochondrial DNA rearrangements. *Nature* 376(6536): 163–165. <https://doi.org/10.1038/376163a0>
- Bouchet P, Frýda J, Hausdorf B, Ponder W, Valdés A, Warén A (2005) Classification and nomenclator of gastropod families. *Malacologia* 47: 1–397.
- Bouchet P, Rocroi JP, Hausdorf B, Kaim A, Kano Y, Nützel A, Parkhaev P, Schrödl M, Strong EE (2017) Revised classification, nomenclator and typification of gastropod and monoplacophoran families. *Malacologia* 61(1–2): 1–526. <https://doi.org/10.4002/040.061.0201>
- Cameron SL (2014) Insect mitochondrial genomics: Implications for evolution and phylogeny. *Annual Review of Entomology* 59(1): 95–117. <https://doi.org/10.1146/annurev-ento-011613-162007>
- Dayrat B, Conrad M, Balayan S, White TR, Albrecht C, Golding R, Gomes SR, Harasewych MG, de Frias Martins AM (2011) Phylogenetic relationships and evolution of pulmonate gastropods (Mollusca): New insights from increased taxon sampling. *Molecular Phylogenetics and Evolution* 59(2): 425–437. <https://doi.org/10.1016/j.ympev.2011.02.014>
- Downton M, Castro L, Austin A (2002) Mitochondrial gene rearrangements as phylogenetic characters in the invertebrates: The examination of genome ‘morphology’. *Invertebrate Systematics* 16(3): 345–356. <https://doi.org/10.1071/IS02003>
- Downton M, Cameron SL, Dowavac JI, Austin AD, Whiting MF (2009) Characterization of 67 mitochondrial tRNA gene rearrangements in the Hymenoptera suggests that mitochondrial tRNA gene position is selectively neutral. *Molecular Biology and Evolution* 26(7): 1607–1617. <https://doi.org/10.1093/molbev/msp072>
- Guzmán LB, Vogler RE, Beltramino AA (2021) The mitochondrial genome of the semi-slug *Omalonyx unguis* (Gastropoda: Succineidae) and the phylogenetic relationships within Stylommatophora. *PLoS ONE* 16(6): e0253724. <https://doi.org/10.1371/journal.pone.0253724>
- Hausdorf B (1998) Phylogeny of the Limacoidea sensu lato (Gastropoda: Stylommatophora). *The Journal of Molluscan Studies* 64(1): 35–66. <https://doi.org/10.1093/mollus/64.1.35>
- He ZP, Dai XB, Zhang S, Zhi TT, Lun ZR, Wu ZD, Yang TB (2016) Complete mitochondrial genome of the giant African snail, *Achatina fulica* (Mollusca: Achatinidae): a novel location of putative control regions (CR) in the mitogenome within Pulmonate species. *Mitochondrial DNA, Part A, DNA Mapping, Sequencing, and Analysis* 27: 1084–1085. <https://doi.org/10.3109/19401736.2014.930833>
- Jin JJ, Yu WB, Yang JB, Song Y, DePamphilis CW, Yi TS, Li DZ (2020) GetOrganelle: A fast and versatile toolkit for accurate de novo assembly of organelle genomes. *Genome Biology* 21(1): 1–31. <https://doi.org/10.1186/s13059-020-02154-5>

- Kalyaanamoorthy S, Minh BQ, Wong TK, Von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods* 14(6): 587–589. <https://doi.org/10.1038/nmeth.4285>
- Kück P, Longo GC (2014) FASconCAT-G: Extensive functions for multiple sequence alignment preparations concerning phylogenetic studies. *Frontiers in Zoology* 11(1): 1–8. <https://doi.org/10.1186/s12983-014-0081-x>
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution* 35(6): 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Laslett D, Canbäck B (2008) ARWEN: A program to detect tRNA genes in metazoan mitochondrial nucleotide sequences. *Bioinformatics (Oxford, England)* 24(2): 172–175. <https://doi.org/10.1093/bioinformatics/btm573>
- Minton RL, Cruz MAM, Farman ML, Perez KE (2016) Two complete mitochondrial genomes from *Praticolella mexicana* Perez, 2011 (Polygyridae) and gene order evolution in Helicoidea (Mollusca, Gastropoda). *ZooKeys* 626: 137–154. <https://doi.org/10.3897/zookeys.626.9633>
- Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* 32(1): 268–274. <https://doi.org/10.1093/molbev/msu300>
- Nordsieck H (1992) Phylogeny and system of the Pulmonata (Gastropoda). *Archiv für Molluskenkunde* 121(1–6): 31–52. <https://doi.org/10.1127/arch.moll/121/1992/31>
- Patel RK, Jain M (2012) NGS QC Toolkit: A toolkit for quality control of next generation sequencing data. *PLoS ONE* 7(2): e30619. <https://doi.org/10.1371/journal.pone.0030619>
- Perna NT, Kocher TD (1995) Patterns of nucleotide composition at fourfold degenerate sites of animal mitochondrial genomes. *Journal of Molecular Evolution* 41(3): 353–358. <https://doi.org/10.1007/BF01215182>
- Pilsbry H (1900) On the zoological position of *Achatinella* and *Partula*. In: *Proceedings of the Academy of Natural Sciences of Philadelphia*, 561–567.
- Ponder WF, Lindberg DR, Ponder JM (2020) *Biology and evolution of the Mollusca. Volume one*. CRC Press, Boca Raton, USA. <https://doi.org/10.1201/9781351115254>
- Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA (2018) Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* 67(5): 901–904. <https://doi.org/10.1093/sysbio/syy032>
- Ramírez R, Borda V, Romero P, Ramírez J, Congrains C, Chirinos J, Ramírez P, Velásquez LE, Mejía K (2012) Biodiversity and endemism of the western Amazonia land snails *Megalobulimus* and *Systrophia*. *Revista Peruana de Biología* 19: 59–74. <https://doi.org/10.15381/rpb.v19i1.798>
- Robinson DG (1999) Alien invasions: The effects of the global economy on non-marine gastropod introductions into the United States. *Malacologia* 41: 413–438.
- Rokas A, Holland PW (2000) Rare genomic changes as a tool for phylogenetics. *Trends in Ecology & Evolution* 15(11): 454–459. [https://doi.org/10.1016/S0169-5347\(00\)01967-4](https://doi.org/10.1016/S0169-5347(00)01967-4)
- Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Rosenberg G, Auffenberg K, Bank R, Bieler R, Bouchet P, Herbert D, Köhler F, Neubauer TA, Neubert E, Páll-Gergely B, Richling I, Schneider S (2022) Adapting mark-recapture

- methods to estimating accepted species-level diversity: A case study with terrestrial Gastropoda. *PeerJ* 10: e13139. <https://doi.org/10.7717/peerj.13139>
- Saadi AJ, Wade CM (2019) Resolving the basal divisions in the stylommatophoran land snails and slugs with special emphasis on the position of the Scolodontidae. *Molecular Phylogenetics and Evolution* 139: 106529. <https://doi.org/10.1016/j.ympev.2019.106529>
- Wade CM, Mordan PB, Clarke B (2001) A phylogeny of the land snails (Gastropoda: Pulmonata). *Proceedings of the Royal Society of London, Series B, Biological Sciences* 268(1465): 413–422. <https://doi.org/10.1098/rspb.2000.1372>
- Wade CM, Mordan PB, Naggs F (2006) Evolutionary relationships among the Pulmonate land snails and slugs (Pulmonata, Stylommatophora). *Biological Journal of the Linnean Society. Linnean Society of London* 87(4): 593–610. <https://doi.org/10.1111/j.1095-8312.2006.00596.x>
- Xie GL, Köhler F, Huang XC, Wu RW, Zhou CH, Ouyang S, Wu XP (2019) A novel gene arrangement among the Stylommatophora by the complete mitochondrial genome of the terrestrial slug *Meghimatium bilineatum* (Gastropoda, Arionoidea). *Molecular Phylogenetics and Evolution* 135: 177–184. <https://doi.org/10.1016/j.ympev.2019.03.002>
- Yang H, Zhang JE, Guo J, Deng Z, Luo H, Luo M, Zhao B (2016) The complete mitochondrial genome of the giant African snail *Achatina fulica* (Mollusca: Achatinidae). *Mitochondrial DNA, Part A, DNA Mapping, Sequencing, and Analysis* 27: 1622–1624.

Supplementary material 1

Taxa included in this study, including GenBank Accession numbers and literature references

Authors: Te Zhao, Nan Song, Xingyu Lin, Yang Zhang

Data type: table (docx. file)

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1173.102786.suppl1>

Supplementary material 2

Relative synonymous codon usage (RSCU) for 13 protein-coding genes of *Ambigolimax valentianus*

Authors: Te Zhao, Nan Song, Xingyu Lin, Yang Zhang

Data type: figure (PDF file)

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1173.102786.suppl2>

Supplementary material 3

The secondary structures of tRNA genes inferred for the mitogenome of *Ambigolimax valentianus*

Authors: Te Zhao, Nan Song, Xingyu Lin, Yang Zhang

Data type: figure (PDF file)

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1173.102786.suppl3>

Supplementary material 4

The secondary structure of *rrnL* inferred for *Ambigolimax valentianus*

Authors: Te Zhao, Nan Song, Xingyu Lin, Yang Zhang

Data type: figure (PDF file)

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1173.102786.suppl4>

Supplementary material 5

The secondary structure of *rrnS* inferred for *Ambigolimax valentianus*

Authors: Te Zhao, Nan Song, Xingyu Lin, Yang Zhang

Data type: figure (PDF file)

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1173.102786.suppl5>