

An updated checklist of *Anopheles* (Diptera, Culicidae) of Colombia with new records and distribution data

Nelson Naranjo-Díaz¹, Margarita M. Correa¹ 

¹ Grupo Microbiología Molecular, Escuela de Microbiología, Universidad de Antioquia, Calle 70 No. 52-21, Medellín, Colombia
Corresponding author: Margarita M. Correa (margarita.correa@udea.edu.co)

Abstract

Several species of *Anopheles* mosquitoes (Arthropoda, Insecta, Diptera, Culicidae) are important in public health due to their role in malaria transmission. Of the more than 500 *Anopheles* species worldwide, 47 have been reported in Colombia, but only nine are known to transmit malaria. Taxonomic classification of these mosquitoes is complicated by the existence of species complexes and groups of closely related species that are difficult to distinguish based on morphological characteristics. However, molecular techniques have contributed to resolving taxonomic uncertainties, definition of molecular variants and facilitated the correction of erroneous taxonomic assignments. This study aimed to update the list of *Anopheles* species reported for Colombia. A species checklist was compiled by reviewing catalogs, publications, databases, and unpublished data. Only formally characterized species were included, along with their geographic range and ecological distribution. The final list includes 44 formally characterized *Anopheles* species belonging to five subgenera. The *Nyssorhynchus* subgenus constituted the largest group with 17 species and the widest distribution, occurring in 18 ecoregions. The *Anopheles* subgenus was the second largest group with 16 species and occurrences in 16 ecoregions. Sixty-six new presence records were added to the checklist. The updated *Anopheles* checklist, encompassing presence records and ecological distributions, enhances our understanding of *Anopheles* mosquito biodiversity. Furthermore, it contributes to improved public health by providing a foundation for targeted vector control interventions.

Key words: *Anopheles*, biodiversity, Colombia, ecological distribution, inventory, malaria vectors

Introduction

The *Anopheles* genus is highly diverse, with 511 formally recognized species belonging to eight subgenera, including various species complexes. Some of these complexes still contain unnamed members (Harbach 2023). *Anopheles* mosquitoes thrive in a wide range of ecosystems, contributing to their broad geographical distribution. Additionally, some species have adapted to anthropically modified habitats (Hiwat and Bretas 2011; Harbach 2023). The primary importance of this genus lies in its role as a vector of human and animal pathogens, particularly, *Plasmodium* parasites, the causative agents of malaria (WHO 2022). Some *Anopheles* species also transmit *Wuchereria bancrofti*, the nematode that



Academic editor: John Soghigian
Received: 2 August 2024
Accepted: 16 January 2025
Published: 12 March 2025

ZooBank: <https://zoobank.org/3D134FAA-6E53-44C7-90FE-1DD02C3B1B46>

Citation: Naranjo-Díaz N, Correa MM (2025) An updated checklist of *Anopheles* (Diptera, Culicidae) of Colombia with new records and distribution data. ZooKeys 1231: 169–189. <https://doi.org/10.3897/zookeys.1231.133711>

Copyright: ©
Nelson Naranjo-Díaz & Margarita M. Correa.
This is an open access article distributed under
terms of the Creative Commons Attribution
License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

causes filariasis in Africa and Asia (Manguin et al. 2010), and O'nyong-nyong virus, which produces polyarthritis and fever in Africa (Brault et al. 2004).

Most of Colombia has ecological conditions that favor the widespread distribution of *Anopheles* mosquitoes (Olano et al. 2001; IGAC 2002; Hernández-Valencia et al. 2020). In the country, 47 species have been identified (Montoya-Lerma et al. 2011; Gómez et al. 2015), and nine of these are considered malaria vectors. Three species from the *Nyssorhynchus* subgenus play a significant role in malaria transmission: *Anopheles darlingi*, *An. nuneztovari*, and *An. albimanus* (Olano et al. 2001; Gutiérrez et al. 2008). *Anopheles darlingi* is the primary malaria vector in Latin America (Hiwat and Bretas 2011) and is predominantly found in the northwest, east and Amazon regions of Colombia. *Anopheles nuneztovari* is more common in the northwest, northeast and east, while *An. albimanus* is primarily present in coastal areas (Olano et al. 2001; Naranjo-Díaz et al. 2016a, b).

Knowledge of *Anopheles* species in Colombia is based primarily on catalogs published in the mid-20th century (Gast 1943; Barreto-Reyes 1955; Stone et al. 1959; Knight and Stone 1977) and older reports from the former government malaria control program "Servicio Nacional de Erradicación de la Malaria-SEM" (SEM 1957). Subsequent contributions by Carrejo and González (1992) and González and Carrejo (2007, 2009), focusing on the taxonomy and control of medically important insects, with an emphasis on population and taxonomic studies, have significantly improved our understanding of *Anopheles* in Colombia. While the number of reports on *Anopheles* species occurrence and distribution has increased, information remains scattered and, in some cases, inaccessible to the public.

Molecular tools have significantly improved the resolution of taxonomic ambiguities, particularly within species complexes and among closely related species; they have also facilitated the correction of erroneous taxonomic assignments (Brochero et al. 2007; Ruiz-Lopez et al. 2012; Escovar et al. 2014; Gómez et al. 2015). These findings underscore the need for an updated list of species occurrence. A comprehensive checklist of *Anopheles* species in Colombia, incorporating current ecological and geographical data, is essential for advancing medical entomological research and biodiversity assessment.

Materials and methods

To update the *Anopheles* species checklist, we reviewed various sources, including mosquito catalogs (Gast 1943; Barreto-Reyes 1955; Knight and Stone 1977; Heinemann and Belkin 1978; Faran 1980; Carrejo and Gonzalez 1992), taxonomic keys (Faran and Linthicum 1981; Gonzalez and Carrejo 2009), and government reports (SEM 1957). Additionally, we obtained data on *Anopheles* species occurrence, including geographical coordinates, from databases (Gaffigan et al. 2014; SIB Database 2020; GBIF 2024), scientific articles, and unpublished new records of specimens collected by members of our research group (named "new occurrence data"). The identities of the new records were previously verified using molecular methods, e.g., COI barcoding or ITS2 analysis (Zapata et al. 2007; Cienfuegos et al. 2011; Gómez et al. 2015) (Suppl. material 1).

To be included in the checklist, an *Anopheles* species was required to have a formal description and validation (Harbach 2023; ITIS 2024). Species variants originally described using only molecular methods were excluded. The checklist includes species listed in alphabetic order. Each entry contains

the genus, subgenus, authorship, and year of description; (new occurrence data): This designation is used to indicate that the records are the result of previous work by our research group. Geographical distribution is presented at the level of Colombian administrative departments. An asterisk (*) indicates species occurrences with associated geographical coordinates.

Notes provide information on the sources of the data, changes in the taxonomic classification of species, molecular species designations, and new occurrence data, including the municipalities where specimens were collected. The checklist is accompanied by a map illustrating the distribution of *Anopheles* species with associated geographical coordinates (Figs 1, 2, Suppl. material 1).

In addition to the list, a summary table of the *Anopheles* records by administrative departments is included (Table 1), and also, a table of species distribution per ecoregion (WWF 2015) (Table 2), which includes the following: Amazon-Orinoco-Southern Caribbean mangroves, Apure-Villavicencio dry forests, Caquetá moist forests, Catatumbo moist forests, Cauca Valley dry forests, Cauca Valley montane forests, Chocó-Darién moist forests, Cordillera Oriental montane forests, Eastern Cordillera Real montane forests, Guajira-Barranquilla xeric scrub, Llanos, Magdalena Valley montane forests, Magdalena-Urabá moist forests, Napo moist forests, Negro-Branco moist forests, Northwestern Andean montane forests, Sinú Valley dry forests, Solimões–Japurá moist forests, South American Pacific mangroves, Southwest Amazon moist forests, and Western Ecuador moist forests.




Figure 1. Maps depicting the number of *Anopheles* species reported in Colombia **A** by administrative area (Department) **B** by ecological region.




Figure 2. Map indicating the location of species records, according to geographic coordinates. Blue: previously reported records; red: new occurrence data.

Table 1. Summary of *Anopheles* records by Colombian administrative departments.

Department	Subgenera	Species	New geographic coordinate registers*	Total geographic coordinate registers [‡]
Amazonas	3	8	-	11
Antioquia	3	22	40	155
Arauca	2	16	-	6
Atlántico	2	10	-	1
Bolívar	3	17	-	-
Boyacá	3	17	-	4
Caldas	3	17	-	1
Caquetá	4	18	-	6
Casanare	2	16	-	2
Cauca	4	18	1	42
Cesar	2	15	-	-
Chocó	4	23	-	31
Córdoba	2	18	21	80
Cundinamarca	5	23	-	2
Guainía	2	7	-	-
Guaviare	3	13	3	5
Huila	3	13	-	-
La Guajira	2	15	-	4
Magdalena	3	17	-	6
Meta	5	30	-	60
Nariño	4	15	-	35
Norte de Santander	4	24	-	14
Putumayo	3	18	-	36
Quindío	2	7	-	-
Risaralda	2	7	-	3
San Andres y Providencia	1	1	-	-
Santander	2	19	-	33
Sucre	2	13	-	-
Tolima	3	18	-	1
Valle del Cauca	4	19	1	33
Vaupés	5	15	-	-
Vichada	2	10	-	9

* Number of new geographic coordinate records described during this study; [‡] Total number of geographic coordinate records, including both previous and new occurrence data.

Table 2. List of *Anopheles* recorded in Colombia by ecoregion.

Ecoregions*	Species
Amazon-Orinoco-Southern Caribbean mangroves	<i>An. (Ano.) apicimacula</i> , <i>An. (Ano.) punctimacula</i> , <i>An. (Ano.) pseudopunctipennis</i> [‡]
Apure-Villavicencio dry forests	<i>An. (Ano.) costai</i> , <i>An. (Ano.) apicimacula</i> , <i>An. (Ano.) pseudopunctipennis</i> [‡] , <i>An. (Ker) bambusicolus</i> , <i>An. (Ker) homunculus</i> , <i>An. (Ker) pholidotus</i> [‡] , <i>An. (Nys.) albitarsis</i> , <i>An. (Nys.) argyritarsis</i> , <i>An. (Nys.) benarrochi</i> [‡] , <i>An. (Nys.) braziliensis</i> , <i>An. (Nys.) darlingi</i> [‡] , <i>An. (Nys.) marajoara</i> , <i>An. (Nys.) nuneztovari</i> [‡] , <i>An. (Nys.) rangeli</i> , <i>An. (Nys.) triannulatus</i> , <i>An. (Ste.) nimbus</i>
Caquetá moist forests	<i>An. (Nys.) darlingi</i> [‡] , <i>An. (Nys.) braziliensis</i>
Catatumbo moist forests	<i>An. (Ano.) malefactor</i> , <i>An. (Ano.) neomaculipalpus</i> , <i>An. (Nys.) albitarsis</i> , <i>An. (Nys.) marajoara</i> , <i>An. (Nys.) nuneztovari</i> [‡]
Cauca Valley dry forests	<i>An. (Ano.) calderoni</i> [‡] , <i>An. (Ano.) punctimacula</i> [‡] , <i>An. (Ano.) pseudopunctipennis</i> [‡] , <i>An. (Ker) neivai</i> [‡] , <i>An. (Nys.) albimanus</i> [‡]
Cauca Valley montane forests	<i>An. (Ano.) apicimacula</i> , <i>An. (Ano.) calderoni</i> [‡] , <i>An. (Ano.) pseudopunctipennis</i> [‡] , <i>An. (Nys.) albimanus</i> [‡] , <i>An. (Nys.) albitarsis</i> , <i>An. (Nys.) nuneztovari</i> [‡] , <i>An. (Nys.) triannulatus</i>

Ecoregions*	Species
Chocó-Darién moist forests	<i>An. (Ano.) apicimacula</i> , <i>An. (Ano.) calderoni</i> [‡] , <i>An. (Ano.) costai/forattinii</i> , <i>An. (Ano.) malefactor</i> , <i>An. (Ano.) punctimacula</i> [‡] , <i>An. (Ker) neivai</i> [‡] , <i>An. (Nys.) albimanus</i> [¥] , <i>An. (Nys.) darlingi</i> [¥] , <i>An. (Nys.) eiseni</i> , <i>An. (Nys.) nuneztovari</i> [¥] , <i>An. (Nys.) triannulatus</i>
Cordillera Oriental montane forests	<i>An. (Ker) homunculus</i> , <i>An. (Nys.) darlingi</i> [¥] , <i>An. (Nys.) rangeli</i>
Eastern Cordillera Real montane forests	<i>An. (Nys.) argyritarsis</i>
Guajira-Barranquilla xeric scrub	<i>An. (Ano.) neomaculipalpus</i> , <i>An. (Ano.) punctimacula</i> [‡]
Llanos	<i>An. (Ano.) costai</i> , <i>An. (Ano.) peryassui</i> , <i>An. (Ano.) pseudopunctipennis</i> [‡] , <i>An. (Ano.) shannoni</i> , <i>An. (Ker) bambusicolus</i> , <i>An. (Nys.) albitarsis</i> , <i>An. (Nys.) argyritarsis</i> , <i>An. (Nys.) braziliensis</i> , <i>An. (Nys.) darlingi</i> [¥] , <i>An. (Nys.) marajoara</i> , <i>An. (Nys.) nuneztovari</i> [¥] , <i>An. (Nys.) oswaldoi</i> , <i>An. (Nys.) rangeli</i> , <i>An. (Nys.) triannulatus</i>
Magdalena Valley montane forests	<i>An. (Ano.) apicimacula</i> , <i>An. (Ano.) neomaculipalpus</i> , <i>An. (Ano.) pseudopunctipennis</i> [‡] , <i>An. (Nys.) argyritarsis</i> , <i>An. (Nys.) darlingi</i> [¥] , <i>An. (Nys.) nuneztovari</i> [¥] , <i>An. (Nys.) parvus</i> , <i>An. (Nys.) rangeli</i> , <i>An. (Nys.) triannulatus</i>
Magdalena-Urabá moist forests	<i>An. (Ano.) apicimacula</i> , <i>An. (Ano.) malefactor</i> , <i>An. (Ano.) mattogrossensis</i> , <i>An. (Ano.) neomaculipalpus</i> , <i>An. (Ano.) peryassui</i> , <i>An. (Ano.) pseudopunctipennis</i> [‡] , <i>An. (Ano.) punctimacula</i> [‡] , <i>An. (Ker) neivai</i> [‡] , <i>An. (Lop) squamifemur</i> , <i>An. (Nys.) albimanus</i> [¥] , <i>An. (Nys.) albitarsis</i> , <i>An. (Nys.) aquasalis</i> , <i>An. (Nys.) argyritarsis</i> , <i>An. (Nys.) benarrochi</i> [‡] , <i>An. (Nys.) braziliensis</i> , <i>An. (Nys.) darlingi</i> [¥] , <i>An. (Nys.) evansae</i> , <i>An. (Nys.) marajoara</i> , <i>An. (Nys.) nuneztovari</i> [¥] , <i>An. (Nys.) oswaldoi</i> , <i>An. (Nys.) rangeli</i> , <i>An. (Nys.) strodei</i> , <i>An. (Nys.) triannulatus</i> , <i>An. (Ste.) nimbus</i>
Napo moist forests	<i>An. (Ano.) costai</i> , <i>An. (Ano.) mattogrossensis</i> , <i>An. (Ano.) neomaculipalpus</i> , <i>An. (Ano.) punctimacula</i> [‡] , <i>An. (Nys.) albitarsis</i> , <i>An. (Nys.) bellator</i> , <i>An. (Nys.) benarrochi</i> [‡] , <i>An. (Nys.) braziliensis</i> , <i>An. (Nys.) darlingi</i> [¥] , <i>An. (Nys.) marajoara</i> , <i>An. (Nys.) oswaldoi</i> , <i>An. (Nys.) rangeli</i> , <i>An. (Nys.) strode</i> , <i>An. (Nys.) triannulatus</i>
Negro-Branco moist forests	<i>An. (Ano.) mattogrossensis</i> , <i>An. (Ano.) peryassui</i> , <i>An. (Nys.) braziliensis</i> , <i>An. (Nys.) darlingi</i> [¥] , <i>An. (Nys.) oswaldoi</i>
Northwestern Andean montane forests	<i>An. (Ano.) calderoni</i> [‡] , <i>An. (Ano.) pseudopunctipennis</i> [‡] , <i>An. (Nys.) albimanus</i> [¥] , <i>An. (Nys.) argyritarsis</i> , <i>An. (Nys.) darlingi</i> [¥] , <i>An. (Nys.) nuneztovari</i> [¥] , <i>An. (Nys.) triannulatus</i>
Sinú Valley dry forests	<i>An. (Ano.) punctimacula</i> [‡]
Solimões–Japurá moist forests	<i>An. (Ano.) costai</i> , <i>An. (Ano.) peryassui</i> , <i>An. (Nys.) darlingi</i> [¥] , <i>An. (Nys.) dunhami</i>
South American Pacific mangroves	<i>An. (Ano.) apicimacula</i> , <i>An. (Ano.) calderoni</i> [‡] , <i>An. (Ker) neivai</i> [‡] , <i>An. (Nys.) albimanus</i> [¥]
Southwest Amazon moist forests	<i>An. (Nys.) darlingi</i> [¥]
Western Ecuador moist forests	<i>An. (Ano.) calderoni</i> [‡] , <i>An. (Nys.) albimanus</i> [¥]

* Ecoregions according to WWF (2015). Subgenera *Ano*: *Anopheles*, *Ker*: *Kerteszia*, *Lop*: *Lophopodomyia*, *Nys*: *Nyssorhynchus*, *Ste*: *Stethomyia*. [¥] Primary malaria vector; [‡] Local malaria vector.

Checklist of *Anopheles* mosquitos in Colombia

Order Diptera Linnaeus, 1758

Infraorder Culicomorpha Hennig, 1948

Superfamily Culicoidea Meigen, 1818

Family Culicidae Meigen, 1818

Subfamily Anophelinae Grassi, 1900

Genus *Anopheles* Meigen, 1818

Subgenus *Anopheles* Meigen, 1818

***Anopheles (Anopheles) apicimacula* Dyar & Knab, 1906**

Distribution. Antioquia*, Arauca, Bolívar, Boyacá, Caldas, Casanare, Cauca, Cesar, Chocó*, Córdoba*, Cundinamarca, Guaviare, Huila, La Guajira, Magdalena, Meta*, Nariño*, Norte de Santander, Putumayo, Risaralda, Santander*, Sucre, Tolima, Valle del Cauca*.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Brochero et al. (2005), González and Carrejo (2009), Parra-Henao et al. (2012), Montoya et al. (2017), SIB (2020), Zuleta-Ruiz et al. (2022).

***Anopheles (Anopheles) calderoni* Wilkerson, 1991**

Distribution. Antioquia, Caldas, Chocó*, Huila, La Guajira, Magdalena, Nariño*, Norte de Santander, Quindio, Risaralda*, Tolima, Valle del Cauca*.

Notes. Local malaria vector. Reported by González and Carrejo (2009), González et al. (2010), Lucumi-Aragón et al. (2011), Orjuela et al. (2015), Montoya et al. (2017), Galeano-Castañeda et al. (2019), SIB Database (2020).

***Anopheles (Anopheles) costai* Da Fonseca & Ramos, 1939**

Distribution. Amazonas*, Antioquia, Arauca, Bolívar, Boyacá, Caldas, Caquetá, Cesar, Chocó, Córdoba, Cundinamarca, Guainía, Guaviare, Huila, Meta*, Nariño, Putumayo*, Santander, Sucre, Valle del Cauca, Vaupés.

Notes. Reported by González and Carrejo (2009), Gutiérrez et al. (2009), Ahumada et al. (2013), Orjuela et al. (2013), SIB (2020). *Anopheles costai* was resurrected from synonymy with *Anopheles mediopunctatus* (Sallum et al. 1999), and it was previously erroneously reported in Colombia as *An. mediopunctatus*, as pointed out by Sallum et al. (1999) and Quiñones et al. (2001).

***Anopheles (Anopheles) eiseni* Coquillett, 1902**

Distribution. Antioquia, Boyacá, Caldas, Casanare, Cauca, Chocó*, Cundinamarca, Huila, La Guajira, Magdalena, Meta, Nariño, Norte de Santander, Quindío, Risaralda, Santander, Tolima, Valle del Cauca*.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), SIB (2020).

***Anopheles (Anopheles) fluminensis* Root, 1927**

Distribution. Norte de Santander.

Notes. Reported by González and Carrejo (2009).

***Anopheles (Anopheles) forattinii* Wilkerson & Sallum, 1999**

Distribution. Meta, Vaupés.

Notes. Reported by Wilkerson and Sallum (1999). It was indicated that *An. forattinii* was previously reported in Colombia as *An. mediopunctatus* (Sallum et al. 1999; Quiñones et al. 2001).

***Anopheles (Anopheles) malefactor* Dyar & Knab, 1907**

Distribution. Antioquia*, Chocó*, Córdoba*, Meta, Norte de Santander*.

Notes. Reported by Wilkerson (1990), Álvarez et al. (2018), SIB Database (2020). New occurrence data from Monteria municipality, Córdoba Department.

***Anopheles (Anopheles) mattogrossensis* Lutz & Neiva, 1911**

Distribution. Amazonas*, Arauca, Caquetá*, Cauca, Guainía, Guaviare, Meta, Norte de Santander, Putumayo, Vaupés, Vichada*.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Brochero et al. (2006), Orjuela et al. (2013), Brochero and Conn (2015), Álvarez et al. (2018), Prado et al. (2019).

***Anopheles (Anopheles) neomaculipalpus* Curry, 1931**

Distribution. Amazonas, Antioquia*, Arauca, Atlántico, Bolívar, Boyacá, Caldas, Casanare, Caquetá, Cauca, Cesar, Chocó, Córdoba*, Cundinamarca, Guaviare, Huila, La Guajira*, Magdalena, Meta, Nariño, Norte de Santander*, Putumayo*, Santander*, Sucre, Tolima, Valle del Cauca, Vaupés.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Brochero et al. (2006), Parra-Henao and Alarcón (2008), González and Carrejo (2009), Parra-Henao et al. (2012), Orjuela et al. (2013), Álvarez et al. (2018). New occurrence data from Cáceres Municipality, Antioquia Department, and Montería and Valencia municipalities, Córdoba Department.

***Anopheles (Anopheles) peryassui* Dyar & Knab, 1908**

Distribution. Amazonas*, Antioquia*, Caldas, Caquetá, Casanare, Cundinamarca, Guainía, Guaviare, Meta*, Putumayo, Santander*, Sucre, Vaupés, Vichada*.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), González and Carrejo (2009), Brochero and Conn (2015), Álvarez et al. (2018), SIB (2020).

***Anopheles (Anopheles) pseudopunctipennis* Theobald, 1901**

Distribution. Antioquia*, Arauca, Atlántico, Bolívar, Boyacá, Caldas, Caquetá, Casanare, Cauca, Cesar, Chocó, Córdoba*, Cundinamarca, Guaviare, Huila, La Guajira, Magdalena*, Meta*, Nariño, Norte de Santander, Putumayo, Quindío, Risaralda, Santander*, Sucre, Tolima, Valle del Cauca*.

Notes. Local malaria vector. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Brochero et al. (2006), Parra-Henao and Alarcón (2008), González and Carrejo (2009), Parra-Henao et al. (2012), Naranjo-Díaz et al. (2013), Montoya et al. (2017), SIB (2020). New occurrence data from Apartadó, Arboletes, Cáceres, Necoclí, San Juan de Urabá and Tarazá Municipalities in Antioquia Department, and Canalete Municipality in Córdoba Department.

***Anopheles (Anopheles) punctimacula* Dyar & Knab, 1906**

Distribution. Antioquia*, Arauca, Atlántico, Bolívar, Boyacá*, Caldas, Caquetá, Casanare, Cauca, Cesar, Chocó*, Córdoba*, Cundinamarca, Guaviare, Huila, La Guajira*, Magdalena*, Meta, Nariño*, Norte de Santander, Putumayo*, Quindío, Risaralda, Santander, Sucre, Tolima, Valle del Cauca*, Vaupés, Vichada.

Notes. Local malaria vector. Reported by Barreto-Reyes (1955), Knight and Stone (1977), González and Carrejo (2009), Gutiérrez et al. (2009), Orjuela et al. (2013), Naranjo-Díaz et al. (2013, 2014), Montoya et al. (2017), Álvarez et al. (2018), Galeano-Castañeda et al. (2019), SIB (2020). New occurrence data from Apartadó, Caucasia and Necoclí Municipalities in Antioquia Department, and Canalete Municipality in Córdoba Department.

***Anopheles (Anopheles) shannoni* Davis, 1931**

Distribution. Vaupés, Vichada*.

Notes. Reported by González and Carrejo (2009), Brochero and Conn (2015).

***Anopheles (Anopheles) vestitipennis* Dyar & Knab, 1906**

Distribution. Cesar, Valle del Cauca.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977).

Subgenus *Kerteszia* Theobald, 1905

***Anopheles (Kerteszia) bambusicolus* Komp, 1937**

Distribution. Caquetá, Meta*.

Notes. Reported by Gast (1943), Barreto-Reyes (1955), Knight and Stone (1977), SIB Database (2020).

***Anopheles (Kerteszia) bellator* Dyar & Knab, 1906**

Distribution. Putumayo*.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), SIB Database (2020).

***Anopheles (Kerteszia) boliviensis* (Theobald, 1905)**

Distribution. Caldas, Cauca, Chocó, Cundinamarca, Huila, Meta, Nariño, Tolima.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), González and Carrejo (2009).

***Anopheles (Kerteszia) homunculus* Komp, 1937**

Distribution. Boyacá, Cauca, Chocó, Cundinamarca, Meta*, Norte de Santander, Tolima.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), González and Carrejo (2009), SIB Database (2020).

***Anopheles (Kerteszia) neivai* Howard, Dyar & Knab, 1913**

Distribution. Antioquia, Bolívar, Boyacá, Cauca, Chocó*, Cundinamarca, Nariño*, Norte de Santander, Tolima, Valle del Cauca*, Vaupés.

Notes. Local malaria vector. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Solarte et al. (1996), Gutiérrez et al. (2008, 2009), González and Carrejo (2009), Naranjo-Díaz et al. (2014, 2023), SIB (2020).

***Anopheles (Kerteszia) pholidotus* Zavortink, 1973**

Distribution. Caquetá, Cundinamarca, Magdalena, Meta*, Norte de Santander, Putumayo*, Tolima*, Valle del Cauca.

Notes. Local malaria vector. Reported by González and Carrejo (2009), Escovar et al. (2014), SIB (2020). Previously reported as *Anopheles lepidotus* (Escovar et al. 2014), a species that is not present in the country.

Subgenus *Lophopodomyia* Antunes, 1937

***Anopheles (Lophopodomyia) gilesi* (Peryassú, 1908)**

Distribution. Meta.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977).

***Anopheles (Lophopodomyia) oiketorakras* Osorno-Mesa, 1947**

Distribution. Cundinamarca, Nariño.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977).

***Anopheles (Lophopodomyia) squamifemur* Antunes, 1937**

Distribution. Antioquia*, Cauca, Chocó, Norte de Santander, Valle del Cauca, Vaupés.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977). New occurrence data from Cáceres Municipality in Antioquia Department.

Subgenus *Nyssorhynchus* Blanchard, 1902

***Anopheles (Nyssorhynchus) albimanus* Wiedemann, 1820**

Distribution. Antioquia*, Atlántico, Bolívar, Cauca*, Cesar, Chocó*, Córdoba*, La Guajira, Magdalena, Nariño*, Risaralda*, Sucre, San Andrés y Providencia, Valle del Cauca*.

Notes. Primary malaria vector. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Faran (1980), Gutierrez et al. (Gutiérrez et al. 2008, 2009), Calle et al. (2008), González and Carrejo (2009), Naranjo-Díaz et al. (2013, 2014, 2023), Montoya et al. (2017), Galeano-Castañeda et al. (2019), Altamiranda-Saavedra et al. (2023), SIB (2020). New occurrence data from Apartadó, Mutatá, Necoclí, San Juan de Urabá and Tarazá Municipalities in Antioquia Department, Guapi Municipality in Cauca Department, Canalete, Monteria and Puerto Escondido Municipalities in Córdoba Department, and El Zarzal Municipality in Valle del Cauca.

***Anopheles (Nyssorhynchus) albitarsis* Lynch Arribálzaga, 1878**

Distribution. Antioquia*, Meta*, Norte de Santander*, Putumayo*, Vichada*.

Notes. Reported by Calle et al. (2008), Gutiérrez et al. (2009), González and Carrejo (2009), Jiménez et al. (2012), Ahumada et al. (2013), Orjuela et al. (2013), Montoya et al. (2017), Galeano-Castañeda et al. (2019), SIB (2020). In Colombia, *Anopheles albitarsis* was previously reported as *An. albitarsis* s.l. or *An. marajonara*, both are part of the Albitaris Complex, which potentially comprises at least ten species, only five have been formally described (Bourke et al. 2021). Only the molecular variants *An. albitarsis* F and *An. albitarsis* I are reported in the country,

An. marajoara could not be confirmed (Ruiz-Lopez et al. 2012). New occurrence data from Arboletes and Cauca Municipality in Antioquia Department, Moñitos and San Antero Municipalities in Córdoba Department.

***Anopheles (Nyssorhynchus) aquasalis* Curry, 1932**

Distribution. Atlántico, Bolívar, Chocó, Córdoba*, La Guajira*, Magdalena

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Calle et al. (2008). New occurrence data from San Antero Municipality in Córdoba Department.

***Anopheles (Nyssorhynchus) argyritarsis* Robineau-Desvoidy, 1827**

Distribution. Antioquia*, Arauca, Atlántico, Bolívar, Boyacá, Caldas, Caquetá, Casanare, Cauca, Cesar, Chocó, Córdoba*, Cundinamarca, Huila, La Guajira, Magdalena, Meta*, Nariño*, Norte de Santander, Putumayo, Quindío, Risaralda, Santander*, Tolima, Valle del Cauca, Vaupés, Vichada*.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), González and Carrejo (2009), Jiménez et al. (2012), Parra-Henao et al. (2012) Naranjo-Díaz et al. (2013), SIB (2020). New occurrence data from Cáceres Municipality in Antioquia Department.

***Anopheles (Nyssorhynchus) benarrochi* Gabaldón, Cova García & Lopez, 1941**

Distribution. Meta*, Putumayo*, Santander*.

Notes. Local malaria vector. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Quiñones et al. (2001), Calle et al. (2008), González and Carrejo (2009), Parra-Henao et al. (2012), Orjuela et al. (2013). In Colombia, a molecular variant denominated *An. benarrochi* B was reported, distributed in the south of the country (Ruiz et al. 2005).

***Anopheles (Nyssorhynchus) braziliensis* Chagas, 1907**

Distribution. Amazonas, Antioquia*, Arauca, Bolívar, Boyacá, Caldas, Caquetá, Casanare, Cesar, Chocó, Córdoba, Cundinamarca, Guainía, Guaviare*, Huila, La Guajira, Magdalena, Meta*, Norte de Santander, Putumayo*, Santander, Tolima, Valle del Cauca, Vaupés, Vichada*.

Notes. Reported by Barreto-Reyes (1955), Brochero et al. (2005), Calle et al. (2008), González and Carrejo (2009), Naranjo-Díaz et al. (2013, 2023), Jiménez et al. (2012, 2014), Ahumada et al. (2013), SIB (2020). New occurrence data from Arboletes, Cáceres, Cauca and Tarazá Municipalities in Antioquia Department, San Jose del Guaviare Municipality in Guaviare Department.

***Anopheles (Nyssorhynchus) darlingi* Root, 1926**

Distribution. Amazonas*, Antioquia*, Arauca, Bolívar, Boyacá, Caldas, Caquetá, Casanare, Cesar, Chocó*, Córdoba*, Cundinamarca, Guainía, Guaviare*, La Guajira, Magdalena, Meta*, Norte de Santander, Putumayo*, Santander, Sucre, Vaupés, Vichada*.

Notes. Primary malaria vector. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Brochero et al. (2005), Calle et al. (2008), González and Carrejo (2009), Jiménez et al. (2012, 2014), Ahumada et al. (2013), Orjuela et al. (2013), Naranjo-Díaz et al. (2013, 2014, 2016a, 2019), Montoya et al. (2017), Pacheco et al. (2017), Galeano-Castañeda et al. (2019), SIB (2020). New occurrence data from Apartadó, Cáceres, Caucasia and Necoclí Municipalities in Antioquia Department, San Carlos and Valencia Municipalities in Córdoba Department, San Jose del Guaviare Municipality in Guaviare Department.

***Anopheles (Nyssorhynchus) dunhami* Causey, 1945**

Distribution. Amazonas*.

Notes. Reported by Ruiz et al. (2010).

***Anopheles (Nyssorhynchus) evansae* Brèthes, 1926**

Distribution. Córdoba*.

Notes. Reported by Knight and Stone (1977), Parra-Henao and Alarcon (2008).

***Anopheles (Nyssorhynchus) marajoara* Galvão & Damasceno, 1942**

Distribution. Antioquia*, Arauca, Atlántico, Bolívar, Boyacá, Caldas, Caquetá*, Casanare, Cauca, Cesar, Chocó, Córdoba, Cundinamarca, Guaviare, Huila, La Guajira, Magdalena*, Meta*, Norte de Santander*, Putumayo, Santander, Sucre, Tolima, Vaupés, Vichada*.

Notes. Reported by Knight and Stone (1977), González and Carrejo (2009), Brochero et al. (2010), Jiménez et al. (2012).

***Anopheles (Nyssorhynchus) nuneztovari* Gabaldón, 1940**

Distribution. Antioquia*, Arauca*, Bolívar, Boyacá, Caldas, Caquetá, Casanare*, Cauca, Cesar, Chocó*, Córdoba*, Cundinamarca, Huila, Magdalena, Meta, Norte de Santander*, Putumayo, Santander*, Sucre, Tolima, Valle del Cauca*.

Notes. Primary malaria vector. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Calle et al. (2002, 2008), Brochero et al. (2006), Fajardo et al. (2008), Parra-Henao and Alarcón (2008), González and Carrejo (2009), Fonseca-González et al. (2009), Ruiz et al. (2010), Parra-Henao et al. (2012), Naranjo-Díaz et al. (2013, 2014, 2016b, 2023), Montoya et al. (2017), Galeano-Castañeda et al. (2019), Altamiranda-Saavedra et al. (2023), SIB (2020). New occurrence data from Apartadó, Caucasia, Necoclí and Tarazá Municipalities in Antioquia Department, Monteria and Moñitos Municipalities in Córdoba Department.

***Anopheles (Nyssorhynchus) oswaldoi* Peryassú, 1922**

Distribution. Amazonas, Antioquia*, Arauca, Atlántico, Bolívar, Boyacá, Caldas, Caquetá, Casanare, Cauca, Cesar, Chocó, Córdoba, Cundinamarca, Guainía, Guaviare*, Magdalena, Meta, Nariño, Norte de Santander, Putumayo*, Santander*, Sucre, Tolima, Valle del Cauca, Vaupés, Vichada*.

Notes. Reported by Barreto-Reyes (1955), González and Carrejo (2009), Ruiz et al. (2010), Parra-Henao et al. (2012), Jiménez et al. (2012, 2014), Orjuela et al. (2013). New occurrence data from Caucasia Municipality in Antioquia Department, San Jose del Guaviare Municipality in Guaviare Department.

***Anopheles (Nyssorhynchus) parvus* Chagas, 1907**

Distribution. Arauca, Casanare, Meta, Santander*, Vichada.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Parra-Henao et al. (2012), SIB (2020).

***Anopheles (Nyssorhynchus) rangeli* Gabaldón, Cova García & Lopez, 1940**

Distribution. Antioquia*, Arauca*, Bolívar, Boyacá*, Caldas*, Caquetá*, Casanare, Cauca, Cesar, Chocó, Córdoba*, Cundinamarca*, Guainía, Guaviare*, La Guajira, Magdalena, Meta*, Nariño, Norte de Santander, Putumayo*, Santander*, Tolima, Valle del Cauca.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Brochero et al. (2005, 2006), Calle et al. (2008), González and Carrejo (2009), Jiménez et al. (2012), Parra-Henao et al. (2012), Orjuela et al. (2013), SIB (2020). New occurrence data from Arboletes and Caucasia Municipalities in Antioquia Department, Moñitos Municipality in Córdoba Department.

***Anopheles (Nyssorhynchus) strodei* Root, 1926**

Distribution. Antioquia, Arauca*, Bolívar, Boyacá, Caquetá, Casanare, Chocó, Córdoba, Cundinamarca, La Guajira, Meta*, Norte de Santander, Putumayo*, Santander*, Valle del Cauca.

Notes. Reported by Barreto-Reyes (1955), Calle et al. (2008), González and Carrejo (2009), Parra-Henao et al. (2012), SIB Database (2020).

***Anopheles (Nyssorhynchus) triannulatus* Neiva & Pinto, 1922**

Distribution. Amazonas, Antioquia*, Arauca, Atlántico, Bolívar, Boyacá, Caldas, Caquetá, Casanare, Cauca, Cesar, Chocó*, Córdoba*, Cundinamarca, Guaviare, Huila, La Guajira, Magdalena*, Meta*, Nariño, Norte de Santander, Putumayo*, Quindío, Santander*, Sucre, Tolima, Valle del Cauca.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), Brochero et al. (2006), Calle et al. (2008), González and Carrejo (2009), Parra-Henao et al. (2012), Ahumada et al. (2013), Naranjo-Díaz et al. (2013, 2023), Orjuela et al. (2013), Montoya et al. (2017), Atencia-Pineda et al. (2018), SIB (2020). New occurrence data from Apartadó, Arboletes, Caucasia, Necoclí, and San Juan de Urabá Municipalities in Antioquia Department, Canalete, Monteria, Moñitos, San Antero, and Valencia Municipalities in Córdoba Department.

***Anopheles (Nyssorhynchus) trinkae* Faran, 1979**

Distribution. Meta*.

Notes. Reported by González and Carrejo (2009), SIB Database (2020).

Subgenus *Stethomyia* Theobald, 1902

***Anopheles (Stethomyia) kompi* Edwards, 1930**

Distribution. Caquetá.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977).

***Anopheles (Stethomyia) nimbus* (Theobald, 1902)**

Distribution. Cundinamarca, Guaviare, Meta, Vaupés, Valle del Cauca.

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977), González and Carrejo (2009).

***Anopheles (Stethomyia) thomasi* Shannon, 1933**

Notes. Reported by Barreto-Reyes (1955), Knight and Stone (1977).

Discussion

The genus *Anopheles* contains eight subgenera, five of which are present in the Neotropical region (*Anopheles*, *Kerteszia*, *Lophopodomys*, *Nyssorhynchus*, and *Stethomyia*; Harbach 2023). All five subgenera are present in Colombia. The current checklist only includes formally described species and excludes molecularly identified variants. Therefore, a total of 44 species are listed as present in Colombia. Species of the subgenera *Nyssorhynchus* and *Anopheles* are the most widely distributed in the country, occurring in 32 and 31 Colombian departments, respectively. *Nyssorhynchus* is the subgenus with the highest species richness, comprising 17 species. *Stethomyia* has a limited distribution, being found in only three departments. *Stethomyia* and *Lophopodomys* subgenera are represented by three species each. The highest species richness was detected in Meta Department, with a total of 30 species, followed by Norte de Santander with 24. San Andres and Providencia Department exhibited the lowest species richness, with only the main Colombian malaria vector, *An. albimanus*, being recorded in this insular region (Table 1) (Fig. 1).

It is well known that factors such as temperature, rainfall, and humidity affect the geographical distribution of *Anopheles* species (Abiodun et al. 2016). Colombia exhibits a wide variety of ecological conditions, and 34 ecoregions are described (WWF 2015). The current checklist reports the presence of *Anopheles* species in 21 of those ecoregions (Table 2). The Magdalena-Urabá moist forests ecoregion exhibits the highest richness with 23 species, dominated by the species of the subgenera *Nyssorhynchus* and *Anopheles*, 14 and 7 species, respectively. The characteristics of this ecoregion include consistently high temperatures exceeding 28 °C and average annual rainfall within the range of 2,000–4,000 millimeters (WWF 2015). This ecoregion forms part of the important Tumbes-Chocó-Magdalena biodiversity hotspot, which spans southern Panamá to northern Perú and encompasses coastal and lowland areas of the Magdalena River basin and the Urabá region in northwest Colombia. The diverse ecological requirements of *Anopheles* species, which thrive on a variety of landscapes ranging from mosaic tropical rainforests to wetlands, mangrove swamps, and coastal plains, contribute to the high species richness detected in these Colombian regions.

Several members of the genus *Anopheles* are important in public health due to their role as malaria vectors. In Colombia, the annual number of malaria cases has exceeded 70,000 in recent years (INS 2021, 2022, 2023). Of the nine *Anopheles* species implicated in malaria transmission (Olano et al. 2001; Gutiérrez et al. 2008; Orjuela et al. 2013; Naranjo-Díaz et al. 2013), at least one malaria vector species was registered in each of the 21 ecoregions with occurrence data, except for the Eastern Cordillera Real montane forests (Table 1). As the Magdalena-Urabá moist forests ecoregion exhibited the highest species richness, it is not surprising that it also contained most of the malaria vectors, including the three primary vectors (*An. albimanus*, *An. darlingi*, and *An. nuneztovari*) and four local vectors (*An. pseudopunctipennis*, *An. punctimacula*, *An. neivai*, and *An. benarrochi*). The Chocó-Darién moist forests ecoregion was second in the number of vectors present, including the three primary vectors and three local vectors (*An. calderoni*, *An. punctimacula*, and *An. neivai*). These two ecoregions encompass the most important Colombian malaria regions, the Urabá, Bajo Cauca, Alto Sinú, and Pacific regions, which report the highest annual number of malaria cases (INS 2023).

Several factors contribute to the discrepancies between previous catalogs and the current checklist. For example, *An. mediopunctatus* may be restricted to Brazil, and in Colombia, it was likely misidentified as *An. costai* or *An. forattinii* (Sallum et al. 1999; Quiñones et al. 2001). Additionally, *Anopheles cruzii*, although reported from Costa Rica to Argentina, has not been confirmed in Colombia (Wilkerson and Peyton 1991). Furthermore, *Anopheles vargasi* has been registered in neighboring Venezuela (Del Ventura et al. 2013), but there is no evidence of its presence in Colombia (Osorno-Mesa 1947). Mistaken taxonomic assignments, often due to the existence of cryptic species, have led to erroneous reports. For instance, *An. pholidotus* was previously reported as *An. lepidotus* (Escovar et al. 2014).

Conclusions

In recent decades, numerous studies and surveys conducted in the country have provided valuable data on the presence and distribution of *Anopheles* species. The current checklist attempts to compile the available information. The current list includes 44 formally characterized *Anopheles* species from five subgenera, with the subgenera *Nyssorhynchus* and *Anopheles* being the largest and most widely distributed species groups. Information for 66 new occurrence data is also provided. The incorporation of presence records and ecological distributions is essential for accurately estimating the *Anopheles* species diversity and assessing the malaria vectors. This is fundamental for the design and implementation of effective control interventions.

An interesting finding is that most of the new species records are concentrated in specific regions, likely reflecting research interest in the most malaria-endemic regions of Colombia. It is noteworthy that after 47 species had been recorded in the country, the current checklist only includes 44 species. For an *Anopheles* species to be included in the list, it was required to have a formal description and validation; species variants originally described using only molecular methods were excluded. Among the factors contributing to the discrepancies between previous catalogs and the current checklist are the existence of problematic species, or in some cases damaged specimens, which can lead to

misidentifications; also, mistaken taxonomic assignments can occur due to the existence of cryptic species. In addition, for some species, there was no evidence of their presence in Colombia. Finally, while the use of molecular techniques has helped to clarify the taxonomic status of several problematic species, this has led to an increase in the number of molecular variants reported. However, it has also facilitated the correction of erroneous taxonomic assignments, which in turn, may lead to a decrease in the number of species formally described.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This work received support from Escuela de Microbiología, Universidad de Antioquia, project code No. 2023-66350.

Author contributions

Nelson Naranjo-Díaz: specimen identification, data analysis, writing, review & editing. Margarita M. Correa: data analysis, project coordinator, critical revisions and editing of manuscript drafts.

Author ORCIDs

Margarita M. Correa  <https://orcid.org/0000-0003-2419-7269>

Data availability

The dataset containing the information on *Anopheles* species occurrence in Colombia is available in: <https://doi.org/10.5281/zenodo.13527884>. (Data will be available upon publication).

References

- Abiodun GJ, Maharaj R, Witbooi P, Okosun KO (2016) Modelling the influence of temperature and rainfall on the population dynamics of *Anopheles arabiensis*. *Malaria Journal* 15: 1–15. <https://doi.org/10.1186/s12936-016-1411-6>
- Ahumada ML, Pareja PX, Buitrago LS, Quiñones ML (2013) Biting behavior of *Anopheles darlingi* Root, 1926 (Diptera: Culicidae) and its association with malaria transmission in Villavicencio (Meta, Colombia). *Biomédica* 33: 241–250. [PMID: 24652134] <https://doi.org/10.7705/biomedica.v33i2.1492>
- Altamiranda-Saavedra M, Naranjo-Díaz N, Conn JE, Correa MM (2023) Entomological parameters and population structure at a microgeographic scale of the main Colombian malaria vectors *Anopheles albimanus* and *Anopheles nuneztovari*. *PLoS ONE* 18: e0280066. <https://doi.org/10.1371/journal.pone.0280066>
- Álvarez N, Gómez GF, Naranjo-Díaz N, Correa MM (2018) Discrimination of *Anopheles* species of the Arribalzagia Series in Colombia using a multilocus approach. *Infection, Genetics and Evolution* 64: 76–84. <https://doi.org/10.1016/j.meegid.2018.06.018>

- Atencia-Pineda M, Toro-Cantillo A, Hoyos-López RO (2018) Diversidad genética y estructura poblacional de *Anopheles triannulatus* s.l. en Córdoba, Colombia, determinadas mediante el método de región de código de barras de ADN. *Biomédica* 38: 117–126. <https://doi.org/10.7705/biomedica.v38i0.4055>
- Barreto-Reyes P (1955) Lista de Mosquitos de Colombia, S. A. (Diptera: Culicidae). *Anales de la Sociedad de Biología de Bogotá* 7: 46–94.
- Bourke BP, Justi SA, Caicedo-Quiroga L, Pecor DB, Wilkerson RC, Linton YM (2021) Phylogenetic analysis of the Neotropical *Albitarsis* Complex based on mitogenome data. *Parasites and Vectors* 14: 1–18. <https://doi.org/10.1186/s13071-021-05090-w>
- Brault AC, Foy BD, Myles KM, Kelly CLH, Higgs S, Weaver SC, Olson KE, Miller BR, Powers AM (2004) Infection patterns of o'nyong nyong virus in the malaria-transmitting mosquito, *Anopheles gambiae*. *Insect molecular biology* 13: 625–635. <https://doi.org/10.1111/j.0962-1075.2004.00521.x>
- Brochero H, Conn JE (2015) New records of *Anopheles* (Diptera: Culicidae) in Puerto Carreño, Vichada, Colombia. *Check List* 11(1): 1518. <https://doi.org/10.15560/11.1.1518>
- Brochero HL, Rey G, Buitrago LS, Olano VA (2005) Biting activity and breeding sites of *Anopheles* species in the municipality Villavicencio, Meta, Colombia. *Journal of the American Mosquito Control Association* 21: 182–186. [https://doi.org/10.2987/8756-971X\(2005\)21\[182:BAABSO\]2.0.CO;2](https://doi.org/10.2987/8756-971X(2005)21[182:BAABSO]2.0.CO;2)
- Brochero H, Pareja P, Ortiz G, Olano V (2006) Breeding places and biting activity of *Anopheles* species in the municipality of Cimitarra, Santander, Colombia. *Biomédica* 26: 269–277. [PMID: 16925099] <https://doi.org/10.7705/biomedica.v26i2.1416>
- Brochero HL, Li C, Wilkerson RC (2007) A newly recognized species in the *Anopheles* (*Nyssorhynchus*) *albitarsis* complex (Diptera: Culicidae) from Puerto Carreño, Colombia. *The American journal of tropical medicine and hygiene* 76: 1113–1117. [PMID: 17556620] <https://doi.org/10.4269/ajtmh.2007.76.1113>
- Brochero H, Li C, Wilkerson R, Conn JE, Ruiz-García M (2010) Genetic structure of *Anopheles* (*Nyssorhynchus*) *marajoara* (Diptera: Culicidae) in Colombia. *The American journal of tropical medicine and hygiene* 83: 585–595. <https://doi.org/10.4269/ajtmh.2010.09-0482>
- Calle L DA, Quiñones ML, Erazo HF, Jaramillo O N (2002) Morphometric discrimination of females of five species of *Anopheles* of the subgenus *Nyssorhynchus* from Southern and Northwest Colombia. *Memórias do Instituto Oswaldo Cruz* 97: 1191–1195. <https://doi.org/10.1590/S0074-02762002000800021>
- Calle DA, Quiñones ML, Erazo HF, Jaramillo N (2008) Discriminación por morfometría geométrica de once especies de *Anopheles* (*Nyssorhynchus*) presentes en Colombia. *Biomedica* 28: 371–385. <https://doi.org/10.7705/biomedica.v28i3.75>
- Carrejo NS, Gonzalez R (1992) Introducción al conocimiento de los Diptera. Centro Editorial Universidad del Valle, Cali, 197 pp.
- Cienfuegos A, Rosero DA, Naranjo N, Luckhart S, Conn JE, Correa MM (2011) Evaluation of a PCR-RFLP-ITS2 assay for discrimination of *Anopheles* species in northern and western Colombia. *Acta tropica* 118: 128–135. <https://doi.org/10.1016/j.actatropica.2011.02.004>
- Del Ventura F, Liria J, Navarro JC (2013) Determinación de áreas de endemismo en mosquitos (Diptera: Culicidae) en Venezuela, mediante criterios explícitos de optimización. *Boletín de Malariología y Salud Ambiental* 53: 165–182.
- Escovar JE, González R, Quiñones ML, Wilkerson RC, Ruiz F, Harrison BA (2014) Morphology of the larvae, male genitalia and DNA sequences of *Anopheles* (*Kerteszia*) *pholidotus* (Diptera: Culicidae) from Colombia. *Memórias do Instituto Oswaldo Cruz* 109: 1–7. <https://doi.org/10.1590/0074-0276130596>

- Fajardo Ramos M, González Obando R, Fidel Suárez M, López D, Wilkerson R, Sallum MAM (2008) Morphological analysis of three populations of *Anopheles* (*Nyssorhynchus*) *nuneztovari* Gabaldón (Diptera: Culicidae) from Colombia. *Memórias do Instituto Oswaldo Cruz* 103: 85–92. <https://doi.org/10.1590/S0074-02762008000100013>
- Faran M (1980) Mosquitos Studies (Diptera, Culicidae) XXXIV. A revision of the *Albimanus* Section of the subgenus *Nyssorhynchus* of *Anopheles*. *Contributions of the American Entomological Institute* 15: 1–215.
- Faran M, Linthicum L (1981) handbook of the Amazonian species of *Anopheles* (*Nyssorhynchus*) (Diptera: Culicidae). *Mosquito Systematics* 13: 1–81.
- Fonseca-González I, Cárdenas R, Quiñones ML, McAllister J, Brogdon WG (2009) Pyrethroid and organophosphates resistance in *Anopheles* (*N.*) *nuneztovari* Gabaldón populations from malaria endemic areas in Colombia. *Parasitology Research* 105: 1399–1409. <https://doi.org/10.1007/s00436-009-1570-2>
- Gaffigan TV, Wilkerson RC, Pecor JE, Stoffer JA, Anderson T (2014) Systematic Catalog of Culicidae - Walter Reed Biosystematics Unit - Home. <http://www.mosquitocatalog.org/> [October 5, 2023]
- Galeano-Castañeda Y, Urrea-Aguirre P, Piedrahita S, Bascuñán P, Correa MM (2019) Composition and structure of the culturable gut bacterial communities in *Anopheles albimanus* from Colombia. *PLoS ONE* 14. <https://doi.org/10.1371/journal.pone.0225833>
- Gast A (1943) Biología y distribución de los anophelinos en Colombia. *Revista de la Facultad de Medicina* 12: 1–55.
- GBIF (2024) GBIF Occurrence Download. <https://doi.org/10.15468/dl.wp8y fz/> [accessed 08 May 2024]
- Gómez GF, Bickersmith SA, González R, Conn JE, Correa MM (2015) Molecular taxonomy provides new insights into *Anopheles* species of the neotropical arribalzagia series. *PLoS ONE* 10: e0119488. <https://doi.org/10.1371/journal.pone.0119488>
- González R, Carrejo N (2007) Introducción al estudio taxonomico de *Anopheles* de Colombia Claves y notas de distribución. Universidad del Valle, Cali, 237 pp.
- González R, Carrejo N (2009) Introducción al estudio taxonomico de *Anopheles* de Colombia Claves y notas de distribución. 2nd edn. Universidad del Valle, Cali, 260 pp.
- González R, Carrejo N, Wilkerson RC, Alarcon J, Alarcon-Ormasa J, Ruiz F, Bhatia R, Loaiza J, Linton Y-M (2010) Confirmation of *Anopheles* (*Anopheles*) *calderoni* Wilkerson, 1991 (Diptera: Culicidae) in Colombia and Ecuador through molecular and morphological correlation with topotypic material. *Memórias do Instituto Oswaldo Cruz* 105: 1001–1009. <https://doi.org/10.1590/S0074-02762010000800009>
- Gutiérrez LA, Naranjo N, Jaramillo LM, Muskus C, Luckhart S, Conn JE, Correa MM (2008) Natural infectivity of *Anopheles* species from the Pacific and Atlantic Regions of Colombia. *Acta tropica* 107: 99–105. <https://doi.org/10.1016/j.actatropica.2008.04.019>
- Gutiérrez LA, González JJ, Gómez GF, Castro MI, Rosero DA, Luckhart S, Conn JE, Correa MM (2009) Species composition and natural infectivity of anthropophilic *Anopheles* (Diptera: Culicidae) in the states of Córdoba and Antioquia, Northwestern Colombia. *Memórias do Instituto Oswaldo Cruz* 104: 1117–1124. <https://doi.org/10.1590/S0074-02762009000800008>
- Harbach RE (2023) Culicidae Classification. Mosquito Taxonomic Inventory. <http://mosquito-taxonomic-inventory.info/simpletaxonomy/term/6223> [July 21, 2023]
- Heinemann SJ, Belkin JN (1978) Collection records of the project "Mosquitoes of Middle America" 12. Colombia (COA, COB, COL, COM). *Mosquito Systematics* 10: 493–540.
- Hernández-Valencia JC, Rincón DS, Marín A, Naranjo-Díaz N, Correa MM (2020) Effect of land cover and landscape fragmentation on anopheline mosquito abundance and

- diversity in an important Colombian malaria endemic region. PLoS ONE 15. <https://doi.org/10.1371/journal.pone.0240207>
- Hiwat H, Bretas G (2011) Ecology of *Anopheles darlingi* Root with respect to vector importance: a review. Parasites & Vectors 4: 177. <https://doi.org/10.1186/1756-3305-4-177>
- IGAC (2002) Atlas de Colombia. 5th edn. Imprenta Nacional de Colombia, Bogota, 167 pp.
- INS (2021) Instituto Nacional de Salud, Boletín epidemiológico Semanal. Estadísticas del sistema de vigilancia en salud pública- SIVIGILA, Casos totales en la Semana Epidemiológica 52 y acumulados del año, Subdirección de Vigilancia y Control en Salud Pública. <https://portalsivigila.ins.gov.co/Paginas/Vigilancia-Rutinaria.aspx> [May 4, 2024]
- INS (2022) Instituto Nacional de Salud, Boletín epidemiológico Semanal. Estadísticas del sistema de vigilancia en salud pública- SIVIGILA, Casos totales en la Semana Epidemiológica 52 y acumulados del año, Subdirección de Vigilancia y Control en Salud Pública. <https://portalsivigila.ins.gov.co/Paginas/Vigilancia-Rutinaria.aspx> [May 5, 2024]
- INS (2023) Instituto Nacional de Salud, Boletín epidemiológico Semanal. Estadísticas del sistema de vigilancia en salud pública- SIVIGILA, Casos totales en la Semana Epidemiológica 52 y acumulados del año, Subdirección de Vigilancia y Control en Salud Pública. <https://portalsivigila.ins.gov.co/Paginas/Vigilancia-Rutinaria.aspx> [May 7, 2024]
- ITIS (2024) Integrated Taxonomic Information System. <https://www.itis.gov/> [June 6, 2024]
- Jiménez P, Conn JE, Wirtz R, Brochero H (2012) *Anopheles* (Diptera: Culicidae) vectors of malaria in Puerto Carreño municipality, Vichada, Colombia. Biomédica 32: 13. <https://doi.org/10.7705/biomedica.v32i0.414>
- Jiménez P, Conn JE, Brochero H (2014) Malaria vectors in San José del Guaviare, Orinoquia, Colombia. Journal of the American Mosquito Control Association 30(2): 91–98. <https://doi.org/10.2987/13-6382.1>
- Knight KL, Stone A (1977) A catalog of the mosquitoes in the world. (Diptera: Culicidae). Entomological Society of America, Maryland, 611 pp. <https://doi.org/10.4182/FXDZ6899>
- Lucumi-Aragón D, González RO, Salas-Quinchucua C (2011) Actividad de picadura de *Anopheles calderoni* (Diptera: Culicidae) en dos localidades del Valle del Cauca, Colombia. Revista Colombiana de Entomología 37: 256–261. <https://doi.org/10.25100/socolen.v37i2.9086>
- Manguin S, Bangs MJ, Pothikasikorn J, Chareonviriyaphap T (2010) Review on global co-transmission of human *Plasmodium* species and *Wuchereria bancrofti* by *Anopheles* mosquitoes. Infection, Genetics and Evolution 10: 159–177. <https://doi.org/10.1016/j.meegid.2009.11.014>
- Montoya C, Bascuñán P, Rodríguez-Zabala J, Correa MM (2017) Abundance, composition and natural infection of *Anopheles* mosquitoes from two malaria-endemic regions of Colombia. Biomedica 37: 98–105. <https://doi.org/10.7705/biomedica.v37i0.3553>
- Montoya-Lerma J, Solarte YA, Giraldo-Calderón GI, Quiñones ML, Ruiz-López F, Wilkerson RC, González R (2011) Malaria vector species in Colombia: a review. Memórias do Instituto Oswaldo Cruz 106(Suppl): 223–238. <https://doi.org/10.1590/S0074-02762011000900028>
- Naranjo-Díaz N, Rosero DA, Rua-Uribe G, Luckhart S, Correa MM (2013) Abundance, behavior and entomological inoculation rates of anthropophilic anophelines from

- a primary Colombian malaria endemic area. *Parasites & Vectors* 6: 61. <https://doi.org/10.1186/1756-3305-6-61>
- Naranjo-Díaz N, Altamiranda M, Luckhart S, Conn JE, Correa MM (2014) Malaria vectors in ecologically heterogeneous localities of the Colombian Pacific region. *PLoS ONE* 9: e103769. <https://doi.org/10.1371/journal.pone.0103769>
- Naranjo-Díaz N, Conn JE, Correa MM (2016a) Behavior and population structure of *Anopheles darlingi* in Colombia. *Infection, genetics and evolution* 39: 64–73. <https://doi.org/10.1016/j.meegid.2016.01.004>
- Naranjo-Díaz N, Sallum MAM, Correa MM (2016b) Population dynamics of *Anopheles nuneztovari* in Colombia. *Infection, Genetics and Evolution* 45: 56–65. <https://doi.org/10.1016/j.meegid.2016.08.019>
- Naranjo-Díaz N, Altamiranda-Saavedra M, Correa MM (2019) *Anopheles* species composition and entomological parameters in malaria endemic localities of North West Colombia. *Acta Tropica* 190: 13–21. <https://doi.org/10.1016/j.actatropica.2018.10.011>
- Naranjo-Díaz N, Hernandez-Valencia JC, Gomez GF, Correa MM (2023) Spatial and Temporal Diversity Variation in the *Anopheles* Communities in Malaria-Endemic Regions of Colombia. *The American Journal of Tropical Medicine and Hygiene* 108: 744–754. <https://doi.org/10.4269/ajtmh.22-0569>
- Olano V, Brochero H, Sáenz R, Quiñones M, Molina J (2001) Mapas preliminares de la distribución de especies de *Anopheles* vectores de malaria en Colombia. *Biomédica* 21: 402–408.
- Orjuela LI, Herrera M, Erazo H, Quiñones ML (2013) Especies de *Anopheles* presentes en el departamento del Putumayo y su infección natural con *Plasmodium*. *Biomédica* 33: 42–52. <https://doi.org/10.7705/biomedica.v33i1.619>
- Orjuela LI, Ahumada ML, Avila I, Herrera S, Beier JC, Quiñones ML (2015) Human biting activity, spatial-temporal distribution and malaria vector role of *Anopheles calderoni* in the southwest of Colombia. *Malaria Journal* 14: 1–9. <https://doi.org/10.1186/s12936-015-0764-6>
- Osorno-Mesa E (1947) Una nueva especie de *Anopheles* de Bogotá, Colombia. *Caldasia* 4(20): 431–446.
- Pacheco MA, González R, Brochero HL (2017) *Anopheles darlingi* (Diptera: Culicidae) Rood 1926: Morphometric variations in wings and legs of populations from Colombia. *Biomedica* 37: 124–134. <https://doi.org/10.7705/biomedica.v37i0.3492>
- Parra-Henao G, Alarcón EP (2008) Observaciones sobre la bionomía de *Anopheles* spp. (Diptera: Culicidae) en el municipio Valencia, departamento Córdoba, Colombia. *Boletín de Malariología y Salud Ambiental XLVIII*: 95–98.
- Parra-Henao G, Delgado D, Alarcón EP (2012) Especies de *Anopheles* en el Canón del Río Sogamoso, Santander, Colombia, al inicio de la puesta en marcha del proyecto hidroeléctrico. *Boletín de Malariología y Salud Ambiental* 52: 287–294.
- Prado CC, Alvarado-Cabrera LA, Camargo-Ayala PA, Garzón-Ospina D, Camargo M, Soto-De León SC, Cubides JR, Celis-Giraldo CT, Patarroyo ME, Patarroyo MA (2019) Behavior and abundance of *Anopheles darlingi* in communities living in the Colombian Amazon riverside. *PLoS ONE* 14(3): e0213335. <https://doi.org/10.1371/journal.pone.0213335>
- Quiñones ML, Harbach RE, Calle DA, Ruiz F, Erazo HF, Linton YM (2001) Variante morfológica de adultos hembras de *Anopheles benarrochi* (Diptera: Culicidae) en Putumayo, Colombia. *Biomédica* 21: 351. <https://doi.org/10.7705/biomedica.v21i4.1128>
- Ruiz F, Quiñones ML, Erazo HF, Calle DA, Alzate JF, Linton YM (2005) Molecular differentiation of *Anopheles* (*Nyssorhynchus*) *benarrochi* and *An. (N.) oswaldoi* from

- southern Colombia. *Memorias do Instituto Oswaldo Cruz* 100: 155–160. <https://doi.org/10.1590/S0074-02762005000200008>
- Ruiz F, Linton YM, Ponsonby DJ, Conn JE, Herrera M, Quiñones ML, Vélez ID, Wilkerson RC (2010) Molecular comparison of topotypic specimens confirms *Anopheles* (*Nyssorhynchus*) *dunhami* Causey (Diptera: Culicidae) in the Colombian Amazon. *Memorias do Instituto Oswaldo Cruz* 105: 899. <https://doi.org/10.1590/S0074-02762010000700010>
- Ruiz-Lopez F, Wilkerson RC, Conn JE, McKeon SN, Levin DM, Quiñones ML, Póvoa MM, Linton Y-M (2012) DNA barcoding reveals both known and novel taxa in the Albitarsis Group (*Anopheles*: *Nyssorhynchus*) of Neotropical malaria vectors. *Parasites & vectors* 5: 44. <https://doi.org/10.1186/1756-3305-5-44>
- Sallum MA, Wilkerson RC, Forattini OP (1999) Taxonomic study of species formerly identified as *Anopheles mediopunctatus* and resurrection of *An. costai* (Diptera: Culicidae). *Journal of medical entomology* 36: 282–300. <https://doi.org/10.1093/jmedent/36.3.282>
- SEM Servicio Nacional de Erradicación de la Malaria (1957) Plan para la erradicación de la malaria en Colombia. Volumen II. Ministerio de Salud Pública y Oficina Sanitaria Panamericana, Bogotá, 365 pp.
- SIB Colombia (2020) Sistema de Información sobre Biodiversidad de Colombia. <https://sibcolombia.net/> [accessed 30 May 2024]
- Solarte Y, Hurtado C, Gonzalez R, Alexander B (1996) Man-biting activity of *Anopheles* (*Nyssorhynchus*) *albimanus* and *An. (Kerteszia) neivai* (Diptera: Culicidae) in the Pacific lowlands of Colombia. *Memórias do Instituto Oswaldo Cruz* 91: 141–146. <https://doi.org/10.1590/S0074-02761996000200002>
- Stone A, Knight KL, Starcke H (1959) A synoptic catalog of the mosquitoes of the world (Diptera, Culicidae). *Entomological Society of America*, 358 pp. <https://doi.org/10.4182/UMJB2446>
- WHO (2022) World Malaria Report 2022. World Health Organization: 293. https://reliefweb.int/report/world/world-malaria-report-2022?gclid=CjwKCAiAzp6eBhByEiwA_gGq5NTTnODwucx32cIn1jrven6oy0qHvmQAoWLOEotgDzgzvcTKsdJrPRoCrbkQAvD_BwE [accessed 18 January 2023]
- Wilkerson RC (1990) Redescriptions of *Anopheles punctimacula* and *An. malefactor* (Diptera: Culicidae). *Journal of medical entomology* 27: 225–247. <https://doi.org/10.1093/jmedent/27.2.225>
- Wilkerson R, Peyton E (1991) The Brazilian malaria vector *Anopheles* (*Kerteszia*) *cruzei*: life stages and biology (Diptera: Culicidae). *Mosquito Systematics* 23: 110–122.
- Wilkerson RC, Sallum MA (1999) *Anopheles* (*Anopheles*) *forattinii*: a new species in Series Arribalzagia (Diptera: Culicidae). *Journal of medical entomology* 36: 345–354. <https://doi.org/10.1093/jmedent/36.3.345>
- WWF (2015) List of Ecoregions. World Wildlife Fund. http://wwf.panda.org/about_our_earth/ecoregions/ecoregion_list/
- Zapata MA, Cienfuegos A V, Quirós OI, Quiñones ML, Luckhart S, Correa MM (2007) Discrimination of seven *Anopheles* species from San Pedro de Urabá, Antioquia, Colombia, by polymerase chain reaction-restriction fragment length polymorphism analysis of its sequences. *The American journal of tropical medicine and hygiene* 77: 67–72. <https://doi.org/10.4269/ajtmh.2007.77.67>
- Zuleta-Ruiz BR, Gómez-Vargas W, Ortiz-Reyes A, Ruiz-López F (2022) Identificación de *Anopheles* spp. (Diptera: Culicidae) presentes en áreas endémicas de transmisión de malaria en Turbo, Antioquia y su infección con *Plasmodium* spp. *Actualidades Biológicas* 44(116): e5. <https://doi.org/10.17533/udea.acbi.v44n116a05>

Supplementary material 1

Data on *Anopheles* species occurrence in Colombia

Authors: Nelson Naranjo-Díaz, Margarita M. Correa

Data type: xlsx

Explanation note: Geographic coordinates are shown in decimal degrees and were taken from database portals, scientific articles, and the research group unpublished new occurrence data.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1231.133711.suppl1>