







Plume moths (Lepidoptera, Pterophoridae) of a recently discovered lepidopteran diversity hotspot in the Mount Cameroon area, with descriptions of four new species

Peter Ustjuzhanin^{1,2}, Vasily N. Kovtunovich³, Sylvain Delabye^{4,5}, Vincent Maicher^{5,6}, Szabolcs Sáfian⁷, Eric B. Fokam⁸, Robert Tropek^{4,5}

1 *Altai State University, Lenina 61, Barnaul, RU-656049, Russia*

2 *Biological Institute, Tomsk State University, Lenina Prospekt 36, Tomsk 634050, Russia*

3 *University of the Free State (UFS), 205 Nelson Mandela Dr., Park West, Bloemfontein, Free State, South Africa*

4 *Department of Ecology, Faculty of Science, Charles University, Viničná 7, CZ-12843 Prague, Czech Republic*

5 *Institute of Entomology, Biology Centre of the Czech Academy of Sciences, Branišovská 31, CZ-37005 České Budějovice, Czech Republic*

6 *The Nature Conservancy Gabon, Impasse Edouard, Libreville, Gabon*

7 *Hungarian Natural History Museum, Department of Zoology, Baross utca 13, H-1088 Budapest, Hungary*

8 *Department of Animal Biology and Conservation, Faculty of Science, University of Buea, P.O. Box 63, Buea, Cameroon*

Corresponding author: Robert Tropek (robert.tropek@gmail.com)



Academic editor: Kevin Keegan

Received: 19 October 2024

Accepted: 4 February 2025

Published: 11 March 2025

ZooBank: <https://zoobank.org/DF0C00D9-04D3-4C2C-8CA0-DCD81ED76D73>

Citation: Ustjuzhanin P, Kovtunovich VN, Delabye S, Maicher V, Sáfian S, Fokam EB, Tropek R (2025) Plume moths (Lepidoptera, Pterophoridae) of a recently discovered lepidopteran diversity hotspot in the Mount Cameroon area, with descriptions of four new species. ZooKeys 1231: 145–168. <https://doi.org/10.3897/zookeys.1231.139530>

Copyright: © Peter Ustjuzhanin et al.
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Abstract

Moth diversity on Mount Cameroon, a critical biodiversity hotspot in the Afrotropics, remains understudied despite the region's rich and unique ecosystems. In this study, 34 species of plume moths (Pterophoridae) were recorded from the Mount Cameroon region, including four species new to science: *Titanoptilus bigoti* Ustjuzhanin & Kovtunovich, **sp. nov.**, *Titanoptilus murkwe* Ustjuzhanin & Kovtunovich, **sp. nov.**, *Hellinsia ekonjo* Ustjuzhanin & Kovtunovich, **sp. nov.**, and *Hellinsia mapanja* Ustjuzhanin & Kovtunovich, **sp. nov.** Images of the adult type specimen and male genitalia for *Titanoptilus melanodonta* Hampson, 1905 are published for the first time. In addition, 25 species are reported as new records for the Cameroonian fauna, raising the number of known Pterophoridae species in the country from 19 to 48. The results significantly expand our understanding of plume moth diversity in the region and extend the known distribution range of several species. These findings emphasise the importance of Mount Cameroon as a biodiversity hotspot within the Afrotropics. Enhanced conservation efforts are essential to preserve the unique biodiversity of Mount Cameroon, especially considering threats such as ongoing habitat degradation in some parts of the region and climate change.

Key words: Afrotropics, biodiversity, Cameroon, microlepidoptera, new species, Pterophoridae, taxonomy, tropical rainforest

Introduction

Mount Cameroon (4,040 m a.s.l.), an active volcano in southwestern Cameroon, is recognised as one of the key centres of insect diversity within the Afrotropics, particularly for lepidopterans (Ustjuzhanin et al. 2018, 2020a, 2024; Larsen 2005). Its position within the biodiversity-rich Guineo-Congolian forest zone,

combined with its gradients in elevation (ranging from sea level to 4,040 metres) and precipitation (from its southwestern foothills, which rank among the three rainiest places in the world, to relatively dry savannahs in rain shadows on its northeastern slopes; Cable and Cheek 1998; Maicher et al. 2020a), supports highly heterogeneous habitats that accommodate large numbers of both widespread and endemic species of insects and other taxa (e.g., Ustjuzhanin et al. 2018; Delabye et al. 2020). Moreover, due to its inaccessibility and, later, its legal protection as part of the Mount Cameroon National Park, large portions of Mount Cameroon's ecosystems have remained relatively well-preserved from the negative effects of human activity (Cable and Cheek 1998).

Several lepidopteran groups, including butterflies (e.g., Sáfián and Tropek 2016; Maicher et al. 2018, 2020a, b; Delabye et al. 2021), various microlepidopterans (Ustjuzhanin et al. 2018, 2020a, 2024), and other moth groups (e.g., Maicher et al. 2016, 2020a, b; Yakovlev and Sáfián 2016; Przybyłowicz et al. 2019a, b; Delabye et al. 2020; Mertens et al. 2021), have been the focus of recent studies on Mount Cameroon, revealing the exceptional diversity of these taxa on the mountain. Notably, prior research has identified Mount Cameroon as a hotspot for many-plumed moths (Alucitidae), significantly advancing our understanding of their Afrotropical biodiversity (Ustjuzhanin et al. 2018, 2020a, 2024). Despite this progress, many lepidopteran groups remain understudied in the Mount Cameroon region, highlighting the need for further exploration of its insect diversity.

Pterophoridae, commonly referred to as plume moths, represent one such understudied family in the region. Characterised by their often deeply cleft wings, which give them their distinctive “plume-like” appearance, this group has a global distribution but remains insufficiently documented in the Afrotropics (De Prins and De Prins 2025). Knowledge of plume moths in Cameroon is still very limited, as no comprehensive studies on this family have been conducted in the country. To date, only 19 species have been recorded from Cameroon (De Prins and De Prins 2025). By summarising plume moth material sampled over several years of ecological studies of moth communities in the Mount Cameroon region, including descriptions of new species and numerous new country records, this study provides the first comprehensive survey of Pterophoridae on Mount Cameroon.

Materials and methods

We sampled plume moths on Mount Cameroon using standardised light trapping methods over multiple field seasons between 2014 and 2018, at localities spanning elevations from 30 to 2,400 m a.s.l. across the mountain's southwestern, southern, and northern slopes. Light traps equipped with white and ultraviolet (UV) emitting bulbs were operated from dusk until dawn to attract moths. Specimens were collected manually from white sheets, euthanised using ammonium vapour, pinned, and preliminarily spread directly in the field for easier later processing, and stored dried with silica gel for transportation. The collected moths were identified based on external morphology and genitalia dissections; dissection and preparation of genitalia followed the standard lepidopteran taxonomic protocols described in Ustjuzhanin et al. (2018). The genitalia were mounted on permanent microscopic slides for further study, with unique identification codes assigned to each specimen for reference in taxonomic work. Holotypes will be

deposited in the Nature Education Centre, Jagiellonian University, Kraków, Poland (**NECJU**), while paratypes and other specimens will be split between NECJU and the personal collections of P. Ustjuzhanin and V. Kovtunovich, located in Novosibirsk and Moscow, Russia (**CUK**). Morphological terminology follows Zagulajev (1986) and the distribution of individual species follows the Afromoths database (De Prins and De Prins 2025), unless specified otherwise.

The sampling localities are listed below in an alphabetical order:

- Bamboo Camp, 350 m a.s.l., Mount Cameroon (SW slope), 4.0879°N, 9.0505°E; a lowland rainforest with historical disturbances from selective logging.
- Bimbia-Bonadikombo, 30 m a.s.l., Bimbia-Bonadikombo Community Forest, Mexico Camp, 3.9818°N, 9.2625°E; littoral forest in the part of the community forest that is officially disturbance-free, but with intensive ongoing illegal logging (Ferenc et al. 2018).
- Crater Lake, 1,500 m a.s.l., Mount Cameroon (SW slope), 4.1443°N, 9.0717°E; submontane rainforest with a sparse canopy layer as a consequence of local disturbances by forest elephants (Maicher et al 2020b).
- Drink Gari, 650 m a.s.l., Mount Cameroon (SW slope), Drink Gari camp (also known as “Drinking Gari”), 4.1014°N, 9.0610°E; lowland rainforest with a dense canopy layer.
- Ekonjo, 1,500 m a.s.l., Mount Cameroon (S slope), 4.0881°N, 9.1168°E; upland closed-canopy rainforest belonging to the Ekonjo village.
- Ekonjo farmland, 800 m a.s.l., Mount Cameroon (S slope), 4.0687°N, 9.1311°E; farmlands and secondary growths surrounding the Ekonjo village.
- Elephant Camp, 1,850 m a.s.l., Mount Cameroon (SW slope), 4.1170°N, 9.0729°E; montane rainforest with a sparse canopy layer as a consequence of local disturbances by forest elephants (Maicher et al 2020b).
- Mann’s Spring, 2,200 m a.s.l., Mount Cameroon (SW slope), 4.1428°N, 9.1225°E; montane forest at the natural timberline.
- Mapanja, Mapanja camp, 1,850 m a.s.l., Mount Cameroon (S slope), 4.1157°N, 9.1315°E; montane forest with a mostly closed canopy layer, supplemented by small openings after fallen trees on a steep slope.
- PlanteCam, 1,100 m a.s.l., Mount Cameroon (SW slope), PlanteCam camp (also misspelled as “PlantiCamp” or “PlanteCamp”), 4.1175°N, 9.0709°E; upland rainforest in the transition between lowland and montane zones, with a sparse canopy layer as a consequence of local disturbances by forest elephants (Maicher et al 2020b).
- P&T antenna station, an area surrounding an antenna station of the Ministry of Post and Communications (P&T; sometimes misspelled as “PNT” by foreigners), 2,400 m a.s.l., Mount Cameroon (N slope), 4.21556°N, 9.2725°E; transition between montane forests and subalpine grasslands.

Results

Altogether, we recorded 34 plume moth species in the Mount Cameroon area, four of which are described here as new to science. Additionally, we report 25 species (marked with *) as new records for Cameroon, as only five of the recorded species had been previously known from the country.

Descriptions of the new species

Titanoptilus bigoti Ustjuzhanin & Kovtunovich, sp. nov.

<https://zoobank.org/8EA59C5A-694E-49B9-9B85-2004189371BE>

Figs 1, 2

Type material examined. *Holotype* ♂ (NECJU No. 241015), CAMEROON, Bamboo Camp, 350 m a.s.l., Mount Cameroon, 4.0879°N, 9.0505°E, 14–23.II.2016, Sz. Sáfián, R. Tropek, V. Maicher leg.

External characters. The wingspan is 31 mm. The head and tegulae are pale brown. The thorax is interspersed with dark-brown filiform scales. The labial palpi are thin, straight, and equal in length to the eye diameter, basally brown, and noticeably paler apically. The antennae are thin and dark brown. The forewings are pale brown, with a small elongated brown discal spot at the base. There is a distinct brown spot at the cleft base. The fringe inside the cleft is yellowish brown. The hindwings are yellowish brown. The hindwing fringes are dark, with the third lobe interspersed with evenly distributed dark spatulate scales on the costal margin and forming three distinct patches on the anal margin, a small cluster at the tip, two triangular patches along the middle, and a cluster at the base. The abdomen is long and pale brown, with creamy scales at the base of the thorax extending to the first abdominal segment. The hind legs are long and pale brown with noticeable darkening at the bases of the spurs.

Description. Male genitalia. The genitalia are symmetrical. The valves each comprise two lobes: the basal lobes arise beyond the sacculus are relatively wide and subquadrate, and narrow apically. The distal lobes are narrower, and longer, and $\sim 2 \times$ the length of the basal lobes. Narrow sclerotized processes extend from the distal lobes and are only slightly shorter than those lobes. The uncus is robust, equal in length to the distal lobes of the valves, basally wide, and apically narrow and rounded. The saccus is wide and triangular. Sternum VIII is forked, with sclerotised distal margins. The aedeagus is long and thin in its distal 2/3, bent at a right angle $\sim 1/3$ along its length. The basal part is noticeably thicker up to the bend.

Diagnosis. In terms of size and wings colour, the new species is similar to *Titanoptilus melanodonta* Hampson, 1905 (Figs 3, 4) but differs in the absence of fringe bundles along the lower margin of the second lobe of the forewing



Figures 1, 2. *Titanoptilus bigoti* Ustjuzhanin & Kovtunovich, sp. nov. 1 adult male, holotype, NECJU 2 male genitalia, holotype, NECJU, preparation slide no. 241015. Scale bar: 10 mm.



Figures 3, 4. *Titanoptilus melanodonta* Hampson, 1905 **3** adult male, holotype, NHMUK, locality: [Kenya], British East Africa, N'dimy, Uganda Railway, mile 469, leg. C. S. Betton **4** male genitalia, holotype, NHMUK, gen.pr. no.21811. Scale bar: 10 mm.

and at the base of the third lobe of the hindwing. In the male genitalia, the new species is similar to *Titanoptilus procerus* Bigot, 1969 (male genitalia described and illustrated in Bigot 1969), in having bilobed valves and the narrow, long, and bent aedeagus. However, in the new species, the valves are basally wide, and the aedeagus is bent at a right angle, while in *T. procerus*, the valves are basally narrow, and the aedeagus is arched.

Distribution. Cameroon, Democratic Republic of Congo.

Flight period. On Mount Cameroon, the species was collected in February.

Etymology. The species is named in honour of Louis Bigot, the first researcher who worked on African plume moths, as a token of gratitude for his significant contributions.

Note. In Bigot (1969), this species was misidentified and illustrated as *T. melanodonta* (collected in the Democratic Republic of Congo, d'Eala et de Bokuma, Equateur, leg. L. Bigot), because L. Bigot did not have an opportunity to study the *T. melanodonta* type specimen, and the species' genitalia were not known at that time. Upon examining the male genitalia of *T. melanodonta* type specimen, collected in Kenya (Kenya, British East Africa, N'dimy, Uganda Railway, mile 469, leg. C.S. Betton; Figs 3, 4) and stored in the Natural History Museum in London, UK (NHMUK), PU and VK found that the male image in Bigot (1969) does not correspond to the type specimen. Therefore, we have illustrated the *T. melanodonta* holotype in this study (Figs 3, 4). Instead, the male genitalia image in Bigot (1969) exactly matches the newly described species. Although we do not have access to the specimens, we expect they belong to *T. bigoti*, described in this study, based on Bigot's illustrations.

***Titanoptilus murkwe* Ustjuzhanin & Kovtunovich, sp. nov.**

<https://zoobank.org/ACF340B5-2310-4B02-8D4D-321652E6AFE9>

Figs 5, 6

Type material examined. Holotype ♀ (NECJU no. 241016), CAMEROON, Crater Lake, 1,500 m a.s.l., Mt. Cameroon, 4.1443°N, 9.0717°E, 23–29.IV.2017, V. Maicher, P. Potocký, S. Delabye leg.

External characters. The wingspan is 36 mm. The head and thorax are dark brown, while the tegulae are pale brown. The labial palpi are thin, straight, and

apically acute, with a length equal to the eye diameter. The antennae are thin and dark brown. The forewings are brown, with a narrow oblique yellowish stripe across the base of the first lobe. The first lobe is distally lightened with white scales, while the second lobe is mottled with brown and pale scales. The fringe inside the cleft is dark brown. Along the outer margin of the second lobe, the fringe is narrow and brownish grey but it widens and becomes darker in its distal part. The hindwings are brown, slightly paler than the forewings (unfortunately, not clearly visible in the specimen picture; Fig. 5). The fringe of the third lobe anal margin features a row of short dark spatulate scales on the basal half. These scales narrow further towards the middle before lengthening into a long, rounded scale bundle, which narrows again towards the apical fringe bundle, forming a rounded scale tooth at the lobe terminus. The hind legs are long and pale brown, with noticeable darkening at the bases of the spurs.

Description. Female genitalia. The papillae anales are elongated and oval. The antrum is narrow, tubular, and sclerotized, with the ostium slightly extended. The ductus is thin, long, and membranous. The bursa copulatrix is long and oval, 2 × as long as the ductus, and contains paired robust sclerotized signa in its distal portion, each ~0.24 × the length of the bursa.

Differential diagnosis. In size and wing colour, the new species is similar to *T. melanodonta*, but it differs in the absence of the wide portion of dark spatulate scales along the fringe of the outer margin under the cleft on the second lobe and the narrow basal row of spatulate scales among the fringes of the third lobe. In *T. melanodonta*, a wide scale tooth within the fringe of the anal margin is present under the forewing cleft, and the basal fringe of the third lobe of the hindwing forms a wide triangular scale tooth. In the female genitalia, the presence of signa in the bursa copulatrix distinguishes *T. murkwe* from other large species of the genus.

Distribution. The species is only known from Cameroon.

Flight period. The species was collected in April.



Figures 5, 6. *Titanoptilus murkwe* Ustjuzhanin & Kovtunovich, sp. nov. 5 adult female, holotype, NECJU 6 female genitalia, holotype, NECJU, preparation slide no. 241016. Scale bar: 10 mm.

Etymology. The species name is a noun in apposition, given in honour of the young and talented Cameroonian entomologist, Mercy Murkwe, who accompanied us during sampling of most specimens reported in this study.

***Hellinsia ekonjo* Ustjuzhanin & Kovtunovich, sp. nov.**

<https://zoobank.org/65DE0D2E-884E-46CE-84DE-83E609AC0572>

Figs 7–9

Type material examined. *Holotype* • ♂ (NECJU no. 241017), CAMEROON, Plante-Cam, 1,100 m a.s.l., Mount Cameroon, 4.1175000°N, 9.0709440°E, 09–14.IV.2015, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; *Paratypes* • 1♂, CAMEROON, Buea, Mount Cameroon (SE slope), 09.XI.1986, G. Bassi leg. • 1♀, CAMEROON, Crater Lake, 21.XI.2016, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 3♀, 17–18.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg.; 1♂, 2♀, 23–29.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 1♂, 1♀, CAMEROON, Ekonjo, 24.X.2017, V. Maicher, S. Delabye leg. • 2♀, CAMEROON, Elephant Camp, 19–24.XI.2014, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 4♂, 2♀, 17–22.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg.; 1♀, 18–26.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 1♀, CAMEROON, Mapanja, 14.V.2017, V. Maicher, P. Potocký, S. Delabye leg.; 2♂, 1♀, 23–28.X.2017, V. Maicher, S. Delabye leg.; 1♀, 23.X.2017, V. Maicher, S. Delabye leg. • 3♀, same data as the holotype; 1♂, 2♀, CAMEROON, PlanteCam, 11–18.XII.2014, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 1♂, 3♀, 29.I.–07.II.2016, Sz. Sáfián, R. Tropek, V. Maicher leg. (CUK, NECJU).

External characters. The wingspan ranges from 15 to 19 mm, with the holotype measuring 17 mm. The thorax and tegulae are yellow. The head is covered with clinging yellow scales between the antennae. The collar is brown. The labial palpi are straight, short, and half the length of the eye diameter. The antennae are yellowish grey. The forewings are greyish brown, with the costal margin distally darkened with brown scales. A distinct oblique brown stroke is present at the base of the cleft, not extending as far as the costa. The fringes inside the cleft include a mix of yellow and dark-grey sections. The second lobe has dark-brown filiform scales on the dorsum, with narrow white patches near the apex, and an apical part darkened with brown scales. The hindwings are grey, noticeably darker than the forewings, with a pale brown fringe. The hind legs are yellowish brown.

Description. Male genitalia. The valves are oval and asymmetric. The sacculus on the left valve is relatively narrow and slightly wider apically, and barely extends beyond the middle of the valve. The harpe on the same valve is slightly shorter than the sacculus, slightly bent, and narrows apically. The right sacculus has a plate extending from a bifurcate sclerotised arm. The uncus is straight, acute, and bent slightly along its length. The saccus is slightly moderately rounded. The anellus arms are asymmetric, with the left arm being slightly shorter. Both arms are apically acute. The aedeagus is thin, apically slightly bent, and 2/3 the length of the right valve.

Female genitalia. The papillae anales are narrow and tapered. The posterior apophyses are thin. The distal margin of sternum VII is convex and rounded. The ostium is funnel-shaped, and the antrum is short, tubular, and shifted to the left. The ductus is as thick as the antrum, but membranous and relatively long. The bursa copulatrix is oval and lacks signa.



Figures 7–9. *Hellinsia ekonjo* Ustjuzhanin & Kovtunovich, sp. nov. 7 adult male, holotype, NECJU 8 male genitalia, holotype, NEJCU, preparation slide no. 241017 9 female genitalia, paratype. Scale bar: 5 mm.

Differential diagnosis. The male genitalia of the new species are similar to *Hellinsia adumbratus* (Walsingham, 1881), but the latter differs in the absence of the tapered harpe on the right valve, having a narrower uncus, and an apically curved aedeagus. In the female genitalia, the convex sternum VII and the left-shifted antrum resemble *Hellinsia madecasseus* (Bigot, 1964), but the new species differs by having a narrower ductus and lacking signa in the bursa copulatrix.

Distribution. The species is only known from Cameroon.

Flight period. The species was collected from February to May, and from October to December.

Etymology. The species is named after Ekonjo, a village on the southern slope of Mount Cameroon, where it was collected. Several villagers contributed as field assistants and supported the project in various ways. This dedication aims to support the protection of Mount Cameroon unique biodiversity.

***Hellinsia mapanja* Ustjuzhanin & Kovtunovich, sp. nov.**

<https://zoobank.org/AA56663D-6C2E-463F-B718-2D41DA942DAB>

Figs 10–12

Type material examined. *Holotype* • ♂, (NECJU No. 241019), CAMEROON, Crater Lake, 1,500 m a.s.l., Mount Cameroon, 4.1443°N, 9.0717°E, 21.XI.2016, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; *Paratypes* • 1♂, 1♀, and 3 other ex. CAMEROON, Crater Lake, 23–29.IV.2017, V. Maicher, P. Potocký, S. Delabye leg.; 3♂, 1♀, 17–25.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. • 3♂, CAMEROON, Elephant camp, 17–22.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg.; 5 ex., 18–26.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 2♂, 1♀, and 7 other ex., Cameroon, Mapanja, 10–15.V.2017, V. Maicher, P. Potocký, S. Delabye leg.; 2♂, 2♀, and 3 other ex., 23–28.X.2017, V. Maicher, S. Delabye leg. (CUK, NECJU).

External characters. The wingspan ranges from 17 to 19 mm, with the holotype measuring 19 mm. The head is brown, while the thorax and tegulae are pale yellow. The labial palpi are brown, thin, and short, being half the length of the eye diameter. The antennae are yellow, interspersed with brown scales. The forewings are pale yellow, with an oblique elongated brown streak at the cleft base, reaching the cos-



Figures 10–12. *Hellinsia mapanja* Ustjuzhanin & Kovtunovich, sp. nov. **10** adult male, holotype, NECJU **11** male genitalia, holotype, NEJCU, preparation slide no. 241019 **12** female genitalia, paratype. Scale bar: 5 mm.

tal margin of the first lobe. A more or less round brown discal spot is present near middle of the forewing. The costal margin of the forewing, up to the cleft, is mixed with tiny brown scales. A brown spot is situated medially along the costal margin on the first lobe. The fringe inside the cleft is yellow, with brown filiform scales distally. The second lobe is apically darkened with brown scales. The fringe along the outer edge of the forewing is pale yellow, except for a patch of pale brown filiform scales in its distal part under cleft. The hindwings are yellow and lack any pattern. The fringe on the hindwings is pale yellow, with alternating pale brown portions. The hind legs are pale yellow, interspersed with tiny brown scales.

Description. Male genitalia. The valves are asymmetric, with the left valve narrower and longer than the right valve. The saccular process on the left valve is well-developed and strongly sclerotised, extending far beyond the middle of the valve, and is widened and curved distally. The harpe is shaped like a short, slightly curved spike. The right valve is wide and slightly shorter than the left valve. The sacculus in the distal portion has a well-expressed sclerotized plate with processes on the sides. The uncus is narrow, slightly curved, and apically acute. The saccus is slightly rounded. The anellus arms are short and wide, with the left arm slightly shorter than the right. Both arms are apically narrow and acute. The aedeagus is thin, straight, and less than half as long as the right valve.

Female genitalia. The papillae anales are short and triangular. The posterior apophyses are thin and long. The distal margin of sternum VII is convex and rounded, with a notch on the inner edge. The ostium is funnel-shaped, and the antrum is short, tubular, and shifted to the left. The ductus is short and membranous, smoothly transitioning into the narrow, elongated bursa copulatrix. Small, narrow, elongated signa are present laterally on the bursa.

Differential diagnosis. Externally and in the genital structure, the species is similar to *Hellinsia ekonjo*. The wings are characterised by a pale yellowish tint and an oblique elongated brown streak that reaches the costal edge of the forewing, whereas in *H. ekonjo*, the wings are greyish brown and the elongated streak does not reach the costal edge of the forewing. In the male genitalia, the new species is distinguished by the valves of unequal length and width, the shape and size of the sacculus on the left valve, the shorter harpe, and the shorter, apically straight aedeagus. In the female genitalia, it is characterised by the signa in the bursa copulatrix and the small triangular papillae anales.

Distribution. The species is only known from Cameroon.

Flight period. The species was collected from February to May, and from October to November.

Etymology. The species is named after Mapanja, a village on the southern slope of Mount Cameroon, where it was collected. Several villagers contributed as field assistants and supported the project in various ways. We strongly believe this dedication will also help protect the unique biodiversity of the region.

Other species recorded on Mount Cameroon

****Agdistis clara* Arenberger, 1986**

Agdistis clara Arenberger, 1986: 189. (Type locality: Botswana).

Material examined. 2♂, 1♀ (CUK, NECJU), Cameroon, Ekonjo farmland, 19.II.2017, P. Potocký leg.

Distribution. Botswana, Namibia, Republic of South Africa, and Cameroon.

Note. New species for Cameroon.

****Deuterocopus socotranus* Rebel, 1907**

Deuterocopus socotranus Rebel, 1907: 115. (Type locality: Yemen, Socotra).

Deuterocopus deltoptilus Meyrick, 1930: 565. (Type locality: Uganda).

Deuterocopus henrioti Bigot & Boireau, 2006: 16–17. (Type locality: Cote d'Ivoire).

Material examined. 1♀ (CUK), Cameroon, PlanteCam, 09–14.IV.2015, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.

Distribution. Widely distributed species, besides several Afrotropical countries, it occurs also in Australasian, Oriental, and Palaeartic regions.

Note. New species for Cameroon.

****Titanoptilus procerus* Bigot, 1969**

Titanoptilus procerus Bigot, 1969: 182. (Type locality: West Kasai, Sankuru, Dimbelenge, Democratic Republic of Congo).

Material examined. 1♀ (CUK), Cameroon, PlanteCam, 09–14.IV.2015, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.

Distribution. Congo and Cameroon.

Note. New species for Cameroon.

****Walsinghmiella prolai* (Gibeaux, 1994)**

Titanoptilus prolai Gibeaux, 1994: 82. (Type locality: Comoros).

Material examined. 2♂ (CUK, NECJU), Cameroon, Bamboo Camp, 17–23.IV.2015, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.

Distribution. Comoros, Republic of South Africa, Eswatini, Zimbabwe, Tanzania, and Cameroon.

Note. New species for Cameroon.

***Platyptilia benitensis* Strand, 1913**

Platyptilia benitensis Strand, 1913: 64. (Type locality: Alén, Equatorial Guinea).

Material examined. 1♀, Cameroon, Crater Lake, 17–25.II.2017, P. Potocký, Sz. Sáfián, J. Mertens, Š. Janeček, R. Tropek leg.; 2♂, 1♀, 23–29.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 1♀, Cameroon, PlanteCam, 09–14.IV.2015, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg. • 1♀, Cameroon, Mapanja, 25.X.2017, V. Maicher, S. Delabye leg. (CUK, NECJUK).

Distribution. Equatorial Guinea, Democratic Republic of Congo, Cote d'Ivoire, Cameroon, Kenya, Nigeria, Republic of South Africa, Sao Tome and Principe, Tanzania, and Uganda.

****Platyptilia daemonica* Meyrick, 1932**

Platyptilia daemonica Meyrick, 1932: 109. (Type locality: Jem-Jem Forest, Ethiopia).

Material examined. 1♂, Cameroon, Crater Lake, 24.XI.2016, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 2♂, 17–25.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg.; 1♂, 23–29.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 1♂, 1♀, Cameroon, Elephant Camp, 19–24.XI.2014; 3♂, 18–26.IV.2017, V. Maicher, P. Potocký, S. Delabye leg.; 2♂, 17–22.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. • 3♂, Cameroon, Mann's spring, 07–09.XI.2016, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 1♂, 31.I.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg.; 1 ex., 13.VIII.2017, P. Potocký leg. • 2♂, Cameroon, Mapanja, 22.X.2017, V. Maicher, S. Delabye leg. (CUK, NECJU).

Distribution. Ethiopia, Rwanda, and Cameroon.

Note. New species for Cameroon.

****Platyptilia fletcheri* Ustjuzhanin & Kovtunovich, 2016**

Platyptilia fletcheri Ustjuzhanin & Kovtunovich, 2016: 2. (Type locality: Ruwenzori Range, Lake Mahoma, Uganda).

Material examined. 4♂, 1♀, Cameroon, Mann's spring, 8–12.XI.2016, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 4♂, 1♀, 31.I–04.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. (CUK, NECJU).

Distribution. Uganda, Rwanda (Ustjuzhanin et al. 2016), and Cameroon.

Note. New species for Cameroon.

****Platyptilia gondarensis* Gibeaux, 1994**

Platyptilia gondarensis Gibeaux, 1994: 424. (Type locality: Gondar Province, Ethiopia).

Material examined. 1♀ (CUK), Cameroon, P&T antenna station, 28.III.2015, Sz. Sáfián leg.

Distribution. Ethiopia, Kenya, Uganda, and Cameroon.

Note. New species for Cameroon.

****Platyptilia morophaea* Meyrick, 1920**

Platyptilia morophaea Meyrick, 1920: 38. (Type locality: station 39, Kenya).

Material examined. 3♂, Cameroon, Mann's spring, 08–11.XI.2016, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 35 ex., 28.I–04.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg.; 3 ex., 16–21.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 4 ex., Cameroon, Mapanja, 22–23.X.2017, V. Maicher, S. Delabye leg. (CUK, NECJU).

Distribution. Kenya, Burundi, Democratic Republic of Congo, Ethiopia, Malawi, Tanzania, and Cameroon.

Note. New species for Cameroon.

****Platyptilia mugesse* Kovtunovich & Ustjuzhanin, 2014**

Platyptilia mugesse Kovtunovich & Ustjuzhanin, 2014: 459. (Type locality: Mugesse Forest, Malawi).

Material examined. 2♀, Cameroon, Crater Lake, 17–25.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. • 1♀, Cameroon, Mann's spring, 31.I.2017 • 1♀, Cameroon, Mapanja, 23.X.2017, V. Maicher, S. Delabye leg. (CUK, NECJU).

Distribution. The species was found in Malawi (Kovtunovich et al. 2014) and Cameroon.

Note. New species for Cameroon.

****Platyptilia sciophaea* Meyrick, 1920**

Platyptilia sciophaea Meyrick, 1920: 40. (Type locality: Kenya).

Material examined. 1♀, Cameroon, Mann's spring, 31.I.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. (CUK, NEJCU).

Distribution. Kenya, Uganda, and Cameroon.

Note. New species for Cameroon.

****Platyptiliodes albisignatula* Strand, 1913**

Platyptiliodes albisignatula Strand, 1913: 65. (Type locality: Alén, Equatorial Guinea).

Material examined. 2♂, Cameroon, Bimbia-Bonadikombo, 27–28.XII.2014, V. Maicher, Sz. Sáfián, R. Tropek, S. Janeček, leg. • 1♂, Cameroon, Crater Lake, 17–25.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. (CUK, NECJU).

Distribution. Equatorial Guinea and Cameroon.

Note. New species for Cameroon.

****Amblyptilia direptalis* (Walker, 1864)**

Oxyptilus direptalis Walker, 1864: 934. (Type locality: Cape Province, Republic of South Africa).

Platyptilia amblydectis Meyrick, 1932: 108. (Type locality: Jem-Jem Forest, Ethiopia).

Material examined. 1♂, Cameroon, Crater Lake, 23–29.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. 1♂, Cameroon, Elephant camp, 17–22.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. • 2♂, Cameroon, Mann's spring, 30.I.2017 • (CUK, NECJU).

Distribution. Republic of South Africa, Burundi, Democratic Republic of the Congo, Malawi, Tanzania, Zimbabwe, Kenya, Ethiopia; India, Sri Lanka, Cameroon, and China (Ustjuzhanin et al. 2021).

Note. New species for Cameroon.

****Stenoptilia natalensis* Ustjuzhanin & Kovtunovich, 2010**

Stenoptilia natalensis Ustjuzhanin & Kovtunovich, 2010: 695. (Type locality: Howick, KwaZulu Natal, Republic of South Africa).

Material examined. 1♂, Cameroon, Crater Lake, 02–17.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. • 1♂, Cameroon, Mann's spring, 12.XI.2016, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 9 ex., 28–31.I.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg.; 1♂, 02.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 1♂, 1♀, Cameroon, 22–23.X.2017, V. Maicher, S. Delabye leg. • 1♂, 2♀, Cameroon, P&T antenna station, 28.III.2015, Sz. Sáfián leg. (CUK, NECJU).

Distribution. Republic of South Africa, Lesotho, Malawi, and Cameroon.

Note. New species for Cameroon.

****Vietteilus borbonica* (Viette, 1957)**

Platyptilia borbonica Viette, 1957: 170. (Type locality: Reunion Island).

Material examined. 1♂, 1♀, Cameroon, Crater Lake, 23–29.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 1♂, 1♀, Cameroon, Mann's spring, 13–16.VIII. 2017, P. Potocký leg.; 2♀, Cameroon, Mapanja, 23–29.X.2017. V. Maicher, P. Potocký, S. Delabye leg.; 1♂, 11.XI.2016, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg. • 1♀, Cameroon, Mapanja, 14.V.2017, V. Maicher, P. Potocký, S. Delabye leg.; (CUK, NECJU).

Distribution. Reunion Island, Tanzania, and Cameroon.

Note. New species for Cameroon.

****Bipunctiphorus dimorpha* (Fletcher, 1910)**

Platyptilia dimorpha Fletcher, 1910: 401. (Type locality: Morne Blanc, Mahé, Seychelles).

Platyptilia patriarcha Meyrick, 1912: 54. (Type locality: Mfongosi, Zululand, Republic of South Africa).

Bipunctiphorus etiennei Gibeaux, 1994: 57. (Type locality: above Cialos, Piton Bleu Forest, Réunion).

Material examined. 1♀, Cameroon, Crater Lake, 23–29.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 1♂, 1♀, Cameroon, Mapanja, 11–12.V.2017, V. Maicher, P. Potocký, S. Delabye leg. (CUK, NECJU).

Distribution. Seychelles, Republic of South Africa, Reunion Island, Tanzania, Uganda, Zimbabwe, Kenya, Madagascar, Malawi, Sierra Leone, and Cameroon.

Note. New species for Cameroon.

****Stenoptilodes taprobanes* (Felder & Rogenhofer, 1875)**

Amblyptilia taprobanes Felder & Rogenhofer, 1875: plate 140, fig. 54. (Type locality: Ceylon [Sri Lanka]).

Platyptilia legrandi Bigot, 1962b: 86. (Type locality: Beau Vallon, Mahé, Seychelles).

Stenoptilodes vittata Service, 1966: 11. (Type locality: Nigeria).

Material examined. 1♂, Cameroon, Bamboo Camp, 17–23.IV.2015, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg. • 2♀, Cameroon, PlanteCam, 11–20. XII.2014, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 4 ex., 29.I.–07. II.2016, Sz. Sáfián, R. Tropek, V. Maicher leg. (CUK, NECJU).

Distribution. Widespread throughout tropical and subtropical regions.

Note. New species for Cameroon.

***Sphenarches anisodactylus* (Walker, 1864)**

Sphenarches anisodactylus Walker, 1864: 934. (Type locality: Ceylon [Sri Lanka]).

Platyptilia pygmaeana Strand, 1913: 64. (Type locality: Cameroon).

Material examined. 1♂, Cameroon, Crater Lake, 25.XI.2016, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg. • 1♀, Cameroon, Ekonjo, 21.X.2017, V. Maicher, S. Delabye leg. • 1♂, 2♀, Cameroon, PlanteCam, 11–18.XII.2014, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 2 ex., 29.I.–07.II.2016, Sz. Sáfián, R. Tropek, V. Maicher leg. (CUK, NECJU).

Distribution. Widely distributed in tropical and subtropical regions.

****Procapperia insomnis* (Townsend, 1956)**

Capperia insomnis Townsend, 1956: 93. (Type locality: Nakuru, Kenya).

Procapperia hackeri Arenberger, 2002: 74. (Type locality: Yemen).

Material examined. 1♀ (CUK), Cameroon, PlanteCam, 11–18.XII.2014, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.

Distribution. Kenya, Democratic Republic of the Congo, Yemen, Namibia, Zimbabwe, Republic of South Africa, Malawi, and Cameroon.

Note. New species for Cameroon.

***Megalorhipida leucodactylus* (Fabricius, 1794)**

Pterophorus leucodactylus Fabricius, 1794: 346. (Type locality: Virgin Islands).

Pterophorus defectalis Walker, 1864: 943. (Type locality: Sierra Leone).

Material examined. 1♂ (CUK), Cameroon, Bamboo Camp, 14–23. II.2016, Sz. Sáfián leg.

Distribution. Widespread throughout tropical and subtropical regions.

****Inferuncus strictiformis* (Meyrick, 1932)**

Platyptilia strictiformis Meyrick, 1932: 251. (Type locality: Kampala, Uganda).

Platyptilia spiculivalva Gielis, 1990: 120. (Type locality: E Usambara Mts, Amani, Tanzania).

Material examined. 1♂ (CUK), Cameroon, PlanteCam, 09–14.IV.2015, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.

Distribution. Uganda, Tanzania, Congo, and Cameroon.

Note. New species for Cameroon.

****Inferuncus toxochorda* (Meyrick, 1934)**

Platyptilia toxochorda Meyrick, 1934: 402. (Type locality: Sao Tome).

Platyptilia pentheres Bigot, 1969: 191. (Type locality: Equateur, Bokuma, Democratic Republic of Congo).

Material examined. 1 ♀ (CUK), Cameroon, Mann's spring, 12.VIII.2017, P. Potocký, S. Janeček leg.

Distribution. Sao Tome and Principe, Congo, Democratic Republic of Congo, Tanzania, Ghana, Liberia, and Cameroon.

Note. New species for Cameroon.

****Exelastis pumilio* (Zeller, 1873)**

Mimeseoptilus pumilio Zeller, 1873 (Type locality: Texas, Dallas, U.S.A.).

Mimaeseoptilus gilvidorsis Hedemann, 1896 (Type locality: Saint Croix, American Virgin Islands).

Exelastis gilvidorsis (Hedemann, 1896).

Marasmarcha liophanes Meyrick, 1886 (Type locality: Saint-Denis, Réunion).

Exelastis liophanes (Meyrick, 1886).

Marasmarcha tenax Meyrick, 1913 (Type locality: Barberton, Mpumalanga, Republic of South Africa).

Exelastis tenax (Meyrick, 1913).

Exelastis bergeri Bigot, 1969: 176. (Type locality: Democratic Republic of Congo).

Material examined. 1 ♂ (CUK), Cameroon, PlanteCam, 11–18.XII. 2014, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.

Distribution. Democratic Republic of Congo, Republic of South Africa, Sierra Leone, Uganda, and Cameroon.

Note. New species for Cameroon. Ustjuzhanin et al. (2020b) mentioned *E. bergeri* and *E. tenax* as separate species. Nevertheless, these changes have never been formalised, and until a detailed revision, they remain synonyms of *E. pumilio*.

****Hellinsia ambo* Ustjuzhanin & Kovtunovich, 2011**

Hellinsia ambo Ustjuzhanin & Kovtunovich, 2011: 356. (Type locality: West She-wa, Ethiopia).

Hellinsia ruhuruinia Gielis, 2011: 51. (Type locality: Aberdare N.P., Ruhuruini Gates, Kenya).

Material examined. 1 ♂, Cameroon, Crater Lake, 17.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. • 1 ♀, Cameroon, Ekonjo, 21.X.2017, V. Maicher, S. Delabye leg. • 1 ♂, Cameroon, Elephant Camp, 19–24.XI.2014, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 1 ♂, 18–26.IV. 2017, V. Maicher, P. Potocký, S. Delabye leg.; 1 ex., 17–22.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. • 1 ♂, Cameroon, Mann's spring, 13.VIII.2017, P. Potocký leg.; 1 ♂, 30.I.2017, P. Potocký, Sz. Sáfián, S. Janeček, R. Tropek, leg.; 1 ♂, 15.VIII.2017, P. Potocký leg. • 1 ♂, 1 ♀, Cameroon, Mapanja, 22–23.X.2017, V. Maicher, S. Delabye leg. (CUK, NECJU).

Distribution. Ethiopia, Kenya, Rwanda, Uganda, Ghana, and Cameroon.

Note. New species for Cameroon.

****Picardia eparches* (Meyrick, 1931)**

Pterophorus eparches Meyrick, 1931: 176. (Type locality: Butandiga, Uganda).

Material examined. 5 ex., Cameroon, Crater Lake, 17.II.2017, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 10 ex., 23–29.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 3 ex., Cameroon, Elephant camp, 18–26.IV.2017 • 3 ex., Cameroon, Mann's spring, 16–21.IV.2016, leg.; 1 ex., 04.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg.; 1 ♀, 15.VIII.2017. P. Potocký leg. • 7 ex., Cameroon, Mapanja, 10–15.V.2017, V. Maicher, P. Potocký, S. Delabye leg.; 1 ♂, 23.X.2017, V. Maicher, S. Delabye leg. • 1 ♂, Cameroon, PlanteCam, 09–14.IV.2015, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg. (CUK, NECJU).

Distribution. Uganda, Kenya, Malawi, Zimbabwe, Tanzania (Kovtunovich et al. 2014), Zambia (Ustjuzhanin et al. 2022), and Cameroon.

Note. New species for Cameroon.

***Picardia tropeki* Ustjuzhanin & Kovtunovich, 2022**

Material examined. 1 ♂ (NECJU), Cameroon, PlanteCam, 09–14.IV.2015, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.

Distribution. Cameroon.

Note. This species was previously described from the same material sampled on Mount Cameroon in a separate global overview of the genus *Picardia* (Ustjuzhanin et al. 2022).

****Pselnophorus busoroensis* Gielis, 2011**

Pselnophorus busoroensis Gielis, 2011: 51. (Type locality: Nyungwe NP, Busoro, Rwanda).

Material examined. 1 ♂, 1 ♀, Cameroon, Crater Lake, 23–29. IV. 2017, V. Maicher, P. Potocký, S. Delabye leg. • 1 ♂, 1 ♀, Cameroon, Elephant camp, 17–22. II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. • 1 ♀, Cameroon, Mann's spring, 28.I.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg. (CUK, NECJU).

Distribution. Rwanda and Cameroon.

Note. New species for Cameroon.

****Adaina microdactyla* (Hübner, [1813])**

Alucita microdactyla Hübner, [1813]: pl. 5, figs 26, 27. (Type locality: Europe).

Oidaematophorus madecasseus Gibeaux, 1994: 130. (Type locality: Madagascar).

Material examined. 1 ♀ (CUK), Cameroon, Mapanja, 12.V.2017, V. Maicher, P. Potocký, S. Delabye leg.

Distribution. Madagascar, Democratic Republic of Congo, Tanzania, Malawi, and Cameroon, together with parts of Palearctic and Palaeotropical regions, and New Guinea.

Note. New species for Cameroon.

****Pterophorus cleronoma* (Meyrick, 1920)**

Alucita cleronoma Meyrick, 1920: 41. (Type locality: Mt Kenya, Kenya).

Material examined. 1♀, Cameroon, Crater Lake, 24.XI.2016, V. Maicher, Sz. Sáfián, S. Janeček, R. Tropek leg.; 8 ex., 17–25.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg.; 1♂, 23–29.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 2♂, Cameroon, Elephant camp, 17–22.II.2017, P. Potocký, Sz. Sáfián, R. Tropek, J. Mertens, S. Janeček leg.; 1♂, 18–26.IV.2017, V. Maicher, P. Potocký, S. Delabye leg. • 2♂, Cameroon, Mann's spring, 8–12.XI.2016, P. Potocký, Sz. Sáfián, S. Janeček, R. Tropek leg. • 2♂, Cameroon, Mapanja, 22.X.2017, V. Maicher, S. Delabye leg. (CUK, NECJU).

Distribution. Kenya and Cameroon.

Note. New species for Cameroon.

***Cosmoclostis schouteni* Gielis, 1990**

Cosmoclostis schouteni Gielis, 1990: 123. (Type locality: Daloa, Cote d'Ivoire).

Pterophorus ashanti Arenberger, 1995: 250. (Type locality: Kumasi, Ghana).

Material examined. 1♂, 1♀ (CUK), Cameroon, Drink Gari, 06–15.II.2016, V. Maicher, Sz. Sáfián, R. Tropek leg.

Distribution. Cote d'Ivoire, Cameroon, Liberia, Nigeria, Ghana.

Discussion

In this study, we report 34 species of Pterophoridae from Mount Cameroon, including four species new to science, and 25 species recorded in the country for the first time. These species also represent 14 genera not recorded in Cameroon previously: *Adaina*, *Agditis*, *Amblyptilia*, *Bipunctiphorus*, *Deuterocopus*, *Inferuncus*, *Platyptiloides*, *Procapperia*, *Pselnophorus*, *Stenoptilia*, *Stenoptilodes*, *Titanoptilus*, *Vietteilus*, and *Walsinghmiella*. These records highlight Mount Cameroon as a significant locality for plume moth diversity. Beyond the newly described species, the addition of numerous new records for Cameroon confirms the importance of Mount Cameroon as a previously overlooked site for studying Afrotropical moths (Maicher et al. 2016; Ustjuzhanin et al. 2018, 2020a, 2024; Delabye et al. 2020).

This study substantially increased the documented diversity of plume moths in this underexplored region, raising the number of Pterophoridae species known from Cameroon from 19 (De Prins and De Prins 2025) to 48. Consequently, Cameroon is now known to host more than 10% of the 429 known species from the Afrotropics (based on De Prins and De Prins 2025, plus the new

species described here). This significant contribution emphasises the need for further surveys in understudied areas, such as Cameroon, to fully elucidate the diversity of Afrotropical Pterophoridae.

In addition to the new species descriptions, this study significantly expands the known distribution of several plume moth species. While some of the recorded species also occur in neighbouring countries, nine species (*P. daemonica*, *P. fletcheri*, *P. gondarensis*, *P. mugesse*, *P. sciophaea*, *V. borbonica*, *P. eparches*, *P. busoroensis*, and *P. cleronoma*) were previously known only from East Africa, *A. clara* was known only from South Africa, and two species (*W. prolai* and *S. natalensis*) were previously recorded from both of these distant regions. Although such scattered distributional patterns are documented in other moth groups (Maicher et al. 2016; Delabye et al. 2020) and better-studied taxa like butterflies (Larsen 2005), these range extensions more likely reflect the insufficient knowledge of plume moth distribution in Sub-Saharan Africa. Mount Cameroon is now the easternmost point in the known distribution for the West African species *C. schouteni*, and the westernmost point for the Central African species *T. procerus* and *P. albisignatula*, reflecting Cameroon's position as a biogeographical crossroads between West and Central Africa. Nevertheless, for any biogeographical conclusions and interpretations, much better faunistic and taxonomic knowledge of the Pterophoridae Afrotropical diversity is needed.

Mount Cameroon continues to demonstrate its status as a critical biodiversity hotspot within the Afrotropics. Although the diversity at the site is not as exceptional for Pterophoridae as for Alucitidae (Ustjuzhanin et al. 2018, 2020a, 2024), the number of endemic plume moth species and the biogeographically significant records align with patterns seen in butterflies (Larsen 2005), other moths (Yakovlev and Sáfián 2016), and some other non-lepidopteran taxa (Cable and Cheek 1998; Damaška et al. 2022; Hlaváč et al. 2023; Janeček and Janečková 2025). This can be attributed to the mountain's wide range of habitats, from littoral forests and lowland rainforests to montane cloud forests and subalpine grasslands, with large areas under minimal anthropogenic pressure. Its position at the border of two biogeographic regions, along with its isolation from other montane ranges, further contributes to the area's high level of endemism.

Our findings confirm the critical need to protect the ecosystems of Mount Cameroon. While parts of the region remain relatively well-preserved inside Mount Cameroon National Park, ongoing threats from habitat degradation and deforestation persist, particularly in lowland areas (Cable and Cheek 1998; Ferenc et al. 2018; Dongmo et al. 2023) and on the more densely populated eastern and southeastern slopes, where the regional capital, Buea, is located (Cable and Cheek 1998). Additionally, climate change poses significant risks to high-elevation species uniquely adapted to cooler, more stable environments (e.g., Ivory et al. 2016; Linck et al. 2021). Continued conservation efforts, particularly in the form of habitat protection within the national park and sustainable land-use practices in its buffer zones, will be key to preserving the biodiversity of this globally important site. We hope that dedicating some of the newly described species to local communities will foster greater appreciation for the biodiversity of their ecosystems. Our findings reinforce the urgent need to prioritise the conservation of the Mount Cameroon region to ensure the long-term survival of its unique ecosystems and the species they harbour.

Acknowledgements

We are grateful to Francis E. Luma, Štěpán Janeček, Pavel Potocký jr., Jan E.J. Mertens, Jennifer T. Kimbeng, Mercy Murkwe, Ishmeal N. Kobe, Congo S. Kulu, and several other assistants for their help in the field, and to the MCNP staff for all their assistance. We used the ChatGPT 4o language model for English proofreading. This study was performed under several authorisations from the Ministries of the Republic of Cameroon for Forestry and Wildlife, and for Research and Innovations.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

R. Tropek and S. Delabye were funded by the Czech Science Foundation (project no. 20-16499S).

Author contributions

RT conceived the study ideas and supervised the project; RT, SD, VM, SzS, and EBF sampled the material; PU and VK identified and compared the material, and prepared the species descriptions; RT, PU, and SD wrote the first draft; all authors contributed to writing and approved the final text.

Author ORCIDs

Peter Ustjuzhanin  <https://orcid.org/0000-0002-5222-2241>

Vasily N. Kovtunovich  <https://orcid.org/0000-0001-5091-4263>

Sylvain Delabye  <https://orcid.org/0000-0003-0911-9721>

Vincent Maicher  <https://orcid.org/0000-0002-9147-3529>

Szabolcs Sáfián  <https://orcid.org/0000-0002-0614-4203>

Eric B. Fokam  <https://orcid.org/0000-0003-4531-3580>

Robert Tropek  <https://orcid.org/0000-0001-7499-6259>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Bigot L (1969) Les Lépidoptères Pterophoridae du musée royal de l'Afrique centrale, à Tervuren. *Revue de Zoologie et de Botanique Africaines* 79: 165–206.
- Cable S, Cheek M (1998) The plants of Mount Cameroon: A conservation checklist. Royal Botanic Gardens, Kew, 198 pp.
- Damaška AF, Konstantinov A, Fikáček M (2022) Multiple origins of moss-inhabiting flea beetles (Coleoptera: Chrysomelidae): molecular phylogeny, overview of genera and a new genus from Africa. *Zoological Journal of the Linnean Society* 196: 647–676. <https://doi.org/10.1093/zoolinnean/zlab112>

- De Prins J, De Prins W (2025) AfroMoths. Online database of Afrotropical moth species (Lepidoptera). <http://www.afromoths.net> [accessed 22 January 2025]
- Delabye S, Maicher V, Sáfián S, Potocký P, Mertens J, Przybyłowicz Ł, Murkwe M, Kobe I, Fokam E, Janeček Š, Tropek R (2020) First records of 31 species of butterflies and moths (Lepidoptera) in Cameroon, with remarks on their elevational ranges. *Biodiversity Data Journal* 8: e50543. <https://doi.org/10.3897/BDJ.8.e50543>
- Delabye S, Maicher V, Sáfián S, Doležal J, Altman J, Janeček Š, Kobe IN, Murkwe M, Šebek P, Tropek R (2021) Butterfly and moth communities differ in their response to habitat structure in rainforests of Mount Cameroon. *Biotropica* 53: 567–580. <https://doi.org/10.1111/btp.12900>
- Dongmo MA, Hanna R, Bonebrake TC (2023) Enhancing scientific and community capacity to conserve Central African Lepidoptera. *Biological Conservation* 279: 109938. <https://doi.org/10.1016/j.biocon.2023.109938>
- Ferenc M, Sedláček O, Tropek R, Albrecht T, Altman J, Dančák M, Doležal J, Janeček Š, Maicher V, Majeský L, Motombi FN, Murkwe M, Sáfián S, Svoboda M, Hořák D (2018) Something is missing at the bottom: Importance of coastal rainforests for conservation of trees, birds and butterflies in the Mount Cameroon area. *African Journal of Ecology* 56: 679–683. <https://doi.org/10.1111/aje.12506>
- Hlaváč P, Baňář P, Fokam EB (2023) Pselaphinae of the Cameroon Volcanic Line. Part 3. A new genus of Goniacerini (Coleoptera: Staphylinidae: Goniaceritae) from Mount Cameroon. *Zootaxa* 5323: 255–267. <https://doi.org/10.11646/zootaxa.5323.2.5>
- Ivory SJ, Early R, Sax DF, Russell J (2016) Niche expansion and temperature sensitivity of tropical African montane forests. *Global Ecology and Biogeography* 25: 693–703. <https://doi.org/10.1111/geb.12446>
- Janeček Š, Janečková P (2025) Field Guide to Plants of Mount Cameroon 1. Flowering plants above the forest line. Janečkovi, Třeboň, 159 pp.
- Kovtunovich VN, Ustjuzhanin PY, Murphy RJ (2014) Plume moths of Malawi (Lepidoptera: Pterophoridae). *Zootaxa* 3847: 451–494. <https://doi.org/10.11646/zootaxa.3847.4.1>
- Larsen TB (2005) Butterflies of West Africa. Apollo Books, Stenstrup, 865 pp. <https://doi.org/10.1163/9789004531093>
- Linck EB, Freeman BG, Cadena CD, Ghalambor CK (2021) Evolutionary conservatism will limit responses to climate change in the tropics. *Biology Letters* 17: 20210363. <https://doi.org/10.1098/rsbl.2021.0363>
- Maicher V, Sáfián S, Ishmeal KN, Murkwe M, Kimbeng TJ, Janeček Š, Tropek R (2016) Two genera and nineteen species of fruit-feeding erbid moths (Lepidoptera: Erebidae) recorded in Cameroon for the first time. *Entomological News* 126: 64–70. <https://doi.org/10.3157/021.126.0108>
- Maicher V, Sáfián S, Murkwe M, Przybyłowicz Ł, Janeček Š, Fokam EB, Pycrz T, Tropek R (2018) Flying between raindrops: Strong seasonal turnover of several Lepidoptera groups in lowland rainforests of Mount Cameroon. *Ecology and Evolution* 8: 12761–12772. <https://doi.org/10.1002/ece3.4704>
- Maicher V, Sáfián S, Murkwe M, Delabye S, Przybyłowicz Ł, Potocký P, Kobe IN, Janeček Š, Mertens JEJ, Fokam EB, Pycrz T, Doležal J, Altman J, Hořák D, Fiedler K, Tropek R (2020a) Seasonal shifts of biodiversity patterns and species' elevation ranges of butterflies and moths along a complete rainforest elevational gradient on Mount Cameroon. *Journal of Biogeography* 47: 342–354. <https://doi.org/10.1111/jbi.13740>

- Maicher V, Delabye S, Murkwe M, Doležal J, Altman J, Kobe IN, Desmist, J, Fokam EB, Pycrz T, Tropek R (2020b) Effects of disturbances by forest elephants on diversity of trees and insects in tropical rainforests on Mount Cameroon. *Scientific Reports* 10: 21618. <https://doi.org/10.1038/s41598-020-78659-7>
- Mertens JEJ, Brisson L, Janeček Š, Klomberg Y, Maicher V, Sáfián S, Delabye S, Po-tocký P, Kobe IN, Pycrz T, Tropek R (2021) Elevational and seasonal patterns of butterflies and hawkmoths in plant-pollinator networks in tropical rainforests of Mount Cameroon. *Scientific Reports* 11: 9710. <https://doi.org/10.1038/s41598-021-89012-x>
- Przybyłowicz L, Maicher V, László GM, Sáfián S, Tropek R (2019a) *Amerila* (Lepidoptera: Erebidae: Arctiinae) of Cameroon with morphological remarks on male and female genitalia. *Zootaxa* 4674: 283–295. <https://doi.org/10.11646/zootaxa.4674.2.8>
- Przybyłowicz L, Tarcz S, László GM, Sáfián S, Zilli A (2019b) Revision of the *Amerila syntomina* species complex with description of a new species from West Africa (Lepidoptera: Erebidae: Arctiinae). *Annales Zoologici* 69: 703–717. <https://doi.org/10.3161/00034541ANZ2019.69.4.006>
- Ustjuzhanin P, Kovtunovich V, Sáfián S, Maicher V, Tropek R (2018) A newly discovered biodiversity hotspot of many-plumed moths (Lepidoptera, Alucitidae) in the Mount Cameroon area: first report on species diversity, with description of nine new species. *Zookeys* 777: 119–139. <https://doi.org/10.3897/zookeys.777.24729>
- Ustjuzhanin PY, Kovtunovich VN, Ustjuzhanina AK (2016) New species of “giant” plume moths of the genus *Platyptilia* (Lepidoptera, Pterophoridae) from Uganda. *European Journal of Taxonomy* 172: 1–6. <https://doi.org/10.5852/ejt.2016.172>
- Ustjuzhanin P, Kovtunovich V, Maicher V, Sáfián S, Delabye S, Streltsov A, Tropek R. (2020a) Even hotter hotspot: Description of seven new species of many-plumed moths (Lepidoptera, Alucitidae) from Mount Cameroon. *Zookeys* 935: 103–119. <https://doi.org/10.3897/zookeys.935.49843>
- Ustjuzhanin P, Kovtunovich V, Udovichenko P, Armstrong A, Streltsov A (2020b) Plume moths in the protected areas of KwaZulu Natal province, Republic of South Africa. *Ecologica Montenegrina* 27: 11–21. <https://doi.org/10.37828/em.2020.27.2>
- Ustjuzhanin PY, Kovtunovich VN, Saldaitis A, Streltsov AN (2021) New species of plume moths (Lepidoptera: Pterophoridae) from China, Sichuan Province. *Russian Entomological Journal* 30: 323–327. <https://doi.org/10.15298/rusentj.30.3.10>
- Ustjuzhanin PY, Kovtunovich VN, Streltsov AN (2022) Overview of the species of the genus *Picardia* Gibeaux, 1994 (Lepidoptera: Pterophoridae) in the world fauna, with description of new species. *Ecologica Montenegrina* 52: 12–23. <https://doi.org/10.37828/em.2022.52.2>
- Ustjuzhanin P, Kovtunovich V, Delabye S, Maicher V, Sáfián Sz, Streltsov A, Tropek R (2024) Magnifying the hotspot: Description of nine new species of many-plumed moths (Lepidoptera, Alucitidae), with an identification key to all species known from Cameroon. *Zookeys* 1193: 25–48. <https://doi.org/10.3897/zookeys.1193.111544>
- Yakovlev RV, Sáfián S (2016) *Geraldocossus* gen. nov. (Lepidoptera, Cossidae) from Mount Cameroon (West Africa). *Zootaxa* 4114: 595–599. <https://doi.org/10.11646/zootaxa.4114.5.8>
- Zagulajev AK (1986) Pterophoridae. In: *Opredelitel Nasekomykh Evropeyskoi Chasti SSSR*. Vol. 4. Cheshuekrylye, Part 3. Nauka, Leningrad, pp. 26–215.

Appendix 1

List of all species of Pterophoridae recorded in Cameroon

The list follows the AfroMoths online database (de Prins and de Prins 2025), with the addition of the species included in this paper (new country records are marked by an asterisk).

- **Adaina microdactyla* (Hübner, [1813])
- **Agdistis clara* Arenberger, 1986
- **Amblyptilia direptalis* (Walker, 1864)
- Antarches tessmanni* (Strand, 1913)
- **Bipunctiphorus dimorpha* (Fletcher, 1910)
- Cosmoclostis schouteni* Gielis, 1990
- **Deuterocopus socotranus* Rebel, 1907
- Exelastis crudipennis* (Meyrick, 1932)
- **Exelastis pumilio* (Meyrick, 1913)
- Fletcherella niphadarcha* (Meyrick, 1930)
- **Hellinsia ambo* Ustjuzhanin & Kovtunovich, 2011
- **Hellinsia ekonjo* Ustjuzhanin & Kovtunovich, sp. nov.
- Hellinsia lienigianus* (Zeller, 1852)
- **Hellinsia mapanja* Ustjuzhanin & Kovtunovich sp. nov.
- **Inferuncus strictiformis* (Meyrick, 1932)
- **Inferuncus toxochorda* (Meyrick, 1934)
- Marasmarcha sisyrodes* Meyrick, 1921
- Megalorhipida leucodactylus* (Fabricius, 1794)
- **Picardia eparches* (Meyrick, 1931)
- Picardia tropeki* Ustjuzhanin & Kovtunovich, 2022
- Platyptilia benitensis* Strand, 1913
- **Platyptilia daemonica* Meyrick, 1932
- Platyptilia farfarellus* (Zeller, 1867)
- **Platyptilia fletcheri* Ustjuzhanin & Kovtunovich, 2016
- **Platyptilia gondarensis* Gibeaux, 1994
- **Platyptilia morophaea* Meyrick, 1920
- **Platyptilia mugesse* Kovtunovich & Ustjuzhanin, 2014
- **Platyptilia sciophaea* Meyrick, 1920
- **Platyptiliodes albisignatula* Strand, 1913
- **Procapperia insomnis* (Townsend, 1956)
- **Pselnophorus busoroensis* Gielis, 2011
- Pterophorus albidus* (Zeller, 1852)
- Pterophorus candidalis* (Walker, 1864)
- **Pterophorus cleronoma* (Meyrick, 1920)
- Pterophorus ceraunia* (Bigot, 1969)
- Pterophorus lampra* (Bigot, 1969)
- Pterophorus rhyparias* (Meyrick, 1908)
- Pterophorus spissa* (Bigot, 1969)
- Pterophorus virgo* (Strand, 1913)
- Sphenarches anisodactylus* (Walker, 1864)

Stenodacma wahlbergi (Zeller, 1852)

**Stenoptilia natalensis* Ustjuzhanin & Kovtunovich, 2010

**Stenoptilodes taprobanes* (Felder & Rogenhofer, 1875)

**Titanoptilus bigoti* Ustjuzhanin & Kovtunovich, sp. nov.

**Titanoptilus murkwe* Ustjuzhanin & Kovtunovich, sp. nov.

**Titanoptilus procerus* Bigot, 1969

**Vietteilus borbonica* (Viette, 1957)

**Walsinghmiella prolai* (Gibeaux, 1994)