

# A new subgenus of *Stigmatophora* Staudinger, 1881 (Lepidoptera, Erebidae, Arctiinae, Lithosiini) from China, with descriptions of three new species

Lu Zhang<sup>1</sup>, Tingting Zhao<sup>1</sup>, Huilin Han<sup>1,2,3</sup>

1 School of Forestry, Northeast Forestry University, Harbin 150040, China

2 Northeast Asia Biodiversity Research Center, Northeast Forestry University, Harbin 150040, China

3 Ministry of Education, Key Laboratory of Sustainable Forest Ecosystem Management, Northeast Forestry University, Harbin 150040, China

Corresponding authors: Huilin Han ([hanhuilin@aliyun.com](mailto:hanhuilin@aliyun.com)); Tingting Zhao ([zhaotingting11@foxmail.com](mailto:zhaotingting11@foxmail.com))

## Abstract

We describe three newly discovered species of the genus *Stigmatophora* Staudinger, 1881 from China, (*S. bucseki* **sp. nov.**, *S. longa* **sp. nov.**, and *S. dianensis* **sp. nov.**) and establish a new subgenus *Bifurcistigma* **subgen. nov.** for the first two. Detailed descriptions of the external morphology and male and female genitalia of the new species are provided, along with those of *S. obratzovi* Daniel, 1951, which is also assigned to the new subgenus. A checklist of all species of *Stigmatophora* is presented including the distribution, type locality, and synonymy of each.

**Key words:** Checklist, distribution, footman moth, morphology, taxonomy



Academic editor: Reza Zahiri

Received: 9 October 2024

Accepted: 22 December 2024

Published: 10 February 2025

ZooBank: <https://zoobank.org/134D1DB9-40F3-4375-831E-90AB8627CFEC>

**Citation:** Zhang L, Zhao T, Han H (2025) A new subgenus of *Stigmatophora* Staudinger, 1881 (Lepidoptera, Erebidae, Arctiinae, Lithosiini) from China, with descriptions of three new species. ZooKeys 1226: 303–317. <https://doi.org/10.3897/zookeys.1226.138892>

Copyright: © Lu Zhang et al.  
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Introduction

The genus *Stigmatophora* Staudinger, 1881 belongs to the tribe Lithosiini of the subfamily Arctiinae and was established with *Setina micans* Bremer & Grey, 1852 as the type species from Beijing (Peking), China (Staudinger 1881). Before 2013, 23 *Stigmatophora* species were known (Bremer and Grey 1852; Walker and Gray 1855; Moore 1878; Meyrick 1894; Swinhoe 1894; Blanford 1898; Leech 1898; Seitz 1911; Wileman 1911; Matsumura 1927, 1930; Daniel 1951; Fang 1991), which are distributed in the northern part of the Oriental region, as well as the central and eastern Palaearctic region (Volynkin et al. 2018; Volynkin 2020). In recent years, *Stigmatophora* has been reviewed and divided into two subgenera based on the characters of male genitalia, *Stigmatophora* and *Pseudomiltochrista* Dubatolov & Bucsek, 2016 (Volynkin et al. 2018; Volynkin 2020; Volynkin et al. 2021), three new species were described. To date, the genus contains 26 species worldwide, and among them, 22 species are found in China.

In this study, we describe two new species (*Stigmatophora bucseki* **sp. nov.**, *Stigmatophora longa* **sp. nov.**) with an external morphology closely resembling *S. obratzovi* Daniel, 1951. The wing pattern and male genitalia of three species do not comply with any known subgenera. Therefore, we propose a third subgenus, *Bifurcistigma* **subgen. nov.**, to accommodate them. A third new species,

*S. dianensis* sp. nov., is placed in the subgenus *Pseudomiltochrista* Dubatolov & Bucsek, 2016. Additionally, the male genitalia of *S. obratzsovi* are illustrated for the first time, and a checklist of all species in the genus, including their type localities and distributions, is provided.

## Materials and methods

Specimens of the new species were collected using a 220V/450W mercury lamp. For identification, standard methods for dissection and preparation of the genitalia slides were used (Kononenko and Han 2007). Images of the specimens were taken with a Nikon D700 camera, while the genitalia slides were photographed with an Olympus photo microscope and imaged-stacked using Helicon Focus v. 7.0; figures were further processed in Adobe Photoshop CS6. The terminology of genitalia follows the guidelines set by Volynkin (2024). All type specimens of the new species are deposited in the Insect Collection of Northeast Forestry University (NEFU), Harbin, China.

## Abbreviations used

<b>NEFU</b>	Northeast Forestry University, Harbin, China
<b>TS</b>	type species
<b>TL</b>	type locality

## Taxonomic account

**Family Erebiidae Leach, 1815**

**Subfamily Arctiinae Leach, 1815**

**Tribe Lithosiini Billberg, 1820**

**Genus *Stigmatophora* Staudinger, 1881**

Common name. 痣苔蛾属

*Stigmatophora* Staudinger, 1881: 399.

**TS.** *Setina micans* Bremer & Grey, 1852.

**TL.** Peking region (China, Beijing).

**Subgenus *Bifurcistigma* subgen. nov.**

<https://zoobank.org/6E96C5DE-EC89-48DC-ACE1-7474AC8B709F>

Common name. 岔带痣苔蛾亚属

**TS.** *Stigmatophora obratzsovi* Daniel, 1951, here designated.

**TL.** Chekiang, “Berge südlich Wenchow” (China, Zhejiang).

**Diagnosis.** Unlike members of subgenus *Stigmatophora* (Figs 7, 8, 19, 25) and *Pseudomiltochrista* (Figs 9–12, 20–22, 26), *Bifurcistigma* subgen. nov. (Figs 1–6, 13–18, 23–24) has the following characters (details for *Stigmatophora* and *Pseudomiltochrista* are in parentheses, respectively): 1) forewing ground color brown, with an orange band (mostly yellow, with a pattern or spots, partially

orange; dark yellow, with rosy veins); 2) ground color of hindwing pale brown to pale blackish brown (mostly dark yellow, partially dark orange; light yellow, with rosy tint); 3) head and patagium orange (mostly yellow, partially red; yellow mixed with red); 4) thorax purple-grey (mostly dark yellow, partially light red and orange; brownish-yellow with red); and 5) abdomen pale brown to pale blackish brown (almost dark yellow, partially light red and orange; light yellow).

Male genitalia: 1) uncus stout (others are thinner, ca 1/2 length of *Bifurcistigma* subgen. nov.; elongated); 2) valva without harpe (harpe small, thorn-like; broad, crest-like); 3) lamella centralis moderately sclerotized (others with no lamella centralis); 4) sacculus with a medial longitudinal sclerotized band at 1/3 of the length (with no medial process, except in *S. flava*; with a setose thorn-like medial process); 5) juxta with two lateral lobes each side (flat, shield-like; with dorsal process); and 6) cornuti in vesica dense, short, and stout, arranged in a U-shaped band (shortest, almost granular; longest, spine-like, and sparsely arranged).

In the female genitalia: 1) posterior margin of 8<sup>th</sup> sternite sclerotized and semicircular (membranous; irregularly dentate); 2) ductus bursae extremely broad, strongly sclerotized, roughened with wrinkles (elongate, thin, and weakly sclerotized, and smooth; thin, only sclerotized on left side); and 3) corpus bursae densely covered with spinules, without appendix bursae (other two with a strong spine-like scobination and narrow appendix bursae).

### ***Stigmatophora (Bifurcistigma) obratzovi* Daniel, 1951**

Figs 1, 2, 13, 14, 23


Common name. 盆带痣苔蛾

*Stigmatophora obratzovi* Daniel, 1951: 297, pl. 1, fig. 3.

**TL.** Chekiang, “Berge südlich Wnchow” (China, Zhejiang).

**Material examined.** CHINA • 3♂♂; Jiangxi, Jinggangshan, Liujiaping; 3–9.VIII.2021; H. L. Han leg.; genit. prep. no. ztt-435-1, ztt-436-1, ztt-437-1; in NEFU • 1♀; Nanling, 1,000–1,400 m elev., 1–6.VI.2006; M. Wang et al. leg.; genit. prep. no. NSMT-4743, in NSMT.

**Redescription. Adults** (Figs 1, 2). Forewing length 10 mm in males, 11.6 mm in female. Head and patagium orange, tegula purple-grey; antenna filiform, brownish-yellow, with a dark brown base; thorax purple-grey; abdomen camel-colored, lighter at 1<sup>st</sup> abdominal segment. Forewing ground color brown, with orange bands along costal margin, inner, and lower margins of cell, and veins  $M_2$  and  $Cu_1$ ; band on  $M_2$  slightly broadens, and on  $Cu_1$ , it widens toward outer margin, connecting fully with costal margin; orange band on inner margin not reaching anal angle. Hindwing ground color brownish-gray. **Male genitalia** (Figs 13, 14). Uncus robust, nearly cylindrical, slightly curved, with a hooked apex. Tegumen weakly sclerotized, thick, ca 1.25 times length of uncus. Valva elongate, narrow, asymmetrical, covered with long hairs, and with a narrowly rounded apex. Costa smooth, slightly raised near base at 1/3 its length. Sacculus weakly sclerotized, broad at base, tapering towards end; ca 7/8 length of valva, middle third rough and band-like, with distal third separated from valva; distal sacculus process curves inward, hook-shaped. Ampulla sclerotized, elongated, with a mound-shaped base. Lamella centralis moderately sclerotized,



**Figures 1–12.** Adults of *Stigmatophora* spp. **1** *S. (Bifurcistigma) obratzsovi*, male, China (Jiangxi) **2** ditto, female, China (Nanling) **3** *S. (B.) bucseki* sp. nov., male, holotype, China (Guizhou) **4** *S. (B.) longa* sp. nov., male, holotype, China (Heilongjiang) **5** ditto, male, paratype, China (Heilongjiang) **6** *S. (B.) bucseki* sp. nov., female, paratype, China (Guizhou) **7** *S. (Stigmatophora) micans*, male, Zavkhan aimak, Mongolia (after Volynkin et al. 2018) **8** ditto, female, East Siberia, Russia (after Volynkin et al. 2018) **9** *S. (Pseudomiltochrista) zolotuhini*, male, India (Assam) (after Volynkin et al. 2018) **10** *S. (P.) dianensis* sp. nov., male, holotype, China (Yunnan) **11** ditto, male, paratype, China (Yunnan) **12** *S. (P.) zolotuhini*, female, holotype, Vietnam (Gia Lai) (after Volynkin et al. 2018). Scale bars: 1 cm.

extending from base of costa to sacculus process. Vinculum strongly sclerotized, wide V-shaped. Juxta with two kidney-shaped lateral lobes on each side, distal half densely covered with small granules, while basal half sclerotized, flat. Aedeagus robust, slightly twisted at middle, tapering towards apex. Carinal region sclerotized. Vesica membranous near base, with a granular field at middle, and a U-shaped band of stout, elongate, triangular cornuti distally.

**Female genitalia.** (Fig. 23). Papillae anales covered with dense setae. Anterior and posterior apophysis almost equal in length. Ostium bursae strongly sclerotized, rough, and trapezoidal. Ductus bursae cylindrical, extremely broad, and strongly sclerotized, with its interior densely covered with small spines. Corpus bursae membranous, ovoid, rough, and densely covered with small spines that are shorter than those on ductus bursae.

**Distribution.** China (Fujian, Zhejiang, Jiangxi, Hunan).



***Stigmatophora (Bifurcistigma) bucseki* sp. nov.**

<https://zoobank.org/2EA1A963-CD22-49B7-85E5-1BB9E1CCDBFA>

Figs 3, 6, 15, 16, 24


Common name. 布痣苔蛾

**Material examined. Holotype:** CHINA • ♂; Guizhou, Zunyi, Suiyang, Huilong; 6–7.VIII.2020; H. L. Han & J. Wu leg.; genit. prep. no. ztt-312-1; in NEFU.


**Paratypes:** 3♂♂; same data as holotype; H. L. Han & J. Wu leg.; genit. prep. no. ztt-311-1, ztt-313-1, ztt-314-1 • 1♀; Guizhou, Qiandongnan, Leishan, Langde; 18–22.VIII.2020; H. L. Han & J. Wu leg.; genit. prep. no. ztt-397-2; in NEFU • 1♀; Yunnan, Mojiang, Hani Autonomous County; 18–19.IX.2008; H. L. Han & J. Wu leg.; genit. prep. no. ztt-375-2, in NEFU.

**Diagnosis.** Externally, the new species is similar to *S. (B.) obraztsovi* (Figs 1, 2, 13, 14, 23), but it can be separated by the following morphological characters (details for *S. (B.) obraztsovi* are in parentheses): in *S. (B.) bucseki*, band on vein  $M_2$  slightly widened toward outer margin and partially connected to costal margin (narrower and fully connected). In the male genitalia, the uncus is long, approximately equal in length to the tegumen (shorter, about 0.75 times); the ampulla is more robust, horn-shaped (slender); and the costa is broader, with the base distinctly enlarged (narrow and smooth). In the female genitalia, the ductus bursae is cylindrical (asymmetrical) and the corpus bursae is approximately circular (elliptical).

**Description. Adults** (Figs 3, 6). Forewing length 9.5 mm in male, 10.7 mm in female. Head and patagium orange; tegula purple-grey; antenna filiform, brown, with dark brown base; thorax purple-grey; abdomen camel-colored, lighter at 1<sup>st</sup> abdominal segment. Forewing ground color brown, fringe orange; costal, inner, and lower margins of cell,  $M_2$  and  $Cu_1$  veins covered with orange band; band on  $M_2$  slightly widening toward outer margin and slightly connected with costal margin. Hindwing ground color brown, terminal line distinct, fringe camel-colored. **Male genitalia** (Figs 15, 16). Uncus robust, moderately sclerotized, nearly sickle, with a spiculate hooked apex. Tegumen weakly sclerotized, thick, same length as uncus. Valva short and wide, with long hairs at apex; costa slightly depressed medially, distinctly widened at base around 1/3 position. Sacculus slightly sclerotized, ca 9/10 length of valva, base wide and tapering towards end, end third separated from valva and hooked; medial third strongly sclerotized, slightly inwardly protruding. Ampulla strongly sclerotized, horn-shaped. Lamella centralis moderately sclerotized, connected with base of ampulla and sacculus process. Vinculum strongly sclerotized, V-shaped. Juxta with two lateral lobes on each side, kidney-shaped, with distal half densely covered small granules. Aedeagus robust and slightly sclerotized. Base of vesica smooth, distal part densely covered in small granules, row of cornuti structures U-shaped; subchamber diverticula with prominent elasma. **Female genitalia** (Fig. 24). Papillae anales slightly sclerotized, covered with setae. Anterior and posterior apophysis slender, almost equal in length. 8<sup>th</sup> sternite sclerotized and semicircular. Ostium bursae slightly sclerotized and rough, scalloped, with a sclerotized, transverse band. Ductus bursae very broad at base and gradually narrowing towards apex, strongly sclerotized, with dense microspines, with conspicuous folds, and a strongly sclerotized, nipple-shaped process on right. Corpus bursae ellipsoidal, rough, and densely covered with small spines.



**Figures 13–22.** Male genitalia of *Stigmatophora* spp. **13** *S. (Bifurcistigma) obrastsovi*, genit. prep. no. ztt-436-1 **14** ditto, genit. prep. no. ztt-435-1 **15** *S. (B.) bucseki* sp. nov., holotype, genit. prep. no. ztt-312-1 **16** ditto, paratype, genit. prep. no. ztt-313-1 **17** *S. (B.) longa* sp. nov., holotype, genit. prep. no. wp-90-1 **18** ditto, paratype, genit. prep. no. wp-89-1 **19** *S. (Stigmatophora) micans* (after Volynkin et al. 2018) **20** *S. (Pseudomiltochrista) zolotuhini* (after Volynkin et al. 2018) **21** *S. (P.) dianensis* sp. nov., holotype, genit. prep. no. ztt-391-1 **22** ditto, paratype, genit. prep. no. ztt-390-1. Scale bars: 1 mm.



Figures 23–26. Female genitalia of *Stigmatophora* spp. **23** *S. (Bifurcistigma) obratzovi*, genit. prep. no. NSMT-4743 **24** *S. (Bifurcistigma) bucseki* sp. nov., paratype, genit. prep. no. ztt-397-2 **25** *S. (Bifurcistigma) micans* (after Volynkin et al. 2018) **26** *S. (Pseudomiltochrista) zolotuhini* (after Volynkin et al. 2018). Scale bar: 1 mm.

**Bionomics.** The type series was collected in early August at an altitude of 958 m in a subtropical mixed forest. The forest floor was densely covered with a variety of shrubs, ferns, and expansive patches of grassland.

**Etymology.** The new species is named after Slovakian entomologist Karol Bucsek for his help with identification.

**Distribution.** China (Guizhou: Zunyi, Qiandongnan; Yunnan: Mojiang) (Fig. 27).

***Stigmatophora (Bifurcistigma) longa* sp. nov.**

<https://zoobank.org/599DDE26-EF3C-43CF-9098-5DCBADE8C0B4>

Figs 4, 5, 17, 18

Common name. 龙痣苔蛾

**Material examined. Holotype:** CHINA • ♂; Heilongjiang, Shangzhi, Changshou, Laoyeling; 5.VIII.2009; P. Wang et.al. leg.; genit. prep. no. wp-90-1, in NEFU.






Figure 27. Collecting sites of three new *Stigmatophora* spp.

**Paratype:** 1 ♂, same data as holotype; genit. prep. no. wp-89-1, in NEFU.

**Diagnosis.** This new species is highly similar to *S. (B.) obratzovi* (Figs 1, 2, 13, 14, 23) and *S. (B.) bucseki* (Figs 3, 6, 15, 16, 24), but can be distinguished by the following morphological characters (details for *S. (B.) obratzovi* and *S. (B.) bucseki* are in parentheses, respectively): in *S. (B.) longa*, band on  $M_2$  not connected with costal margin (completely connected; slightly connected). In the male genitalia, ampulla sclerotized and crutch-shaped (narrow and slender; shorter and horn-shaped).

**Description. Adults** (Figs 4, 5). Forewing length 9.8 mm in male. Antenna filiform. Thorax dark brown; tegula chocolate; abdomen taupe-brown, lighter on last abdominal segment. Forewing ground color brownish-black, with orange bands along costal, inner, and lower margins of cell and veins  $M_2$  and  $Cu_1$ ; bands along  $M_2$  and  $Cu_1$  slightly widening toward outer margin, band along inner margin not reaching anal angle. Hindwing ground color brown, terminal line distinct, fringe brown. **Male genitalia** (Figs 17, 18). Uncus sclerotized, robust, and blunt at apex. Tegumen bell-shaped, ca 1.21 times as long as valva. Valva wide, with cucullus rounded. Costa relatively smooth, mid-region slightly concaved. Sacculus sclerotized, wide at base, tapering towards end, sacculus process hooked, separated from valva. Ampulla strongly sclerotized, elongated, crutch-shaped. Lamella centralis moderately sclerotized, extending from base of ampulla to sacculus process. Vinculum strongly sclerotized, V-shaped. Juxta with two weakly sclerotized lateral lobes, kidney-shaped. Aedeagus robust, slightly bent near vesica. Carinal plate weakly sclerotized. Vesica nearly spherical, thin membranous near base, granulated at middle, and with a row of short, distal cornuti distributed. **Female genitalia.** Unknown.



**Bionomics.** The type series was collected in early August at an altitude of 1318 m in a mixed coniferous and broadleaf forest. The forest floor was covered with dense growth of shrubs and trees.

**Distribution.** China (Heilongjiang: Shangzhi) (Fig. 27).

**Etymology.** The species is named after “loong”, in honor of the current year of the Chinese zodiac, and it also represents the species collection location, Heilongjiang.

### **Subgenus *Pseudomiltochrista* Dubatolov & Bucsek, 2016**

Common name. 拟美痣苔蛾亚属

*Pseudomiltochrista* Dubatolov & Bucsek, 2016: 233.

**TS.** *Pseudomiltochrista zolotuhini* Dubatolov & Bucsek, 2016, by original designation.

**TL.** Central Vietnam, Gia Lai Prov., K' Bang Distr., Dak Roong Comm., vill. [age] Kon Loc, Kon Ka Kinh NP [National Park], 14°42.602'N, 108°39.062'E, 1050 m.

### ***Stigmatophora (Pseudomiltochrista) dianensis* sp. nov.**

<https://zoobank.org/1BF307C4-3F94-4FE1-A43D-57FF285D439B>

Figs 10, 11, 21, 22

Common name. 云南痣苔蛾

**Material examined. Holotype:** CHINA • ♂; Yunnan, Pu' er, Simao, Manxieba; 3.VIII.2018; H. L. Han et. al. leg.; genit. prep. no. ztt-391-1; in NEFU.

**Paratypes:** 1♂, same data as holotype; genit. prep. no. ztt-390-1 • 1♂, Yunnan, Xishuangbanna, Jinghong, Mengyang, Wild Elephant Valley; 4–5.VIII.2018; H. L. Han et. al. leg., genit. prep. no. ztt-321-1, in NEFU.

**Diagnosis.** The new species is similar to *S. (P.) zolotuhini* (Figs 9, 12, 20, 26), but it can be separated by the following morphological characters (details for *S. (P.) zolotuhini* are in parentheses): in *S. (P.) dianensis*, forewing 8 mm long, ovoid, costal and outer margins more arched (10 mm, slightly narrow, smooth). In the male genitalia, uncus short and spindle-shaped (long and slender); apex of apical sacculus process hook-shaped (bluntly finger-shaped); harpe elongate conical, moderately sclerotized, covered with short setae (crest-like, weakly sclerotized); juxta with a flat triangular dorsal process (with a clavate dorsal process); and vesica elongate (fist-shaped, with multiple diverticula).

**Description. Adults** (Figs 10, 11). Forewing length 8.7 mm in males. Head, patagium, tegula, and thorax dark yellow, with rose veins; antenna filiform, yellow; thorax yellow-brown; abdomen pale yellow. Forewing ovoid, costal and outer margins more arched, with yellow ground color and rose veins, with irregular submarginal line and medial curved spots. Rose-colored pattern completely reaches outer margin. Hindwing pale yellow, scattered with rosy tint. **Male genitalia** (Figs 21, 22). Uncus robust, rhombic, with bristle. Tegumen membranous, triangular. Valva narrow and symmetrical; costa membranous and smooth; cucullus rather sharp, covered with rough hairs.

Sacculus sclerotized, wide at base, gradually narrowing toward apex, with a strongly sclerotized medial process, sharkfin-shaped, directed downwards. Harpe elongate conical, moderately sclerotized, covered with short setae. Vinculum moderately sclerotized, thick, U-shaped. Juxta flat, with a triangular dorsal process. Aedeagus slender tubular, with swollen caecum. Vesica elongate; central region covered with small graniculi and a cluster of cornuti. **Female genitalia.** Unknown.

**Bionomics.** Wild Elephant Valley in Xishuangbanna is within the Mengyangzi Protection Area of the Xishuangbanna National Nature Reserve. It is north of Jinghong City and flanked by the Lancang River to the west and the Xiaohei River to the east. The terrain comprises low hills and mountain basins. The valley has abundant natural resources, harboring tropical rainforests, South Asian tropical evergreen broadleaf forests, and a diverse array of rare flora and fauna. Specimens were collected in a dense broadleaf forest covered with ferns.

**Distribution.** China (Yunnan: Pu' er, Xishuangbanna) (Fig. 27).

**Etymology.** The species name "dian" is derived from the Chinese character "滇" (Diān), which is the abbreviation for Yunnan Province.

### **Checklist of species in the genus *Stigmatophora* Staudinger, 1881, with synonyms, type locality, and distribution**

Subgenus *Stigmatophora* Staudinger, 1881

The *palmata* species-group

1. *S. (S.) cernyi* Volynkin, 2020.  
TL: Thailand, Mount Doi Phahompok.  
Distribution: China (Yunnan); Thailand.
2. *S. (S.) chekiangensis* Daniel, 1951.  
TL: China, Chekiang (Zhejiang: W. Tien-Mu-Shan).  
Distribution: China (Gansu, Hunan, Guangxi, Sichuan).
3. *S. (S.) danieli* Volynkin, Huang & Dubatolov, 2021.  
TL: China, Jiangxi: Wuyi Shan.  
Distribution: China (Zhejiang, Jiangxi, Hubei, Guangdong).
4. *S. (S.) flavogrisea* Leech, 1899.  
TL: China, Sichuan, N of Kangding.  
= *Hypeugoa flavogrisea* Leech, 1899  
TL: China, Sichuan, N of Kangding.  
Distribution: China (Gansu, Sichuan).
5. *S. (S.) hainanensis* Fang, 1991.  
TL: China, Hainan.  
Distribution: China (Guangdong, Hainan, Jiangxi, Guangxi).

6. *S. (S.) inanis* Seitz, 1913.  
TL: Pakistan, Punjab "Murree".  
= *S. palliduspalmata* N. Singh, Kirti & Joshi in Kirti and Singh 2016.  
TL: India "Jammu & Kashmir, Patnitop".  
Distribution: Pakistan; India; Nepal.
7. *S. (S.) karenkonis* Matsumura, 1930.  
TL: Formosa, Karenko (China: Taiwan).  
= *Miltochrista karenkonis* Matsumura, 1930.  
TL: Formosa, Karenko (China: Taiwan).  
Distribution: China (Taiwan); Japan.
8. *S. (S.) orientalis* Daniel, 1951.  
TL: China, Chekiang (Zhejiang).  
= *Hypeugoa flavogrisea orientalis* Daniel, 1951.  
TL: China, Chekiang (Zhejiang: West-Tien-Mu-Shan).  
= *Hypeugoa flavogrisea flavogrisea* Daniel, 1951.  
TL: China, Chekiang (Zhejiang).  
Distribution: China (Shaanxi, Hebei, Jiangsu, Zhejiang, Jiangxi, Sichuan); Japan; Korea.
9. *S. (S.) palmata* Moore, 1878.  
TL: Northern India, Himachal Pradesh.  
= *Lyclene palmata* Moore, 1878.  
TL: northern India, Himachal Pradesh "Dharmasala N.W. Himalaya".  
Distribution: China (Yunnan, Xizang); India.
10. *S. (S.) ranruna* Matsumura, 1927.  
TL: China, Taiwan.  
Distribution: China (Taiwan).
11. *S. (S.) rhodophila* Walker, 1864.  
TL: China, Shanghai.  
= *Barsine rhodophila* Walker, 1865.  
TL: China, Shanghai.  
Distribution: China (Shanghai).  
*S. (S.) rhodophila rhodophila* Butler, 1879.  
TL: China, Shanghai.  
= *Miltochrista torrens* Butler, 1879.  
TL: Japan, Yokohama.  
Distribution: China (Heilongjiang, Jilin, Beijing, Shaanxi, Shanxi, Shandong, Jiangsu, Shanghai, Zhejiang, Henan, Hubei, Hunan, Guangxi, Sichuan); Japan; Korea; Russia.  
*S. (S.) rhodophila zeyana* Dubatolov, 2013.  
TL: Russia, Amurskaya Oblast, Zeyskii Nature Reserve (the status is unclear).  
Distribution: Russia.

12. *S. (S.) roseivena* (Hampson, 1894).  
TL: Burma, Momeit.  
Distribution: China (Jiangxi, Fujian, Hainan); Myanmar; Thailand; Vietnam.

Other species with unclear status

13. *S. (S.) acerba* Leech, 1899.  
TL: China, Moupin (Sichuan).  
= *Miltochrista acerba* Leech, 1899.  
TL: China, Moupin (Sichuan).  
Distribution: China (Gansu, Hunan, Guangxi, Sichuan).
14. *S. (S.) confusa* Daniel, 1951.  
TL: China, Yunnan, Li-Kiang.  
Distribution: China (Yunnan).
15. *S. (S.) conjuncta* Fang, 1991.  
TL: China, Gansu.  
Distribution: China (Gansu, Beijing, Shanxi, Hubei, Guangxi).
16. *S. (S.) disticha* Meyrick, 1894.  
TL: Burma, Shán States, Koni.  
= *Ammatho disticha* Meyrick, 1894.  
TL: Burma, Shán States, Koni.  
= *Asura disticha* Hampson, 1900.  
TL: Burma, Shán States, Koni.  
Distribution: Burma; Bhutan.
17. *S. (S.) flava* Bremer & Grey, 1852.  
TL: China, Shanghai.  
= *Setina sinensis* Walker, 1854.  
TL: China, Shanghai.  
= *Setina ochracea* Kindermann, 1855.  
TL: China, Altai.  
Distribution: China (Heilongjiang, Jilin, Liaoning, Xinjiang, Hebei, Shanxi, Shandong, Jiangsu, Zhejiang, Fujian, Jiangxi, Hubei, Hunan, Guangdong, Guangxi, Sichuan, Guizhou, Yunnan); Kazakhstan; Russia; Japan; North Korea.
18. *S. (S.) grisea* Hering, 1936.  
TL: China, Gansu.  
Distribution: China (Gansu).
19. *S. (S.) leacrita* Swinhoe, 1894.  
TL: Japan, Yokohama.  
= *Setina leacrita* Swinhoe, 1894.  
TL: Japan, Yokohama.  
Distribution: China (Taiwan); Russia; Japan; Korea.



20. *S. (S.) likiangensis* Daniel, 1951.  
TL: China, N. Yunnan, Li-Kiang.  
Distribution: China (Yunnan).
21. *S. (S.) micans* Bremer & Grey, 1852.  
TL: China, Peking (Beijing).  
= *Setina micans* Bremer & Grey, 1852.  
TL: China, Peking (Beijing).  
= *Setina albosericea* Moore, 1877.  
TL: China, Shanghai.  
Distribution: China (Heilongjiang, Jilin, Liaoning, Shaanxi, Inner Mongolia, Beijing, Hebei, Shanxi, Shandong, Jiangsu, Shanghai, Anhui, Henan, Hubei, Sichuan); East Kazakhstan, South Siberia, Amurland, Russia, Mongolia, North Korea.
22. *S. (S.) rubivena* Fang, 1991.  
TL: China, Yunnan.  
Distribution: China (Yunnan).
23. *S. (S.) strigivenata* Hampson, 1894.  
TL: Burma, E. Pegu.  
= *Eugoa strigivenata* Hampson, 1894.  
TL: Burma, E. Pegu.  
Distribution: Burma, Bhutan.
24. *S. (S.) tridens* Wileman, 1910.  
TL: China, Kanshirei.  
= *Miltochrista tridens* Wileman, 1910.  
TL: China, Kanshirei.  
Distribution: China (Taiwan).  
Subgenus *Pseudomiltochrista* Dubatolov & Bucsek, 2016.
25. *S. (P.) dianensis* sp. nov.  
TL: China, Yunnan.  
Distribution: China (Yunnan).
26. *S. (P.) zolotuhini* Dubatolov & Bucsek, 2016.  
TL: central Vietnam.  
Distribution: India; Myanmar; Thailand; Laos; Vietnam.  
Subgenus *Bifurcistigma* subgen. nov.
27. *S. (B.) bucseki* sp. nov.  
TL: China, Guizhou.  
Distribution: China (Guizhou, Yunnan).
28. *S. (B.) obratzovi* Daniel, 1951.  
TL: China, Chekiang (Zhejiang).  
Distribution: China (Zhejiang, Fujian, Jiangxi, Hunan).

29. *S. (B.) longa* sp. nov.

TL: China, Heilongjiang.

Distribution: China (Heilongjiang).

## Acknowledgements

We express our sincere gratitude to Dr Károl Bucsek for his invaluable assistance and guidance throughout this research. We also sincerely thank Mamoru Owada (NSMT, Japan) for providing the female adult and genitalia slide of *S. (B.) obratzovi*. Additionally, we appreciate the support of lab members Mr Peng Wang, Dr Jun Wu, and Dr Ming-Rui Li in collection specimens. We greatly appreciate Dr Christopher Glasby's help with English language editing. We are also deeply grateful to reviewer Siyao Huang and editors Reza Zahiri and Robert Forsyt for their time and efforts in enhancing the quality and clarity of the manuscript.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Funding

This study is supported by the project of National Nature Science Foundation of China (No.31572294), and Northeast Asia Biodiversity Research Center (2572022DS09, NABRI202303).

### Author contributions

Investigation: LZ, TTZ. Writing - original draft: LZ. Writing - Review and Editing: HHL, TTZ.

### Author ORCIDs

Lu Zhang  <https://orcid.org/0009-0007-5269-3426>

Tingting Zhao  <https://orcid.org/0000-0003-4139-6120>

Huilin Han  <https://orcid.org/0000-0002-2045-6182>

### Data availability

All of the data that support the findings of this study are available in the main text.

## Reference

- Blanford WT (1898) The Fauna of British India, Including Ceylon and Burma. Moths. Vol. 2. Taylor & Francis, London, 609 pp.
- Bremer O, Grey W (1852) Diagnoses de lépidoptères nouveaux, trouvés par MM. Tatarinoff et Gaschkewitsch aux environs de Pékin. Études entomologiques, Imprimerie de la Société de Littérature Finnoise, Helsingfors 1: 58–67.
- Daniel F (1951) Beiträge zur Kenntnis der Arctiidae Ostasiens unter besonderer Berücksichtigung der Ausbeuten von Dr. h.c. H. Höne aus diesem Gebiet (Lep.-Het.). III. Teil: Lithosiinae. Bonner zoologische Beiträge 2(3–4): 291–327.

- Dubatolov VV, Bucsek K (2016) New lichen-moth taxa (Lepidoptera, Arctiidae, Lithosiinae) from Vietnam. *Euroasian Entomological Journal* 15(3): 228–238.
- Fang C (1991) A study on the Chinese *Stigmatophora* Staudinger (Lepidoptera: Arctiidae: Lithosiinae). *Sinozoologia* 8: 377–382.
- Kononenko V, Han H (2007) Atlas Genitalia of Noctuidae in Korea (Lepidoptera). In: Park KT (Ed.) *Insects of Korea* (Series 11). Junhaeng-Sa, Seoul, 464 pp.
- Leech JH (1898) Lepidoptera Heterocera from northern China, Japan, and Corea. *Transactions of the Royal Entomological Society of London* 1898: 261–379. <https://doi.org/10.5962/bhl.title.22195>
- Matsumura S (1927) New species and subspecies of moths from the Japanese Empire. *Journal of the College of Agriculture, Hokkaido Imperial University, Sapporo, Japan* 19: 1–5.
- Matsumura S (1930) New species and forms of Arctiidae from Japan. *Insecta Matsu-murana* 5: 31–40.
- Meyrick E (1894) On a collection of Lepidoptera from Upper Burma. *Transactions of the Royal Entomological Society of London* 1894: 16–19.
- Moore P (1878) A revision of certain genera of European and Asiatic Lithosiidae, with characters of new genera and species. *Proceedings of the Zoological Society of London* 1878: 3–37. <https://doi.org/10.1111/j.1469-7998.1878.tb07927.x>
- Seitz A (1911) Subfamilie: Lithosiinae, Flechtenbärchen (partim: 1. Darantoides – 28. Scaphidriotis). In: Seitz A (Ed.) *Die Gross-Schmetterlinge der Erde*. Vol. 10. Spinner und Schwärmer des indo-australischen Gebiets. Alfred Kernen, Stuttgart, 118–134.
- Staudinger O (1881) Beitrag zur Lepidopteren-Fauna Central-Asiens. *Stettiner Entomologische Zeitung* 42: 393–424.
- Swinhoe C (1894) New species of eastern Lepidoptera. *Journal of Natural History* 14: 429–443. <https://doi.org/10.1080/00222939408677830>
- Volynkin AV (2020) On the taxonomy of the genera *Stigmatophora* Staudinger, 1881 and *Miltochrista* Hübner, [1819], with description of a new species from North Thailand and China (Lepidoptera, Erebidae, Arctiinae, Lithosiini). *Zootaxa* 4789: 508–522. <https://doi.org/10.11646/zootaxa.4789.2.8>
- Volynkin AV (2024) On the terminology of the genitalia structures of lichen moths (Lepidoptera: Erebidae: Arctiinae: Lithosiini) with some references to Noctuidae. *Ecologica Montenegrina* 73: 176–207. <https://doi.org/10.37828/em.2024.73.18>
- Volynkin AV, Dubatolov VV, Kishida Y (2018) On the taxonomy of the genera *Planovalvata* Dubatolov & Kishida, *Stigmatophora* Staudinger and *Pseudomiltochrista* Dubatolov & Bucsek, with description of one new species (Lepidoptera, Erebidae, Arctiinae, Lithosiini). *Zootaxa* 4394: 128–136. <https://doi.org/10.11646/zootaxa.4394.1.7>
- Volynkin AV, Huang S-Y, Dubatolov VV (2021) A review of the *Stigmatophora palmata* (Moore, 1878) species-group (Lepidoptera: Erebidae: Arctiinae) with description of a new species from China. *Zootaxa* 4999: 243–257. <https://doi.org/10.11646/zootaxa.4999.3.4>
- Walker F, Gray JE (1855) List of the specimens of Lepidopterous insects in the collection of the British Museum. Vol. 3. Trustees of the British Museum, London, 583–775.
- Wileman AE (1911) New Lepidoptera–Heterocera from Formosa. *The Entomologist* 44: 148–152.