

Pharmacological properties of selenium and its preparations: from antioxidant to neuroprotector

Igor F. Belenichev¹, Nadia O. Gorchakov², Iryna B. Samura¹,
Natalia V. Savchenko², Nina V. Bukhtiyarova¹, Olena O. Popazova¹

¹ Zaporizhzhia State Medical University, 26 Mayakovsky Ave., Zaporizhzhia 69000, Ukraine

² Bogomolets National Medical University, 13 Taras Shevchenko Blvd., Kyiv 01601, Ukraine

Corresponding author: Olena O. Popazova (popazova.ea@gmail.com)

Academic editor: Oleg Gudyrev ♦ Received 15 August 2021 ♦ Accepted 24 October 2021 ♦ Published 3 December 2021

Citation: Belenichev IF, Gorchakova NO, Samura IB, Savchenko NV, Bukhtiyarova NV, Popazova OO (2021) Pharmacological properties of selenium and its preparations: from antioxidant to neuroprotector. *Research Results in Pharmacology* 7(4): 29–40. <https://doi.org/10.3897/rrpharmacology.7.73051>

Abstract

Selenium is an essential component of more than two dozen enzymes and other selenoproteins that play critical roles in reproduction, DNA synthesis, thyroid hormone metabolism, and protection from oxidative damage and infection. Selenium has a protective action against some forms of cancer and cardiovascular diseases, modulates levels of inflammatory mediators, promotes to maintain bone homeostasis and protects against bone loss. Selenium significance as a cardioprotective agent may be associated not only with its antioxidant properties, but also with its ability to prevent inflammation, autophagy, as well as the intrinsic and extrinsic apoptosis pathways. Signaling pathways, such as p-AMPK, PARP, Nrf2, STAT, are involved in the protective effects of selenium. Selenium protects against cardiovascular damage by increasing the survival rate of cardiomyocytes, including a mitochondria-dependent pathway and autophagy through peroxisome proliferator-activated receptors. Research demonstrating neuroprotective and cardioprotective effects of selenium preparations – selenoline, selenocysteine and selenomethionine – is growing at a rapid rate. It has been established that these compounds are able to normalize the levels of heat shock proteins (HSP70), which limit the cytotoxic effects of free radicals, produce energotropic action, prevent a decrease in the membrane mitochondria charge, and the opening of the mitochondrial pore. Also regulate the expression of transmembrane factors NF- κ B, c-fos, which is associated with their main biological function of chaperone proteins, providing protection of neurons from damage. In this review, we want to emphasize pharmacological role of Selenium and its derivatives on human health is very complex and has yet to be fully understood.

Keywords

selenium, heat shock proteins, cardioprotective, neuroprotective, antioxidant action.

Introduction

Selenium, the 34th element of the periodic table placed in the subgroup VIA (also called chalcogen group), is the electronic and chemical analogue of sulfur, a multifunc-

tional (e.g. antioxidant, antineoplastic, antidiabetic, antimicrobial) nutrient element. Selenium is one of the most important trace elements in the human body, nutritionally essential as a component of more than two dozen enzymes and other selenoproteins that play critical roles in

reproduction, DNA synthesis, thyroid hormone metabolism, and protection from oxidative damage and infection (Rakov and Muz 2019). The human body absorbs more than 90% of selenium from plant and animal food, and about 10% – with drinking water (Krelsek et al. 2021). Dietary selenium deficiency can lead to the development of severe and dangerous diseases, such as cardiomyopathy, arrhythmias, and increased susceptibility to infection (Yovanovich and Ermakov 2020). Conversely, a selenium-excessive diet can cause acute poisoning and chronic symptoms. Classical symptoms of dietary selenium deficiency in farm and laboratory animals were characterized in the 1950s to 1960s, and selenium supranutrition was found to have anti-cancer properties without affecting healthy cells and cancer mortality in the 1990s (Alhasan et al 2019). Its anti-cancer properties are related to antioxidant and prooxidant activity. Selenium-loaded bone frameworks have potential for cancer treatment and regeneration. Many studies report that selenium has a protective effect against some forms of cancer, cardiovascular disease mortality, regulates the inflammatory mediators in asthma, maintains bone homeostasis and protects against bone loss. Antioxidant properties of selenium preparations have been widely studied and discussed in a number of recent review articles (Rahmanto and Davies 2012; Steinbrenner and Sils 2013; Solovyev 2015; Gandin et al. 2018; Talbi et al. 2019; Chen et al 2020; Kuria et al. 2020; Wang et al. 2020). Selenium is widely used in industry for the manufacture of several products, such as semiconductors, photovoltaic cells (due to its ability to convert light energy into electrical energy), rectifiers, corrosion-resistant alloys, pigments (red and orange color) for paints, ceramics, glass-making process and pharmaceutical substances for the creation of mono-drugs and complex drugs, which have a wide spectrum of action (Rahmanto and Davies 2012).

Selenium was discovered in 1817 by the Swedish chemist J.J. Berzelius, who named the new element from the Greek word ‘Selene’, meaning Goddess of the Moon (Duntas and Benvenega 2015). In terms of chemical properties, it is similar to tellurium, discovered in 1782 by M.H. Klaproth (Rakov and Muz 2019). The history of the study of the biological role of selenium includes 3 periods. Selenium was considered highly toxic to the body until 1857 (Golubkina et al. 2017). The reason for this statement was the massive poisoning of livestock in some areas of the Great Plains in the United States. The next stage, characterized by the emergence of interest in selenium, was the discovery by K. Schwartz and S. Foltz in 1857 that established the prevention of developing liver cirrhosis in rats and exudative diathesis in chickens by sodium selenite administered in very low doses. This was the beginning of the use of selenium for the treatment of white muscle disease and other diseases in animals. The third stage started with the discovery of the selenium-containing enzyme, glutathione peroxidase, regulating the antioxidant state of the body and also contributing to the identification of the areas where the lack of selenium

in the environment and food products were established (Golubkina et al. 2017). However, it was not until 1873 when all the aspects of the biochemical mechanisms of action of selenium on organs and tissues were disclosed. In 1873, J. Rotrak with co-authors and a group of German scientists led by L. Flos (Yovanovich and Ermakov 2020) demonstrated the effect of selenium on the metabolic parameters of living organisms. Selenium supports the function of several systems, including the endocrine, immune, and cardiovascular systems. The thyroid is the organ with the highest concentration of selenium per weight of organ tissue. About 25 proteins were discovered to contain a selenocysteine in place of the more common sulfurated amino acid. In these proteins, selenocysteine, present in the active site of the enzyme, is essential for its catalytic activity. Selenium acts in the catalytic center playing a principal role in several major metabolic pathways, such as thyroid hormone metabolism, immune functions, and antioxidant defense systems.

Nowadays, much evidence demonstrated the role of glutathione peroxidase, and that this enzyme, like catalase, with its antioxidant properties, protects erythrocyte membranes from oxidation by destroying hydrogen peroxide. This work (Wichman et al. 2016) became a trigger for the discovery of other selenium-containing enzymes, such as 3 types of oxidoreductases – iodothyronine deiodinases (D1, D2, D3), which catalyze the activation and inactivation of thyroxine with the formation of active triiodothyronine and reverse triiodothyronine in deiodination reactions.

Deiodinases have tissue and organ specificity, which is determined by their different localization. D1 is predominantly expressed in the liver, kidneys, thyroid gland and pituitary glands; D2 – in the gastrointestinal tract, heart, central nervous system, pituitary gland, skeletal muscle, brown adipose tissue, and placenta; D3 – in the in pregnant uterus and placenta, in the liver, brain, and skin of the embryo.

The important role of selenium and its derivatives

Selenium and its derivatives have pronounced antioxidant properties, even if they are a part of complex compounds. Feeding healthy rats with chlorella algae, which contains a significant amount of selenium, has lowered the levels of lipid peroxidation indicators in the organs and plasma of animals, such as malondialdehyde and diene conjugates, while increasing the levels of indicators of antioxidant protection, such as the activity of superoxide dismutase, succinate dehydrogenase, catalase, and cytochrome-C-oxidase (Krelsek et al. 2021).

Selenoproteins, due to their enzymatic role, have antioxidant properties that are more marked inside cells. It is selenoproteins that play a decisive role in maintaining the levels of selenium in the body and preventing organ

dysfunctions associated with changes in homeostasis. **Selenocysteine** and **selenomethionine** are key forms of selenoproteins and are found in various tissues of the body. **Selenocysteine** is found in animal tissues, while **selenomethionine** is obtained from yeast, algae, and some plants (Kreliszek et al. 2021).

Selenoprotein P, a selenium-rich plasma protein, is a glycoprotein that plays an important role in selenium metabolism. Decreased **selenoprotein P** expression is associated with dietary selenium deficiency. The fall in **selenoprotein P** concentration in plasma is associated with dietary selenium deficiency, as a result of which oxidative stress may develop. Selenium administration first leads to an increase in the levels of **selenoprotein P**, and then of other proteins. **Selenoprotein P** is mainly produced in hepatocytes, although it is found in plasma and other tissues (Steinbrenner and Sils 2013). **Selenoprotein P** consists of two domains: the larger N-terminal domain responsible for maintaining the redox potential in the cell, and the smaller C-terminal domain, which provides selenium transport.

Selenoprotein P is expressed in testes, brain tissues, and liver. It is considered a marker of selenium in the body. **Selenoprotein P** maintains homeostasis, protects lipoproteins from oxidation by removing the peroxynitrite molecule, which is formed due to oxidative stress inside the cells as a result of the reaction between superoxide ions and nitric oxide molecules at the sites of inflammation (Di Giuseppe et al. 2019). **Selenoprotein P** enters the circulatory system and is a source of selenium for peripheral tissues (Talbi et al. 2019). With a sufficient intake of selenium with food, its content increases more significantly in the nervous system and testes, while the levels of selenium increase insignificantly in the kidneys and other organs. It should be noted that recent advances in the selenium field have not only contributed to our better understanding of how selenium functions at the molecular level, but also provided complicated details on general protein translation mechanisms and evolution of the genetic code.

Selenoprotein W, a small size of 9 kDa, one of the most widespread selenoproteins found in the cytosol, is detected in high concentrations in muscles and the brain. It is essential for muscle metabolism, and skeletal muscle calcification. Selenium supplements are prescribed prophylactically in cattle and sheep for enzootic muscular dystrophy, commonly known as white muscle disease (Yildirim et al. 2019). Besides, the level of selenoprotein W deficiency may be correlated with the occurrence of stress, since the severity of stress is strongly influenced by the bioavailability of selenium.

All thioredoxin D1, D2, D3 enzymes act as intracellular antioxidant molecules, regulating redox reactions; therefore they are the main factors of apoptosis and DNA synthesis. It is believed that thioredoxin acts as a key enzyme in selenium metabolism, providing the synthesis of selenoproteins (Weekley and Harris 2013). Thioredoxin activates the p53 tumor suppressor gene, so D1 is strongly

expressed in all carcinoma cells. In addition, inhibition of thioredoxin D1 enzyme in cancer cell lines contributes to a decrease in cell proliferation and cancer progression by administering knockdown D1 cells to mice. D1 may have two opposite functions in the development and progression of cancer. On the one hand, D1 can help prevent cancer by maintaining the redox balance of cells and reducing the frequency of mutations that cause tumor development. On the other hand, D1 is also important for tumor progression because cancer cells are very susceptible to oxidative damage (Kieliszek et al. 2019).

Glutathione peroxidase performs several functions in the body, regulating detoxification processes and redox homeostasis (Jiao et al. 2017). Glutathione peroxidase is expressed in the erythrocytes, plasma, extracellular fluid, kidneys, lungs, liver, thyroid gland, heart, testes, and placenta (Liu et al. 2021). Glutathione peroxidase is detected in mitochondria, cytosol, and nucleus (Alhasan et al. 2019). The enzyme is essential for the maturation of sperm motility and effectively improves male fertility (Alhasan et al. 2019).

The number of reports that trace the relationship between pathogenesis of some diseases and prolonged dietary selenium deficiency or selenium excess is growing annually (Hadrup and Ravn-Haren 2020), and it is proposed to use new selenium preparations and other selenium-containing sources, such as algae (Viniarska et al. 2017).

Organoselenium compounds behaving very similarly to their organosulfur analogs have been synthesized in recent years; however, selenium single bondcarbon bonds are typically weaker. The selenium functionality to see broad application in organic synthesis is the selenide, analogous to the sulfide. The attention was paid again to ebselen, and then diselenides appeared, which are united by similar antioxidant and anti-inflammatory properties. It is assumed that these properties are realized due to the activation of transcription factors that regulate the expression of antioxidant genes (Nogueira et al. 2021).

Thiol-containing compounds can be lithium mimetics and used as protease inhibitors in SAPS-COV-2 (Heller et al. 2020). Selenium has been compared to vitamin E as a protector of cell membranes from oxidation. Vitamin E directly absorbs phospholipid hydroperoxide radicals, while selenium in hydroxy peroxidase reduces the levels of phospholipid peroxides, which can generate reactive peroxide radicals (Chen et al 2020).

Selenium is widespread not only in the biosphere, but also in the atmosphere. Hydrogen selenide, dimethyl selenide, and selenium oxide are determined in the atmosphere (Hossain et al 2020). Selenium can be found in plants in the inorganic form, such as selenites, selenates, elemental selenium and others. In plants, these forms are converted into organic compounds, mainly selenocysteine and selenomethionine. In human and plant biological systems, selenomethionine replaces methionine. Among plants that accumulate selenium are lucerne, cruciferous species, such as cabbage, broccoli and cauliflower,

as well as yeasts. Most selenium in animal tissue in the form of selenomethionine is primarily stored in the skeletal muscle that stores anywhere from 28% to 46% of the body's selenium. Seafood and organ meats are noted as the sources that contain the highest levels of selenium, while other good products include muscle meats, cereals, and dairy products (Wichman et al. 2016).

A significant proportion of selenium is contained in the thyroid gland in the form of deiodinases; therefore, hypothyroidism and goiter are common in areas with soil low in selenium (Skowronska-Joozwtak 2015). In autoimmune thyroiditis, selenium reduces the titer of antibodies to thyroid peroxidase and thyroglobulin (Wichman et al. 2016).

Selenium also improves the visual function in patients with endocrine ophthalmopathy, although glucocorticoids have been used to manage severe ophthalmopathy (Bartalena et al. 2016). The anti-inflammatory effect of selenium was first discovered in diabetic rats (Tong et al. 2020). It was shown that selenium can influence inflammatory reactions inducing inhibition of NF- κ B, which promotes the production of tumor necrosis factor and interleukins (Lv et al. 2020).

Selenium influences the process of gene transcription responsible for the synthesis of immunoglobulins, which play an important role in the regulation of the activity of the immune system. Selenium levels also determine macrophage differentiation.

More pro-inflammatory macrophages M1 are produced under conditions of selenium deficiency, whereas anti-inflammatory macrophages M2 are mainly formed under conditions of selenium excess.

This model of selenium action makes it possible to induce a proinflammatory cellular immune response against

viral and bacterial pathogens using Th1 lymphocytes (Tinggi 2014). The release of inflammatory mediators in the body can trigger the occurrence of various painful conditions, whereas a decrease in selenoprotein activity may be associated with a disruption in the functioning of the antiradical system of the body.

Levels of proteins containing selenocysteine are regulated by markers of inflammation and extracellular sugar levels in liver cells. Selenium-containing proteins have antitumor action in systemic macrophages and brain astrocytes, reducing stress. The protective and antitumor effect of selenium is associated not only with its antioxidant properties, but also with the effect on the immune system, since lipid peroxidation activation reduces the potential of immune reactivity.

Selenium may play a role in the regulation of the redox states of proteins, such as NF- κ B transcription factors. Transcription factors regulated by redox mechanisms can have two states, active and inactive, which differ in the oxidation state of the cysteine residue. This is due to the presence or absence of an intramolecular disulfide bond between two cysteine residues. Disulfide formation by oxidation is reversed by cellular systems of disulfide-reducing enzymes, such as glutaredoxin and thioredoxin. Se-derivatives react with protein thiols or disulfides and form more reactive intermediates. In this mechanism, Se acts as a redox catalyst that binds the redox equilibrium of the cell. Se reacts with the cysteine of the non-catalytic thioredoxin site to form the selenenyl sulfide derivative. Modulation of the transcription factor NF- κ B by Se can be related to gene expression and regulate the synthesis of enzymes (antioxidants, participants in energy metabolism, glutathione synthesis). An important link in the neuroprotective action of selenium derivatives is the ac-




Figure 1. Hypothetical mechanism of action of selenium derivatives. **Note:** TRX – thioredoxin; TRXR – thioredoxin reductase; MGH – malate dehydrogenase; SOD – superoxide dismutase; GS – glutamine synthetase; GPX – glutathione peroxidase; NF κ B – nuclear transcription factor; HSP70 – heat shock protein 70 kDa; HIF – hypoxia-induced factor.

tivation of GSH-dependent mechanisms of endogenous neuroprotection, realized by increasing the expression of HSP70 (Fig. 1).

Selenium is necessary for the functioning of the immune system, both the humoral innate and cellular adaptive immune subtypes: it stimulates the function of natural killer cells, increases the production of interleukin-1, interleukin-2, suppresses hypersensitivity of the immediate and delayed types, modulates the phagocytic function of polymorphonuclear leukocytes, antibody response, has antiapoptogenic and radioprotective effects (Saburov et al. 2016).

Selenium preparations exhibit the protective effect on brain tissue, liver tissue, the cardiovascular, endocrine, and reproductive systems. In brain tissue, selenium not only exhibits antioxidant properties, but also takes part in signal transmission in neurons, which helps to maintain cognitive function and prevents the development of Alzheimer's disease (Pillai et al. 2014; Solovyev 2015; Solovyev et al. 2018). Alzheimer's disease is the most common type of dementia with subsequent development of progressive loss of memory, self-expression, face and object recognition, as well as sudden or dramatic changes in behavior.

The levels of selenium-containing compounds affects the levels of mediators, the metabolic activity of neurotransmitters and their structure. That is, impairment of neurotransmitters plays a role in the pathogenesis of the disease on the one hand, and impairment of prooxidant-antioxidant homeostasis, on the other (Loef et al. 2011). The hereditary factors of Alzheimer's disease include the presence of risk factors, the interaction between selenium, apolipoprotein E and presenilin 2.

At the same time, no correlation has been found between the activity of glutathione peroxidase in the cerebrospinal fluid and the occurrence of cerebrovascular diseases. Changes in calcium metabolism have also been implicated in the pathogenesis of Alzheimer's disease. A decrease in selenoproteins levels can lead to a rapid increase in cytosolic free calcium concentration. The data obtained open up the prospect of using selenium preparations in treating Alzheimer's disease (Reddy et al. 2017; Reeves et al. 2019; Tamtaji et al. 2019; De Silva Leme 2020).

Selenium levels in the body

Insufficient content of selenium, zinc, magnesium, and disturbance of the ratio between them leads to depression (Wang et al. 2018). This is primarily due to the fact that selenium is involved in the antioxidant defense of brain tissue and the nervous system. The daily dose of selenium is 55 mg/kg; it is possible that the serum selenium content is 70–90 mg/L. Such a daily intake of selenium is advertised for the treatment of depression, given that selenium deficiency is noted in pregnancy and kidney diseases when it is used in a milk dose.

The recommended daily intake of selenium is 55 mcg / day, is 55 mg/kg, the content of selenium in the blood serum may be 70–90 mg/L. Such a daily intake of selenium is recommended for the management of depression, taking into account that selenium deficiency is noted in pregnancy and kidney disease when it should be administered in a double dose (Bleys et al 2009). Selenium deficiency was established during the use of neuromodulators and a decrease in the levels of brain-derived neurotrophic factor (Iglesias et al. 2013). Beyond mood and general well-being, the role of selenium on mental health is complex and has yet to be fully understood (Conner et al. 2014), and research linking the two is growing at a rapid rate. In recent years, evidence has shown that a drop in blood selenium levels correlates with the development of mental health conditions, including depression and anxiety disorders (Myniec et al. 2015).

The administration of selenium compound m-trifluoromethyl-diphenyl-diselenide to rats relieved depressive symptoms (Bruning et al 2011), and vice versa, a decrease in the selenium content in the diet of women increased the risk of depression (Pasco et al. 2012).

At the same time, it was noted that depression had not disappeared after the intake of sufficient selenium doses in geriatric patients (Gao et al. 2012), as well as during hemodialysis (Ekramzadeh et al. 2015). It should be noticed that depression may develop even during selenium intake, when its blood levels exceeded 82–83 mg/L (Colangelo et al. 2014).

The mechanism of the antidepressant effect of selenium is associated with its influence on the dopaminergic and serotonergic systems, possibly with blockade of MAO activity, and increased sensitivity of 5-HT receptors (Bruning et al. 2009; Solovyev 2015). It is considered that selenium due to its antioxidant effect can be recommended for the management of cardiovascular diseases and hyperlipidemia (Gharipour et al. 2017).

A decrease in selenium intake in some countries correlated with an increased risk of cardiomyopathies and cardiovascular diseases in general (Jenkins et al. 2020; Kuria et al. 2020).

Studies investigating the role of selenium deficiency in the development of cardiovascular diseases have shown that a correlation between insufficient selenium intake and the development of cardiovascular diseases may result from the presence of oxidative stress, as well as its complications (Al-Mubarak et al 2020; Gac et al. 2020).

Animal experiments carried out on animals with the decreased activity of glutathione peroxidase have shown that the restoration of glutathione peroxidase activity after the administration of selenium compounds occurs with a decrease in the content of active oxygen radicals. Selenium deficiency decreases the number of receptors for low density lipoprotein, which are necessary for cholesterol level regulation (Toppo et al. 2009; Talbi et al. 2019; Tong et al. 2020).

It should be noted that the number of low density lipoprotein receptors, which are necessary for the regulation

of cholesterol levels, decreases with selenium deficiency. A drop in selenium levels is associated with the risk of coronary heart disease when oxidative stress develops (Joseph and Weber 2013).

Selenium is also involved in the synthesis of an important cardioprotective agent coenzyme Q, a powerful antioxidant (Rees et al. 2013). In addition, selenium has hypolipidemic effects, especially when combined with tocopherol (Jin et al. 2011).

After the combination of selenium with amino acids, selenocysteine, selenomethionine, methylselenocysteine, and selenocystathionine are formed, which have organoprotective effects and play an important role in enzymatic reactions. Selenocysteine is present in the active site of glutathione peroxidase, which reduces the levels of hydrogen peroxide and organic peroxides. These compounds can regulate apoptosis, proliferation, and reproductive processes (Mehdi et al. 2013; Gac et al. 2020).

The development of cardiomyopathies, Keshan disease, and heart failure are associated with selenium deficiency in humans. In the first place, the levels of selenoproteins and, what is especially important, the activity of glutathione peroxidase, and succinate dehydrogenase are significantly reduced in such patients (Van der Pol et al. 2019; Bomer et al. 2020; Al-Mubarak et al. 2021). Recent studies have shown that selenium significance as a protective agent in heart failure is associated not only with its antioxidant properties, but also with its ability to prevent inflammation, autophagy, as well as the intrinsic and extrinsic apoptosis pathways (Mirdamadi et al. 2019; Al-Mubarak et al. 2020; Belenichev et al. 2020). Signaling pathways, such as p-AMPK, PARP, Nrf2, STAT, are involved in the protective effects of selenium (Zhang et al. 2019; Rayman 2020; Shalihat et al. 2021).

In addition, selenium protects against cardiovascular damage by increasing the survival rate of cardiomyocytes, including a mitochondria-dependent pathway and autophagy through peroxisome proliferator-activated receptor (PPAR) (Ren et al. 2016; Wang et al. 2020). It is established that the occurrence of oxidative and nitrosative stress plays a crucial pathophysiological role in the development of cardiovascular and nervous system disorders. One of the main groups of drugs that prevent stress development is selenium preparations. Sodium selenite (selenase, selenoline) is one of the registered selenium medicines.

The content of components of the enzymatic and non-enzymatic links of the thiol-disulfide system changes in acute disturbance of cerebral circulation. Disturbance of the non-enzymatic link is manifested by a decrease in the content of reduced glutathione, and an increase in oxidized glutathione. At the same time, the levels of cysteine and methionine also fall, against the background of an increase in homocysteine and nitrotyrosine.

It has been shown that cysteine selenite increases the levels of reduced glutathione, cysteine, methionine, and reduces the content of homocysteine and tyrosine to a greater extent than methionine selenite and sodium selenite. Similar alterations in the indices of the non-enzymat-

ic link of the thiol-disulfide system were noted in the myocardium after administration of selenium compounds.

The data obtained indicate the presence of neuro- and cardioprotective activity in compounds of selenium with amino acids (Gorchakova et al. 2017). A decrease in the activity of glutathione peroxidase, glutathione reductase, glutathione-S-transferase has been observed in brain tissue of rats under conditions of acute insufficiency of cerebral circulation.

Preliminary administration of selenium preparations may prevent the alterations in the parameters of the enzymatic link of the thiol-disulfide system (Notsek et al. 2015).

Neuroprotective and cardioprotective action of selenium preparations have also been established in relation to their action on heat shock proteins (HSP70). It was also noted that only 40% of rats with neurological disturbances survived on the 4th day after bilateral ligation of the common carotid arteries. The levels of heat shock proteins, which have a regulatory effect on the processes of cell death, decreased in the homogenate of brain tissue (Belenichev 2020).

Introduction of selenium preparations – selenoline, selenocysteine or selenomethionine – for 4 days significantly relieved acute neurological symptoms. Selenoline reduced animal mortality on the 4th day by 30%, methionine selenite – by 35%, and cysteine selenite – by 45%. Course administration of selenium preparations led to the expression of mRNA, to a greater extent with the introduction of cysteine selenite (Belenichev 2020).

The neuroprotective properties of selenium preparations are realized due to their ability to normalize the levels of heat shock proteins that limit the cytotoxic effects of free radicals, produce energotropic action, prevent a decrease in the membrane mitochondria charge, and the opening of the mitochondrial pore, as well as regulate the expression of transmembrane factors NF- κ B, c-fos, which is associated with their main biological function of chaperone proteins and providing protection of neurons from damage.

Cysteine selenite demonstrated the most significant impact due to its ability to positively influence on the synthesis of glutathione, modulate expression of selenium glutathione peroxidase, as well as influence on GSH-dependent mechanisms of neuroprotection (Belenichev et al. 2020)

The available literature data indicate that selenium is involved in hepatoprotective mechanisms. A decrease in the levels of blood selenium was noted due to decreased levels of transport selenoprotein P. When modeling liver lesions by the administration of lipoproteins, a disturbance of amino acid transport and production of cysteine selenite have been noted. Therefore, selenium may serve as a marker of liver diseases (Kozeniecki et al. 2020; Sherlock et al. 2020; Gul-Klein et al. 2021).

Selenium is involved in the body's metabolism in several ways. It can be involved in protein synthesis as selenomethionine instead of methionine (easily acylates met-tRNA). Also, through the mechanism of trans-sulfonation, it is converted to selenocysteine, which, through

information MEDLINE, EMBASE, Medline (PubMed), the Web of Science, and the Cochrane Central were searched for publications in English having the key words of this study. We used the following key words: [Selenium]; [antioxidants]; [neuroprotection], and [transcription factors]. All the authors independently selected articles, evaluated the quality of the data, presented and interpreted the data corresponding to the main idea of the study, and made up the final list of the references.

Conclusions

In recent years, the importance of selenium has been established for the prevention and management of viral infections. A relationship has been established between a decrease in selenoprotein levels and the incidence of coronavirus and HIV infection since the expression of pro-inflammatory interleukin-6 is increased in these diseases. Selenium is recommended for viral infections at a dose of 55 mcg per day. The non-pathogenic Coxsackie virus may turn into a pathogenic one when mice are kept on a low-selenium diet. Selenium deficiency can also significantly affect the survival of patients with HIV infection (Zhang and Liu 2020; Zhang et al. 2020).

A decrease in selenium levels in the body and the environment is known to increase mortality from AIDS and COVID-19 (Rayman 2012; Song et al. 2020; Zhang et al. 2020; Zhou et al. 2020), which is associated with the role of selenium as an antioxidant agent, as well as its action on gene activity in response to HIV or other viral infection.

Selenoproteins, first of all glutathione peroxidase, play an important role in the regulation of HIV manifestation. Proper selenium levels can delay the spread of the viral infections. A hypothetical HIV gene was cloned to prove the role of glutathione peroxidase in the prevention of viral infections, and that increased the synthesis of glutathione peroxidase. AIDS is often accompanied by lack of selenium, decreased numbers of T-helpers, or CD4 cells, and it is these lymphocytes that are the main target for HIV.

When searching for anticancer drugs, attention was drawn to organic and inorganic selenium compounds, which are liable to transformation. The mechanism of action of anticancer agents is mainly based on the induction

of apoptosis in tissues and organs (Siegel et al. 2020). They can affect gene expression and various signaling pathways, DNA repair or damage (Tan et al. 2019), and the processes of angiogenesis and metastasis (Gandin et al. 2018). In this case, the metabolites of the compounds have high activity, which is associated with the formation of reactive oxygen species, which in turn lead to oxidative stress with the oxidation of the thiol groups of proteins in the cell (Harmanci et al 2017). Selenium and its compounds may have prooxidant and antioxidant properties. Besides apoptosis, selenium compounds may cause other types of cell death. The manifestation of oxidative stress is promoted by iron ions, which accumulate in the cell, undergo the Fenton reaction and contribute to the formation of free radicals. Iron also increases the activity of the enzyme lipoxygenase (Chen et al. 2020; Subburayan et al. 2020).

Selenium-dependent glutathione peroxidase is considered the main enzyme that prevents tumor development and is the main antitumor regulator. Ferroptosis can occur when the antioxidant defense of the cell is exceeded. Excessive production of free radicals damages DNA and activates the p53 protein. Cancer cell proliferation can be inhibited by suppressing NF-kB. Binding to DNA inhibits protein kinase P or histone deacetylase. Selenium can stop the cell cycle. There is evidence suggesting that selenite can induce necrosis in breast cancer cells (Chen et al. 2020; Zhang et al 2020). These data indicate that the trace element selenium is a significant factor for the health and normal metabolism of vital systems and organs. At the same time, abnormalities in its content in the body, pharmacokinetics and pharmacodynamics can contribute to the development of a wide range of diseases that requires adherence to certain doses when using selenium preparations.

Funding/Support

This research was financially supported by Zaporizhzhia State Medical University (Grant No.2301020).

Conflict of interests

The authors declare no conflict of interests.

References

- Alhasan R, Kharma A, Leroy P, Jacob, Gaucherer C (2019) Selenium donors at the junction of inflammatory diseases. *Current Pharmaceutical Design* 35(15): 1707–1716. <https://doi.org/10.2174/1381612825666190701153903> [PubMed]
- Al-Mubarak Ali A, Van der Meer P, Bomer N (2021) Selenium, selenoproteins and heart failure: Current knowledge and future perspective. *Comorbidities of Heart Failure Reports* 18(3): 122–131. <https://doi.org/10.1007/s11897-021-00511-4> [PubMed] [PMC]
- Al-Mubarak AA, Grote Beverborg N, Anker SD, Samani NJ, Dickstein K, Filippatos G, van Veldhuisen DJ, Voors AA, Bomer N, van der Meer P (2020) A clinical tool to predict low serum selenium in patients with worsening heart failure. *Nutrients* 12(9): 35–41. <https://doi.org/10.3390/nu12092541> [PubMed] [PMC]
- Ahsan U, Kamran Z, Raza I, Ahmad S, Babar W, Riaz MH, Iqbal Z (2014) Role of selenium in male reproduction—a review. *Animal Reproduction Science* 146 (1–2): 55–62. <https://doi.org/10.1016/j.anireprosci.2014.01.009> [PubMed]
- Bartalena L, Baldeschi L, Boboridis K, Eckstein A, Kahaly GJ, Marcocci C, Perros P, Salvi M, Wiersinga WM, European Group on Graves' Orbitopathy [EUGOGO] (2016) The 2016 European thyroid

- association. European group on graves orbicopathy guidelines for the management of graves orbitopathy. *European Thyroid Journal* 5(1): 9–26. <https://doi.org/10.1159/000443828> [PubMed] [PMC]
- Belenichev IF, Gorchakova NO, Bukhtiyarova NV, Samura IB, Savchenko NV, Nefedov OO, Bak PG (2020) Modulation of HSP70-dependent mechanisms of endogenous neuroprotection with selenium derivatives under conditions of ischemic-type acute cerebrovascular accident modeling. *Pedagogy and Psychology of Sport* 6(4): 99–108. <http://dx.doi.org/10.12775/PPS.2020.06.04.009>
 - Bleys J, Navas-Achien A, Guallar E (2008) Serum selenium levels and all-cause, cancer and cardiovascular mortality among US adults. *Jama Internal Medicine* 168(4): 404–410. <https://doi.org/10.1001/archinternmed.2007.74> [PubMed]
 - Bomer N, Grote Beverborg N, Hoes MF, Streng KW, Vermeer M, Dokter MM, IJmker J, Anker SD, Cleland JGF, Hillege HL, Lang CC, Ng LL, Samani NJ, Tromp J, van Veldhuisen DJ, Touw DJ, Voors AA, van der Meer P (2020) Selenium and outcome in heart failure. *European Journal Heart Failure* 22(8): 1415–1423. <https://doi.org/10.1002/ejhf.1644> [PubMed] [PMC]
 - Bruning CA, Prigol M, Roehrs JA, Nogueira CW, Zeni CV (2009) Involvement of the serotonergic system in the anxiolytic-like effect caused by m-trifluoromethyl-diphenyl diselenide in mice. *Behavioural Brain Research* 205(2): 511–517. <https://doi.org/10.1016/j.bbr.2009.08.010> [PubMed]
 - Bruning CA, Souza AC, Gai BM, Zeni G, Nogueira CW (2011) Antidepressant-like effect of m-trifluoromethyl-diphenyl diselenide in the mouse forced swimming test involves opioid and serotonergic systems. *European Journal of Pharmacology* 658(2–3): 145–149. <https://doi.org/10.1016/j.ejphar.2011.02.039> [PubMed]
 - Chauychu-Noo N, Thananurak P, Boonkum W, Vongpralub T, Chankitisakudi V (2021) Effect of organic selenium dietary supplementation on quality and fertility of cryopreserved chicken sperm. *Cryobiology* 98: 57–62. <https://doi.org/10.1016/j.cryobiol.2020.12.008> [PubMed]
 - Chen Y, Liu S, Li J, Li Z, Quan J, Liu X, Tang Y, Liu B (2020) The latest view on the mechanism of ferroptosis and its research progress in spinal cord injury. *Oxidative Medicine and Cellular Longevity* 2020: 6375038. <https://doi.org/10.1155/2020/6375938> [PubMed] [PMC]
 - Chen Z, Lai H, Hou L, Chen T (2020) Rational design and action mechanism of chemically innovative organoselenium in cancer therapy. *Chemical Communication* 56(2): 179–196. <https://doi.org/10.1039/C9CC07683B> [PubMed]
 - Colangelo LA, He K, Whooley MA, Daviglius ML, Morris S, Liu K (2014) Selenium exposure and depressive symptoms. The coronary artery risk development in young adults' trace element study. *NeuroToxicology* 41: 167–174. <https://doi.org/10.1016/j.neuro.2014.02.003> [PubMed] [PMC]
 - Conner TS, Richardson AC, Miller JC (2014) Optimal serum selenium concentrations are associated with lower depressive symptoms and negative mood among young adults. *The Journal of Nutrition* 145(1): 59–65. <https://doi.org/10.3945/jn.114.198010> [PubMed]
 - De Silva Leme AGH, Cardoso BR (2020) Selenium and Alzheimer's disease. *Genetics, Neurology, Behavior, and Diet in Dementia* 2: 739–748. <https://doi.org/10.1016/B978-0-12-815868-5.00047-5>
 - Di Giuseppe R, Koch M, Nothlings U, Kastenmuller G, Artati A, Adamski J, Gunnar J, Lieb W (2019) Metabolomics signature associated with circulating serum selenoprotein P levels. *Endocrine* 64(3): 486–495. <https://doi.org/10.1007/s12020-018-1816-9> [PubMed]
 - Duntas LH, Benvenega S (2015) Selenium: an element for life. *Endocrine* 18(3): 756–775. <https://doi.org/10.1007/s12020-014-0477-6> [PubMed]
 - Ekramzadeh M, Mazloom Z, Sagheb M (2015) Association of depression with selenium deficiency and nutritional markers in the patients with end-stage renal disease on hemodialysis. *Journal Renal Nutrition* 25(4): 381–387. <https://doi.org/10.1053/j.jrn.2014.12.005> [PubMed]
 - Fontenelle LC, Feitosa MM, Morais JBS, Severo JS, de Freitas T, Beserra JB, Henriques GS, Marreiro DN (2018) The role of selenium in insulin resistance. *Brazilian Journal of Pharmaceutical Sciences* 54(1). <https://doi.org/10.1590/s2175-97902018000100139>
 - Gandin V, Khallar P, Braude J, Fernandes AP (2018) Organic selenium compounds as potential chemotherapeutic agents for improved cancer treatment. *Free Radical Biology and Medicine* 127: 80–97. <https://doi.org/10.1016/j.freeradbiomed.2018.05.001>
 - Gac P, Czerwinska K, Macek P, Jaremkov A, Mazur G, Pawlas K, Poreba R (2020) The importance of selenium and zinc deficiency in cardiovascular disorders. *Environment Toxicology and Pharmacology* 82: 1035–1053. <https://doi.org/10.1016/j.etap.2020.103553> [PubMed]
 - Gao S, Jin Y, Unverzagt FW, Liang C, Hall KS, Cao J, Ma F, Murrel JA, Cheng Y, Li P, Bian J, Hendrie HC (2012) Selenium level and depressive symptoms in a rural elderly Chinese cohort. *BMC Psychiatry* 12: 72–74. <https://doi.org/10.1186/1471-244X-12-72> [PubMed] [PMC]
 - Gharipour M, Sadeghi M, Behmanesh M, Salehi M, Nezafati P, Gharipour A (2017) Selenium homeostasis and clustering of cardiovascular risk factors: a systemic review. *Acta Biomed* 88(3): 263–270. <https://doi.org/10.23750/abm.v88i3.5701> [PubMed] [PMC]
 - Gül-Klein S, Haxhiraj D, Seelig J, Kästner A, Hackler J, Sun Q, Heller RA, Lachmann N, Pratschke J, Schmelzle M, Schomburg L (2021) Serum selenium status as a diagnostic marker for the prognosis of liver transplantation. *Nutrients* 13(2): 619. <https://doi.org/10.3390/nu13020619> [PubMed] [PMC]
 - Golubkina NA, Sindireva AV, Zaitsev VF (2017) Intraregional variability of the selenium status of the population. *Medical Ecology* 12(1): 107–127. <https://doi.org/10.18470/1992-1098-2017-1-107-127>
 - Gorchakova NO, Belenichev IF, Bukhtiyarova NB (2017) Influence of selenium-containing agents on the state of the non-enzymatic part of the thiol-disulfide system of rats in acute cerebrovascular insufficiency. *The National Congress of Pharmacologists: abstracts add. Zaporizhzhia*, pp. 28–29. [in Ukrainian]
 - Hadrup N, Ravn-Haren G (2020) Acute human toxicity and mortality after selenium ingestion a review. *Journal of Trace Elements in Medicine and Biology* 58: 126–435. <https://doi.org/10.1016/j.jtemb.2019.126435> [PubMed]
 - Harmanci D, Erbayraktar Z, Sayin O, Guner GA (2017) In vitro effects of selenium on human glioblastoma multiforme cells lines: A preliminary study. *Acta Clinica Croatia* 56(1): 48–57. <https://doi.org/10.20471/acc.2017.56.01.08> [PubMed]
 - Heller RA, Sun Q, Hackler J, Seelig J, Seibert L, Cherkezov A, Minich WB, Seemann P, Diegmann J, Pilz M, Bachmann M, Ranjbar A, Moghaddam A, Schomburg L (2020) Prediction of survival odds in COVID-19 by zinc, age and selenoprotein P as composite biomarker. *Redox Biology* 38: 101764. <https://doi.org/10.1016/j.redox.2020.101764> [PubMed] [PMC]

- Hofstee P, McKeating DR, Bartho LA, Anderson ST, Perkins AV, Cuffe J (2020) Maternal selenium deficiency in mice alters offspring glucose metabolism and thyroid status in a sexually dimorphic manner. *Nutrients* 12(1): 267. <https://doi.org/10.3390/nu12010267> [PubMed] [PMC]
- Hossain A, Skalicky M, Brestic M, Maitra S, Sarkar S, Ahmad Z, Vemuri H, Garai S, Mondal M, Bhatt R, Kumar P, Banerjee P, Saha S, Islam T, Laing AM (2021) Selenium biofortification: roles, mechanisms, responses and prospects. *Molecules* 26(4): 881. <https://doi.org/10.3390/molecules26040881> [PubMed] [PMC]
- Iglesias P, Selgas R, Romero S, Dies JI (2013) Selenium and kidney disease. *Journal of Nephrology* 26(2): 266–272. <https://doi.org/10.5301/jn.5000213> [PubMed]
- Jenkins DJA, Kitts D, Giovannucci EL, Sahye-Pudurath S, Paquette M, Blanco Mejia S, Patel D, Kavanagh M, Tsirakis T, Kendall CWC, Pichika SC, Sievenpiper JL (2020) Selenium, antioxidants, cardiovascular disease, and all-cause mortality: a systematic review and meta-analysis of randomized controlled trials. *The American Journal of Clinical Nutrition* 112(6): 1642–1682. <https://doi.org/10.1093/ajcn/nqaa245> [PubMed] [PMC]
- Jiao Y, Wang Y, Guo S, Wang G (2017) Glutathione peroxidases as oncotargets. *Oncotarget* 8(45): 80093–80102. <https://doi.org/10.18632/oncotarget.20278> [PubMed] [PMC]
- Jin RC, Mahoney CE, Coleman Anderson L, Ottaviano F, Croce K, Leopold JA, Zhang YY, Tang SS, Handy DE, Loscalzo J (2011) Glutathione peroxidase-3 deficiency promotes platelet dependent thrombosis in vivo. *Circulation* 123(18): 1963–1973. <https://doi.org/10.1161/CIRCULATIONAHA.110.000034> [PubMed] [PMC]
- Joseph J, Weber KT (2013) Selenium and cardiometabolic health: inconclusive yet intriguing evidence. *The American Journal of the Medical Sciences* 346(3): 216–220. <https://doi.org/10.1097/MAJ.0b013e3182638716> [PubMed] [PMC]
- Kieliszek M, Blasejak S, Bzducha-Wrobel A, Kot AM (2019) Effect of selenium on growth and antioxidative system of yeast cells. *Molecular Biology Reports* 46(2): 1797–1808. <https://doi.org/10.1007/s11033-019-04630-z> [PubMed]
- Kim J, Chung HS, Choi MK, Roh YK, Yoo HJ, Park JH, Kim DS, Yu JM, Moon S (2019) Association between serum selenium and the presence of diabetes mellitus; meta-analysis of observational studies. *Diabetes & Metabolism Journal* 43(4): 447. <https://doi.org/10.4093/dmj.2018.0123> [PubMed] [PMC]
- Kohler LN, Foote J, Kelley CP, Florea A, Shelly C, Chow HS, Hsu P, Batai K, Ellis N, Saboda K, Lance P, Jacobs ET (2018) Selenium and type 2 diabetes: systematic review. *Nutrients* 10(12): 1924. <https://doi.org/10.3390/nu10121924> [PubMed] [PMC]
- Kozeniecki M, Ludke R, Kerner J, Patterson B (2020) Micronutrients in liver disease: roles, risk factors for deficiency, and recommendations for Supplementation. *Nutrition in Clinical Practice* 35: 50–62. <https://doi.org/10.1002/ncp.10451> [PubMed]
- Krelszek M, Bano I, Zare HA (2021) Comprehensive review on selenium and its effects on human health and distribution in Middle Eastern Countries. *Biological Trace Element Research*, 17 pp. <https://doi.org/10.1007/s12011-021-02716-z>
- Kuria A, Tian H, Li M, Wang Y, Aaseth JO, Zang J, Cho Y (2020) Selenium status in the body and cardiovascular disease: a systematic review and meta-analysis. *Critical Reviews in Food Science and Nutrition* 1–10. <https://doi.org/10.1080/10408398.2020.1803200> [PubMed]
- Liu C, Yan Q, Gao C, Lin L, Wei J (2021) Study on antioxidant effect of recombinant glutathione peroxidase 1. *International Journal of Biological Macromolecules* 170(15): 503–513. <https://doi.org/10.1016/j.ijbiomac.2020.12.183> [PubMed]
- Loef M, Schrauzer GN, Walach H (2011) Selenium and Alzheimer's disease: a systematic review. *Journal of Alzheimer's Disease* 26(1): 81–104. <https://doi.org/10.3233/JAD-2011-110414> [PubMed]
- Lv J, Ai P, Lei S, Zhou F, Chen S, Zhang Y (2020) Selenium levels and skin diseases systematic review and meta-analysis. *Journal of Trace Elements in Medicine and Biology* 62: 126548. <https://doi.org/10.1016/j.jtemb.2020.126548> [PubMed]
- Mehdi Y, Hornick JL, Istasse E, Dufrasne I (2013) Selenium in the environment, metabolism and involvement in body functions. *Molecules* 18(3): 3292–3311. <https://doi.org/10.3390/molecules18033292> [PubMed] [PMC]
- Mintziori G, Mousiolis A, Duntas LH, Goulis DG (2020) Evidence for a manifold role of selenium in infertility. *Hormones* 19(1): 55–59. <https://doi.org/10.1007/s42000-019-00140-6> [PubMed]
- Mirdamadi A, Rafiei R, Kahazaipour G, Fouladi L (2019) Selenium level in patients with heart failure versus normal individuals. *International Journal of Preventive Medicine* 10: 210–220. https://doi.org/10.4103/ijpvm.IJPVM_45_18 [PubMed] [PMC]
- Młyniec K, Gawel M, Doboszevska U, Strawicz J, Pytka K, Davies CI, Budziszewska B, Nowak G (2015) Essential elements in depression and anxiety. Part II. *Pharmacological Reports* 67(2): 187–194. <https://doi.org/10.1016/j.pharep.2014.09.009> [PubMed]
- Nogueira CW, Barbosa NV, Rocha JB (2021) Toxicology and pharmacology of synthetic organoselenium compounds: an update. *Archives of Toxicology* 95: 1179–1226. <https://doi.org/10.1007/s00204-021-03003-5> [PubMed] [PMC]
- Notsek MS, Gorchakova NO, Belenichev IF, Puzyrenko AM, Chekman IS (2015) Influence of selenium preparations on indicators of enzymatic link of thiol-disulfide system in brain tissues of animals with acute cerebral insufficiency blood circulation. *Bulletin of Problems of Biology and Medicine* 2(4): 202–205. [in Ukrainian]
- Pasco JA, Jacka FN, Williams LJ, Evans-Cleverdon M, Brennan SL, Kolowicz MA, Nicholson GC, Ball MJ, Berk M (2012) Dietary selenium and major depression: a nested case-control study. *Complementary Therapies in Medicine* 20(3): 119–123. <https://doi.org/10.1016/j.ctim.2011.12.008> [PubMed]
- Piagntini M, Silva DC, Dell'Aqua CPF, Moya-Araujo CF, Codognoto VM, Ramos AA, Oba E (2017) Effect of selenium supplementation on semen characteristics of Brazil's ram. *Reproduction in Domestic Animals* 52(3): 355–358. <https://doi.org/10.1111/rda.12903> [PubMed]
- Pillai R, Uyehara-Lock JH, Bellinger FP (2014) Selenium and selenoprotein function in brain disorders. *IUBMB Life* 66(4): 229–239. <https://doi.org/10.1002/iub.1262> [PubMed]
- Pogotova GA, Gorchakova NO, Belenichev IF, Chekman IS (2014) The effect of selenase on energy metabolism and prooxidant-antioxidant system in the organs of rats with toxic hepatitis. *Bulletin of Problems of Biology and Medicine* 2(3): 216–220. [in Ukrainian]
- Rahmanto AS, Davies MJ (2012) Selenium containing amino acids as direct and indirect antioxidants. *IUBMB Life* 46(11): 86–87. <https://doi.org/10.1002/iub.1084> [PubMed]
- Rakov OV, Muz VA (2019) Selenium and human health. *Endocrinology* 24(2): 178–186. <https://doi.org/10.31793/1680-1466.2019.24-2.178>

- Rayman MP (2012) Selenium and human health. *The Lancet* 379(9822): 1256–1263. [https://doi.org/10.1016/S0140-6736\(11\)61452-9](https://doi.org/10.1016/S0140-6736(11)61452-9) [PubMed]
- Rayman MP (2020) Selenium intake, status and health: a complex relationship. *Hormones (ATHens)* 19(1): 9–14. <https://doi.org/10.1007/s42000-019-00125-5> [PubMed] [PMC]
- Rayman MP, Stranges S (2013) Epidemiology of selenium and type 2 diabetes: Can we make sense of it? *Free Radical Biology and Medicine* 65: 1557–1564. <https://doi.org/10.1016/j.freeradbiomed.2013.04.003> [PubMed]
- Reddy VS, Bukke S, Dutt N, Rana P, Pandey AK (2017) A systematic review and meta-analysis of the circulatory, erythrocytic and CSF selenium levels in Alzheimer's disease: A meta-analysis (AMMA study-I). *Journal of Trace Elements in Medicine and Biology* 42: 58–75. <https://doi.org/10.1016/j.jtemb.2017.04.005> [PubMed]
- Rees K, Hartley L, Day C, Flowers N, Clarke A, Stranges S (2013) Selenium supplementation for the primary prevention of cardiovascular disease. *Cochrane Database of Systematic Reviews* 346(3): CD009671. <https://doi.org/10.1002/14651858.CD009671.pub2> [PubMed] [PMC]
- Reeves MA, Bellinger PP, Berry MJ (2019) The neuroprotective functions of selenoprotein M and its role in cytosolic calcium regulation. *Antioxidants and Redox Signaling* 13(7): 809–818. <https://doi.org/10.1089/ars.2009.2883> [PubMed] [PMC]
- Ren H, Mu J, Ma J, Gong J, Li J, Wang J, Gao T, Zhu P, Zheng S, Xie J, Yuan B (2016) Selenium inhibits homocysteine-induced endothelial dysfunction and apoptosis via activation of AKT. *Cellular Physiology and Biochemistry* 38 (3): 871–882. <https://doi.org/10.1159/000443041> [PubMed]
- Saburov KA, Kazybekova AA, Zakharov TA (2016) Influence of selenium on the state of immunity and oxidative homeostasis in the ruffed populations of the Central Tien Shan. *TSU Bulletin* 21(6): 2290–2299. [in Russian]
- Saitar T (2021) Selenium role in reproduction, pregnant/postpartum women and neonates: a current study. *Current Nutrition and Food Science* 17(1): 28–37. <https://doi.org/10.2174/1573401316999200423121620>
- Shalihat A, Hasanah AN, Lesmana R, Budiman A, Gozali D (2021) The role of selenium in cell survival and its correlation with protective effects against cardiovascular disease: A literature review. *Biomedicine and Pharmacotherapy* 134: 111–125. <https://doi.org/10.1016/j.biopha.2020.111125> [PubMed]
- Sherlock LG, Sjostrom K, Sian L, Delaney C, Tipple TE, Krebs NF, Nozik-Grayck E, Wright CJ (2020) Hepatic-specific decrease in the expression of selenoenzymes and factors essential for selenium processing after endotoxemia. *Frontiers in Immunology* 11: 595282. <https://doi.org/10.3389/fimmu.2020.595282> [PubMed] [PMC]
- Siegel KI, Miller KD, Jemal A (2020) Cancer statistics, 2020. *CA: A Cancer Journal for Clinicians* 70(1): 7–30. <https://doi.org/10.3322/caac.21590> [PubMed]
- Skowronska-Joozwtak E (2015) The effect of selenium on thyroid physiology and pathology. *Thyroid Research* 8: A23. <https://doi.org/10.1186/1756-6614-8-S1-A23>
- Solovyev ND (2015) Importance of selenium and selenoprotein for brain function. From antioxidant protection to neuronal signaling. *Journal of Inorganic Biochemistry* 153: 1–12. <https://doi.org/10.1016/j.jinorgbio.2015.09.003> [PubMed]
- Solovyev ND, Drobishev E, Bjorklund G, Dubrovskii Y, Lysiuk R, Rayman MP (2018) Selenium, selenoprotein P and Alzheimer's disease. Is there a link? *Free Radical Biology and Medicine* 127: 124–133. <https://doi.org/10.1016/j.freeradbiomed.2018.02.030> [PubMed]
- Song P, Li W, Xie J, Hou Y, You C (2020) Cytokine storm induced by STARS-Cov-2. *Clinica Chimica Acta* 509: 280–287. <https://doi.org/10.1016/j.cca.2020.06.017> [PubMed] [PMC]
- Steinbrenner H, Sies H (2013) Selenium homeostasis and antioxidant selenoproteins in brain: implications for disorders in the central nervous system. *Archives of Biochemistry and Biophysics* 536(2): 152–157. <https://doi.org/10.1016/j.abb.2013.02.021> [PubMed]
- Stoedter M, Renko K, Hog A, Schomburg (2010) Selenium controls the sex-specific immune response and selenoprotein expression during the acute phase response in mice. *Biochemical Journal* 429(1): 43–51. <https://doi.org/10.1042/BJ20091868> [PubMed]
- Stoffaneiler R, Morse NL (2015) A review of dietary selenium intake and selenium status in Europe and the Middle East. *Nutrients* 7(3): 1494–1537. <https://doi.org/10.3390/nu7031494> [PubMed] [PMC]
- Subburayan K, Thaayulathil F, Pallichankandy S, Cheratta AR, Galadari S (2020) Superoxide-mediated ferroptosis in human cancer cells induced by sodium selenite. *Translational Oncology* 13(11): 100843. <https://doi.org/10.1016/j.tranon.2020.100843> [PubMed] [PMC]
- Talbi W, Ghazouani T, Braconi D, Ben Abdallah R, Raboudi F, Santucci A, Fattouch S (2019) Effects of selenium on oxidative damage and antioxidant enzymes of eukaryotic cells, wine *Saccharomyces cerevisiae*. *Journal of Applied Microbiology* 126(2): 555–566. <https://doi.org/10.1111/jam.14150> [PubMed]
- Tan HW, Mo HY, Lau ATY, Xu YM (2019) Selenium species: current status and potentials in cancer prevention and therapy. *International Journal of Molecular Sciences* 20(1): 75. <https://doi.org/10.3390/ijms20010075> [PubMed] [PMC]
- Tamtaji OR, Heidari-Soureshjani R, Mirhosseini N, Kouchaki E, Bahmani F, Aghadavod E, Tajabadi-Ebrahimi M, Asemi Z (2019) Probiotic and selenium co-supplementation, and the effects on clinical, metabolic and genetic status in Alzheimer's disease: A randomized, double-blind, controlled trial. *Clinical Nutrition* 38(6): 2569–2375. <https://doi.org/10.1016/j.clnu.2018.11.034> [PubMed]
- Tinggi U (2014) Essentiality and toxicity of selenium and its status in Australia: a review. *Toxicology Letters* 137(1–2): 103–110. [https://doi.org/10.1016/S0378-4274\(02\)00384-3](https://doi.org/10.1016/S0378-4274(02)00384-3) [PubMed]
- Tong C, Li P, Yu LH, Li L, Li K, Chen E, Yang SH, Long M (2020) Selenium-rich yeast attenuates ochratoxin A-induced small intestinal injury in broiler chickens by activating the Nrf2 pathway and inhibiting NF-KB activation. *Journal of Functional Foods* 66: 103784. <https://doi.org/10.1016/j.jff.2020.103784>
- Toppo S, Flohe L, Ursini F, Vanin S, Matorino M (2009) Catalytic mechanisms and specificities of glutathione peroxidases: Variations of a basic scheme. *Biochimica et Biophysica Acta (BBA) – General Subjects* 1790(11): 1486–1500. <https://doi.org/10.1016/j.bbagen.2009.04.007> [PubMed]
- Van der Pol A, van Gilst WH, Voors AA, van der Meer P (2019) Treating oxidative stress in heart failure: past, present and future. *European Journal Heart Failure* 21(1): 425–435. <https://doi.org/10.1002/ejhf.1320> [PubMed] [PMC]
- Viniarska HB, Bodnar OI, Vasilenko OV, Stanislavichuk AV (2017) Accumulation and effects of selenium and zinc on *Chlorella vulgaris* beij. (chlorophyta) metabolism. Heavy Metals and other pollutants in the Environment. In: Zalkov GE (Ed.) *Biological Aspects*. Apple Academic Press, 293–315.

- Wang G, Um P, Dickerman BA, Liu J (2018) Zinc, magnesium, selenium and depression: a review of the evidence, potential mechanism and implications. *Nutrients* 10(5): 584. <https://doi.org/10.3390/nu10050584> [PubMed] [PMC]
- Wang YG, Liu CZ, Li YZ, Peng Y, Yan S (2020) Cotreatments with Dex and Na₂SeO₃ further improved antioxidant and anti-inflammatory protection of myocardial cells from I/R injury compared to their individual treatments. *Free Radical Research* 54(1): 76–90. <https://doi.org/10.1080/10715762.2019.1707198> [PubMed]
- Wang Y, Rijntjes E, Wu Q, Ly H, Gao C, Shi B, Schomburg L (2020) Selenium deficiency is linearly associated with hypoglycemia in healthy adults. *Redox Biology* 37: 101709. <https://doi.org/10.1016/j.redox.2020.101709> [PubMed] [PMC]
- Weekley CM, Harris HH (2013) Which form is that? The importance of selenium speciation and metabolism in the prevention and treatment of disease. *Chemical Society Reviews* 42(23): 8870–8894. <https://doi.org/10.1039/c3cs60272a> [PubMed]
- Wichman J, Winther KH, Bonnema SJ, Negedus L (2016) Selenium supplementation significantly reduces thyroid autoantibody levels in patients with chronic autoimmune thyroiditis: a systematic review and meta-analysis. *Thyroid* 26(12): 1681–1692. <https://doi.org/10.1089/thy.2016.0256> [PubMed]
- Yildirim S, Ozkan C, Huyut Z, Cinar A (2019) Detection of Se, vit E, vit A, MDA, 8-OHdG and CoQ10 levels and histopathological changes in heart tissue in sheep with white muscle disease. *Biological Trace Element Research* 188(2): 419–423. <https://doi.org/10.1007/s12011-018-1434-7> [PubMed]
- Yovanovich LN, Ermakov VV (2020) The value of selenium and zinc in the prevention and treatment of certain diseases. In: *Materials of the International Biochemical Symposium, Tiraspol*, pp. 71–83. [in Russian]
- Zhang J, Taylor EW, Bennett K, Saad R, Rayman MP (2020) Association between regional selenium status and reported of COVID-19 cases in China. *The American Journal of Clinical Nutrition* 111(6): 1297–1299. <https://doi.org/10.1093/ajcn/nqaa095> [PubMed] [PMC]
- Zhang L, Gao Y, Feng H, Zou N, Wang K, Sun D (2019) Effects of selenium deficiency and low protein intake on the apoptosis through a mitochondria-dependent pathway. *Journal of Trace Elements in Medicine and Biology* 56: 21–30. <https://doi.org/10.1016/j.jtemb.2019.06.019> [PubMed]
- Zhang L, Liu Y (2020) Potential interventions for novel coronavirus in China: a systematic review. *Journal of Medical Virology* 63: 479–490. <https://doi.org/10.1002/jmv.25707> [PubMed] [PMC]
- Zhang Y, Cartland SP, Henriquez R, Patel S, Gammelgaard B., Flouda K, Hawkins CL, Rayner BS (2020) Selenomethionine supplementation reduces lesion burden, improves vessel function and modulates the inflammatory response within the setting of atherosclerosis. *Redox Biology* 29: 401–411. <https://doi.org/10.1016/j.redox.2019.101409> [PubMed] [PMC]
- Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, Xiang J, Wang Y, Song B, Gu X, Guan L, Wei Y, Li H, Wu X, Xu J, Tu S, Zhang Y, Chen H, Cao B (2020) Clinical course and risk factors for mortality adult in patients with COVID-19 in Wuhan China: a retrospective cohort study. *The Lancet* 395(10229): 1054–1063. [https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3) [PubMed] [PMC]

Author contributions

- **Igor F. Belenichev**, Dr. habil. of Biology, Full Professor, head at the Department of Pharmacology and Medical Formulation with Course of Normal Physiology, Zaporizhzhia State Medical University, e-mail: ifb1914@mail.ru, ORCID ID <https://orcid.org/0000-0003-1273-5314>. The author of the work concept and design of the article was responsible for critical review and final approval of the article.
- **Nadia O. Gorchakova**, Dr. habil. of Medicine, Full Professor, Bogomolets National Medical University, e-mail: gorchakova.na21@gmail.com, ORCID ID <https://orcid.org/0000-0002-7240-5862>. The author gave critical review and the final approval of the version to be submitted.
- **Nina V. Bukhtiyarova**, PhD of Medicine, Assistant Professor, Zaporizhzhia State Medical University, e-mail: hce.181901@gmail.com, ORCID ID <https://orcid.org/0000-0003-3499-3111>. The author was engaged in collection, analysis and interpretation of the data for publication, and in writing the article.
- **Iryna B. Samura**, PhD of Medicine, Assistant Professor, Zaporizhzhia State Medical University, e-mail: pea.160119@gmail.com, ORCID ID <https://orcid.org/0000-0001-5352-3209>. The author was engaged in collection, analysis and interpretation of the data for publication, and in their of the data critical review.
- **Natalia V. Savchenko**, PhD of Medicine, Assistant Professor, Bogomolets National Medical University, e-mail: zhc.9416@gmail.com, ORCID ID <https://orcid.org/0000-0003-3392-6638>. The author was engaged in collection, analysis and interpretation of the data for publication, and in their of the data critical review.
- **Olena O. Popazova**, teaching assistant, Master of Biology, Zaporizhzhia State Medical University, e-mail: popazova.ea@gmail.com, ORCID ID <https://orcid.org/0000-0001-7740-460X>. The author was engaged in collection, analysis and interpretation of the data for publication, and in their of the data critical review.