

Springtails and Silverfishes (Apterygota) Chapter 13.5

Jürg Zettel

University of Bern, Institute of Ecology and Evolution Baltzerstrasse 6, CH-3012 Bern, Switzerland

Corresponding author: *Jürg Zettel* (juersi.zettel@bluewin.ch)

Academic editor: *David Roy* | Received 27 January 2010 | Accepted 25 May 2010 | Published 6 July 2010

Citation: Zettel J (2010) Springtails and Silverfishes (Apterygota). Chapter 13.5. In: Roques A et al. (Eds) Alien terrestrial arthropods of Europe. *BioRisk* 4(2): 851–854. doi: 10.3897/biorisk.4.47

Abstract

The alien fauna of Apterygota is still limited in Europe. Springtails (Collembola) alien to Europe include only three species to which add a cryptogenic one. Two nowadays cosmopolitan species of silverfishes may originate from Central America. The reasons of this limited colonisation of Europe are briefly discussed.

Keywords

Apterygota, Collembola, springtails, Zygentoma, silverfishes

13.5.1. Diplura alien to Europe

No introductions of alien species into Europe are known.

13.5.2. Collembola (Springtails) alien to Europe

Worldwide ca. 6500 collembolan species are listed, belonging to 18 families (Hopkin 1997). For Europe, there are estimated to be ca. 1500 species, belonging to 16 families (taxonomic work is still progressing).

Collembola are the most abundant terrestrial arthropods, colonising all soil habitats that provide enough humidity and food, such as organic matter or microorganisms. Example habitats include root rosettes of high alpine plants, plant debris on the shore, natural soils, as well as microhabitats such as flower pots. Most species are soil or

litter dwellers, whilst only few species live on the surface or in the vegetation (mainly Entomobryidae and Symphyleona). In mature soil, abundances may attain values of 50–100'000 individuals/m². Local gradations in abundance are a well known phenomenon in many Collembola.

As detritivores, Collembola are not generally considered as pest species. Exceptions are two species of Symphyleona living above ground in the vegetation layer: the European *Sminthurus viridis* which became a severe pest in Australia on alfalfa, clover etc, and the ubiquitous *Bourletiella hortensis* is known to feed on vegetable seedlings when natural food (weed seedlings) is absent. The ubiquitous onychiurid *Protaphorura armata* also switches food source in the absence of weeds, but only as a secondary pest when feeding on wounds infected by microorganisms. In Europe, no Collembola are declared as agronomic pests (e.g. in CABI Crop Protection Compendium) (CABI 2009).

A 100 or more Collembola species may occur in the same soil habitat, and through occupying all available niches are believed to preclude establishment of alien species. To date, alien Collembola have only been observed to become invasive and replace indigenous species in isolated microhabitats and in extreme climates such as two species on sub-Antarctic islands (Convey et al. 1999, Greenslade 2002).

Identifying alien Collembola is difficult due to the limited number of specialists investigating soil fauna. No intentional introductions to Europe have occurred. Unintentional transport within soil of ornamental plants, with vegetables, dirty equipment and vehicles easily moves Collembola over large distances. Short life cycles and parthenogenetic development of a number of species may also increase the chances to colonise new sites. Therefore the distribution ranges of alien species within Europe may increase steadily.

Only three records of alien Collembola introduced to Europe have been published. These concern two species in the family Isotomidae, *Proisotoma filifera*, originating from Central America but found in Dutch greenhouses (Ellis 1970), and *Desoria trispinata*, that originates from North America but has appeared in anthropogenic habitats, mainly towns (Christian 1987, Christian and Kindl-Stomatopulos 1999, Kindl-Stomatopulos). A third species in the family Onychiuridae, *Onychiurus folsomi*, originating from Australia, is restricted to earthworm cultures in Spain (Arbea and Jordana 1988). In addition, we considered a cryptogenic species, *Sminthurinus trinotatus* (Katiannidae), which presents a very disjunct known distribution (southern Europe, eastern Asia).

13.5.3. *Zygentoma* (Silverfishes) alien to Europe

Zygentoma or silverfishes comprise five families (in Europe Lepismatidae only) with 12 genera (three in Europe) and ca. 370 species (ten in Europe). The two (today) cosmopolitan species *Ctenolepisma longicaudata* and *Thermobia domestica* (both Lepismatidae), may originate from central parts of America. Once moved from western Mediterranean regions to central and northern Europe, they mainly colonise anthropogenic habitats, where they may become pests by destroying paper or stored products.

References

- Arbea JL, Jordana R (1988) Nota sobre la presencia masiva de *Onychiurus folsomi* Schaeffer (Collembola, Onychiuridae) en lechos de *Eisenia andrei* (Oligochaeta, Lumbricidae). *Boletín de Sanidad Vegetal Plagas* 14: 535–540.
- CABI Crop Protection Compendium. <http://www.cabicompendium.org/cpc>.
- Christian E (1987) *Collembola (Springschwänze)*. *Catalogus Faunae Austriae XIIIa*, Wien: Österreichische Akademie der Wissenschaften. 80 pp.
- Christian E, Kindl-Stamatopolos L (1999) Arthropods on plastered riverbanks in the builtup area of Vienna. In: Tajovský K, Pižl V (Eds) *Soil Zoology in Central Europe. Proceedings of the 5th Central European Workshop on Soil Zoology, April 1999*. České Budějovice: Institute of Soil Biology, 27–30.
- Convey P, Greenslade P, Arnold RJ, Block W (1999) Collembola of subantarctic South Georgia. *Polar Biology* 22: 1–6.
- Ellis WN (1970) *Proisotoma filifera* Denis in Holland, with a note on its classification (Collembola, Isotomidae). *Entomologische Berichten* 30: 18–24.
- Essl F, Rabitsch W (2002). *Neobiota in Österreich*. Wien: Umweltbundesamt. 432 pp.
- 8-Greenslade P (2002) Assessing the risk of exotic Collembola invading subantarctic islands: prioritising quarantine management. *Pedobiologia* 46: 338–344.
- Hopkin SP (1997). *Biology of the springtails (Insecta: Collembola)*. Oxford: Oxford University Press. 330 pp.
- Hopkin SP. *The Natural World in Close Up*. available at [www.stevehopkin.co.uk].
- Kindl-Stamatopolos L (2001) Arthropoden des Wienflussufers im dicht bebauten Stadtgebiet Wiens. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Österreich* 138: 1–15.

Table 13.5.1. List and characteristics of the Collembola and Zygentoma species alien to Europe. Status: **A** Alien to Europe **C** cryptogenic species. Country codes abbreviations refer to ISO 3166 (see appendix I). Habitat abbreviations refer to EUNIS (see appendix II). Last update 11/01/2010;

Order-Family Species	Status	Regime	Native range	1st record in Europe	Invaded countries	Habitat	References
Collembola- Onychiuridae							
<i>Onychiurus folsomi</i> (Schäffer, 1900)	A	detritivorous	Australia	1988 ?, ES	ES, GB	I (Vermiculture), J100	Arbea and Jordana (1988), Hopkin (2009)
Collembola- Isotomidae							
<i>Proisotoma filifera</i> (Denis, 1931)	A	detritivorous	Central America	1968, NL	NL	J100	Ellis (1970)
<i>Desoria trispinata</i> (MacGillivray, 1896)	A	detritivorous	North America	ca 1900, AT	AT, DE, IT, NO, PT, RU	J	Christian (1987), Christian and Kindl- Stomatopulos (1999), Kindl-Stamatopulos (2001)
Collembola- Katiannidae							
<i>Smimthurinus trinotatus</i> (Axelson, 1905)	C	detritivorous	South Europe, East Asia?	1925, GB	AT, DE, FR, GB, IT	I2, J100	Essl and Rabitsch (2002), Hopkin (2009)
Zygentoma- Lepismatidae							
<i>Ctenolepisma longicaudata</i> (Escherich, 1905)	C	detritivorous	Central America?	Unknown	CY, FR, IT, MT, PT	J1 (stored products)	Essl and Rabitsch (2002)
<i>Thermobia domestica</i> (Packard, 1873)	C	detritivorous	Central America?	Unknown	CY, DE, DK, FR, GB, IT, PT	J1 (stored products)	Essl and Rabitsch (2002)