

# Management of invasive alien species in Spain: A bibliometric review

Rafael Muñoz-Mas<sup>1</sup>, Martina Carrete<sup>2</sup>, Pilar Castro-Díez<sup>3</sup>,  
Miguel Delibes-Mateos<sup>4</sup>, Josep A. Jaques<sup>5</sup>, Marta López-Darias<sup>6</sup>,  
Manuel Nogales<sup>6</sup>, Joan Pino<sup>7,8</sup>, Anna Traveset<sup>9</sup>, Xavier Turon<sup>10</sup>,  
Montserrat Vilà<sup>11,12</sup>, Emili García-Berthou<sup>1</sup>

**1** GRECO, Institute of Aquatic Ecology, University of Girona, 17003 Girona, Spain **2** Departamento de Sistemas Físicos, Químicos y Naturales, Universidad Pablo de Olavide, Ctra. de Utrera, km. 1 41013, Seville, Spain **3** Biological Invasions Research Group (BioInv), Departamento de Ciencias de la Vida, Universidad de Alcalá, Pza. San Diego, s/n, 28801, Alcalá de Henares, Madrid, Spain **4** Instituto de Estudios Sociales Avanzados (IESA-CSIC), Plaza Campo Santo de los Mártires, 7, 14004 Córdoba, Spain **5** Departament de Ciències Agràries i del Medi Natural, Universitat Jaume I, Av. Vicent Sos Baynat, s/n, 12071 Castelló de la Plana, Spain **6** Island Ecology and Evolution Research Group, Instituto de Productos Naturales y Agrobiología (IPNA-CSIC), Avda. Astrofísico Francisco Sánchez, 3, 38206 - San Cristóbal de La Laguna Santa Cruz de Tenerife - Islas Canarias, Spain **7** CREA, E08193 Bellaterra (Cerdanyola del Vallès), Catalonia, Spain **8** Universitat Autònoma de Barcelona, E08193 Bellaterra (Cerdanyola del Vallès), Catalonia, Spain **9** Instituto Mediterráneo de Estudios Avanzados (IMEDEA, CSIC-UIB), C/ Miquel Marquès, 21 - 07190 Esporles, Mallorca, Illes Balears, Spain **10** Departamento de Ecología Marina, Centro de Estudios Avanzados de Blanes (CEAB-CSIC), Accés a la Cala St. Francesc, 14, 17300 Blanes, Spain **11** Estación Biológica de Doñana (EBD-CSIC), Avda. Américo Vespucio 26, 41092 Sevilla, Spain **12** Department of Plant Biology and Ecology, University of Sevilla, 41012 Sevilla, Spain

Corresponding author: Rafael Muñoz-Mas ([rafa.m.mas@gmail.com](mailto:rafa.m.mas@gmail.com))

---

Academic editor: Moritz von der Lippe | Received 3 May 2021 | Accepted 22 October 2021 | Published 14 December 2021

---

**Citation:** Muñoz-Mas R, Carrete M, Castro-Díez P, Delibes-Mateos M, Jaques JA, López-Darias M, Nogales M, Pino J, Traveset A, Turon X, Vilà M, García-Berthou E (2021) Management of invasive alien species in Spain: a bibliometric review. NeoBiota 70: 123–150. <https://doi.org/10.3897/neobiota.70.68202>

---

## Abstract

Scientific and grey literature on invasive alien species (IAS) is conditioned by social, economic and political priorities, editorial preferences and species and ecosystem characteristics. This leads to knowledge gaps and mismatches between scientific research interests and management needs. We reviewed the literature on IAS management in Spain found in Scopus, Web of Science, Google Scholar and Dialnet to identify key deficiencies and priority research areas. The collected literature was classified, employing features

describing formal aspects and content. We used bibliometric and keyword co-occurrence network analyses to assess the relationship between features and reveal the existence of additional topics. Most of the compiled documents ( $n = 388$ ) were focused on terrestrial ecosystems and inland waters, whereas marine and urban ecosystems were under-represented. The literature was largely generic and not species-specific, focusing on raising awareness and proposing changes on current regulation as prominent approaches to prevent further introductions. The compiled authors exhibited many clear publishing preferences (e.g. language or document type), but less regarding target taxa. In addition, there was a strong association between species and the different features considered, especially between the methodological approach (e.g. review, field experiment) and the primary emphasis of study (i.e. basic/theoretical, applied or interdisciplinary). This indicates that research on IAS has had a strong species-specific focus. References about terrestrial species focused mainly on vascular plants, whereas references about inland waters were mostly on fishes and the giant reed (*Arundo donax*), which has been managed with partial success. Animal culling and plant removal were the most frequent eradication and small-scale control treatments, whereas the documents addressing wider spatial scales were largely theoretical. Consequently, the success of described treatments was largely uncertain. Spanish invasion science research has been occasionally innovative, incorporating novel technologies (e.g. species distribution modelling) and engaging society with citizen-science approaches. However, the ratio between basic/theoretical and applied studies indicates that more applied research/management is needed, especially in inland waters and marine ecosystems. We call for increasing effort in the effective dissemination of experience in IAS management to enhance current practical knowledge, including that of schemes undertaken by public agencies.

### Keywords

Biological invasions, eradication, control, Iberian Peninsula, Mediterranean, Spanish archipelagos, keyword co-occurrence analysis, prevention

## Introduction

Biological invasions are human-assisted global phenomena with ravaging effects, not only on biodiversity and ecosystem services, but also on human well-being (McGeoch et al. 2010; Vilà and Hulme 2017). Although transport and introduction of alien species into novel ecosystems is inherent to humankind's expansion (Crees and Turvey 2015), the number of new introductions has increased exponentially since the mid-twentieth century (Seebens et al. 2018, 2019). Invasive alien species (IAS) can reshape ecosystem processes, decrease native species richness and abundance (e.g. McGeoch et al. 2010 and references therein) and cause impact on the economy and human health (Zenni et al. 2021). Thus, preventative, eradication and control actions are required to impede their entry and establishment or minimise their long-term impacts (Robertson et al. 2020).

The incidence of biological invasions and their related costs have led to substantial management efforts worldwide to prevent new introductions and control those already established, by means of eradication or containment (García-de-Lomas and Vilà 2015; Diagne et al. 2021). However, to offer efficient responses is challenging and, regrettably, the specific literature on IAS management is often neglected during the decision-making process (Walsh et al. 2015). In Spain, managers, consultants and assessors face obstacles to find and access suitable references, as a large proportion of papers are written in English, are too scientifically orientated or narrowly focused to be directly

applied or are not open-access (Andreu and Vilà 2007; Mungi et al. 2019; Copp et al. 2021). In addition, many applied studies are scattered amongst the grey literature (often in many different languages), tending to be largely inaccessible to international readers, which further limits the transfer of knowledge on both local and international levels (Haddaway and Bayliss 2015, Jeschke et al. 2019).

In addition to accessibility barriers, the content of scientific literature is also taxonomically and geographically biased (Hulme et al. 2013). This is accentuated by the fact that reviews and positive rather than negative results are more likely to be published (Fanelli 2012). Thus, several characteristics inherent to IAS and recipient ecosystems utterly favour the availability of literature on specific taxa, regions or ecosystems that are easier to study and/or manage (Thomsen et al. 2014; Nghiem et al. 2016; Shackleton et al. 2019). Moreover, trends in invasion science are also affected by social and political priorities, which are, in turn, influenced by communication media and the outcomes of previous research (Gläser and Laudel 2016; Geraldi et al. 2019; Shackleton et al. 2019). This can lead to changes in the importance of research topics over time, for instance through fostering a given species over others or by abandoning certain research areas.

Bibliometric and keyword co-occurrence network analyses are useful to reveal the main knowledge components of any discipline, such as areas with deep insights, outstanding gaps and peripheral research areas (Aria and Cuccurullo 2017, Radhakrishnan et al. 2017). Despite some recent contributions (Enders et al. 2019, 2020), such analyses are scarce in previous reviews of IAS literature, particularly related to management. Consequently, we chose to use them to identify the main patterns in the scientific literature shown by IAS management in Spain. Our specific aims were to: (i) characterise the species, topics and approaches from a management viewpoint, (ii) detect relationships between them and (iii) identify research areas deserving further attention. We focused on Spain due to its diversity of climates and ecosystems, along with its insular and continental territories, which have favoured the establishment of a large and diverse number of IAS (Muñoz-Mas and García-Berthou 2020). Mainland Spain is part of the Mediterranean biodiversity hotspot (Williams et al. 2011) and encompasses an enormous range of ecosystem types, from arid coastal regions to mountain ranges and woodlands. Spain includes two markedly-different populated archipelagos: the Mediterranean Balearic Islands and the Macaronesian Canary Islands off the north-western African coast (Andreu et al. 2009; Benito-Calvo et al. 2009). Islands are in themselves biodiversity hotspots, but they have especially suffered from the establishment of IAS, which have caused numerous extinctions (Lenzner et al. 2020). In addition, Spain has two autonomous cities located on the Mediterranean coast of the African continent, which may require different IAS management approaches. Former reviews on invasion management in the country were sectorial and focused on stakeholder perceptions and management of alien plants (Andreu et al. 2009) or were based on the most common deficiencies in IAS management (Dana et al. 2019). Thus, reviewing past and present experience may help to improve current IAS management actions and identify future research lines in Spain and similar territories. Our results may help funding agencies to target as yet unidentified research needs.

## Methods

### Literature review

We performed our literature search using Scopus, Web of Science (WoS) and Google Scholar. The first two mainly focus on English language literature, largely scientific papers, whereas the references compiled within Google Scholar are more heterogeneous and less structured (Haddaway et al. 2015). Monolingual searches have been shown to limit and bias results in global literature reviews (Angulo et al. 2021; Nuñez and Amano 2021). To avoid such potential bias, in addition to Google Scholar, we also consulted Dialnet. This is a multidisciplinary reference database launched in 2001 that focuses on scientific literature published in Spanish, including books, theses and other documents.

We used the following search terms in Scopus and WoS (both accessed 28 April 2020): (Spain OR Iberian Peninsula OR Canary Islands OR Balearic Islands OR Ceuta OR Melilla) AND (alien species OR exotic species OR non-native species OR invasive species OR introduced species OR species introduction OR translocated species OR species translocation OR species spread OR naturalised species OR casual species OR species of concern OR noxious species OR pernicious species OR harmful species) (Rytwinski et al. 2020). We also used the equivalent Spanish terms for searches in Google Scholar and Dialnet, the latter accessed 30 May 2020. The results were downloaded from Scopus, WoS and Dialnet and duplicates removed (Rytwinski et al. 2020) (Table 1). Following Rytwinski et al. (2020), we modified this general workflow for searches in Google Scholar to deal with the inferior specificity of the searching engine. Between 29 April and 30 May 2020, we retrieved up to 1,000 documents, including duplicates, starting the search using the Spanish keywords describing the toponymy (e.g. España or península Ibérica) and terms used to name IAS (e.g. especie exótica or especie invasora) with one of the following terms: gestión (management), erradicación (eradication) or control. Starting with gestión, we selected documents up to the point when they were clearly irrelevant or duplicated. We then repeated the query using the following keyword (first erradicación and finally control) and repeated the entire process using the English keywords. Once 1,000 documents were compiled, the full text was checked to discard irrelevant documents and, only then, we added the document to the references obtained from Scopus and WoS (Table 1).

Only documents focusing on direct IAS management (i.e. eradication and control) or with explicit management implications in Spain were included (e.g. risk assessment and prevention of future invasions, regulations or education). Studies exclusively focusing on the biology or the ecology of IAS, with no management implications or with no examples, were discarded. We only counted introduced and established species or species able to spread in the wild (e.g. Blackburn et al. 2011), excluding those in captivity or cultivated. We also considered microorganisms and parasites (potentially) affecting other species in the wild (e.g. the crayfish plague *Aphanomyces astaci* or the amphibian chytrid fungus *Batrachochytrium dendrobatidis*), but excluding agricultural pests or pathogens of captive animals or humans.

In general, no agriculture-related studies were retained, nor weeds interfering with crops or plantations. However, studies on the red weevil, *Rhynchophorus ferrugineus*, were not excluded, although they focused on the date palm *Phoenix dactylifera*, because this weevil also affects the endemic Canary Islands date palm *Phoenix canariensis*. We compiled a total of 388 documents (Table 1).

## Literature characterisation and keyword compilation

We selected relevant features of the retrieved documents for our analysis, following previous bibliographic studies on IAS management (mostly Bayliss et al. 2013; Matzek et al. 2014) (Table 2). The selected features described formal aspects (e.g.

**Table 1.** Total number of references retrieved before scrutiny (e.g. including duplicates) and final number of references retained for the database. Web of Science and Scopus were managed simultaneously because they show a great overlap of hosted references.

Bibliographic database	Number of references retrieved	Number of references retained
Scopus	1569	214
Web of Science	1152	
Dialnet	75	62
Google Scholar	1000	112

**Table 2.** Features and categories used to characterise the compiled literature (based on Andreu and Vilà 2007; Bayliss et al. 2013; Matzek et al. 2014). Specific definitions for each category of the selected features are provided in Supplementary Material (Table S1).

Feature	Categories
Authors	-
Document type	Scientific article, book, book chapter, conference proceedings, report or thesis
Ecological organisation level	Population/autoecology, community/assemblage, ecosystem or generic/not applicable
Ecosystem type	Terrestrial, inland waters, marine or urban
Insular territory	Yes or no
Language	English or Spanish
Main topic	Management (eradication and control), prediction, prevention, prioritisation (including risk management), regulation/decision-making, risk assessment, social, biology/ecology, climate change and other synergisms, conservation, definitions, economics, education and awareness-raising, impacts, information sharing, communication and collaboration, introduction/spread or survey/monitoring
Management topic	Prevention, eradication, control or generic/not applicable
Methodological approach	Review/meta-analysis, observational field study, field experiment, greenhouse/laboratory experiment, opinion paper or theory
Primary emphasis	Basic (without direct manipulation of the target taxa), applied (with direct manipulation of the target taxa) or interdisciplinary (encompassing social and policy issues)
Protected area	Yes or no
Spatial scale	Local, regional, national, supranational/global or other/undefined (e.g. greenhouse/lab experiment)
Species	Target species, taxon or generic/multiple taxa
Success	Yes, partial, no, unknown or not applicable
Treatment	Regulation, awareness raising, culling, biocontrol agent, poisoning, hydrology manipulation, hydrogeomorphological restoration, survey and monitoring, mechanical removal, manual removal, herbicide, genetic selection, enclosure, not applicable, DNA metabarcoding, re-afforestation, heating, baiting, containment, mulching, prevention, restocking, incineration, insecticide or sterilisation
Year	-

document type or text language) and content (e.g. methodological approach, main topic, spatial scale). To better characterise documents with multiple topics and sections, the categories within the different features were non-exclusive. That is, a document could focus on the biology/ecology, introduction/spread and survey/monitoring of IAS or on prevention, eradication and control. However, during subsequent analyses, we downweighed each reference in the frequency-related calculations to sum up to one and ensure the equal contribution of all documents (Muñoz-Mas and García-Berthou 2020).

We downloaded the keywords from documents retrieved from Scopus and WoS and manually scrutinised those documents compiled from Google Scholar and Dialnet to compile the available keywords. Then, we inspected the keywords to detect mistakes and misspellings and translated those words into Spanish using Google Translator, adjusting English keywords to well-established terms when necessary (e.g. caña común/common reed to giant reed). The references and features can be found in Supplementary Material.

## Data analyses

We used the cumulative sum of number of documents per year to compare the publishing trends in management of invasive alien species in Spain with more general science publishing trends (in Spain and worldwide). The total number of documents published worldwide and in Spain were those included in the Journal Citation Reports (JCRs). The series were retrieved from: [www.scimagojr.com/countryrank.php](http://www.scimagojr.com/countryrank.php). Frequency bar plots enabled scrutiny of the prevalence of the different categories of each additional feature, except species and ecosystem types. We investigated these two features simultaneously using the function `comparison.cloud` of the R (R Core Team 2021) package `wordcloud` (Fellows 2018), but without graphical scrutiny for language, insular territory or protected area, due to their binary nature.

The association amongst features was analysed, except year (Table 2), employing Cramér's V Index (Cramér 1946), implemented in the R package `oii` (Hale et al. 2017). This Index ranges from 0 (no association) to 1 (perfect association) and is based on a corrected  $\chi^2$  statistic. To graphically describe the association patterns, we built a network using the R package `igraph` (Csardi and Nepusz 2006), employing the values of the Index only when statistically significant ( $P < 0.05$ ).

Cramér's V Index evaluates the association between features, but provides no information about the relationship amongst categories. Therefore, we built two alluvial diagrams to graphically scrutinise the relationship between the categories of the features: methodological approach, ecosystem type, management topic and spatial scale and ecosystem type, treatment and success (Table 2). Alluvial diagrams are a kind of Sankey diagram that group together observations of the same category and visualise them as flows across the considered set of features (Rosvall and Bergstrom 2010). We used the function `SankeyDiagram` in the R package `flipPlots` (Displayr 2019) to build the alluvial diagram.

To investigate the existence of additional topics and research areas not described by the features and categories detailed in Table 2, we analysed the literature using a


keyword co-occurrence network (Radhakrishnan et al. 2017). As customary, we first systematically lemmatised/stemmed the resulting keywords in R (i.e. inflected or derived words were reduced to their root form) to reduce the variability within the collected keywords. For example, by applying this procedure, the word *biolog* would result from the words *biological* and *biology*. For this, we used the function *wordStem* of the R package *SnowballC* (Bouchet-Valat 2020) to allow reproducible results. Compound keywords were split (e.g. *invasive species* was divided into *invasive* and *species*), each word was lemmatised independently (e.g. *invas* and *speci*) and the resulting words were reassembled (e.g. *invas speci*) to build the co-occurrence network. The co-occurrence matrix cross-product was obtained via the function *cocMatrix* of the R package *bibliometrix* (Aria and Cuccurullo 2017) and we built the co-occurrence network with *igraph* (Csardi and Nepusz 2006). We delineated the relevant research areas or clusters employing the function *cluster\_edge\_betweenness* (Newman and Girvan 2004) in *igraph* (Csardi and Nepusz 2006). Finally, the most frequent keyword of each cluster was used to simplify the complete network into a smaller network and facilitate its interpretation. To avoid oversimplification of the network, we kept the most frequent keywords ( $> Q_{95}$  or number of occurrences  $\geq 4$  occurrences), while the less frequent keywords were collapsed to the most frequent keyword in the corresponding research area (i.e. the cluster centre). We depicted the resulting network with the most important keywords in each research area (i.e. cluster centres encompassing the most frequent and less frequent keywords into single vertices and additionally those keywords whose frequency of occurrence was  $\geq 4$ ) as two-level circular treemaps (Zhao and Lu 2015) using the R package *ggraph* (Pedersen 2021).

## Results

The number of documents published on IAS management has grown steadily since 1995 (Figure 1A). However, those published in Spain are under-represented compared to the scientific production trends both globally and in Spain, although during the last decade, the scientific production accelerated significantly. Most documents were scientific articles (72.4%) (Figure 1B). Review/meta-analysis was the most common methodological approach (54.9% out of the 388 documents), followed by field experiment (19.0%) and observational field study (18.9%) (Figure C). The total number of authors was 1,280. The most prolific author was Montserrat Vilà (13 documents/3.4%), followed by Pilar Castro-Díez, Elías D Dana and Juan García de Lomas (7 documents each/1.8%), whereas 1,067 authors appeared in one single document (Figure 1D). Most documents were written in English (59.3%) and the remainder in Spanish.

Management (i.e. eradication and control) was the most frequent topic (31.8% out of 388 documents), followed by risk assessment (9.6%), survey/monitoring (9.3%), biology/ecology (7.6%), and impacts (7.4%) (Figure 2A). Studies focusing on populations and the autoecology of a single species were the most frequent (45.2%), followed by studies at the community/assembly level (24.0%) (Figure 2B). The dominant






**Figure 2.** Numbers of documents by main topics are represented in (A), ecological organisation levels in (B), management topics in (C), spatial scales in (D), primary emphasis in (E), treatment in (F) and success in (G).

sustained management actions to control the target IAS (Figure 2G). About 13% of the studies focused exclusively on insular territories and 17% on protected areas.


The compiled studies involved 159 species or higher taxa (e.g. vascular plants). The largest number dealt with terrestrial ecosystems (50.4%), followed by inland waters (36.2%), whereas marine ecosystems and urban environments were the least common (8.3% and 5.0%, respectively) (Figure 3). Terrestrial vascular plants (10.4%) were the most frequent, especially eucalypts (*Eucalyptus* spp.) (1.6%), prickly pear species (*Opuntia* spp.) (1.3%), Hottentot figs (*Carpobrotus* spp.) (1.3%), Monterey pine (*Pinus radiata*) (1.3%) and cordgrasses (*Spartina* spp.) (1.0%). Amongst terrestrial animals, the most frequent species were the yellow-legged hornet (*Vespa velutina*) (1.6%), the pinewood nematode *Bursaphelenchus xylophilus* (1.3%), American mink *Neovison vison* (1.0%) and generic studies on vertebrates (3.3%), mainly birds (1.3%). In inland waters, generic studies were also the most common (7.5%), followed by those on fish management (7.0%). The most studied species in inland waters were the giant reed (*Arundo donax*) (3.1%) and the red swamp crayfish (*Procambarus clarki*) (1.8%), followed by generic studies on riparian vegetation (1.6%). Most studies on marine






**Figure 4.** Association network displaying Cramér's V between features characterising the documents on IAS management in Spain. Connections are only depicted when the  $\chi^2$  statistic is significant ( $P < 0.05$ ). Connection width and colour are based on Cramér's V and range from 0 (no association) to 1 (perfect association).

The most frequent studies were reviews and meta-analyses focusing on terrestrial ecosystems and providing general guidelines to manage invasive species at the national (6.5% out of the 388 documents), supranational/global (5.3%) and regional scales (4.3%) (Figure 5). They were followed by field experiments (manipulative) on terrestrial ecosystems addressed to locally eradicate specific taxa (3.3%). Reviews and meta-analyses on inland waters and those providing general guidelines to manage IAS at higher scales (i.e. regional to supranational/global) were also frequent (3.2% and 2.8%, respectively). Field experiments (manipulative) in inland waters addressed to eradicate (3.0%) or control (2.8%) specific taxa were in all cases local. The most frequent observational field studies on both terrestrial ecosystems (2.3%) and inland waters (2.0%) addressed generic aspects and were local. In contrast, documents focusing on marine ecosystems were scarce and studied generic aspects of marine invasion science with supranational/global (1.0%) or local (0.9%) perspectives. Manipulative field experiments to locally eradicate marine species were markedly rare (0.3%). Studies in urban environments were mostly reviews and meta-analyses addressing generic aspects at both supranational/global and local scales were the most abundant (0.5% and 0.5%, respectively). The proportion of field manipulative experiments to eradicate or control species was negligible.




**Figure 5.** Alluvial diagram relating the methodological approach, ecosystem type, management topic and spatial scale of the compiled documents on invasive alien species (IAS) management in Spain. Connection width is proportional to the number of documents (An interactive version of this figure can be downloaded from <https://doi.org/10.6084/m9.figshare.16547790.v1>).

The most frequent approaches, described within terrestrial studies, focused on awareness-raising, survey/monitoring and regulation (7.3%, 6.8% and 3.1%, respectively out of the 388 documents). Most of them had no quantifiable (not applicable/17.2%) or unknown success (2.1%). This was followed by the use of biocontrol agents whose efficacy has not yet been tested (i.e. unknown, 1.6%) and unsuccessful culling of IAS (none and partial, 1.7%). The successful approaches most frequently reported combined herbicides and mechanical and manual plant removal (2.9% in total). In inland waters, awareness-raising and regulation were amongst the most frequently indicated approaches (3.9% and 3.0%, respectively), although with no quantifiable success (not applicable, 6.9%). Water level and flow regime manipulation was the most frequent management approach, but it was not tested (unknown, 2.3%) and occasionally turned out useless (0.7%). Culling and poisoning seldom worked (0.8% and 0.5%, respectively) and the success of most reported experiences was unknown or partial (1.6% and 0.9%, respectively). Hydrogeomorphological restoration, re-afforestation, plant removal and herbicide use were the most common approaches to control invasive riparian vegetation (1.2%, 1.1%, 1.6% and 0.7%, respectively). However, success of these treatments was partial or uncertain and only 0.6% reported successful experiences. The treatments for marine ecosystems followed a similar pattern and focused on preventative approaches: awareness-raising, survey/monitoring and regulation (1.9%, 1.0% and 0.9%, respectively), most of them with no quantifiable (not applicable/2.0%) or unknown success (1.6%). The only successful study involved raising awareness and DNA metabarcoding to confirm the elimination of the pygmy mussel (*Xenostrobus securis*) (Miralles et al. 2016). Awareness-raising, regulation and survey/monitoring were the most common approaches for urban environments (3.7% in total), but the few applied experiences indicated partial success through culling and plant removal and subsequent herbicide spraying (0.3%).



**Figure 6.** Alluvial diagram relating the ecosystem type, treatment and success of the compiled documents on invasive alien species (IAS) management in Spain. Connection width is proportional to the number of documents. The category Other includes mulching, prevention, sterilisation, heating, incineration, containment, baiting, restocking, DNA metabarcoding, insecticide and genetic selection (Alternative static and interactive versions of this figure relating species/taxa, treatment and success can be downloaded from <https://doi.org/10.6084/m9.figshare.16547790.v1>).

The collected keywords encompassed 1,145 different terms. The aggregation algorithm revealed 39 different research areas or clusters (Figure 7A), with 67 keywords occurring on  $\geq 4$  occasions (Figure 7B). The largest research area highlighted the importance of invasive plants amongst the Spanish literature on IAS management, the Mediterranean nature of much of the territory and the numerous studies carried out on this taxon in the Balearic Islands. The simplified network reflected the main topics of the study: invasive species and management, eradication and control. It also reflected research carried out on specific taxa, such as on the elimination of American mink (*Neovison vison*), which clustered with invasive species. Exclusion experiments on the European rabbit (*Oryctolagus cuniculus*) and feral cats (*Felis silvestris catus*) appeared in different research areas. The studies on feral cats were undertaken mainly in the Canary Islands as their research areas were connected. In addition, the resulting network highlighted the impacts on freshwater biodiversity caused by giant reed (*A. donax*) and aquaculture activities. The simplified network reflected the importance of the yellow-legged hornet (*V. velutina*) and mosquitoes (mainly the tiger mosquito *A. albopictus*) and the extensive use of species distribution models (SDMs), such as MaxEnt (Phillips et al. 2004), to foresee expansion trends and suitable regions. Specific topics (propagule pressure and ecosystem services) and introduction vectors (ballast waters, aquaculture and hunting) appeared in separate research areas. The simplified network reflected specific management approaches, such as general forest management, use of herbicides or emergence of citizen science. It also reflected studies addressing the interaction be-



**Figure 7.** (A) Complete keyword co-occurrence network developed to visualise the importance of the research areas. Vertex colours are based on research areas or clusters and vertex size is proportional to the frequency of keyword occurrence. (B) Simplified network based on cluster centres and most frequent keywords ( $> Q_{95}$  or number of occurrences  $\geq 4$ ). Overall vertex sizes are proportional to the log-transformed number of occurrences and inner circles to the proportion within each research area. Label sizes have been rescaled to avoid overlapping.

tween invasions and climate change. In addition to the main toponymy used during the bibliography search, the network reflected specific regions and environments, such as the Strait of Gibraltar and Galicia (NW Spain) and the importance of wetlands and salt marshes. Moreover, the simplified network highlighted the importance of the archipelagos and islands within Spanish invasion science research, given that island and Canary Islands appear in differentiated research areas. It also revealed the multiplicity of terms used to name similar concepts, such as the terms alien species and exotic species that appear scattered throughout different research areas.

## Discussion

Spanish literature on IAS management has been mainly theoretical (55% review/meta-analysis), with a balance between theoretical and applied studies similar to that reported in other studies on biological and ecological aspects of biological invasions (Andreu and Vilà 2007; Bayliss et al. 2013; Matzek et al. 2014, 2015). Nonetheless, IAS man-

agement literature has been under-represented compared to the Spanish and global trends on scientific production, although recently its share has grown. Altogether, this suggests that Spanish invasion science may also suffer from a knowing-doing gap caused by a preponderance of theoretical studies (Matzek et al. 2014, 2015). Moreover, scientific papers written in English constituted a large proportion of the compiled literature, which suggests that most of it may be too scientifically orientated to be directly applied (Andreu and Vilà 2007; Mungi et al. 2019; Copp et al. 2021). Nevertheless, scientific activity in Spain has also incorporated new forecasting technologies (e.g. SDMs/MaxEnt, de Medeiros et al. 2018) along with new approaches to engage society in IAS control (e.g. citizen science, Clusa et al. 2018) and for biomonitoring (e.g. DNA metabarcoding, Borrell et al. 2017). In addition, risk assessments and horizon scanning studies for decision-making formed a notable part of the literature, with direct implications for IAS regulation (Bayón and Vilà 2019). There was also discussion regarding the inefficiency of current codes of conduct and laws (Maceda-Veiga et al. 2013), aimed at enforcing policies to overcome the highlighted deficiencies.

The proportion of applied studies and field experiments was markedly low, which impeded rating the efficacy of most of the described approaches (66.2%). Moreover, studies on novel biocontrol agents, such as the use of pathogens (McColl and Sunarto 2020) were infrequent. In part, this is because these agents require rigorous risk assessment studies and are, hence, subject to strict regulations (Loomans 2021). Thus, species eradication and control experiences often relied on the use of herbicides, mechanical elimination or culling (e.g. Melero et al. 2010; Mateos-Naranjo et al. 2012) and successful approaches were restricted to small-scale areas, such as islets and ponds (e.g. Ferreras-Romero et al. 2016; Maceda-Veiga et al. 2017). Consequently, concerns of Spanish managers about the problem that too much research focuses generally on the ecological aspects of alien plants, rather than on specific cost-efficient management strategies (Andreu et al. 2009), can be considered, to some extent, applicable to most invasive alien taxa.

The proportion of documents per species and ecosystem type in Spain was similar to that estimated in other countries (Thomsen et al. 2014). Terrestrial species, mainly plants, attracted the bulk of the literature, followed by studies involving species of inland waters. By contrast, marine and urban environments were infrequent in the collected literature. The preponderance of terrestrial ecosystems and the associated species can be justified by the primary introduction pathways of terrestrial species (i.e. release and/or escape), which are largely related to forestry, livestock, agriculture and wildlife trade (Essl et al. 2015). Vascular plants are the most frequently introduced taxon worldwide; consequently, they were expected to receive the largest proportion of studies. However, the feasibility of managing terrestrial invaders or aliens, especially sessile species and stages of their life history (e.g. nests, Enríquez et al. 2013), is greater compared to aquatic species, due to the lower accessibility of these environments. Therefore, a publication bias towards successful studies with positive results on tractable terrestrial species is not discernible (Fanelli 2012; Booy et al. 2017).

Studies on terrestrial ecosystems involved all spatial scales (i.e. local, regional, national and supranational). However, applied experiments and experience were mostly

local interventions, such as management of prickly pear species (*Opuntia* spp.) and the sentry plant (*Agave americana*) (Arévalo et al. 2015). By contrast, eradication and control of vagile terrestrial organisms proved to be economically unaffordable (e.g. American mink *N. vison*, Melero et al. 2010; Mañas et al. 2016), especially in a context of multilevel overlapping or competing public administrations with ill-defined jurisdictions and pervasive budget shortages (Tollington et al. 2017; Dana et al. 2019). In this regard, island territories were well covered by the collected literature, with several successful management experiences in these territories, such as the eradication of American mink (*N. vison*) in the Atlantic Islands National Park (Velando et al. 2017) or of the red palm weevil (*R. ferrugineus*) from the Canary Islands (Fajardo et al. 2019). Impacts of IAS on islands are likely to increase in the future, especially on oceanic islands, such as the Canaries and, to a lesser extent, on the continental Balearic Archipelago. Indeed, insular terrestrial ecosystems are generally the most threatened (Lenzner et al. 2020). Therefore, these two successful examples shed hope on the future management of IAS in Spanish insular territories.

The published research on terrestrial invertebrates and microorganisms appeared to be concentrated on a few species with direct impacts on economics and human health (e.g. yellow-legged hornet *V. velutina*, pinewood nematode *B. xylophilus* or tiger mosquito *A. albopictus*). However, further applied research and knowledge transfer is particularly needed to control invertebrates, due to the rising number of introductions worldwide (Saul et al. 2017; Seebens 2019). Such a task often requires strategies based on prevention and prompt eradication (e.g. ballast water sanitation) (Booy et al. 2017, 2020), but these were the least frequent of the categories amongst those implying direct manipulation of the target taxa. From a theoretical viewpoint, the compiled literature raises awareness and proposes changes to current regulations as its main approach, directed at preventing further introductions at a higher level of organisation. Legislation regarding IAS has become more restrictive over the years worldwide and Spain is also immersed in this useful trend (Turbelin et al. 2017; Maceda-Veiga et al. 2019). However, in light of the number of recent introductions (Muñoz-Mas and García-Berthou 2020), it can be concluded that the real effective capability of Spain to impede the establishment of further species is limited. This general pattern is shared with other European countries and is unlikely to change substantially in the near future (Seebens et al. 2021). Tackling the establishment of further terrestrial invertebrates and microorganisms will require further and stronger innovative and well-funded preventative approaches.

The number of studies conducted in inland waters was notably high due to the enormous number of established species and their associated economic costs (e.g. Durán et al. 2012; Muñoz-Mas and García-Berthou 2020). Some of this research was promoted by the Water Framework Directive (European Parliament & Council 2000), even if IAS are not explicitly mentioned therein (Boon et al. 2020). Our keyword co-occurrence network reflected the numerous studies recommending flow management as a way to control IAS (Sabater et al. 2008; Fornaroli et al. 2020). However, despite the schemes conducted in other countries (Kiernan et al. 2012), no applied examples

in large and intermediate regulated river systems were found in the literature. River basin management plans increasingly account for the presence of IAS, but more emphasis on applied management of medium-to-large river systems is necessary (Boon et al. 2020). Nonetheless, applied studies of inland water ecosystems dealt with control of the giant reed *A. donax* (the most managed species appearing in the compiled literature) in relatively small areas (Bruno et al. 2019) or described experience in relatively small lentic environments (i.e. common carp *Cyprinus carpio* in ponds and lakes; Ferreras-Romero et al. 2016) and small streams (signal crayfish *Pacifastacus leniusculus*; Dana et al. 2010). Unfortunately, more research is needed to optimise water allocation schemes because climate change is facilitating the establishment of further IAS. Meanwhile, the increased demands of agriculture will reduce the availability of water resources to undertake the aforementioned actions (Rahel and Olden 2008; Escribano Francés et al. 2017).

In Spain, stowaway introductions in brackish and marine environments have also gained prominence (García-Gómez et al. 2020; Painting et al. 2020), causing a shift in the type of introduced species that can be framed within the current increase in global maritime traffic (Saul et al. 2017; Seebens 2019). In this regard, Spain enacted in 2004 the International Convention for the Control and Management of Ships' Ballast Water and Sediments (BWM Convention), which has been addressed in local studies (e.g. Moreno-Andrés et al. 2017), dissertations (e.g. Bartolomé Lamarca 2014) and is highlighted in the keyword co-occurrence network. However, new IAS records are being reported frequently. For example, the gastropod *Mitrella psilla* was recently found thriving on western coasts of Spain (Martínez-Ortí et al. 2020). The effectiveness of measures to fulfil the BWM Convention remains limited, which underscores the difficulties faced in managing marine IAS (Thomsen et al. 2014; Cuesta et al. 2016). Indeed, the number of first records whose most probable introduction pathways are ballast waters or biofouling (Davidson et al. 2018) and the increasing importance of aquaculture and related introductions (Nunes et al. 2015; Garlock et al. 2020) suggest these introduction pathways must be taken into account. It can, therefore, be concluded that marine invasion science should move towards our central focus in the future.

Worldwide, urban environments are becoming active introduction hubs (Gaertner et al. 2017). Indeed, there are several examples of initial IAS establishment in urban zones that have spread outwards over natural environments in Spain, such as *Lippia filiformis* (Casasayas i Fornell 1989) or the black-headed weaver *Ploceus melanocephalus* (Grundy et al. 2014). Ornamental plants and alien exotic birds kept as pets are perhaps the most striking and troublesome introductions in city surroundings (Riera et al. 2021), but other less-known taxa have been found on numerous occasions (e.g. *Pseudosuccinea columella* (Mollusca) Martínez-Ortí 2013). Besides the introduction of ornamental plants and tortoises and terrapins (*Trachemis* spp.), which have already spread over natural environments and prompted specific studies (Muñoz-Mas and García-Berthou 2020), the majority of Spanish management literature on urban environments has focused on birds (e.g. the monk parakeet *Myiopsitta monachus*

and the rose-ringed parakeet *Psittacula krameri* (Álvarez-Pola and Muntaner 2009; Maceda-Veiga et al. 2019; Hernández-Brito et al. 2020; Saavedra and Medina 2020). Therefore, in the light of increasing urbanisation of the Spanish population and the relatively low number of specific studies, we conclude that prevention protocols and further studies, specifically addressed to urban environments, should be strengthened to encompass the full spectrum of potential introductions.

The compiled literature on IAS management in Spain does not particularly reflect budget reductions related to the Great Recession of 2008 (Catanzaro 2018). However, IAS management literature has been under-represented compared to overall scientific production trends and the current rise in publication rates has not compensated for this historical delay. Nonetheless, our compiled references only represent a small fraction of the total number of reviewed documents ( $388/3796 = 10.22\%$ ). Moreover, a significant number of documents were written in Spanish ( $n = 158; 40.7\%$ ). This finding deserves special attention as it handicaps knowledge transfer (Di Bitetti and Ferreras 2017) and biases conclusions inferred by international agents (Konno et al. 2020). However, it is not problematic from a national point of view because texts, documents and software packages assisting the process of decision-making by administrators and functionaries have proven to be most efficient when presented in local languages (Copp et al. 2021). Nevertheless, despite the proliferation of public repositories and open access publications, a wealth of information is still hidden away, not easily accessible to risk assessors, managers and researchers through standard search engines. For example, it is known that more than a hundred plant species have been managed by Spanish regional administrations (Andreu and Vilà 2007), but only the most frequent species generated accessible documents (e.g. reports). Altogether, it indicates that public agencies produce insufficient literature (*sensu lato*) as they focus on other tasks. Neither communication between managers and scientists beyond undertaking management action schemes nor protocols to evaluate their success are common practices in Spain. It would be beneficial for public agencies to encourage and facilitate such interguild contact, perhaps using legislative and labour changes, to disseminate applied experience in accessible ways.

Despite these recommendations and the highlighted deficiencies, Spanish literature on IAS management should not be considered completely defective. Recent studies on alien animal species, currently thriving in Spanish inland waters, indicate that no single management protocol can be applied to every taxonomic group, due to marked differences amongst species, introduction pathways and invaded habitats (Muñoz-Mas and García-Berthou 2020). Likewise, our study shows a strong association between species, taxon or group of taxa and the features/categories used to describe the compiled literature. This indicates that species-specific studies are often needed, which highlights how difficult and complex the task of IAS management is (Woodford et al. 2016; Portela et al. 2020; Yelenik et al. 2020). Our results should help to properly drive future research efforts towards IAS management in Spain. We recommend more research into applied techniques to shift the balance between theoretical and empirical studies, especially in inland waters and marine ecosystems due to their lower accessibility. The same need for

more studies applies to urban environments, as they are often the bridgehead of IAS introductions. Renewed effort in prevention and prompt eradication should be made to fulfil, for example, the BWM Convention and impede further introductions into marine ecosystems. Finally, we encourage public agencies to support and strengthen the dissemination of applied experience and thus enhance know-how and knowledge transfer in the field.

## Acknowledgements

We thank Dr. Núria Roura-Pascual, an anonymous reviewer and the Editor Dr. Moritz von der Lippe for their helpful comments about the manuscript.

This research was funded by the Spanish Ministry of Science and Innovation (InvaNET network, RED2018-102571-T). Additional financial support was provided by the Spanish Ministry of Science and Innovation, Spanish State Research Agency (AEI) and European Regional Development Fund (FEDER, UE) (grants PID2020-118550RB, PID2019-103936GB-C21, RTI2018-093504-B-I00) and the Government of Catalonia (ref. 2017 SGR 548). RMM benefitted from a postdoctoral Juan de la Cierva Fellowship from the Spanish Ministry of Science and Innovation (FJCI-2016-30829).

## References

- Álvarez-Pola C, Muntaner J (2009) Control de aves invasoras en las islas Baleares. *Anuari Ornitològic de les Balears: Revista d'observació estudi i conservació dels aucells* 2009: 67–71. <https://www.raco.cat/index.php/AnuariOrnitologic/article/download/244672/327694>.
- Andreu J, Vilà M (2007) Análisis de la gestión de las plantas exóticas en los espacios naturales españoles. *Ecosistemas* 16: 109–124. <https://www.revistaecosistemas.net/index.php/ecosistemas/article/view/118> (May 30, 2020).
- Andreu J, Vilà M, Hulme PE (2009) An assessment of stakeholder perceptions and management of noxious alien plants in Spain. *Environmental Management* 43: 1244–1255. <https://doi.org/10.1007/s00267-009-9280-1>
- Angulo E, Digne C, Ballesteros-Mejía L, Adamjy T, Ahmed DA, Akulov E, Banerjee AK, Capinha C, Dia CAKM, Dobigny G, Duboscq-Carra VG, Golivets M, Haubrock PJ, Heringer G, Kirichenko N, Kourantidou M, Liu C, Nuñez MA, Renault D, Roiz D, Taheri A, Verbrugge LNH, Watari Y, Xiong W, Courchamp F (2021) Non-English languages enrich scientific knowledge: The example of economic costs of biological invasions. *Science of The Total Environment* 775: 144441. <https://doi.org/10.1016/j.scitotenv.2020.144441>
- Arévalo JR, Fernández-Lugo S, Mellado M, de la Concepción T (2015) Experimental management control of *Opuntia dillenii* Haw. and *Agave americana* L. in Teno Rural Park, Canary Islands. *Plant Species Biology* 30: 137–146. <https://doi.org/10.1111/1442-1984.12049>

- Aria M, Cuccurullo C (2017) bibliometrix: An R-tool for comprehensive science mapping analysis. *Journal of Informetrics* 11: 959–975. <https://doi.org/10.1016/j.joi.2017.08.007>
- Bartolomé Lamarca I (2014) Sistemas de gestión de agua del lastre: Fundamentos jurídicos y esquemas operativos. Universitat Politècnica de Catalunya. <http://hdl.handle.net/2099.1/24626>
- Bayliss HR, Stewart G, Wilcox A, Randall N (2013) A perceived gap between invasive species research and stakeholder priorities. *NeoBiota* 19: 67–82. <https://doi.org/10.3897/neobiota.19.4897>
- Bayón A, Vilà M (2019) Horizon scanning to identify invasion risk of ornamental plants marketed in Spain. *NeoBiota* 52: 47–86. <https://doi.org/10.3897/neobiota.52.38113>
- Benito-Calvo A, Pérez-González A, Magri O, Meza P (2009) Assessing regional geodiversity: The Iberian Peninsula. *Earth Surface Processes and Landforms* 34: 1433–1445. <https://doi.org/10.1002/esp.1840>
- Di Bitetti MS, Ferreras JA (2017) Publish (in English) or perish: The effect on citation rate of using languages other than English in scientific publications. *Ambio* 46: 121–127. <https://doi.org/10.1007/s13280-016-0820-7>
- Blackburn T, Pyšek P, Bacher S, Carlton J, Duncan R, Jarošík V, Wilson J, Richardson D (2011) A proposed unified framework for biological invasions. *Trends in Ecology and Evolution* 26: 333–339. <https://doi.org/10.1016/j.tree.2011.03.023>
- Boon PJ, Clarke SA, Copp GH (2020) Alien species and the EU Water Framework Directive: A comparative assessment of European approaches. *Biological Invasions* 22: 1497–1512. <https://doi.org/10.1007/s10530-020-02201-z>
- Booy O, Mill AC, Roy HE, Hiley A, Moore N, Robertson P, Baker S, Brazier M, Bue M, Bullcock R, Campbell S, Eyre D, Foster J, Hatton-Ellis M, Long J, Macadam C, Morrison-Bell C, Mumford J, Newman J, Parrott D, Payne R, Renals T, Rodgers E, Spencer M, Stebbing P, Sutton-Croft M, Walker KJ, Ward A, Whittaker S, Wyn G (2017) Risk management to prioritise the eradication of new and emerging invasive non-native species. *Biological Invasions* 19: 2401–2417. <https://doi.org/10.1007/s10530-017-1451-z>
- Booy O, Robertson PA, Moore N, Ward J, Roy HE, Adriaens T, Shaw R, Valkenburg J, Wyn G, Bertolino S, Blight O, Branquart E, Brundu G, Caffrey J, Capizzi D, Casaer J, De Clerck O, Coughlan NE, Davis E, Dick JTA, Essl F, Fried G, Genovesi P, González-Moreno P, Huysentruyt F, Jenkins SR, Kerckhof F, Lucy FE, Nentwig W, Newman J, Rabitsch W, Roy S, Starfinger U, Stebbing PD, Stuyck J, Sutton-Croft M, Tricarico E, Vanderhoeven S, Verreycken H, Mill AC (2020) Using structured eradication feasibility assessment to prioritise the management of new and emerging invasive alien species in Europe. *Global Change Biology* 26: 6235–6250. <https://doi.org/10.1111/gcb.15280>
- Borrell YJ, Miralles L, Do Huu H, Mohammed-Geba K, Garcia-Vazquez E (2017) DNA in a bottle – Rapid metabarcoding survey for early alerts of invasive species in ports. *PLoS ONE* 12: e0183347. <https://doi.org/10.1371/journal.pone.0183347>
- Bouchet-Valat M (2020) SnowballC: Snowball stemmers based on the C “libstemmer” UTF-8 Library. <https://cran.r-project.org/package=SnowballC>
- Bruno D, Zapata V, Guareschi S, Picazo F, Dettori E, Carbonell JA, Millán A, Velasco J, Robledano F (2019) Short-term responses of aquatic and terrestrial biodiversity to riparian restoration measures designed to control the invasive *Arundo donax* L. *Water (Switzerland)* 11: 2551. <https://doi.org/10.3390/w11122551>

- Casasayas i Fornell T (1989) La flora al·lòctona de Catalunya. Catàleg raonat de les plantes vasculares exòtiques que creixen sense cultiu al NE de la Península Ibèrica. Universitat de Barcelona. <http://hdl.handle.net/10803/969>.
- Catanzaro M (2018) Spain's biggest-ever science petition decries "abandonment" of research. *Nature* 556: 285. <https://doi.org/10.1038/d41586-018-04523-4>
- Clusa L, Miralles L, Fernández S, García-Vázquez E, Dopico E (2018) Public knowledge of alien species: A case study on aquatic biodiversity in North Iberian rivers. *Journal for Nature Conservation* 42: 53–61. <https://doi.org/10.1016/j.jnc.2018.01.001>
- Copp GH, Vilizzi L, Wei H, Li S, Piria M, Al-Faisal AJ, Almeida D, Atique U, Al-Wazzan Z, Bakiu R, Bašić T, Bui TD, Canning-Clode J, Castro N, Chaichana R, Çoker T, Dashinov D, Ekmekçi FG, Erős T, Ferincz Á, Ferreira T, Giannetto D, Gilles AS, Głowacki Ł, Gouilletquer P, Interesova E, Iqbal S, Jakubčinová K, Kanongdate K, Kim JE, Kopecký O, Kostov V, Koutsikos N, Kozic S, Kristan P, Kurita Y, Lee HG, Leuven RSEW, Lipinskaya T, Lukas J, Marchini A, González Martínez AI, Masson L, Memedemin D, Moghaddas SD, Monteiro J, Mumladze L, Naddafi R, Năvodaru I, Olsson KH, Onikura N, Paganelli D, Pavia RT, Perdikaris C, Pickholtz R, Pietraszewski D, Povž M, Preda C, Ristovska M, Rosíková K, Santos JM, Semenchenko V, Senanan W, Simonović P, Smeti E, Števoe B, Švolíková K, Ta KAT, Tarkan AS, Top N, Tricarico E, Uzunova E, Vardakas L, Verreycken H, Zięba G, Mendoza R (2021) Speaking their language – Development of a multilingual decision-support tool for communicating invasive species risks to decision makers and stakeholders. *Environmental Modelling and Software* 135: 104900. <https://doi.org/10.1016/j.envsoft.2020.104900>
- Cramér H (1946) *Methods of mathematical statistics*. Princeton University Press., Princeton, NJ (USA), 500 pp.
- Crees JJ, Turvey ST (2015) What constitutes a "native" species? Insights from the Quaternary faunal record. *Biological Conservation* 186: 143–148. <https://doi.org/10.1016/j.biocon.2015.03.007>
- Csardi G, Nepusz T (2006) The igraph software package for complex network research. *InterJournal Complex Systems* 1695: 1–9. <http://igraph.org>.
- Cuesta JA, Almón B, Pérez-Dieste J, Trigo JE, Bañón R (2016) Role of ships' hull fouling and tropicalization process on European carcinofauna: New records in Galician waters (NW Spain). *Biological Invasions* 18: 619–630. <https://doi.org/10.1007/s10530-015-1034-9>
- Dana ED, García-de-Lomas J, Verloove F, Vilà M (2019) Common deficiencies of actions for managing invasive alien species: A decision-support checklist. *NeoBiota*: 97–112. <https://doi.org/10.3897/neobiota.48.35118>
- Dana ED, López-Santiago J, García-de-Lomas J, García-Ocaña DM, Gámez V, Ortega F (2010) Long-term management of the invasive *Pacifastacus leniusculus* (Dana, 1852) in a small mountain stream. *Aquatic Invasions* 5: 317–322. <https://doi.org/10.3391/ai.2010.5.3.10>
- Davidson IC, Scianni C, Minton MS, Ruiz GM (2018) A history of ship specialization and consequences for marine invasions, management and policy. *Journal of Applied Ecology* 55: 1799–1811. <https://doi.org/10.1111/1365-2664.13114>
- Diagne C, Leroy B, Vaissière A-C, Gozlan RE, Roiz D, Jarić I, Salles J-M, Bradshaw CJA, Courchamp F (2021) High and rising economic costs of biological invasions worldwide. *Nature* 592: 571–576. <https://doi.org/10.1038/s41586-021-03405-6>

- Displayr (2019) *flipPlots: Creates Plots* (R package version 1.2.0).
- Durán C, Lanao M, Pérez LPY, Chica C, Anadón A, Touya V (2012) Estimación de los costes de la invasión del mejillón cebra en la cuenca del Ebro (periodo 2005-2009). *Limnetica* 31: 213–230.
- Enders M, Havemann F, Jeschke JM (2019) A citation-based map of concepts in invasion biology. *NeoBiota* 47: 23–42. <https://doi.org/10.3897/neobiota.47.32608>
- Enders M, Havemann F, Ruland F, Bernard-Verdier M, Catford JA, Gómez-Aparicio L, Haider S, Heger T, Kueffer C, Kühn I, Meyerson LA, Musseau C, Novoa A, Ricciardi A, Sagouis A, Schittko C, Strayer DL, Vilà M, Essl F, Hulme PE, van Kleunen M, Kumschick S, Lockwood JL, Mabey AL, McGeoch MA, Palma E, Pyšek P, Saul W-C, Yannelli FA, Jeschke JM (2020) A conceptual map of invasion biology: Integrating hypotheses into a consensus network. Belmaker J (Ed.). *Global Ecology and Biogeography* 29: 978–991. <https://doi.org/10.1111/geb.13082>
- Enríquez ML, Abril S, Díaz M, Gómez C (2013) Nest site selection by the Argentine ant and suitability of artificial nests as a control tool. *Insectes Sociaux* 60: 507–516. <https://doi.org/10.1007/s00040-013-0317-3>
- Escribano Francés G, Quevauviller P, San Martín González E, Vargas Amelin E (2017) Climate change policy and water resources in the EU and Spain. A closer look into the Water Framework Directive. *Environmental Science & Policy* 69: 1–12. <https://doi.org/10.1016/j.envsci.2016.12.006>
- Essl F, Bacher S, Blackburn TM, Booy O, Brundu G, Brunel S, Cardoso A-C, Eschen R, Gallardo B, Galil B, García-Berthou E, Genovesi P, Groom Q, Harrower C, Hulme PE, Katsanevakis S, Kenis M, Kühn I, Kumschick S, Martinou AF, Nentwig W, O’Flynn C, Pagad S, Pergl J, Pyšek P, Rabitsch W, Richardson DM, Roques A, Roy HE, Scalera R, Schindler S, Seebens H, Vanderhoeven S, Vilà M, Wilson JRU, Zenetos A, Jeschke JM (2015) Crossing frontiers in tackling pathways of biological invasions. *BioScience* 65: 769–782. <https://doi.org/10.1093/biosci/biv082>
- European Parliament & Council (2000) Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy. *Official Journal of the European Union* 327: 1–73. <http://data.europa.eu/eli/dir/2000/60/oj>.
- Fajardo M, Rodríguez X, Hernández CD, Barroso L, Morales M, González A, Martín R (2019) The Canary Islands success story in eradicating red palm weevil. In: Al-Dobai S, Elkahky M, Faleiro R (Eds), *Proceedings of the “Scientific Consultation and High-Level Meeting on Red Palm Weevil Management.”* FAO and CIHEAM, Rome (Italy), 29–31.
- Fanelli D (2012) Negative results are disappearing from most disciplines and countries. *Scientometrics* 90: 891–904. <https://doi.org/10.1007/s11192-011-0494-7>
- Fellows I (2018) *wordcloud: Word clouds*. <https://cran.r-project.org/package=wordcloud>.
- Ferreras-Romero M, Márquez-Rodríguez J, Fernández-Delgado C (2016) Long-time effect of an invasive fish on the Odonata assemblage in a Mediterranean lake and early response after rotenone treatment. *Odonatologica* 45: 7–21. <https://doi.org/10.5281/zenodo.50846>
- Fornaroli R, Muñoz-Mas R, Martínez-Capel F (2020) Fish community responses to antecedent hydrological conditions based on long-term data in Mediterranean river basins (Iberian

- Peninsula). *Science of the Total Environment* 728: 1–16. <https://doi.org/10.1016/j.scitotenv.2020.138052>
- Gaertner M, Wilson JRU, Cadotte MW, MacIvor JS, Zenni RD, Richardson DM (2017) Non-native species in urban environments: patterns, processes, impacts and challenges. *Biological Invasions* 19: 3461–3469. <https://doi.org/10.1007/s10530-017-1598-7>
- García-de-Lomas J, Vilà M (2015) Lists of harmful alien organisms: Are the national regulations adapted to the global world? *Biological Invasions* 17: 3081–3091. <https://doi.org/10.1007/s10530-015-0939-7>
- García-Gómez JC, Sempere-Valverde J, González AR, Martínez-Chacón M, Olaya-Ponzzone L, Sánchez-Moyano E, Ostalé-Valriberas E, Megina C (2020) From exotic to invasive in record time: The extreme impact of *Rugulopteryx okamurae* (Dictyotales, Ochrophyta) in the strait of Gibraltar. *Science of the Total Environment* 704: 135408. <https://doi.org/10.1016/j.scitotenv.2019.135408>
- Garlock T, Asche F, Anderson J, Bjørndal T, Kumar G, Lorenzen K, Ropicki A, Smith MD, Tveterås R (2020) A global blue revolution: Aquaculture growth across regions, species, and countries. *Reviews in Fisheries Science and Aquaculture* 28: 107–116. <https://doi.org/10.1080/23308249.2019.1678111>
- Geraldi NR, Anton A, Lovelock CE, Duarte CM (2019) Are the ecological effects of the “worst” marine invasive species linked with scientific and media attention? *PLoS ONE* 14: e0215691. <https://doi.org/10.1371/journal.pone.0215691>
- Gläser J, Laudel G (2016) Governing science: How science policy shapes research content. *European Journal of Sociology* 57: 117–168. <https://doi.org/10.1017/S0003975616000047>
- Grundy JPB, Franco AMA, Sullivan MJP (2014) Testing multiple pathways for impacts of the non-native Black-headed Weaver *Ploceus melanocephalus* on native birds in Iberia in the early phase of invasion. *Ibis* 156: 355–365. <https://doi.org/10.1111/ibi.12144>
- Haddaway NR, Bayliss HR (2015) Shades of grey: Two forms of grey literature important for reviews in conservation. *Biological Conservation* 191: 827–829. <https://doi.org/10.1016/j.biocon.2015.08.018>
- Haddaway NR, Collins AM, Coughlin D, Kirk S (2015) The role of google scholar in evidence reviews and its applicability to grey literature searching. *PLoS ONE* 10: e0138237. <https://doi.org/10.1371/journal.pone.0138237>
- Hale S, Bright J, Blank G (2017) oii: Crosstab and statistical tests for OII MSc stats course. <https://cran.r-project.org/package=oi>
- Hernández-Brito D, Blanco G, Tella JL, Carrete M (2020) A protective nesting association with native species counteracts biotic resistance for the spread of an invasive parakeet from urban into rural habitats. *Frontiers in Zoology* 17: 13. <https://doi.org/10.1186/s12983-020-00360-2>
- Hulme PE, Pyšek P, Jarošík V, Pergl J, Schaffner U, Vilà M (2013) Bias and error in understanding plant invasion impacts. *Trends in Ecology and Evolution* 28: 212–218. <https://doi.org/10.1016/j.tree.2012.10.010>
- Jeschke JM, Börner K, Stodden V, Tockner K (2019) Open access journals need to become first choice, in invasion ecology and beyond. *NeoBiota* 52: 1–8. <https://doi.org/10.3897/neo-biota.52.39542>

- Kiernan JD, Moyle PB, Crain PK (2012) Restoring native fish assemblages to a regulated California stream using the natural flow regime concept. *Ecological Applications* 22: 1472–1482. <https://doi.org/10.1890/11-0480.1>
- Konno K, Akasaka M, Koshida C, Katayama N, Osada N, Spake R, Amano T (2020) Ignoring non-English-language studies may bias ecological meta-analyses. *Ecology and Evolution* 10: 6373–6384. <https://doi.org/10.1002/ece3.6368>
- Lenzner B, Latombe G, Capinha C, Bellard C, Courchamp F, Diagne C, Dullinger S, Gollivets M, Irl SDH, Kühn I, Leung B, Liu C, Moser D, Roura-Pascual N, Seebens H, Turbelin A, Weigelt P, Essl F (2020) What will the future bring for biological invasions on islands? An expert-based assessment. *Frontiers in Ecology and Evolution* 8: 280. <https://doi.org/10.3389/fevo.2020.00280>
- Loomans AJM (2021) Every generalist biological control agent requires a special risk assessment. *BioControl* 66: 23–35. <https://doi.org/10.1007/s10526-020-10022-1>
- Maceda-Veiga A, López R, Green AJ (2017) Dramatic impact of alien carp *Cyprinus carpio* on globally threatened diving ducks and other waterbirds in Mediterranean shallow lakes. *Biological Conservation* 212: 74–85. <https://doi.org/10.1016/j.biocon.2017.06.002>
- Maceda-Veiga A, Escribano-Alacid J, de Sostoa A, García-Berthou E (2013) The aquarium trade as a potential source of fish introductions in southwestern Europe. *Biological Invasions* 15: 2707–2716. <https://doi.org/10.1007/s10530-013-0485-0>
- Maceda-Veiga A, Escribano-Alacid J, Martínez-Silvestre A, Verdaguer I, Mac Nally R (2019) What's next? The release of exotic pets continues virtually unabated seven years after enforcement of new legislation for managing invasive species. *Biological Invasions* 21: 2933–2947. <https://doi.org/10.1007/s10530-019-02023-8>
- Mañas S, Gómez A, Palazón S, Pödra M, Minobis B, Alarcia OE, Casal J, Ruiz-Olmo J (2016) Are we able to affect the population structure of an invasive species through culling? A case study of the attempts to control the American mink population in the Northern Iberian Peninsula. *Mammal Research* 61: 309–317. <https://doi.org/10.1007/s13364-016-0277-x>
- Martínez-Ortí A (2013) Nuevo hallazgo del limnéido exótico *Pseudosuccinea columella* (Say, 1817) (Gastropoda: Pulmonata) en la península ibérica. *Noticiario Sociedad Española de Malacología* 60: 41–42.
- Martínez-Ortí A, Nappo A, Escutia V (2020) Nuevos hallazgos de los gasterópodos *Aplus assimilis* (Reeve, 1846) (F. Pisaniidae) y *Mitrella psilla* (Duclos, 1846) (F. Columbellidae) en la costa mediterránea española. *Arxius de Miscel·lània Zoològica* 18: 51–57. <https://doi.org/10.32800/amz.2020.18.0051>
- Mateos-Naranjo E, Cambrollé J, De Lomas JG, Parra R, Redondo-Gómez S (2012) Mechanical and chemical control of the invasive cordgrass *Spartina densiflora* and native plant community responses in an estuarine salt marsh. *Journal of Aquatic Plant Management* 50: 106–111. [http://www.apms.org/japm/vol50/2-17716\\_p106-124\\_APMdj.pdf](http://www.apms.org/japm/vol50/2-17716_p106-124_APMdj.pdf)
- Matzek V, Pujalet M, Cresci S (2015) What managers want from invasive species research versus what they get. *Conservation Letters* 8: 33–40. <https://doi.org/10.1111/conl.12119>
- Matzek V, Covino J, Funk JL, Saunders M (2014) Closing the knowing-doing gap in invasive plant management: Accessibility and interdisciplinarity of scientific research. *Conservation Letters* 7: 208–215. <https://doi.org/10.1111/conl.12042>

- McCull KA, Sunarto A (2020) Biocontrol of the Common Carp (*Cyprinus carpio*) in Australia: A Review and Future Directions. *Fishes* 5: 17. <https://doi.org/10.3390/fishes5020017>
- McGeoch MA, Butchart SHM, Spear D, Marais E, Kleynhans EJ, Symes A, Chanson J, Hoffmann M (2010) Global indicators of biological invasion: Species numbers, biodiversity impact and policy responses. *Diversity and Distributions* 16: 95–108. <https://doi.org/10.1111/j.1472-4642.2009.00633.x>
- de Medeiros CM, Hernández-Lambrano RE, Sánchez Agudo JA (2018) How reliable is the untrained eye in the identification of an invasive species? The case of alien bee-hawking yellow-legged hornet in Iberian Peninsula. *Contemporary Problems of Ecology* 11: 666–681. <https://doi.org/10.1134/S1995425518060136>
- Melero Y, Palazón S, Bonesi L, Gosálbez J (2010) Relative abundance of culled and not culled American mink populations in northeast Spain and their potential distribution: Are culling campaigns effective? *Biological Invasions* 12: 3877–3885. <https://doi.org/10.1007/s10530-010-9778-8>
- Miralles L, Dopico E, Devlo-Delva F, Garcia-Vazquez E (2016) Controlling populations of invasive pygmy mussel (*Xenostrobus securis*) through citizen science and environmental DNA. *Marine Pollution Bulletin* 110: 127–132. <https://doi.org/10.1016/j.marpolbul.2016.06.072>
- Moreno-Andrés J, Romero-Martínez L, Acevedo-Merino A, Nebot E (2017) Tratamientos basados en luz ultravioleta para aguas de lastre como opción viable hacia el control de especies invasoras en la bahía de Algeciras. *Almoraima. Revista de Estudios Campogibraltareños* 47: 159–172. <https://dialnet.unirioja.es/servlet/articulo?codigo=7213382&orden=0&info=link>
- Mungi NA, Kaushik M, Mohanty NP, Rastogi R, Antony Johnson J, Qureshi Q (2019) Identifying knowledge gaps in the research and management of invasive species in India. *Biologia* 74: 623–629. <https://doi.org/10.2478/s11756-018-00186-8>
- Muñoz-Mas R, García-Berthou E (2020) Alien animal introductions in Iberian inland waters: An update and analysis. *Science of the Total Environment* 703: 134505. <https://doi.org/10.1016/j.scitotenv.2019.134505>
- Newman MEJ, Girvan M (2004) Finding and evaluating community structure in networks. *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics* 69, 026113. <https://doi.org/10.1103/PhysRevE.69.026113>
- Nghiem LTP, Papworth SK, Lim FKS, Carrasco LR (2016) Analysis of the capacity of Google trends to measure interest in conservation topics and the role of online news. *PLoS ONE* 11: e0152802. <https://doi.org/10.1371/journal.pone.0152802>
- Nunes AL, Tricarico E, Panov VE, Cardoso AC, Katsanevakis S (2015) Pathways and gateways of freshwater invasions in Europe. *Aquatic Invasions* 10: 359–370. <https://doi.org/10.3391/ai.2015.10.4.01>
- Núñez MA, Amano T (2021) Monolingual searches can limit and bias results in global literature reviews. *Nature Ecology and Evolution* 5: 64. <https://doi.org/10.1038/s41559-020-01369-w>
- Painting SJ, Collingridge KA, Durand D, Grémare A, Créach V, Bernard G (2020) Marine monitoring in Europe: Is it adequate to address environmental threats and pressures? *Ocean Science* 16: 235–252. <https://doi.org/10.5194/os-16-235-2020>
- Pedersen TL (2021) ggraph: An implementation of grammar of graphics for graphs and networks. <https://cran.r-project.org/package=ggraph>
- Phillips SJ, Dudík M, Schapire RE (2004) A maximum entropy approach to species distribution modeling. In: *Proceedings of the Twenty-First International Conference on Machine Learn-*

- ing, ICML '04. Association for Computing Machinery, New York, NY, USA, 83. <https://doi.org/10.1145/1015330.1015412>
- Portela R, Vicente JR, Roiloa SR, Cabral JA (2020) A dynamic model-based framework to test the effectiveness of biocontrol targeting a new plant invader – the case of *Alternanthera philoxeroides* in the Iberian Peninsula. *Journal of Environmental Management*. <https://doi.org/10.1016/j.jenvman.2020.110349>
- R Core Team (2021) R: A language and environment for statistical computing. Version 4. <https://www.r-project.org>
- Radhakrishnan S, Erbis S, Isaacs JA, Kamarthi S (2017) Novel keyword co-occurrence network-based methods to foster systematic reviews of scientific literature. *PLoS ONE* 12: 1–16. <https://doi.org/10.1371/journal.pone.0172778>
- Rahel FJ, Olden JD (2008) Assessing the effects of climate change on aquatic invasive species. *Conservation Biology* 22: 521–533. <https://doi.org/10.1111/j.1523-1739.2008.00950.x>
- Riera M, Pino J, Melero Y (2021) Impact of introduction pathways on the spread and geographical distribution of alien species: Implications for preventive management in Mediterranean ecosystems. *Diversity and Distributions* 27: 1019–1034. <https://doi.org/https://doi.org/10.1111/ddi.13251>
- Robertson PAPA, Mill A, Novoa A, Jeschke JM, Essl F, Gallardo B, Geist J, Jarić I, Lambin X, Musseau C, Smith K, Booy O, Pergl J, Pyšek P, Rabitsch W, von Schmalensee M, Shirley M, Strayer DL, Stefansson RA, Smith K, Booy O (2020) A proposed unified framework to describe the management of biological invasions. *Biological Invasions* 22: 2633–2645. <https://doi.org/10.1007/s10530-020-02298-2>
- Rosvall M, Bergstrom CT (2010) Mapping change in large networks. *PLoS ONE* 5: e8694. <https://doi.org/10.1371/journal.pone.0008694>
- Rytwinski T, Harper M, Taylor JJ, Bennett JR, Donaldson LA, Smokorowski KE, Clarke K, Bradford MJ, Ghamry H, Olden JD, Boisclair D, Cooke SJ (2020) What are the effects of flow-regime changes on fish productivity in temperate regions? A systematic map. *Environmental Evidence* 9: 7. <https://doi.org/10.1186/s13750-020-00190-z>
- Saavedra S, Medina FM (2020) Control of invasive ring-necked parakeet (*Pittacula krameri*) in an island Biosphere Reserve (La Palma, Canary Islands): Combining methods and social engagement. *Biological Invasions* 22: 3653–3667. <https://doi.org/10.1007/s10530-020-02351-0>
- Sabater S, Artigas J, Durán C, Pardos M, Romaní AM, Tornés E, Ylla I (2008) Longitudinal development of chlorophyll and phytoplankton assemblages in a regulated large river (the Ebro River). *Science of the Total Environment* 404: 196–206. <https://doi.org/10.1016/j.scitotenv.2008.06.013>
- Saul W-C, Roy HE, Booy O, Carnevali L, Chen H-J, Genovesi P, Harrower CA, Hulme PE, Pagad S, Pergl J, Jeschke JM (2017) Assessing patterns in introduction pathways of alien species by linking major invasion data bases. *Journal of Applied Ecology* 54: 657–669. <https://doi.org/10.1111/1365-2664.12819>
- Seebens H (2019) Invasion ecology: Expanding trade and the dispersal of alien species. *Current Biology* 29: R120–R122. <https://doi.org/10.1016/j.cub.2018.12.047>
- Seebens H, Bacher S, Blackburn TM, Capinha C, Dawson W, Dullinger S, Genovesi P, Hulme PE, Kleunen M, Kühn I, Jeschke JM, Lenzner B, Liebhold AM, Pattison Z, Pergl J, Pyšek P, Winter M, Essl F (2021) Projecting the continental accumulation of alien species through to 2050. *Global Change Biology* 27: 970–982. <https://doi.org/10.1111/gcb.15333>

- Seebens H, Blackburn TM, Dyer EE, Genovesi P, Hulme PE, Jeschke JM, Pagad S, Pyšek P, van Kleunen M, Winter M, Ansong M, Arianoutsou M, Bacher S, Blasius B, Brockerhoff EG, Brundu G, Capinha C, Causton CE, Celesti-Grapow L, Dawson W, Dullinger S, Economo EP, Fuentes N, Guénard B, Jäger H, Kartesz J, Kenis M, Kühn I, Lenzner B, Liebhold AM, Mosena A, Moser D, Nentwig W, Nishino M, Pearman D, Pergl J, Rabitsch W, Rojas-Sandoval J, Roques A, Rorke S, Rossinelli S, Roy HE, Scalera R, Schindler S, Štajerová K, Tokarska-Guzik B, Walker K, Ward DF, Yamanaka T, Essl F (2018) Global rise in emerging alien species results from increased accessibility of new source pools. *Proceedings of the National Academy of Sciences* 115: E2264-LP-E2273. <https://doi.org/10.1073/pnas.1719429115>
- Seebens H, Briski E, Ghabooli S, Shiganova T, MacIsaac HJ, Blasius B (2019) Non-native species spread in a complex network: the interaction of global transport and local population dynamics determines invasion success. *Proceedings of the Royal Society B: Biological Sciences* 286: 20190036. <https://doi.org/10.1098/rspb.2019.0036>
- Shackleton RT, Larson BMH, Novoa A, Richardson DM, Kull CA (2019) The human and social dimensions of invasion science and management. *Journal of Environmental Management* 229: 1–9. <https://doi.org/10.1016/j.jenvman.2018.08.041>
- Thomsen M, Wernberg T, Olden J, Byers JE, Bruno J, Silliman B, Schiel D (2014) Forty years of experiments on aquatic invasive species: Are study biases limiting our understanding of impacts? *NeoBiota* 22: 1–22. <https://doi.org/10.3897/neobiota.22.6224>
- Tollington S, Turbé A, Rabitsch W, Groombridge JJ, Scalera R, Essl F, Shwartz A (2017) Making the EU legislation on invasive species a conservation success. *Conservation Letters* 10: 112–120. <https://doi.org/10.1111/conl.12214>
- Turbelin AJ, Malamud BD, Francis RA (2017) Mapping the global state of invasive alien species: Patterns of invasion and policy responses. *Global Ecology and Biogeography* 26: 78–92. <https://doi.org/10.1111/geb.12517>
- Velando A, Morán P, Romero R, Fernández J, Piorno V (2017) Invasion and eradication of the American mink in the Atlantic Islands National Park (NW Spain): A retrospective analysis. *Biological Invasions* 19: 1227–1241. <https://doi.org/10.1007/s10530-016-1326-8>
- Vilà M, Hulme PE [Eds] (2017) *Impact of biological invasions on ecosystem services*. Springer International Publishing, Cham (Switzerland), 354 pp. <https://doi.org/10.1007/978-3-319-45121-3>
- Walsh JC, Dicks L V., Sutherland WJ (2015) The effect of scientific evidence on conservation practitioners' management decisions. *Conservation Biology* 29: 88–98. <https://doi.org/10.1111/cobi.12370>
- Williams KJ, Ford A, Rosauer DF, De Silva N, Mittermeier R, Bruce C, Larsen FW, Margules C (2011) Forests of East Australia: The 35th Biodiversity Hotspot. In: *Biodiversity Hotspots*. Springer Berlin Heidelberg, 295–310. [https://doi.org/10.1007/978-3-642-20992-5\\_16](https://doi.org/10.1007/978-3-642-20992-5_16)
- Woodford DJ, Richardson DM, MacIsaac HJ, Mandrak NE, van Wilgen BW, Wilson JRU, Weyl OLF (2016) Confronting the wicked problem of managing biological invasions. *NeoBiota* 31: 63–86. <https://doi.org/10.3897/neobiota.31.10038>
- Yelenik SG, D'Antonio CM, Rehm EM, Caldwell IR (2020) Multiple feedbacks due to biotic interactions across trophic levels can lead to persistent novel conditions that hinder restoration. In: *Plant invasions: the role of biotic interactions*. CABI, 402–420. <https://doi.org/10.1079/9781789242171.0402>

- Zenni RD, Essl F, García-Berthou E, McDermott SM (2021) The economic costs of biological invasions around the world. *NeoBiota* 67: 1–9. <https://doi.org/10.3897/neobiota.67.69971>
- Zhao H, Lu L (2015) Variational circular treemaps for interactive visualization of hierarchical data. In: Liu S, Scheuermann G, Takahashi S (Eds) 2015 IEEE Pacific Visualization Symposium (PacificVis). Institute of Electrical and Electronics Engineers (IEEE), Hangzhou (China), 81–85. <https://doi.org/10.1109/PACIFICVIS.2015.7156360>

## Supplementary material 1

### Complete reference list and the features used to characterise the references

Authors: Rafael Muñoz-Mas, Martina Carrete, Pilar Castro-Díez, Miguel Delibes-Mateos, Josep A. Jaques, Marta López-Darias, Manuel Nogales, Joan Pino, Anna Traveset, Xavier Turon, Montserrat Vilà, Emili García-Berthou.

Data type: references and features

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/neobiota.70.68202.suppl1>

## Supplementary material 2

### Table S1. Table of features, categories and definitions used to characterise the compiled literature

Authors: Rafael Muñoz-Mas, Martina Carrete, Pilar Castro-Díez, Miguel Delibes-Mateos, Josep A. Jaques, Marta López-Darias, Manuel Nogales, Joan Pino, Anna Traveset, Xavier Turon, Montserrat Vilà, Emili García-Berthou

Data type: definitions

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/neobiota.70.68202.suppl2>