

# The reproductive biology of *Saccharum spontaneum* L.: implications for management of this invasive weed in Panama

Graham D. Bonnett<sup>1,2</sup>, Josef N. S. Kushner<sup>2</sup>, Kristin Saltonstall<sup>2</sup>

**1** CSIRO Plant Industry, QBP, 306 Carmody Road, St Lucia, Qld 4067, Australia **2** Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancon, Panamá

Corresponding author: *Graham D. Bonnett* (graham.bonnett@csiro.au)

---

Academic editor: *R. Hufbauer* | Received 28 August 2013 | Accepted 2 December 2013 | Published 24 January 2014

---

**Citation:** Bonnett GD, Kushner JNS, Saltonstall K (2014) The reproductive biology of *Saccharum spontaneum* L.: implications for management of this invasive weed in Panama. NeoBiota 20: 61–79. doi: 10.3897/neobiota.20.6163

---

## Abstract

*Saccharum spontaneum* L. is an invasive grass that has spread extensively in disturbed areas throughout the Panama Canal watershed (PCW), where it has created a fire hazard and inhibited reforestation efforts. Currently physical removal of aboveground biomass is the primary means of controlling this weed, which is largely ineffective and does little to inhibit spread of the species. Little is known about reproduction of this species, although it is both rhizomatous and produces abundant seed. Here we report a series of studies looking at some of the basic reproductive mechanisms and strategies utilised by *S. spontaneum* to provide information to support development of better targeted management strategies.

We found that seed produced between September and November was germinable both in the lab and *in situ*. Genetic diversity of mature stands was assessed using microsatellite markers and found to be high, even at small scales. Studies of vegetative reproduction showed that buds on stems that had been dried for up to six weeks were still capable of sprouting. Separate experiments showed that stem fragments could sprout when left on the surface or buried shallowly and that larger pieces sprouted more readily than smaller pieces.

Collectively these results demonstrate that *S. spontaneum* in the PCW has the capability to produce many propagules that can successfully recruit and it is likely that seed dispersal drives the spread of the species. Timing of management actions to reduce flowering would significantly reduce the seed load into the environment and help to prevent spread to new sites. Similarly, where biomass is cut, cutting stems into smaller pieces will allow the stems to dry out and reduce the ability of buds to sprout. Additionally, attention should be paid to prevent accidental transport to new sites on machinery.

## Keywords

Invasive species, seed germination, asexual reproduction, microsatellite

## Introduction

Many of the “World’s Worst Weeds” are perennial species with the ability to spread with both seeds and vegetative structures (Holm et al. 1977) and perennial grasses are notorious invaders of both agricultural and natural areas (D’Antonio and Vitousek 1992). Of these, large-statured grasses are particularly difficult to manage due to their high biomass and rapid dominance both above- and belowground (Lambert et al. 2010). Grasses can produce copious amounts of seeds, which are often wind-dispersed and may remain dormant until conditions are right for germination (Donohue et al. 2010). In addition, many are rhizomatous and can spread vegetatively both to new sites and rapidly within a site once established (e.g. *Arundo donax* L., *Phragmites australis* (Cav.) Trin. ex Steud.; Boose and Holt 1999, Bart and Hartmann 2003). Stem fragments containing axillary buds may also play a role in dispersal to new sites (Boose and Holt 1999).

*Saccharum spontaneum* L. (wild sugarcane; Poaceae) is a polymorphic species believed to have evolved in India (Mukherjee 1957). It is a highly adaptable polyploid that grows in a wide range of habitats across southern Asia and east Africa to the Mediterranean, spanning the tropics to temperate regions from latitudes 8° S to 40° N (Daniels and Roach 1987; Tai and Miller 2001). Although it is both wind-pollinated and -dispersed, it also reproduces vegetatively, through clonal spread either from underground rhizomes or culm fragments. Because of *S. spontaneum*’s potential to propagate via seeds and its propensity for aggressive rhizomatous spread, it is considered a weed, even in countries to which it is native such as India (Panje 1970, Yadav et al. 2007) and Thailand (Pichitkul 2009). It has also been introduced into other countries for use in sugarcane breeding programs (Bonnett et al. 2008) and is considered a noxious weed in many countries including the USA (USDA 2010). Research in India has shown that flowering in *S. spontaneum* is controlled by photoperiod, with vegetative growth occurring in the wet monsoon season, followed by a dormant phase during the dry season. In India, flowering is initiated at the end of the rainy season, with diurnal variations in humidity and temperature likely influencing both pollen and seed dispersal (Panje and Srinivisan 1959).

In the Republic of Panamá, *S. spontaneum* (Paja Canalera) has spread extensively in the Panama Canal Watershed (PCW) since the first herbarium specimen was collected in 1960 (MO1824369 J.E. Ebinger 490). It now dominates in abandoned agricultural lands and along human transportation corridors, such as roads and railroad tracks, encompassing over 3 percent of the watershed (ACP-ANAM 2006). It grows to an average height of 3–4 m in dense, impenetrable stands which impede growth of other plants and provide little useful habitat for wildlife. Efforts towards reforestation of the PCW are seriously hindered by this species, as it can inhibit germination, establishment, and growth of native tree species (Hammond 1999, Hooper et al. 2002,

Jones et al. 2004) and increases the vulnerability of reforestation projects and nearby forests to devastating fires in the dry season (Saltonstall and Bonnett 2012).

Flowering of *S. spontaneum* across the landscape in Panama typically begins in August, midway through the rainy season which normally occurs from May through December. Inflorescences can be seen year-round, but are typically restricted to certain clones on a small scale during other times of the year (K. Saltonstall, pers. obs.). Flowering densities vary, averaging 4-5 stems/m<sup>2</sup> during the peak flowering season (Saltonstall and Bonnett 2012), and hundreds of seeds are produced by each inflorescence. However, the reproductive biology of *S. spontaneum* as an invasive species has not yet been studied. The relatively recent establishment and spread of *S. spontaneum* in Panama makes this an ideal environment to understand the relative contribution of sexual vs vegetative dispersal between and within sites.

The case for understanding the biology of invasive plants as a precursor to developing control strategies has been well made. Zamora et al. (1989) identified knowledge of the population dynamics and life cycle as key to finding the stage of development most vulnerable to control measures. In many cases the relative contributions of sexual and vegetative reproduction to invasive spread are largely unknown. Here we present a series of studies which evaluate reproductive and genetic factors that may contribute to the invasive success of *S. spontaneum* in Panama. The hypotheses that we tested were: 1) Seeds collected *in situ* are viable through the period of production, as tested under control conditions; 2) seeds can germinate *in situ*; 3) stands of mature plants are genetically diverse; 4) vegetative buds are able to produce plants under a range of conditions. We hope that a better understanding of these reproductive factors will assist in improving management strategies to prevent further spread throughout the country and beyond.


## Methods

### Study area

The Republic of Panamá is located at approximately 8–9° N latitude (Fig. 1), which is a latitude central to the native range of *S. spontaneum* (Mukherjee 1957). Day length varies only about one hour from its maximum at the June summer solstice to December and average daily temperatures range between 19°C and 35°C. During the rainy season months of May to December, rainfall averages around 240 mm per month, with rainfall occurring frequently (World Meteorological Organization 2010). All studies occurred during 2009, which was a year with typical rainfall patterns.

### Germination ability

We collected seeds of *S. spontaneum* from 12 sites (Sites 1–12 Fig. 1; Appendix 1) in the PCW weekly between September and December. Each week three mature inflo-



**Figure 1.** Map of Panama showing the locations of 22 sites where *S. spontaneum* samples were collected. Seeds were collected weekly at Sites 1-12 for germination trials. Lines in the larger map indicate district boundaries. In the inset, the canal is shown between Panama City and Gamboa and major roads are indicated by grey lines.

rescences, from different plants within a few meters of each other, were sampled and assayed separately. Seeds that had disarticulated from the inflorescence but remained loosely attached were collected, and kept at room temperature until assayed within two days of harvest. When wet at the time of harvest due to rain, seeds were dried under ambient conditions prior to conducting germination assays.

One hundred seeds were taken at random from those collected from each inflorescence and germinated on moist paper towel in petri dishes ( $n=3$  inflorescences per site per week). Seeds were germinated at  $36^{\circ}\text{C}$  as *Saccharum* seeds have a very high temperature optimum for germination of around  $36^{\circ}\text{C}$  (Skinner 1959, Singh 1988).

Assays were conducted in the dark, as there is no evidence of a light requirement for the germination of *Saccharum* seeds (G. Bonnett, unpublished).

Germination was defined as root or radicle emergence visible to the naked eye. Seed germination was checked every three to four days and counted seedlings were removed from the plates. Total number of seeds germinated after two weeks is presented. For analysing the effect of date of collection on germination percentage we carried out a one-way ANOVA with date as a factor and log-transformed ( $\log_{10}(\text{value}+1)$ ) germination data as the dependent variable (Sigmaplot 11, Systat Software Inc. San Jose, CA). Post-hoc comparisons of means were conducted using the Tukey method, for this and all subsequent analyses significance was tested at the 0.05 level.

### **In situ germination**

To test if seeds would germinate *in situ*, vegetation was cleared and the soil surface raked on 16 September exposing areas of bare soil between two *S. spontaneum* stands (Site 6, Fig. 1, Appendix 1). The experiment comprised 10 replicates of three treatments, each in 60 by 120 cm plots. Treatment one, bare soil, was designed to observe any seedlings emerging from seeds deposited by wind. The second treatment comprised six mature inflorescences placed on each plot and covered with nylon flywire to prevent seeds from blowing away. Plots in treatment three were bare soil covered by nylon flywire, with no additional seeds added. One replicate of each treatment was randomly placed in each of 10 evenly spaced blocks. After 19 days we removed the mesh to allow continued growth of seedlings. The number of germinated *S. spontaneum* seedlings per plot was recorded one week after, and then every two weeks until nine weeks after the start of the experiment.

To test for differences in seedling numbers between treatments, we conducted a repeated measures analysis with time (within subject) and treatment (between subjects) as factors and the number of seedlings as the dependent variable (Sigmaplot 11). The number of seedlings was log-transformed ( $\log_{10}(\text{value}+1)$ ) prior to analysis and post-hoc comparisons of means were conducted using the Holm-Sidak method (a variation of the Holm multi-comparison test using the Sidak correction (Sidak 1967)).

### **Genetic diversity of mature stands**

To assess levels of genetic diversity within stands, plants were collected at 22 sites across central Panama, with a focus on the PCW but including the eastern- and westernmost edges of the invasion (Fig. 1, Appendix 1). Green leaves were harvested from individual culms spaced up to 10 m apart ( $n = 3$  per site). To examine if the patterns we found at these sites were consistent across large stands, we also sampled parallel transects separated by 20 m at two independent stands in Soberania National Park within an area dominated by *S. spontaneum* (Sites 5 and 6, Fig. 1, Appendix 1; see Saltonstall and

Bonnett 2012 for a description of the stands). Five transects were laid out in Stand 1 and three in Stand 2 and samples collected every 10 m along the transects (n=26 for Stand 1, n=9 for Stand 2). All samples were kept frozen at -20°C until DNA was extracted using a CTAB extraction protocol (Doyle and Doyle 1987).

We assessed multilocus allele profiles using 12 microsatellite loci (msscir14, msscir17, msscir53, msscir58, SMC28, SMC221, SMC334, SMC336, SMC597, SMC1047, SMC1237, SMC1493; Brown et al. 2007; Pan 2006) targeting different regions of the genome. PCR amplifications were performed in 7  $\mu$ L reaction volumes containing approximately 1 ng DNA, 0.056 U Amplitaq (Applied Biosystems, Inc., Carlsbad, CA), 1x reaction buffer (Applied Biosystems, Inc.), 3 mM MgCl<sub>2</sub>, 200  $\mu$ M dNTPS, 0.5 pmol of forward primer including an M13 tag, 2.5 pmol of reverse primer, 2 pmol of FAM, PET, VIC, or NED labelled M13 primer, and 1 mg mL<sup>-1</sup> BSA. All PCR reactions were conducted with 2 min of denaturation at 94°C, 35 cycles of denaturation at 94°C for 45 s, annealing at 56°C for 25 s, and 25 s of extension at 72°C, and a final extension period of 72°C for 5 min in a thermal cycler (Eppendorf Multimax, Eppendorf, Hamburg, Germany). Primers were not multiplexed due to the high ploidy level of the species, but PCR reactions with differing florescent tags were multiloading for capillary electrophoresis on an ABI prism 3730 automated sequencer. Fragment sizes were analysed using GeneMapper 3.7 (Applied Biosystems Inc.) and individuals were scored for presence or absence of identified alleles for each locus. All scored alleles had consistent results over replicate amplifications. Due to polyploidy in this species, allele frequencies are unknown thus multilocus allele profiles are hereafter referred to as allele phenotypes. We compared all multilocus allele phenotypes found within and among sites to identify repeated allele phenotypes, which we assumed to result from asexual reproduction (e.g. clonal expansion).

### **Sprouting of vegetative propagules**

*Effect of drying:* To assess the ability of buds to remain viable and sprout after different periods of drying, culms were collected from Site 4 (Fig. 1, Appendix 1) on 3 September. Culms, approximately 2 m tall, that had not flowered were cut at the base and randomly assigned to five treatments (drying durations) spread out on benches that were shielded from rainfall but otherwise open. Drying durations were 0 days (day of harvest), one, two, four, and six weeks. After drying, stem pieces with a node at either end were cut from the culms and ten were placed into each of five replicates and covered with commercial peat. Each replicate comprised two plastic seedling trays (53 cm by 26 cm). Plants were grown under an open structure with a transparent roof, watered daily, and the plants that sprouted recorded four weeks after planting. To assess changes in moisture content, plant material was harvested from the same site on 19 November. Fresh and dry weights of stem pieces cut from culms at the time of harvest and after one, two and four weeks were recorded. Dry mass was taken after drying the stem pieces to constant weight at 40°C.

Data from the different times of drying were compared by ANOVA and times of drying were compared with Tukey multiple comparison tests (Sigmaplot 11). The proportion of buds that sprouted was square root transformed, other data was not transformed prior to analysis.

### **Effects of propagule size and planting depths on sprouting**

Fresh culms and rhizomes were collected at Site 6 (Fig. 1, Appendix 1) on July 3 and propagules were cut to the appropriate size and planted the same day. Both experiments were performed under a covered shelter, open on the sides. Unmodified soil representative of the area (Oxisol, collected in Rio Hato, Panama) was used and all pots were watered daily or as needed. All propagules were harvested on September 12 when the presence of a sprout above-ground, a sprout below-ground but not emerged, or the presence of roots was recorded.

*Propagule size:* Culm pieces with one node (and bud) were cut into either 2, 5, 10, or 20 cm lengths or two node (and bud) pieces ranging in length from 9.9 to 12.7 cm. Rhizomes were cut into pieces ranging from 15–40 mm in length, either with or without visible axillary buds. There were 10 replicates for each culm length (single and two node pieces) and 25 pieces of each rhizome type. Culm pieces shorter than 10 cm were planted individually in 10 cm diameter plastic nursery sacks while longer pieces were planted in 12 L pots. Rhizomes were planted individually in 10 cm sacks. All propagules were planted at a depth of 5 cm.

*Planting depth:* Individual propagules (5 cm single node stem pieces or single node rhizome pieces (ranging from 11–44 mm in length)) were planted at different depths in 10 cm nursery sacks. Planting depths were surface, 5 cm, 10 cm, and 20 cm for stem pieces, and surface, 10 cm, and 20 cm for rhizome pieces; there were ten replicates per planting depth. Time to sprouting was recorded for each propagule.

Our original intent was to analyse growth of sprouts using analysis of variance appropriate to the experimental design. However, because of the frequency of failure to sprout, and thus zero growth, we decided to analyse only the proportion sprouted either above- or belowground, as the percentage of the 10 replicates that sprouted for each treatment.

To test if soil type influenced the results of this experiment, growth was also tested using commercial potting mix in nursery sacks. Single node stem pieces (5 cm) were planted at the surface or 2 cm depth, in either commercial potting mix or unmodified soil from Rio Hato, with six replicates per treatment. Presence of sprouting was recorded after four weeks.

We further distinguish between propagules that produced roots at the node but did not initiate other growth, propagules that sprouted but died before they reached the soil surface, and propagules that produced visible growth above the soil surface. Results were analysed using generalized linear models (Poisson GLM for propagule depth and Binomial GLM for propagule size and soil type) as implemented in R 2.13.1 (R Development Core Team 2010). The proportion of variance explained by each model ( $r^2$ ) was calculated from the null and residual variances.

## Results

### Sexual reproduction

#### Germination ability

The proportion of seeds germinating and the temporal patterns of germination varied greatly across the 12 sites (Appendix 2). The range of maximum germination percentages were 72%–30% (Appendix 2), with an average level of germination around 35% across sites until November (Fig. 2). Significantly fewer seeds germinated after 16 November (ANOVA,  $F_{12,403} = 25.5$ ,  $P < 0.001$ , Fig. 2).


#### In situ germination

The numbers of seedlings germinating *in situ* showed a significant effect of treatment ( $F_{2,18} = 21.6$ ,  $p < 0.001$ ), week ( $F_{4,36} = 7.4$ ,  $p < 0.001$ ) and treatment by week interaction ( $F_{8,72} = 29.9$ ,  $p < 0.001$ ), so the effect of treatment was dependent upon time. One week after the experiment was initiated, germination was evident in all plots (Fig. 3) and there were significant differences in the number of seedlings between each treatment, with the plots with added seeds having many more seedlings than the controls. The average numbers of seedlings in plots with added seeds decreased from week to week, while those in control plots increased over time. Within the flywire treatment, there was a significant increase in the number of seedlings between weeks 3 and 5, corresponding with the removal of the flywire, whereas the seeds added and bare soil control plots did not show significant changes after week 3. After nine weeks there were no significant differences in the numbers of seedlings remaining between any of the treatments.


#### Genetic diversity of mature stands

High levels of genetic diversity were found both within and across sites. A total of 227 alleles were observed across the twelve microsatellite loci, of which 215 were shared between sites. The average number of alleles per locus ( $A_o$ ) was 18.4 and up to ten alleles per individual were found within a locus (range 1–10).

In the two stands sampled with transects, the majority of culms sampled were unique (Stand 1 at Site 5 = 89% and Stand 2 at Site 6 = 65%) and all replicate allele phenotypes were found in adjacent plots. Similarly, in the 22 sites where only 3 samples were collected, 86% of them contained multiple allele phenotypes (19 of 22 sites). Fifty four multilocus allele profiles were found across the 22 sites, with three unique phenotypes sampled at each of 13 sites (Sites 3, 5, 6, 11, 13–16, and 18–22), two phenotypes at each of six sites (2, 4, 7, 9, 10, and 17), and a single phenotype at each of three sites (1, 8, and 12; Appendix 1). The same multilocus profile was not found at more than one site.



**Figure 2.** Weekly average germination percentage of *S. spontaneum* seeds from 12 sites. Error bars represent standard error of the mean. Dates of collection with different letters are significantly different.



**Figure 3.** Average numbers of *S. spontaneum* seedlings germinating each week per plot in the *in situ* germination experiment. “*Saccharum* added” plots had mature inflorescences placed in them and were initially covered with flywire, “flywire control” plots were initially covered with flywire but no seeds were added, and control plots were similar sized plots of bare soil. Error bars represent standard error of the mean (n=10). Weeks with different capital letters are significantly different from each other. Treatments with different small letters are significantly different from each other within that week.

**Table 1.** Moisture content of stems and the proportion of buds that sprouted after various times of drying. Prior to testing for sprouting, buds were cut from culms immediately after harvesting (0) and 1–4 weeks of drying. Sprouting of buds was recorded after 4 weeks. Results are presented as the mean with the standard error in parenthesis. Values within a column with different letters are significantly different.

Time (weeks)	Moisture content (%)	Proportion of buds sprouting (%)
0	80.2 (1.08) <sup>a</sup>	70.9 (4.39) <sup>ab</sup>
1	76.7 (1.21) <sup>ab</sup>	84.0 (2.92) <sup>a</sup>
2	77.2 (0.68) <sup>ab</sup>	57.0 (6.63) <sup>bc</sup>
4	73.7 (0.74) <sup>b</sup>	45.0 (3.16) <sup>c</sup>
6	N.D.	38.8 (3.75) <sup>c</sup>

N.D. = not determined

## Vegetative propagation

### Effects of drying


There was a significant effect of time of drying of the culm on the proportion of buds that sprouted (ANOVA  $F_{4,19} = 16.992$ ,  $P < 0.001$ , Table 1), less than half of the buds sprouting after 4 weeks. However while significant over time, the reduction in moisture content was only 8% lower after 4 weeks of drying (ANOVA  $F_{3,16} = 7.819$ ,  $P < 0.01$ , Table 1).

### Effects of propagule size


Increased size of stem fragments increased sprouting ability ( $p < 0.01$ ,  $r^2 = 0.34$ ; Fig. 5). While roots were produced in the majority of fragments of 5, 10 and 20 cm lengths, no fragments of 2 cm, 5 cm, or 10 cm produced active sprouts from the node. One 20 cm (10%) and one 2-node fragment (10%) sprouted belowground but died before reaching the soil surface. Three (30%) 20 cm fragments and seven (70%) 2-node fragments sprouted aboveground. All 2-node fragments that grew sprouted at only one node, but had roots present at both nodes. No rhizome fragments, with or without buds, sprouted.

### Effects of planting depth

There was a significant negative effect of planting depth on the ability of stem cuttings to sprout, with lower depths having an inhibitory effect on sprouting ( $p < 0.05$ ,  $r^2 = 0.29$ , Fig. 4). While 100% of stem cuttings laid at the soil surface sprouted, only 40% successfully grew aboveground from a depth of 5 cm and only 20% from 10 cm. Stems at a depth of 20 cm failed to grow aboveground, but 20% of propagules at this depth sprouted and died before reaching the soil surface. All rhizome pieces, at any depth, failed to sprout suggesting that single node pieces of *S. spontaneum* rhizomes may have low sprouting ability (data not shown). One rhizome piece, at 10 cm depth, produced



**Figure 4.** Proportion of buds sprouting from different depths of planting. Proportion of buds that produced visible growth above the soil surface (Above ground), that sprouted but died before they reached the soil surface (Below ground), or produced roots at the node (Roots), but did not initiate other growth as determined by depth of planting.



**Figure 5.** Proportion of buds sprouting from different sizes of culm piece planted. Proportion of buds that produced visible growth above the soil surface (above ground), that sprouted but died before they reached the soil surface (below ground), or produced roots at the node (roots), but did not initiate other growth as determined by size of culm piece.

roots but did not produce a sprout. Rhizome pieces laid at the soil surface dried out quickly, despite daily watering, and did not even produce roots.

Depth trials comparing commercial potting soil (PS) to the unmodified soil (US) used in the previous trial showed similar results, with 83.3% (PS) and 67.7% (US) of cuttings laid at the soil surface sprouting aboveground and 67.5% (PS) and 50.0% (US) sprouting when planted at a depth of 2 cm, respectively. No significant differences were seen between depths or soil types ( $p > 0.05$ ,  $r^2 = 0.05$ ).

## Discussion

The spread of *S. spontaneum* in the PCW has been rapid and dramatic, occurring largely in recent decades. Flowering across the landscape is extensive and our results suggest that seed dispersal clearly plays a major role in this spread. Seed germination rates were high when tested under optimal conditions, and we found seedlings readily establish *in situ*. Although we did not follow the fate of these seedlings to maturity, 79% of our samples collected in mature stands had unique allele phenotypes, with all replicated genotypes occurring in adjacent samples within a site. Such high genotypic diversity cannot result from vegetative spread and can only be the result of sexual reproduction. Our trials with vegetative propagules also showed that growth can occur from stem fragments, although they can be highly sensitive to the conditions under which they are grown.

Although we did not test viability directly, our germination trials indicate that on a landscape scale large numbers of germinable seeds are produced annually. Variability in germination within and between sites was high but seeds germinated from all sites, particularly those collected during the peak flowering months of September and October. Other invasive grasses, such as *Phragmites australis* and *Spartina alterniflora* in North America, have also been shown to have highly variable production of viable seeds across sites (Daehler and Strong 1994, Kettenring and Whigham 2009). Although the average viability of these other species is typically much lower than the rates we have observed (e.g. less than 20 % for *Phragmites australis* (Kettenring and Whigham 2009)), they are aggressive invaders in North American wetlands and seeds play an important role in their dispersal (Belzile et al. 2010; Daehler and Strong 1994; McCormick et al. 2010).

Seeds which germinated *in situ* showed high mortality in plots where they were added, but control plots which received seeds only through natural dispersal increased their numbers of seedlings from week to week. In the case of the bare soil control plots, seedling densities were ~27 seedlings/plot (37 seedlings/m<sup>2</sup>) by the end of the experiment, which is similar to densities counted in plots measured in a *S. spontaneum* stand which had burned earlier in the year (Saltonstall and Bonnett 2012). While we did not follow the fate of individual seedlings, it is likely that mortality occurred in the control plots as well and it was a constant rain of novel seeds landing and germinating

in the plots that kept their numbers increasing. The high mortality seen in the seed-added plots could be due to negative interactions on a seed-seed or seed-seedling basis, such as competitive or allelopathic effects which were compounded by the high densities of seeds and seedlings. Such effects have been shown in sugarcane (Sampietro et al. 2007) and several other species (Inderjit and Streibig 2001; Murray 1998) but not yet demonstrated for *S. spontaneum*. As the fate of the seedlings established in this experiment was not followed after the nine weeks of observations, additional research regarding the long-term survivorship of *S. spontaneum* seedlings is needed.

Genetic diversity across the landscape is high. We found high numbers of alleles both within individuals and between sites, and nearly all samples tested were genetically unique. This suggests that stands of *S. spontaneum* typically form from the coalescence of multiple individuals recruited from seeds rather than vegetative propagation of a single clone across a large area. Consequently, though we have not demonstrated recruitment of adult plants from seedlings, flowering is a life-history stage that needs to be targeted to effectively manage the spread of the species.

The effects of disconnection of stems and rhizomes on vegetative reproduction has been studied extensively in rhizomatous grasses, and disconnection from the parent plant has been shown to release buds from inhibition by other buds on the stem or rhizome. This response may be influenced by the length of the segment containing the buds as well as which the conditions to which propagules may be exposed to following disconnection from the parent plant (Kigel and Koller 1985, Boose and Holt 1999). *Saccharum spontaneum* appears to regenerate better from larger cuttings, with larger stem fragments having a greater chance of sprouting and reaching the soil surface. Dessication reduced sprouting ability to some extent but 39% of buds still sprouted after six weeks of drying. Similar results have been seen in experiments with other rhizomatous grasses, such as *Agropyron repens* (Turner 1968) and *Pennisetum macrourum* (Harradine 1980) where successful emergence and establishment decreased with planting depth and increased with size of the fragment. It is likely that larger segments are able to sustain buds in a dormant condition for longer periods of time and that once growing is initiated by the dominant bud, heterotrophic growth (e.g. underground) can be sustained for longer periods of time (Kigel and Koller 1985).

Ability of buds to sprout also appears to be strongly affected by planting depth. Planting stem cuttings at depths as shallow as 5 cm reduced the number of aboveground sprouts by 60%, and deeper plantings inhibited even the sprouting of fine roots from the nodes. The lack of growth in our initial studies prompted concerns about the soil that we used inhibiting growth in this experiment. Although the bags we used had drainage holes, the Oxisols used here have high clay content and waterlogging may have been an issue preventing growth of propagules. However, our secondary trials comparing growth in the Oxisol versus commercial potting mix showed similar results, with reduced growth when fragments were planted below the soil surface suggesting that the inhibition of growth could be a light response rather than an effect of the growth substrate.

## Management of *S. spontaneum*

Management of *S. spontaneum* in Panama is restricted to controlling aboveground biomass, and generally involves physical cutting of mature biomass, and in some cases, chemical control where reforestation is a goal (Craven et al. 2009). Fire is also used during the dry season months to remove biomass. When cut, biomass is typically left on site to dry and degrade naturally. These activities occur year round, and regrowth of stands following biomass removal occurs frequently and rapidly (Saltonstall pers. Obs). Like many other weeds (Holm et al. 1977), the remaining buds and rhizome system thus presents the greatest barrier to control of established stands of *S. spontaneum*. However, little thought goes into the effect of these management efforts on seed production and further spread of the species.

Timing and method of control thus become important when considering management of this aggressive invader in Panama. While intermittent flowering of *S. spontaneum* occurs year round, the peak season of flowering is from August – October. Our studies have shown that seed germinability is highest in the months of September, October, and early November with a rapid decline in germinability in December. We suspect that germinability of seeds produced at other times of the year is also much lower. Timing of biomass removal can be important, with removal in June or July, prior to inflorescence emergence and seed development but late enough in the year that reproduction will not occur in the months of August – November, possibly being a method to reduce spread by seed. This may be particularly important in areas on the edges of the distribution where *S. spontaneum* does not yet dominate and spread can be minimized on a local scale. As herbicides are not commonly used in Panama except during the initial stages of reforestation projects, repeated cutting of stems throughout the year to deplete below-ground carbohydrate resources in rhizomes is another approach which may help to prevent flowering at optimal times of the year. However, as our studies have shown that cut stems are still capable of sprouting after six weeks of desiccation, cut biomass should also be handled to minimize sprouting and spread to new sites.

Better control of dry-season fires is also needed to help control spread of *S. spontaneum* by seed, as rapid regrowth following the onset of the rainy season typically leads to stems with mature inflorescences developing in the peak months of seed production. Further, burned areas have higher densities of flowering stems and comparable seed germination rates as unburned areas (Saltonstall and Bonnett 2012) suggesting that the potential for spread to new sites is increased by fire.

As *S. spontaneum* in Central Panama is widespread and many thousands of seeds are produced by each plant, a strategy to control seed numbers will have to be co-ordinated over a large area to be truly effective. While the extent of seed dispersal has not been experimentally quantified, research conducted on Barro Colorado Island (BCI) in the Panama Canal, utilising 200 seed rain traps (Wright and Calderón 2006), provides some evidence. *Saccharum spontaneum* is found only in small, isolated patches on BCI,

mostly within treefall gaps in the forest interior (S.J. Wright, personal communication). However, between 1990 and 2011 over 7500 *S. spontaneum* seeds were trapped (annual median 229); S.J. Wright unpublished data). These seeds were likely wind dispersed from mainland areas and would have travelled at least 2 km across the Canal from the nearest sources of *S. spontaneum* plants. Information regarding the longevity of *S. spontaneum* seed remaining in the seed bank is also lacking, but a concerted effort over several years to eliminate *S. spontaneum* seed from the seed bank will be required to stop spread by seed. The closely related invasive *Andropogon gayanus* has been shown to retain 1% of germination of seed after one year in burial experiments (Flores et al. 2005), suggesting that establishment of new plants could continue to occur for some time even if annual seed production is reduced across the landscape.

In conclusion we have presented evidence that the spread of *S. spontaneum* in Panama has been driven by seed production, but that vegetative propagules derived from stem fragments are also robust and can be the source of new plants. Seed production from *S. spontaneum* can be reduced through the timing of management actions to reduce biomass that prevents re-growth of flowering stems. However the mobility of the seed means that to be effective, such actions would have to be conducted over large areas. Once established, the persistence of individual stands through re-growth from buds lower on the stems and rhizomes will continue to impede control and eradication over large areas so further research to determine a weak point in the vegetative growth stage is needed.

## **Acknowledgements**

GDB thanks the Queensland Government (Queensland Smithsonian Fellowship), CRC Sugar Industry Innovation through Biotechnology and the Australian Government and the Australian Sugarcane Industry as provided by the Sugar Research and Development Corporation for financial support for an extended visit to the Smithsonian Tropical Research Institute. KS was funded by a Smithsonian Postdoctoral Fellowship. We thank the Smithsonian Tropical Research Institute and Jefferson Hall for logistical support, Eldredge Bermingham for the use of his laboratory, and Milton Salano for assistance with Fig. 1. Deanne Haffner is thanked for extensive input to the germination assays. J. Hall and K. Juneau provided helpful comments on earlier drafts of the manuscript.

## **References**

ACP-ANAM (2006) Programa de Vigilancia de la Cobertura Vegetal, Región Oriental de la Cuenca del Canal. Panama City, Panama, Panama Canal Authority. <http://www.pancanal.com/esp/cuenca/cobertura-vegetal.pdf>

- Bart D, Hartman JM (2003) The role of large rhizome dispersal and low salinity windows in the establishment of common reed, *Phragmites australis*, in salt marshes: New links to human activities. *Estuaries* 26: 436–443. doi: 10.1007/BF02823720
- Belzile F, Labbe J, Leblanc M-C, Lavoie C (2010) Seeds contribute strongly to the spread of the invasive genotype of the common reed (*Phragmites australis*). *Biological Invasions* 12: 2443–2250. doi: 10.1007/s10530-009-9634-x
- Bonnett GD, Nowak E, Olivares-Villegas JJ, Berding N, Morgan T, Aitken KS (2008) Identifying the risks of transgene escape from sugarcane crops to related species, with particular reference to *Saccharum spontaneum* in Australia. *Tropical Plant Biology* 1: 58–71. doi: 10.1007/s12042-007-9002-x
- Boose AB, Holt JS (1999) Environmental effects on asexual reproduction in *Arundo donax*. *Weed Research* 39: 117–127. doi: 10.1046/j.1365-3180.1999.00129.x
- Brown JS, Schnell RJ, Power EJ, Douglas SL, Kuhn DN (2007) Analysis of clonal germplasm from five *Saccharum* species: *S. barberi*, *S. robustum*, *S. officinarum*, *S. sinense* and *S. spontaneum*. A study of inter- and intra species relationships using microsatellite markers. *Genetic Resources and Crop Evolution* 54: 627–648. doi: 10.1007/s10722-006-0035-z
- Craven D, Hall J, Verjans JM (2009) Impacts of herbicide application and mechanical cleanings on growth and mortality of two timber species in *Saccharum spontaneum* grasslands of the Panama Canal watershed. *Restoration Ecology* 17: 751–761. doi: 10.1111/j.1526-100X.2008.00408.x
- D’Antonio CM, Vitousek PM (1992) Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annual Review of Ecology and Systematics* 23: 63–87.
- Daehler CC, Strong DR (1994) Variable reproductive output among clones of *Spartina alterniflora* (Poaceae) invading San Francisco Bay, California: the influence of herbivory, pollination, and establishment site. *American Journal of Botany* 81: 307–313. doi: 10.2307/2445457
- Daniels J, Roach BT (1987) Taxonomy and evolution. In: D Heinz (Ed) *Sugarcane improvement through breeding*. Elsevier Science Publishers B.V. (Amsterdam): 7–84. doi: 10.1016/B978-0-444-42769-4.50007-2
- Donohue K, Rubio de Casas R, Burghardt L, Kovach K, Willis CG (2010) Germination, postgermination adaptation, and species ecological ranges. *Annual Review of Ecology and Systematics* 41: 293–319.
- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19:11–15.
- Flores TA, Setterfield SA, Douglas MM (2005) Seedling recruiting of the exotic grass *Andropogon gayanus* (Poaceae) in Northern Australia. *Australian Journal of Botany* 53: 243–249. doi: 10.1071/BT03154
- Hammond BW (1999) *Saccharum spontaneum* (Gramineae) in Panama: The physiology and ecology of invasion. *Journal of Sustainable Forestry* 8: 23–38. doi: 10.1300/J091v08n03\_03
- Harradine AR (1980) The biology of African feather grass (*Pennisetum macrourum* Trin.) in Tasmania. II. Rhizome behaviour. *Weed Research* 20: 171. doi: 10.1111/j.1365-3180.1980.tb00064.x
- Holm LG, Plucknett D, Pancho J, Herberger J (1977) *The World’s Worst Weeds: distribution and biology*. (University of Hawaii Press, Honolulu, HI): 609 pp.

- Hooper ER, Condit R, Legendre P (2002) Responses of 20 native tree species to reforestation strategies for abandoned farmland in Panama. *Ecological Applications* 12: 1626–1641. doi: 10.1890/1051-0761(2002)012[1626:RONTST]2.0.CO;2
- Inderjit MO, Streibig JC (2001) Wheat (*Triticum aestivum*) interference with seedling growth of ryegrass (*Lolium perenne*): influence of density and age. *Weed Technology* 15: 807–812. doi: 10.1614/0890-037X(2001)015[0807:WTAIWS]2.0.CO;2
- Jones ER, Wishnie MH, Deago J, Sautu A, Cerazo A (2004) Facilitating natural regeneration in *Saccharum spontaneum* (L.) grasslands within the Panama Canal watershed: effects of tree species and tree structure on vegetation recruitment patterns. *Forest Ecology and Management* 191: 171–183. doi: 10.1016/j.foreco.2003.12.002
- Kettenring KM, Whigham DF (2009) Seed viability and seed dormancy of non-native *Phragmites australis* in suburbanized and forested watersheds of the Chesapeake Bay, USA. *Aquatic Botany* 91: 199–204. doi: 10.1016/j.aquabot.2009.06.002
- Kigel J, Koller D (1985) Asexual reproduction in weeds. In: Duke SO (Ed) *Weed Physiology*, Vol. I. Reproduction and Ecophysiology. CRC Press (Boca Raton, FL.), 65–100.
- Lambert AM, Dudley TL, Saltonstall K (2010) Ecology and impacts of the large-statured invasive grasses *Arundo donax* and *Phragmites australis* in North America. *Invasive Plant Science and Management* 3: 489–494. doi: 10.1614/IPSM-D-10-00031.1
- Lambrinos JG (2001) The expansion history of a sexual and asexual species of *Cortaderia* in California, USA. *Journal of Ecology* 89: 88–98. doi: 10.1046/j.1365-2745.2001.00524.x
- McCormick MK, Kettenring KM, Brown HM, Whigham DF (2010) Extent and reproductive mechanisms of *Phragmites australis* spread in brackish wetlands in Chesapeake Bay, Maryland (USA). *Wetlands* 30: 67–74. doi: 10.1007/s13157-009-0007-0
- Mukherjee SK (1957) Origin and distribution of *Saccharum*. *Botanical Gazette* 119: 55–61. doi: 10.1086/335962
- Murray BR (1998) Density-dependent germination and the role of seed leachate. *Australian Journal of Ecology* 23: 411–418. doi: 10.1111/j.1442-9993.1998.tb00748.x
- Pan YB (2006) Highly polymorphic microsatellite DNA markers for sugarcane germplasm evaluation and variety identity testing. *Sugar Technology* 8: 246–256. doi: 10.1007/BF02943564
- Panje RR (1970) The evolution of a weed. *PANS* 16: 590–595.
- Panje RR, Srinivasan K (1959) Studies in *Saccharum spontaneum*. The flowering behavior of latitudinally displaced populations. *Botanical Gazette* 120: 193–202. doi: 10.1086/336026
- Pichitkul P (2009) Aquatic plants in Chumphon and Surat Thani. *Proceedings of the 47<sup>th</sup> Kaetsart University Annual Conference*, 262–272.
- Poulin J, Weller SG, Sakai AK (2005) Genetic diversity does not affect the invasiveness of fountain grass (*Penisetum setaceum*) in Arizona, California, and Hawaii. *Diversity and Distributions* 11: 241–247. doi: 10.1111/j.1366-9516.2005.00136.x
- Saltonstall K, Bonnett GD (2012) Fire promotes growth and reproduction of *Saccharum spontaneum* (L.) in Panama. *Biological Invasions* 14: 2479–2488. doi: 10.1007/s10530-012-0245-6
- Sampietro DA, Soberon JR, Sgariglia MA, Quiroga EN, Vattuone MA (2007) Allelopathic Plants. 17. Sugarcane (*Saccharum officinarum* L.). *Allelopathy Journal* 20: 243–250.

- Setterfield SA, Rossiter-Rachor NA, Hutley LB, Douglas MM, Williams RJ (2010) Turning up the heat: the impacts of *Andropogon gayanus* (gamba grass) invasion on fire behaviour in northern Australian savannas. *Diversity and Distributions* 16: 854–861. doi: 10.1111/j.1472-4642.2010.00688.x
- Sidak ZK (1967) Rectangular confidence regions for the means of multivariate normal distributions. *Journal of the American Statistical Association* 62: 626–633.
- Singh S (1988) Effect of temperature on germination of sugar cane seeds (fluff). *Sugarcane* 1: 11–12.
- Skinner JC (1959) Fuzz germination in the photoperiod room in Meringa. *Sugarcane Breeder's Newsletter* 20: 2–6.
- Tai PYP, Miller JD (2001) A core collection for *Saccharum spontaneum* L. from the world collection of sugarcane. *Crop Science* 41: 879–885. doi: 10.2135/cropsci2001.413879x
- Turner DJ (1968) *Agropyron repens* (L.) Beauv. - some effects of rhizome fragmentation, rhizome burial and defoliation. *Weed Research* 8: 298. doi: 10.1111/j.1365-3180.1968.tb01434.x
- USDA (2010) <http://plants.usda.gov/java/noxious?rptType=federal>
- World Meteorological Organisation (2010) World Weather Information Service, Panama. <http://www.worldweather.org/076/c01221.htm>
- Wright SJ, Calderón O (2006) Seasonal, El Niño and longer term changes in flower and seed production in a moist tropical forest. *Ecology Letters* 9: 35–44.
- Yadav A, Balyan RS, Malik RK, Malik RS, Singh S, Banga RS, Singh S (2007) Efficacy of glyphosate, MON-8793 and MON-8794 for general weed control under non-cropped situations. *Environment and Ecology* 25: 636–639.
- Zamora DL, Thill DC, Eplee RE (1989) An eradication plan for plant invasions. *Weed Technology* 3: 2–12.

## Appendix I

Locations, description and diversity of the sites used for the collection of *Saccharum spontaneum* seeds. (doi: 10.3897/neobiota.20.6163.app1) File format: Adobe PDF file (pdf).

**Explanation note:** The table gives the latitude and longitude, description and number of genotypes found among the 3 plants of *Saccharum spontaneum* tested from each of 22 Sites. Sites 1–12 were used to assess seed germinability through time.

**Copyright notice:** This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

---

**Citation:** Bonnett GD, Kushner JNS, Saltonstall K (2014) The reproductive biology of *Saccharum spontaneum* L.: implications for management of this invasive weed in Panama. *NeoBiota* 20: 61–79. doi: 10.3897/neobiota.20.6163  
Locations, description and diversity of the sites used for the collection of *Saccharum spontaneum* seeds. *NeoBiota* 20: 61–79. doi: 10.3897/neobiota.20.6163.app1

---

## Appendix II

Germinability of *Saccharum spontaneum* seed through time from 12 sites. (doi: 10.3897/neobiota.20.6163.app2) File format: Adobe PDF file (pdf).

**Explanation note:** Proportion of seeds that germinated each week between September and December from samples taken at 12 sites. 100 seeds were germinated from each of three replicate samples and tested for germination in laboratory conditions. Results are presented as the mean and the error bar represents the standard error of the mean.

**Copyright notice:** This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

---

**Citation:** Bonnett GD, Kushner JNS, Saltonstall K (2014) The reproductive biology of *Saccharum spontaneum* L.: implications for management of this invasive weed in Panama. *NeoBiota* 20: 61–79. doi: 10.3897/neobiota.20.6163  
Germinability of *Saccharum spontaneum* seed through time from 12 sites. *NeoBiota* 20: 61–79. doi: 10.3897/neobiota.20.6163.app2

---