

# Chromosome numbers for the Italian flora: 14

Antonio Giacò<sup>1</sup>, Angelino Carta<sup>1</sup>, Jacopo Franzoni<sup>1</sup>, Alessio Mo<sup>1</sup>,  
Mario Pentassuglia<sup>1</sup>, Lorenzo Peruzzi<sup>1</sup>

<sup>1</sup> PLANTSEED Lab, Dipartimento di Biologia, Università di Pisa, Via Derna 1, 56126 Pisa, Italy

Corresponding author: Antonio Giacò ([antonio.giacco@biologia.unipi.it](mailto:antonio.giacco@biologia.unipi.it))

---

Academic editor: Giannantonio Domina | Received 4 June 2024 | Accepted 12 June 2024 | Published 24 June 2024

---

**Citation:** Giacò A, Carta A, Franzoni J, Mo A, Pentassuglia M, Peruzzi L (2024) Chromosome numbers for the Italian flora: 14. Italian Botanist 17: 69–75. <https://doi.org/10.3897/italianbotanist.17.128997>

---

## Abstract

In this contribution, new chromosome data obtained on material collected in Italy are presented. It includes the first counts for four subspecies of the Italian endemic *Centaurea aplolepa* Moretti, i.e. *C. aplolepa* subsp. *aplolepa*, *C. aplolepa* subsp. *bertolonii*, *C. aplolepa* subsp. *levantina*, and *C. aplolepa* subsp. *parvula*. In addition, the first chromosome count for an Italian population of *Silene canescens* (Caryophyllaceae) is provided.

## Keywords

Asteraceae, Caryophyllaceae, *Centaurea*, cytotaxonomy, endemism, *Silene*

## How to contribute


Texts concerning new chromosome data should be submitted electronically to Antonio Giacò ([antonio.giacco@biologia.unipi.it](mailto:antonio.giacco@biologia.unipi.it)), including indications on voucher specimens and methods used.

## Chromosome counts

*Centaurea aplolepa* Moretti subsp. *aplolepa* (Asteraceae)

**Chromosome number.**  $2n = 18$  (Fig. 1).

**Voucher specimen.** ITALY. Liguria. Capo Noli (Savona), sulle pareti rocciose (WGS84: 44.196751°N, 8.424978°E), 12 July 2023, A. Giacò, L. Peruzzi (PI066169).



**Figure 1.** *Centaurea aplolepa* Moretti subsp. *aplolepa* from Capo Noli (Savona),  $2n = 18$ . Scale bar: 10  $\mu\text{m}$ .

**Method.** Squash preparations were made on root tips obtained from germinating seeds. Root tips were pre-treated with 0.4% colchicine for 3 h and then fixed in Carnoy solution for 1 h. After hydrolysis in 1N HCl at 60 °C for 7.5 minutes, the tips were stained with leuco-basic fuchsin.

**Observations.** *Centaurea aplolepa* is endemic to the Tyrrhenian portion of central-northern Italy and includes ten subspecies (Greuter 2008). The autonymic subspecies is endemic to western Liguria, where it grows on garrigues and cliffs along the coastline (Arrigoni 2003). The chromosome number, obtained from seeds, is reported here for the first time, and is in accordance with chromosome counts previously published for other subspecies of *C. aplolepa* (Viegi et al. 1972; Viegi and Cela Renzoni 1976).


A. Giacò, L. Peruzzi

***Centaurea aplolepa* Moretti subsp. *bertolonii* (Arrigoni) Greuter (Asteraceae)**

**Chromosome number.**  $2n = 18$  (Fig. 2).

**Voucher specimen.** ITALY. Liguria. Italia, Genova, nei pressi di Forte Sperone (WGS84: 44.437325°N, 8.931887°E), 12 July 2023, A. Giacò, L. Peruzzi (PI066182).

**Method.** Squash preparations were made on root tips obtained from germinating seeds. Root tips were pre-treated with 0.4% colchicine for 3 h and then fixed in Carnoy



**Figure 2.** *Centaurea aplolepa* Moretti subsp. *bertolonii* (Arrigoni) Greuter from Genova,  $2n = 18$ . Scale bar: 10  $\mu\text{m}$ .

solution for 1 h. After hydrolysis in 1N HCl at 60 °C for 7.5 minutes, the tips were stained with leuco-basic fuchsin.

**Observations.** *Centaurea aplolepa* subsp. *bertolonii* is endemic to the hills over and west of Genova, showing a distribution range that is approximately in-between those of *C. aplolepa* subsp. *aplolepa* to the west and *C. aplolepa* subsp. *levantina* and *C. aplolepa* subsp. *lunensis* to the east (Arrigoni 2003). The chromosome number, obtained from seeds, is reported here for the first time.


A. Giacò, M. Pentassuglia, L. Peruzzi

### ***Centaurea aplolepa* subsp. *levantina* (Arrigoni) Greuter (Asteraceae)**

**Chromosome number.**  $2n = 18$  (Fig. 3).

**Voucher specimen.** ITALY. Liguria. Sestri Levante (Genova) (WGS84: 44.269873°N, 9.38648°E), piccoli affioramenti rocciosi presso il molo, A. Giacò, L. Peruzzi (PI066170).

**Method.** Squash preparations were made on root tips obtained from germinating seeds. Root tips were pre-treated with 0.4% colchicine for 3 h and then fixed in Carnoy solution for 1 h. After hydrolysis in 1N HCl at 60 °C for 7.5 minutes, the tips were stained with leuco-basic fuchsin.



**Figure 3.** *Centaurea aplolepa* Moretti subsp. *levantina* (Arrigoni) Greuter from Sestri Levante (Genova),  $2n = 18$ . Scale bar: 10  $\mu\text{m}$ .

**Observations.** According to Arrigoni (2003), *Centaurea aplolepa* subsp. *levantina* is a coastal ecotype endemic to eastern Liguria. The chromosome number, obtained from seeds, is reported here for the first time.

A. Giacò, M. Pentassuglia, L. Peruzzi


***Centaurea aplolepa* subsp. *parvula* (Ces.) Arcang. (Asteraceae)**

**Chromosome number.**  $2n = 18$  (Fig. 4).

**Voucher specimen.** ITALY. Piemonte. Acqui Terme, su calanchi (WGS84: 44.58492°N, 8.35745°E), 12 August 2023, A. Mo (PI066166).

**Method.** Squash preparations were made on root tips obtained from germinating seeds. Root tips were pre-treated with 0.4% colchicine for 3 h and then fixed in Carnoy solution for 1 h. After hydrolysis in 1N HCl at 60 °C for 7.5 minutes, the tips were stained with leuco-basic fuchsin.

**Observations.** Albeit *Centaurea aplolepa* subsp. *parvula* was not included by Arrigoni (2003) in his taxonomic revision of the *Centaurea paniculata* L. group, according to Bartolucci et al. (2024), this subspecies is endemic to Liguria and Piemonte. While in Liguria it is only reported for Rossiglione (Genova) (Gola 1912), the only record for Piemonte is represented by the type locality, at Acqui Terme



**Figure 4.** *Centaurea aplolepa* Moretti subsp. *parvula* (Ces.) Arcang. from Acqui Terme (Alessandria),  $2n = 18$ . Scale bar: 10  $\mu\text{m}$ .

(Alessandria). During a field trip, we found there several populations growing on badlands. The chromosome number, obtained from seeds, is reported here for the first time.

A. Giacò, A. Mo


### ***Silene canescens* Ten. (Caryophyllaceae)**

**Chromosome number.**  $2n = 24$  (Fig. 5).

**Voucher specimen.** ITALY. Toscana. Marina di Torre del Lago (Viareggio, Lucca), retroduna (WGS84: 43.829054°N, 10.253978°E), 15 Jun 2023, A. Mo (seeds collected and deposited at the germplasm bank of the Department of Biology, University of Pisa; a herbarium specimen collected from the same area in 2012 is conserved at PI under the barcode PI063832).

**Method.** Squash preparations were made on root tips obtained from germinating seeds. Root tips were pre-treated with 0.4% colchicine for 3 h and then fixed in Carnoy solution for 1 h. After hydrolysis in 1 N HCl at 60 °C for 8 minutes, the tips were stained with leuco-basic fuchsin.

**Observations.** *Silene canescens* Ten. is a psammophyte belonging to a species complex distributed throughout the sandy shores of southern Europe (Chater et al. 1993).



**Figure 5.** *Silene canescens* Ten. from Torre del Lago (Toscana),  $2n = 24$ . Scale bar: 10  $\mu\text{m}$ .

Difficulties to delineate its distribution range are due to the confusion with the closely related *S. colorata* Poir. (Valsecchi 1995). In Italy, *S. canescens* is considered taxonomically doubtful, but it occurs in most of the Thyrrhenian and Adriatic coasts (Bartolucci et al. 2024). Here we document the first chromosome count ( $2n = 2x = 24$ ) for an Italian population of this species, which is line with previous karyological data (Blackburn and Morton 1957), also concerning the related *S. colorata* (Rice et al. 2015).

J. Franzoni, A. Carta, A. Mo

## Acknowledgements

AG is thankful to G. Casazza for having provided precise locations of occurrence of *Centaurea* populations in Liguria.

## References

Arrigoni PV (2003) Le centauree italiane del gruppo "*Centaurea paniculata* L.". *Parlatorea* 6: 49–78.

- Bartolucci F, Peruzzi L, Galasso G, Alessandrini A, Ardenghi NMG, Bacchetta G, Banfi E, Barberis G, Bernardo L, Bouvet D, Bovio M, Calvia G, Castello M, Cecchi L, Del Guacchio E, Domina G, Fascetti S, Gallo L, Gottschlich G, Guarino R, Gubellini L, Hofmann N, Iberite M, Jiménez-Mejías P, Longo D, Marchetti D, Martini F, Masin RR, Medagli P, Peccenini S, Prosser F, Roma-Marzio F, Rosati L, Santangelo A, Scoppola A, Selvaggi A, Selvi F, Soldano A, Stinca A, Wagensommer RP, Wilhalm T, Conti F (2024) A second update to the checklist of the vascular flora native to Italy. *Plant Biosystems* 158(2): 219–296. <https://doi.org/10.1080/11263504.2024.2320126>
- Blackburn KB, Morton JK (1957) The incidence of polyploidy in the Caryophyllaceae of Britain and of Portugal. *New Phytologist* 56: 344–351. <https://doi.org/10.1111/j.1469-8137.1957.tb06979.x>
- Chater AD, Walters SM, Akeroyd JR [Eds] (1993) *Silene* L. In: Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, Webb DA (Eds) *Flora Europaea* (Vol. 4). Cambridge University Press, Cambridge, 191–218.
- Gola G (1912) La vegetazione dell'Appennino piemontese. *Annali di Botanica* 10: 189–338.
- Greuter W (2008) *Centaurea* L. In: Greuter W, von Raab-Straube E (Eds) *Med-checklist: a critical inventory of vascular plants of the circum-Mediterranean countries*. Organization for the Phyto Taxonomic Investigation of the Mediterranean Area (OPTIMA), Genève, 81–152.
- Hausknecht C (1894) Floristische Beiträge. *Mitteilungen des Thüringischen Botanischen Vereins* 6: 13–14.
- Rice A, Glick L, Abadi S, Einhorn M, Kopelman NM, Salman-Minkov A, Mayzel J, Chay O, Mayrose I (2015) The Chromosome Counts Database (CCDB) – a community resource of plant chromosome numbers. *New Phytologist* 206: 19–26. <https://doi.org/10.1111/nph.13191>
- Valsecchi F (1995) Indagini sistematiche, tassonomiche e corologiche nel gruppo “*Silene colorata* Poir., *S. sericea* All., *S. canescens* Ten.”. *Bollettino della Società Sarda di Scienze Naturali* 30: 447–476.
- Viegi L, Cela Renzoni G, Corsi G, Garbari F (1972) Numeri cromosomici per la flora italiana: 135–144. *Informatore Botanico Italiano* 4(3): 229–236.
- Viegi L, Cela Renzoni G (1976) Numeri cromosomici per la flora italiana: 276–282. *Informatore Botanico Italiano* 8(3): 276–281.