

Two new species of *Pambolus* (Hymenoptera, Braconidae) from Jamaica

Juan José Martínez^{1,†}, Fadia Sara Ceccarelli^{1,‡}, Alejandro Zaldívar-Riverón^{1,§}

I Colección Nacional de Insectos, Instituto de Biología, Universidad Nacional Autónoma de México, 3er. circuito exterior s/n, Cd. Universitaria, Copilco Coyoacán, A. P. 70-233, C. P. 04510., D. F., México

† [urn:lsid:zoobank.org:author:D27E443D-CBAE-4229-BB98-E9F375604407](https://zoobank.org/D27E443D-CBAE-4229-BB98-E9F375604407)

‡ [urn:lsid:zoobank.org:author:E0FC81B0-2F57-4253-9608-C970E5570BB8](https://zoobank.org/E0FC81B0-2F57-4253-9608-C970E5570BB8)

§ [urn:lsid:zoobank.org:author:B6E09990-6CA5-403F-AC3F-9498DBB5BEDF](https://zoobank.org/B6E09990-6CA5-403F-AC3F-9498DBB5BEDF)

Corresponding author: Juan José Martínez (jjmartinez80@hotmail.com)

Academic editor: G. Broad | Received 31 October 2011 | Accepted 8 December 2011 | Published 10 January 2012

[urn:lsid:zoobank.org:pub:B1A8A488-CBD2-4152-9C7C-E9294FC6B942](https://zoobank.org/pub/B1A8A488-CBD2-4152-9C7C-E9294FC6B942)

Citation: Martínez JJ, Ceccarelli FS, Zaldívar-Riverón A (2012) Two new species of *Pambolus* (Hymenoptera, Braconidae) from Jamaica. Journal of Hymenoptera Research 24: 85–93. doi: 10.3897/JHR.24.2300

Abstract

Pambolus albospina sp. n. and *Pambolus rastafari* sp. n. are described from Jamaica. *Pambolus rastafari* can be distinguished from all other Neotropical species of the genus by its antennal color pattern and the smooth sculpture of the head and mesoscutum. *Pambolus albospina* is morphologically close to *P. hemitaeniatus* van Achterberg, from which it can be distinguished by the number of white antennal segments and the sculpture on the first metasomal tergite. These two new species constitute the first records of the genus for the Caribbean Islands.

Keywords

Caribbean, Hormiinae, Pambolinae, Taxonomy

Introduction

Pambolus Haliday is a braconid genus with 42 currently valid species from all biogeographic regions (Yu et al. 2005). It has been included in the subfamily Hormiinae (Whitfield and Wharton 1997) or in the small subfamily Pambolinae (van Achterberg 1995; Braet and van Achterberg 2003; van Achterberg and Braet 2004; Yu et al. 2005). In this paper we follow the latter classification since Whitfield and Wharton's (1997)

concept of Hormiinae includes a heterogeneous array of genera and does not represent a natural group as revealed by a previous phylogenetic study of the cyclostome subfamilies of Braconidae (Zaldívar-Riverón et al. 2006). Currently, two subgenera are recognized within *Pambolus*, the typic subgenus includes all species with reduced wings, whereas *Phaenodus* gathers all macropterous forms (Braet and van Achterberg 2003).

In the Neotropical Region, *Pambolus* includes 12 described mainland species that are distributed from Honduras to northern Argentina (Yu et al. 2005). Specimens of *Pambolus* are fairly commonly observed in hymenopteran samples, though its species are not particularly abundant (Shaw and Huddelston 1991). Nine of the twelve recognized Neotropical species were described based on single females, and only one, *P. hebes* Papp, is known from both sexes (Papp 1996). The biology of species of *Pambolus* is largely unknown. Shaw and Huddelston (1991) report that a European species of *Pambolus* has been associated with chrysomelids (Coleoptera), although this information needs to be confirmed. The aim of this work is to describe two new species of *Pambolus* recently collected in Jamaica, which represent the first records of the genus for the Caribbean Islands.

Methods

Specimens were collected using yellow pan traps at two localities in Jamaica.

Morphological terminology follows Sharkey and Wharton (1997), surface sculpture terminology follows Harris (1979). Descriptions of the new species are organized following the basic format of recent descriptions of *Pambolus* species (Braet and van Achterberg 2003; van Achterberg and Braet 2004) in order to facilitate comparisons. Photographs were taken and edited using a Leica® Z16 APO-A stereoscopic microscope, a Leica® DFC295/DFC290 HD camera, and the Leica Application Suite® program. DNA Barcode sequences [~650 bp of the cytochrome oxidase I (COI) mitochondrial DNA gene; Hebert et al. 2003] were generated for specimens of the two new species using the same DNA extraction and amplification protocols employed by Ceccarelli et al. (in press).

Specimens are deposited at Colección Nacional de Insectos (CNIN), Instituto de Biología, Universidad Nacional Autónoma de México, and at Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Buenos Aires, Argentina (MACN).

Results


Pambolus albospina sp. n.

urn:lsid:zoobank.org:act:19BA7908-B656-4B6E-A661-F9C68582B370

http://species-id.net/wiki/Pambolus_albospina

Figures 1–6

Diagnosis. Following the key to Neotropical species (Braet and van Achterberg 2003), the female of this new species runs to *P. hemitaeniatus* van Achterberg, but differs from



Figures 1–6. *Pambolus albospina* sp. n. **1** Habitus of female (lateral view) **2** head (anterior view) **3** head, mesosoma and first metasomal segment (dorsal view) **4** head, mesosoma and first metasomal segment (lateral view) **5** antenna (dorsal view) **6** wings.

the latter and all other Neotropical species of the subgenus *Phaenodus* by having an entirely smooth first metasomal tergite without striations, and by its entirely white propodeal spines. Males, however, are difficult to relate to those of the other species. They can be distinguished from *P. longicornis* (Enderlein)-the only neotropical species of the subgenus *Phaenodus* with known males- also by the color of the propodeal spines and the smooth first metasomal tergite.

Female. Color: Mostly reddish brown (fig. 1); scape and base of pedicel pale yellow, apex of pedicel and first five flagellomeres honey yellow, sixth to eleventh flagellomeres gradually darkening from brownish yellow to black, further flagellomeres (12th to 24th) entirely white. Palpi white; lower area of temple, gena, face, clypeus and labrum reddish brown; frons, vertex and upper area of temple orange brown, ocellar triangle darkened. Mesoscutum, scutellum and propodeum except propodeal spines, orange brown; pronotum, mesopleuron and metapleuron darker, rusty brown; propodeal spines entirely white. First metasomal tergite reddish brown, remainder of metasoma brown. Coxae, trochanters and trochantelli white, femora white basally and brown on apical third fourth, tibiae whitish, tarsi light brown. Fore wing with faint infusate bands, hind wing hyaline, pterostigma and veins brown, except vein r-m and junction of veins 2RS and 2M, which are not pigmented.

Head: Antenna (fig. 5) with 24 flagellomeres, first flagellomere 1.1 times longer than second, first and second flagellomeres 5.5 and 5.0 times longer than their maximum width respectively. Scapus oblique apically; in dorsal view (fig. 3) length of eyes 1.9 times temple; POL 1.5 times OD and 0.7 times OOL; in dorsal view width of head 1.7 times its maximum length; face flattened and clypeus weakly convex in lateral view; face (fig. 2) weakly coriaceous near toruli, turning smooth near clypeus; clypeus largely smooth; anterior tentorial pit big and circular; frons, vertex and temple rugulose; gena smooth and shining (fig. 4); malar space 1.5 times basal width of mandible; occipital carina complete, meeting hypostomal carina ventrally.

Mesosoma: Length of mesosoma 1.4 times its maximum height and 1.6 times its maximum width; mesosoma setose; pronotum largely smooth, pronotal furrow deep and scrobiculate; propleuron convex and weakly coriaceous; mesopleuron smooth (fig. 4), precoxal sulcus deep and scrobiculate, occupying two thirds length of mesopleuron, posterior mesopleural furrow strongly scrobiculate; subalar area rugose; episternal scrobe deep; prepectal and postpectal carinae strongly developed; metapleuron reticulate; mesoscutum (fig. 3) acinose-coriaceous; notauli wide and scrobiculate anteriorly, rather obscured in a median area posteriorly; scutellum coriaceous and slightly convex; scutellar sulcus with five carinae, its median length 0.6 times as long as scutellum; propodeum areolate rugose, with a long and narrow areola medially (fig. 3), areola irregularly rugose anteriorly and with four transverse carinae posteriorly, spines of propodeum 0.8 times as long as fore basitarsus.

Wings: Wings (fig. 6) densely setose. Fore wing: veins r and 3RSa basally widened; r:3RSa:3RSb = 2:5:16; 2RS:2M:r-m = 6:10:3; vein (RS+M)a straight; vein cu-a just interstitial and almost indistinct; first subdiscal cell open. Hind wing: vein cu-a oblique, directed towards wing base; M+CU:1-M = 2:5.

Legs: Hind coxa unsculptured dorsally; length of femur, tibia and basitarsus of hind leg 4.4, 7.6 and 6.0 times their width, respectively; hind femur very weakly coriaceous.

Metasoma: Length of first tergite 0.7 times its apical width, its median area well delimited by carinae but entirely smooth, lateral areas also smooth (fig. 3); second metasomal tergum with a few basal striations, otherwise smooth, remaining terga completely smooth; combined length of second and third tergites as long as their maximum width; length of ovipositor sheath 0.4 times the length of metasoma.

Male Similar to female, except for its slightly lighter body color and its much longer antennae, with 31 flagellomeres, five apical flagellomeres entirely white.

Etymology. The specific epithet refers to the white propodeal spines of this species.

Material examined. Holotype female: Jamaica, Trelawny, Windsor, 18.35752, -76.66406, 82m, 19–21.xi.2010, yellow pan traps, F. S. Ceccarelli col. GenBank accession no. JQ268751. (CNIN IB-UNAM) Paratypes: three males, Jamaica, Saint Andrew, New Castle, 18.06840, -76.7119, 860m, 11.xi.2010, F. S. Ceccarelli col. (CNIN IB-UNAM, MACN); GenBank accession no. JQ268752.

Pambolus rastafari sp. n.

[urn:lsid:zoobank.org:act:76DBA72C-1920-4BF3-860E-860B88A1739F](http://www.zoobank.org/act:76DBA72C-1920-4BF3-860E-860B88A1739F)


http://species-id.net/wiki/Pambolus_rastafari

Figures 7–15

Diagnosis. Following the keys to Neotropical species of *Pambolus* (Braet and van Achterberg 2003; van Achterberg and Braet 2004), *P. rastafari* runs to couplet three due to the presence of a white basal antennal band; however, it can easily be distinguished from *P. duplotaeniatus* van Achterberg, *P. granulatus* van Achterberg and *P. microstriatus* van Achterberg by the basal white band being composed of a single white flagellomere. The three aforementioned species have a white band composed of at least three flagellomeres. This new species also differs from the above three species by its entirely smooth and polished head and mesoscutum.

Female. *Color*: Mostly chestnut brown (fig. 7); palpi, pedicel, second flagellomere, coxae, trochanters and trochantelli, all femora and all tibiae basally ivory white; flagellomeres beyond 12th and fore tarsus light brown; annellus of first flagellomere and apex of all trochantelli distinctly orange-brown; wings almost entirely infusate, with subhyaline areas at junction of veins 2RS and 2M and along vein r-m; remaining veins light brown.

Head: Antenna broken, 13 flagellomeres remaining, first flagellomere 1.2 times longer than second, first and second flagellomeres 5.0 and 3.9 times longer than their maximum width, respectively. Scapus oblique apically. In dorsal view (fig. 9) length of eyes 1.9 times the length of temple; POL as long as OD and 0.4 times OOL; in dorsal view width of head 0.6 times its median length; face rather flattened and clypeus weakly convex in lateral view (fig. 10); face smooth and setose (fig. 8); clypeus smooth; anterior tentorial pit big and circular; frons smooth and slightly concave; vertex, tem-



Figures 7–13. *Pambolus rastafari* sp. n. **7** habitus of female (lateral view) **8** head (anterior view) **9** head (dorsal view) **10** head and basal antennomeres (lateral view) **11** mesosoma (lateral view) **12** mesoscutum and sternalum (dorsal view) **13** propodeum and first metasomal segment (dorsal view).



Figures 14–15. *Pambolus rastafari* sp. n. **14** fore wing **15** hind wing.

ple and gena also smooth and shining; malar space 2.0 times basal width of mandible; occipital carina complete, strongly developed, meeting hypostomal carina ventrally.

Mesosoma: Length of mesosoma 1.4 times its maximum height and 1.6–1.7 times its maximum width; mesosoma setose; pronotum largely smooth, slightly coriaceous anteriorly, pronotal furrow present only medially and scrobiculate; propleuron convex and smooth; mesopleuron smooth (fig. 11), precoxal sulcus deep and scrobiculate, occupying two thirds the length of the mesopleuron, subalar and posterior mesopleural furrows scrobiculate; prepectal and postpectal carinae strongly developed; metapleuron irregularly rugose; mesoscutum smooth and shining (fig. 12); notauli wide and crenulate on anterior edge of mesoscutum, indistinct posteriorly; mesoscutum with a median V-shaped posterior pit; scutellar disc smooth; scutellar sulcus with five coarse and oblique carinae, its median length 0.5 times as long as scutellar disc; propodeum (fig. 13) areolate rugose, with a long and narrow areola medially, areola with a median longitudinal carina anteriorly and with three transverse carinae posteriorly, spines of propodeum 0.7 times as long as fore basitarsus.

Wings: Wings densely setose. Fore wing (fig. 14): veins r and 3-SR basally widened; $r:3RSa:3RSb = 3:8:12$; $2RS:2M:r-m = 9:17:5$; veins (RS+M)b and r-m unpigmented; vein (RS+M)a straight; vein cu-a interstitial; first subdiscal cell open. Hind wing (fig. 15): vein cu-a vertical; $M+CU:1-M = 2:5$.

Legs: Hind coxa largely finely striate dorsally; length of femur, tibia and basitarsus of hind leg 4.4, 8.4 and 5.3 times their width, respectively; hind femur very weakly coriaceous.

Metasoma: Length of first tergite 0.8 times its apical width, its median area wide and coarsely longitudinally striate (fig. 13), lateral areas smooth; remaining terga completely smooth and shining; combined length of second and third tergites as long as their maximum width; length of ovipositor sheath 0.4–0.5 times the length of metasoma.

Male. Unknown.

Etymology. The specific epithet refers to the religious and social movement from Jamaica.

Material examined. Holotype female: Jamaica, Saint Andrew, New Castle, 18.06840, -76.71119, 860m, 11.xi.2010, F. S. Ceccarelli col. (CNIN IB-UNAM); GenBank accession no. JQ268750. Paratype: one female, same data as holotype (MACN).

Acknowledgments

Grateful thanks are extended to Susana Guzmán Gómez (UNIBIO, IB- UNAM, Mexico), for her assistance in taking the digital images. To Dwight Robinson from UWI Mona, Kingston, Jamaica, for his assistance during our collecting trip. We also thank two anonymous reviewers for their helpful comments on an earlier version of the manuscript. This work was supported by postdoctoral fellowships given by the Universidad Nacional Autónoma de México (UNAM; DGAPA fellowships) to JJM, FSC and SFC, and by grants given by the Consejo Nacional de Ciencia y Tecnología [proyecto Ciencia Básica CONACyT no. 511 and Programa de Cooperación Bilateral México-Argentina (CONACyT-CONICET)] and by the Ministerio de Ciencia e Innovación (CGL2010-15786; Spain) to AZR.

References

- Achterberg C van (1995) Generic revision of the subfamily Betylobraconinae (Hymenoptera: Braconidae) and other groups with modified fore tarsus. *Zoologische Verhandelingen* 298: 1–242.
- Achterberg C. van, Braet Y (2004) Two new species of *Pambolus* Haliday (Hymenoptera: Braconidae: Pambolinae) from Argentina. *Zoologische Mededelingen* 78: 337–344.
- Braet Y, Achterberg C van (2003) New species of *Pambolus* Haliday and *Phaenocarpa* Foerster (Hymenoptera: Braconidae: Pambolinae, Alysiinae) from French Guyana, Suriname and Panama. *Zoologische Mededelingen Leiden* 77: 153–179.
- Ceccarelli FS, Sharkey MJ, Zaldivar-Riverón A (in press) Species identification in the taxonomically neglected, highly diverse, neotropical parasitoid wasp genus *Notiospathius* (Braconidae: Doryctinae) based on an integrative molecular and morphological approach. *Molecular Phylogenetics and Evolution*.
- Harris RA (1979) A glossary of surface sculpturing. *Occasional Papers of Entomology, California Department of Food and Agriculture*, no. 28, 31 pp.

- Hebert PDN, Ratnasingham S, deWaard JR (2003) Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society B* 270: S96–S99. doi: 10.1098/rsbl.2003.0025
- Papp J (1996) On the genus *Pambolus* (Hymenoptera: Braconidae: Pambolinae), with description of four new tropical species. *Acta Zoologica Academiae Scientiarum Hungaricae* 42: 41–57.
- Sharkey MJ, Wharton RA (1997) Morphology and terminology. In: Wharton RA, Marsh PM, Sharkey MJ (Eds) *Manual of the New World genera of the family Braconidae (Hymenoptera)*. Special Publication of the International Society of Hymenopterists no 1, p. 19–37.
- Shaw MR, Huddleston T (1991) Classification and biology of braconid wasps (Hymenoptera: Braconidae). *Handbooks for the Identification of British Insects* 7: 1–126.
- Whitfield JB, Wharton RA (1997) Subfamily Hormiinae. In: Wharton RA, Marsh PM, Sharkey MJ (Eds) *Manual of the New World genera of the family Braconidae (Hymenoptera)*. Special Publication of the International Society of Hymenopterists no. 1, p. 291–307.
- Yu DSK, Achterberg C van, Horstmann K (2005) *World Ichneumonoidea 2004. Taxonomy, Biology, Morphology and Distribution [CD/DVD]*. Taxapad, Vancouver.
- Zaldívar-Riverón A, Mori M, Quicke DLJ (2006) Systematics of the cyclostome subfamilies of braconid parasitic wasps (Hymenoptera: Ichneumonoidea): a simultaneous molecular and morphological Bayesian approach. *Molecular Phylogenetics and Evolution* 38: 130–145. doi: 10.1016/j.ympev.2005.08.006