

RESEARCH ARTICLE

Effects of parasites and predators on nociception: decreases analgesia reduces overwinter survival in root voles (Rodentia: Cricetidae)

Yuan-Gang Yang^{1,2}, Guo-Zhen Shang^{1,2}, Xue-Qin Wu^{1,2}, Hui-Qing Chen^{1,2},
 Yan Wu³, Yi-Fan Cao¹, Jiang-Hui Bian^{1,4}

¹Key Laboratory of Adaptation and Evolution of Plateau Biota, Northwest Institute of Plateau Biology, Chinese Academy of Sciences. Xining 810001, China.

²Graduate University of Chinese Academy of Sciences. Beijing 100049, China.

³School of Life and Environment Sciences, Hangzhou Normal University. Hangzhou 310012, China.

⁴Qinghai Key Laboratory of Animal Ecological Genomics. Xining 810001, China.

Corresponding authors: Jiang-hui Bian (bjh@nwipb.cas.cn), Yan Wu (wuyanqh@163.com)

<http://zoobank.org/A3EEA8A3-3CC2-461B-B351-CD39BE778C3B>

ABSTRACT. Growing evidence suggests that parasite-infected prey is more vulnerable to predation. However, the mechanism underlying this phenomenon is obscure. In small mammals, analgesia induced by environmental stressors is a fundamental component of the defensive repertoire, promoting defensive responses. Thus, the reduced analgesia may impair the defensive ability of prey and increase their predation risk. This study aimed to determine whether coccidia infection increases the vulnerability to predation in root voles, *Microtus oeconomus* (Pallas, 1776), by decreased analgesia. Herein, a predator stimulus and parasitic infection were simulated in the laboratory via a two-level factorial experiment, then, the vole nociceptive responses to an aversive thermal stimulus were evaluated. Further, a field experiment was performed to determine the overwinter survival of voles with different nociceptive responses via repeated live trapping. The coccidia-infected voles demonstrated reduced predator-induced analgesia following exposure to predator odor. Meanwhile, pain-sensitive voles had lower overwinter survival than pain-inhibited voles in enclosed populations throughout the duration of the experiment. Our findings suggest that coccidia infection attenuates predator-induced analgesia, resulting in an increased vulnerability to predation.

KEY WORDS. Analgesic response, coccidian infection, predation effect, small mammal.

INTRODUCTION

In nature, predators and parasites constitute the two primary extrinsic population regulators and play important roles in prey/host population dynamics (Sundell 2006, Tompkins et al. 2011). Predation can increase parasite infection by changing the phenotypic traits of the prey (i.e., morphological and physiologic traits) (Duffy et al. 2011, Caetano et al. 2014, Shang et al. 2019). In turn, increased parasite infection results in hosts vulnerability to predation (Carreon and Faulkes 2014, Shang et al. 2019, Gooding et al. 2020). For example, Møller and Nielsen (2007) showed that prey species with high malaria prevalence have higher predation risk than those with a low prevalence. Moose, *Alces alces* (Linnaeus, 1758), and voles, *Microtus townsendii* (Bachman, 1839), are very different in size, but both are more prone to predation when they have heavy parasite burden (Steen et al. 2002, Joly and Messier 2004). Several possible mechanisms

for the synergistic effects of parasites and predators on host/prey mortality have been proposed, such as deteriorated body condition (Wirsing et al. 2002, Hoey and McCormick 2004), reduced escape ability (Alzaga et al. 2008), and increased metabolism and energy output (Haye and Ojeda 1998, Krams et al. 2013). However, the underlying mechanism by which parasite-infected prey are more vulnerable to predation is obscure.

Animals respond to the threat of predation via a series of defensive responses, including flight, freezing, risk assessment, increased alertness and fear, or analgesia (Lima and Dill 1990, Kavaliers and Colwell 1994). In small mammals, analgesia induced by environmental stress factors is a fundamental component of the defensive repertoire, promoting the coordinated expression of other defensive behaviors (Colwell and Kavaliers 1993, Caio 2011). Thus, the decreased analgesia may impair the defensive ability of small mammals and then increase their vulnerability to predation (Ives and Dobson 1987, Tambeli et al. 2012, Lamana et al. 2018).

Experimental evidence from laboratory has shown that predator or predator cues could activate the analgesic system in mice and rats (Kavaliers and Colwell 1991, 1994). Meanwhile, endoparasitic infections such as coccidia or nematodes reduce this analgesia when the rodents are exposed to predator stimuli (Kavaliers et al. 2000). Notably, most studies on nociceptive responses have been performed on mice and rats under laboratory conditions. Whether parasitic infections decrease analgesia in wild rodents exposed to predators and if this affects their population survival remains unclear.

Our previous study showed that the combined effects of coccidia infection and predators decrease the overwinter survival of root voles, *Microtus oeconomus* (Pallas, 1776) (Shang et al. 2019). Coccidia are the most prevalent parasites in root voles (Cao et al. 2014, Nie et al. 2014). The current study aims to evaluate whether coccidia infection increases the predation vulnerability of root voles via decreased analgesia. This work builds on the previous (Shang et al. 2019) to test two hypotheses: 1) coccidia infection reduces predator-induced analgesia in root voles; 2) individuals with reduced analgesia have lower overwinter survival.

MATERIAL AND METHODS

Statement of animal right

The use of animals in this study was in accordance with the guidelines of the regulations of experiments on animals and was approved by the animal Ethics and Welfare committee of the Northwest Institute of Plateau Biology, Chinese Academy of Science.

Laboratory experiments

The laboratory experiments were conducted at the Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining, China. Root voles were housed singly in clear polyethylene cages ($36 \times 20 \times 17 \text{ cm}^3$) with wood shavings and maintained at $20 \pm 2 \text{ }^\circ\text{C}$ under a 12:12 h light: dark cycle. Food and water were available ad libitum. Twenty voles, six months and older, of each sex from a laboratory colony were divided into two groups: coccidia-infected (hereafter PA+) and parasite-free groups (hereafter PA-). Half of the PA+ and PA- groups were exposed to predator odor (hereafter PR+PA+ or PR+PA-), and the other half to a control odor (hereafter PR-PA+ or PR-PA-). Each of the four treatments involved five voles per sex, and the initial vole body mass of the four treatments did not differ ($F_{3,36} = 0.187, p = 0.904$).

Parasite infection

Voies in the PA+ group were once orally administered with 2000 coccidia oocysts suspended in 0.1 mL distilled water on June 3rd, 2019. Their oocyte levels were comparable to the oocysts per gram in the feces of coccidia-infected root voles studied by Shang et al. (2019). Meanwhile, each vole in the PA- group was treated with a single orogastric gavage dose of 0.1 mL combina-

torial anthelmintic comprising 6.25×10^{-4} mL diclazuril solution (Weierkong, Sichuan) and a 2 mg ivermectin tablet (Weierkong). Combinatorial anthelmintics can effectively expel nematodes, cestodes, and coccidia (Yang et al. 2018).

Our pilot study found that the latency period for coccidia infection in root voles was 6–7 days, and the maximum oocyst output occurred 9–10 days post-infection. Accordingly, we measured nociceptive responses on June 13th, 2019.

Predator odor exposure

Voies were exposed to predator or control odors on June 4–13th, 2019. Silver fox, *Vulpes vulpes* (Linnaeus, 1758), odor was used to stimulate predation risk, while the rabbit odor, *Oryctolagus cuniculus f. domesticus* (Linnaeus, 1758) was used as control (Wang and Liu 2002a, Bian et al. 2005b). Since the major predators in the study area – *Buteo hemilasius* Temminck & Schlegel, 1844 or *Mustela altaica* Pallas, 1811 – are protected animals in China, capturing and collecting fresh feces and urine for 15 consecutive days was challenging. Wang and Liu (2002a, 2002b) found that the silver fox odor could change the behavioral responses of root voles. Thus, our study used silver foxes for predator stimulation instead of natural predators.

Fresh silver fox and rabbit feces and urine were collected in trays under the animal cages daily. Each tray was washed with 500 mL distilled water, and the washing water strained through a filter with a 100 mesh screen (Bian et al. 2005a). Filtered solutions from each animal species collected at different times were thoroughly mixed. At the onset of the laboratory experiment, filter papers infused with predator or control odors were randomly placed in the vole cages three to four times a day between 8:00 am and 11:00 pm. This period was chosen because root voles are primarily diurnal (Sun et al. 1982). Each exposure to predator or control odor lasted 30–60 s.

Nociceptive responses

The nociceptive responses of voies were measured on June 3rd, 2019, prior to parasitic infection. The initial nociceptive response latency did not differ among voies in the four treatments ($F_{3,36} = 0.165, p = 0.919$). Nociception was measured based on the latency of foot-lifting or licking responses to an aversive thermal stimulus (“hot plate,” CAT.NO.T-91-S, CT, USA). Each measurement was replicated thrice in each individual. The individual was immediately removed from the heated surface following the response display and returned to its cage. If no response was observed within 60 s, the test was terminated, and the vole returned to its cage (Kavaliers and Colwell 1994). In the present study, all voies displayed nociceptive responses within 60 s.

Field experiments

Field experiments were conducted at the Haibei Alpine Meadow Ecosystem Research Station, Menyuan County, approximately 155 km north of Xining, Qinghai Province, China (37°37'N, 101°12'E). The station has an elevation of 3200 m,

is surrounded by mountains, and has an average annual temperature and precipitation of -1.6°C and 560 mm, respectively (Li et al. 2004).

Root vole populations in this area fluctuate annually, usually with relatively low numbers in late winter and spring, increasing throughout the breeding season, and declining after the breeding season; multi-year cycles are weak or absent (Jiang et al. 1991). In this study area, root voles prefer dense vegetation, mainly *Elymus nutans* (Poaceae), in habitat selection. The average population in the habitat ranged from 217 to 280 voles ha^{-1} in autumn, even up to 400 ha^{-1} where grazing activities were limited (Bian et al. 1994, Sun et al. 2002). The breeding season typically lasts from April to October. Juveniles reach puberty and breeding age at approximately 50 and 70 days, respectively (Liang et al. 1982). The primary predators in the study area are falcons, *Falco tinnunculus* Linnaeus, 1758, buzzards, *Buteo hemilasius* Temminck & Schlegel, 1844, and weasels, *Mustela altaica* Pallas, 1811.

Experimental facility

The field experiments were carried out in four 0.15 ha (50×30 m) outdoor enclosures located in an old *E. nutans* meadow. Major plants included *E. nutans*, *Poa* spp., *Thalictrum alpinum*, and *Kobresia humilis*. The vegetative cover provided a dense leaf layer, forming a natural refuge for root voles. The enclosures were constructed using galvanized steel panels (1.5 and 0.5 m above and below ground, respectively) but without wire mesh roofs. Further, the enclosures had a series of low panels (~ 0.3 m high) along the exterior walls every 10 m, allowing terrestrial and avian predators to enter but prevented voles from exiting the enclosures. The vegetation conditions are similar in each enclosure. Each enclosure was equipped with 60 laboratory-made wooden traps (Bian et al. 2011), spaced in 5×5 m grids.

Founder populations

Forty-eight voles of each sex, six months or older, from a laboratory colony were used to establish founder populations on October 16, 2017. The voles were divided into two nociception levels according to thermal response latency ("hot plate," CAT. NO.T-91-S, CT, USA); high response latency group (hereafter group H; 53.92 ± 0.11) and low response latency group (hereafter group L; 49.3 ± 0.09). The nociceptive response latency of group H was significantly higher than group L ($F_{1,94} = 1011.93$, $p < 0.001$). Earmarked voles from group H were released into two enclosures, while earmarked voles from group L were released into the other two enclosures. Each enclosure contained 12 voles per sex, and each treatment was conducted in duplicate. The density of the founder population (160 voles ha^{-1}) was in line with natural densities in autumn (Bian et al. 1994, Sun et al. 2002).

Prior to the experiment, all voles were treated with a combinatorial anthelmintic to eliminate parasites and ensure homogeneity. Besides, all enclosures were trapped for two weeks to remove small resident mammals. We also ensured the

initial vole body mass did not differ between the enclosures ($F_{1,94} = 0.004$, $p = 0.95$).

Vole trapping

Live trapping began on October 28th, 2017, after the voles had acclimated to the enclosures for two weeks, and lasted for 141 days (at the end of March 17th, 2018). Standard capture-recapture methods were used throughout the present study. Six trapping sessions were conducted, each consisting of three trapping days. The time interval between two trapping sessions was approximately one month. Each trap was baited with carrots, set between 8:00 am and 5:30 pm, checked every two hours and closed when trapping did not occur. Following capture, the individual was identified and their sex recorded.

Survival rate and population size estimations

We estimated the apparent survival (hereafter "survival") and recapture probability (hereafter "recapture") using the standard open population Cormack-Jolly-Seber model (Lebreton et al. 1992) in the MARK program (White and Burnham 1999). The recapture probability was evaluated under the assumption that the individual was alive and in the sample (Cooch and White 2006). The data comprised a capture history of 96 voles in six trapping sessions from October 28th, 2017, to March 17th, 2018. First, RELEASE in the MARK program was used to conduct a goodness-of-fit test for the global models, namely $\Phi\text{TR} \times \text{T}$, with both vole survival and recapture dependent on treatment, "TR," and time, "T." The goodness-of-fit of the global model was assessed by testing the assumptions of independence and homogeneity of individuals in the enclosures. The goodness-of-fit tests were not significant for voles (tests 2 and 3, RELEASE: $\chi^2 = 12.64$, $\text{df} = 18$, $p = 0.81$), suggesting that voles in the enclosures were independent and that the model fits were acceptable. We then used a bootstrap-based goodness-of-fit test to estimate the \hat{c} -hat value (a variance inflation factor; 1.92), which was adjusted to 1.92 in the global models.

Second, we selected the models as described in our previous study (Yang et al. 2018). Briefly, parsimonious models were selected based on the QAICc values, which allows a compromise between bias and precision when the global model does not fit the data (Anderson et al. 1994) and incorporates a variance inflation factor. Third, we tested the hypothesis that nociception influences vole overwinter survival. For the test, a parsimonious model containing the treatment factor was compared with neighboring populations without the factor, using QAICc values. Subsequently, the model average was estimated from the mean monthly apparent survival probability.

We used the minimum number known alive method to estimate population sizes across trapping sessions in each enclosure. Mark-recapture sampling trials of known populations in the enclosures showed that the minimum number known alive was the best estimate of the actual population size relative to other estimators (Chambers et al. 1999, Bian et al. 2011). The rate of population change for each enclosure was calculated using the

following equation: $r_t = (1/T) \ln (N_{t+1}/N_t)$, where N_t is the population density at time t , N_{t+1} is the population density during the subsequent trapping session, i.e., at time $t + 1$, while T is the time interval between trapping sessions (Klemola et al. 2002).

Statistical analysis

The vole population size (Poisson distribution) was analyzed using generalized linear mixed models, with log link functions in the SPSS v. 20 program (IBM, Armonk, NY, USA). Continuous variables were analyzed using a linear model. Data sampled repeatedly were analyzed using the repeated measures method, and all models were simplified by eliminating non-significant ($p > 0.05$) interactions. Post hoc comparisons of significant effects were computed using the sequential Bonferroni post hoc procedure.

In the analyses of nociceptive responses, treatments were input into the models as fixed factors, and individual IDs were put as the random factors. In the analyses of population change rate and density, treatments and trapping sessions were input as fixed factors to test the primary and interaction effects separately. Meanwhile, enclosures were input as random factors. Since no sex differences were found for any parameter, the data for males and females were pooled during analyses.

RESULTS

Laboratory experiments

We found an effect of treatment on vole nociceptive responses ($F_{3,34} = 8.89, p < 0.001$). Compared with uninfected voles exposed to the control odor (50.70 ± 0.41 ; range from 49.07 to 53.7), uninfected voles exposed to the predator odor had increased nociceptive latencies (53.65 ± 0.41 ; range from 51.73 to 55.77), indicating the induction of analgesia (PR-PA- vs. PR+PA-, $p < 0.001$). However, the response latencies of infected voles after exposure to the predator odor (52.11 ± 0.41 ; range from 51.40 to 53.23) were lower than those of uninfected voles (PR+PA+ vs. PR+PA-, $p < 0.05$; Fig. 1). Thus, coccidia infection altered the vole response to predator odor.




Figure 1. Vole nociceptive latencies among the four treatments. $N=10, 10, 10$ and 10 for the PR+PA+, PR+PA-, PR-PA+ and PR-PA- groups, respectively. Different letters represent significant difference between four treatments ($p < 0.05$). Data were expressed as the mean \pm SE.

Field experiments

Among the various models describing survival, model 1, 2 and 3 were parsimonious models (Table 1). The differences of QAICc value between models 1, 2 and 3 are less than 2 (the differences between model 1 and 2, 3 is 0.11 and 1.71, respectively), thus these models are considered equally valid models. The model 1, 2 and 3 included the effect of time, treatment and interaction between time and treatment, indicating that time and treatment affected vole overwinter survival (Table 1). The average overwinter survival rates in group H and group L were 0.772 ± 0.01 and 0.755 ± 0.01 , respectively (Fig. 2).

The population change rate was affected by time ($F_{5,12} = 10.277, p < 0.05$) and the interaction between time and treatment ($F_{5,12} = 0.785, p < 0.05$). However, no effect of treatment alone was found ($F_{1,12} = 0.06, p = 0.81$), indicating that only

Table 1. Best model structures for modeling survival of the root vole population. The model with the lowest QAICc is reported for the first time. The model structure for recapture remained the best model {P(TR + TR . S + TR . T)}. The effect of treatment is abbreviated TR; time effect, T; sex effect, S. The main effects are symbolized by a plus sign (+) and specific interactions are symbolized by a dot (.), and models including all combinations of additive and interaction effects are represent by an asterisk (*).

	Model number	Model	Number of parameters	QAICc	QAICc weight	QDeviance
General models	1	$\Phi T + TR . T$	12	237.91	0.3696	56.26
	2	ΦT	10	238.02	0.3494	61.06
	3	$\Phi TR + T + TR . T$	13	239.62	0.1573	55.57
	4	$\Phi TR + S + T + TR . T$	14	241.96	0.0488	55.47
Global models	5	$\Phi TR * T * S$	27	269.26	0	47.52


time and its interaction with the treatment affected population change rate. The average population change rates in the group H and group L were -0.008 ± 0.01 and -0.012 ± 0.01 , respectively (Fig. 3). Meanwhile, the population density was affected by time ($F_{5,12} = 53.203$, $p < 0.001$) and treatment ($F_{1,12} = 14.735$, $p < 0.05$), but not the interaction between time and treatment ($F_{5,12} = 0.736$, $p = 0.611$). The group L vole populations demonstrated a higher density than group H ($p < 0.05$). Although the former had a 23.7 % higher density than the latter during the adaptive phase (the first 13 days), its population declined sharply to a size similar to group H at the end of the experiments (March 2018, Fig. 4).

DISCUSSION

The primary finding of this research was that coccidian infection in voles reduces analgesia induced by predator risk, resulting in a lower overwinter survival in root voles.

In small mammals, analgesia can promote defensive responses to stimuli and is advantageous in real-time or potentially dangerous situations (Kavaliers 1988, Rodgers 1995, Kavaliers et al. 2000). Laboratory studies have shown that mice and rats display analgesic response when exposed to predator or predator odors (Kavaliers 1990, Rebecca 2003). For instance, Furuya-da-Cunha et al. (2016) found that mice reduce pain reactivity when exposed to predators. However, parasitic infections in mice reduce the analgesia following exposure to predators (Kavaliers et al. 1997). Our results supported our first hypothesis that coccidia infection reduces predator-induced analgesia in root voles. To our knowledge, the present study is the first to test this hypothesis in wild rodents.

Although various reports have highlighted the positive role of analgesia induced by stress in animal defense responses (Amit and Galina 1986, Butler and Finn 2009, Thomson et al. 2020), the causal correlation between nociception and survival at the population level is obscure. Our study is the first to examine whether nociceptive responses affect the overwinter survival of small mammal populations. We found that pain-sensitive voles have lower overwinter survival than pain-inhibited voles. Firstly, the average overwinter survival rate of pain-sensitive voles was lower than it of pain-inhibited voles. Secondly, some pain-inhibited voles died during the two weeks of acclimatization (October 16–28th, 2017) prior to the mark-recapture experiment. Subsequently, the population of pain-sensitive voles had a higher density than pain-inhibited voles during the first trapping session. However, the population of pain-sensitive populations declined sharply at the end of the experiment (March 17th, 2018) to a size comparable to that of the pain-inhibited populations. This result indicates that pain-sensitive voles had lower survival throughout the experiment, which last approximately five months. Finally, the pain-sensitive voles had a higher population change rate than pain-inhibited voles. Root voles do not breed in winter (Liang et al. 1982) and were prevented from entering or leaving the enclosures throughout the experimental period.



Figures 2–4. Monthly apparent survival probability (2), population change rate (3) and population size (4) of root voles during the live-trapping sessions under two different groups. H signifies that root voles with high thermal responses latency; L signifies that root voles with low thermal responses latency. $n = 48$ and 48 for H and L groups. Data were expressed as the mean \pm SE.

Thus, the higher population change rate of pain-sensitive voles was only due to lower overwinter survival. These findings support our second hypothesis that voles with reduced analgesia have lower overwinter survival.

Numerous studies have found that extrinsic factors, including parasites (Ryberg et al. 2020), predators (Sheriff et al. 2020), climate (Rödel et al. 2004), and food (Pedersen and Greives 2008), can directly or indirectly affect animal mortality. In our field experiments, parasites are removed in both treatments, and food and climatic conditions are consistent in all four enclosures. Moreover, some laboratory studies have shown that increased analgesia can enhance the anti-predator responses, increasing the survival probability (Ornstein and Shimon 1981, Lichtman and Fanselow 1990, Tambeli et al. 2012). For instance, predator-induced analgesia promoted anti-predator behaviors in mice, which decreased mortality when exposed to cats (Ornstein and Shimon 1981). Notably, voles prefer routes with higher vegetation cover to avoid predation risk (Merken et al. 1991, Taraborelli et al. 2008). In this study, vegetation cover was low in winter, which may have enhanced predation risk by decreasing natural shelter. Therefore, in the present study, the lower survival of voles with reduced analgesia was due to the increased vulnerability to predator, which may relate to the decreased anti-predator behaviors.

Growing evidence suggests that predators and parasites can have non-additive effects on a shared group of prey or hosts, which can influence the population dynamics (Ramirez and Snyder 2009, Krkošek et al. 2011, Duffy et al. 2011, Marino and Werner 2013). Shang et al. (2019) found that predators increase both the prevalence and intensity of coccidian infection in voles through immune suppression induced by predation stress. Meanwhile, the increased coccidia infection increases the predation risk, reducing the overwinter survival and population density of voles. The number of individuals present at the beginning of the spring breeding period depends on overwinter survival. Thus, the reduced overwinter survival plays a key crucial role in subsequent population fluctuations (Shang et al. 2020). The present study provides possible insights on how coccidia infection increases the infection vulnerability of root voles. We demonstrate that coccidian infection attenuates predator-induced analgesia, and the reduced analgesia increases the susceptibility to predation.

ACKNOWLEDGEMENTS

This work was Funded by the National Natural Science Foundation of China (Grant 31570421, 31870397), Strategic Priority Research Program of Chinese Academy of Sciences (Grant XDA2005010406), the Natural Science Foundation of Qinghai Province (Grant 2018-ZJ-906), Joint Grant From Chinese Academy of Sciences – People's Government of Qinghai Province on Sanjiangyuan National Park (Grant LHZX-2020-01) and Sanjiangyuan Animal Genome Project. We thank Yan-Bin Yang for his assistance in the field work and biochemical assays.

LITERATURE CITED

Alzaga V, Vicente J, Villanua D, Acevedo P, Casas F, Gortazar C (2008) Body condition and parasite intensity correlates with

- escape capacity in Iberian hares (*Lepus granatensis*). Behavioral Ecology and Sociobiology 62(5): 769–775. <https://doi.org/10.1007/s00265-007-0502-3>
- Amit Z, Galina ZH (1986) Stress-induced analgesia: Adaptive pain suppression. Physiological Reviews 66(4): 1091–1120. <https://doi.org/10.1152/physrev.1986.66.4.1091>
- Anderson DR, Burnham KP, White GC (1994) AIC model selection in overdispersed capture-recapture data. Ecology 75(6): 1780–1793. <https://doi.org/10.2307/1939637>
- Bian JH, Fan NC, Jing ZC, Shi YZ (1994) Studies on the successive relation between small mammal community and plant community in alpine meadow. Acta Theriologica Sinica 14(3): 209–216 <https://doi.org/10.16829/j.slxb.1994.03.009>
- Bian JH, Wu Y, Getz LL, Cao YF, Chen F, Yang L (2011) Does maternal stress influence winter survival of offspring in root voles *Microtus oeconomus*? A field experiment. Oikos 120(1): 47–56. <https://doi.org/10.1111/j.1600-0706.2010.18165.x>
- Bian JH, Wu Y, Liu JK (2005a) Breeding behavior under temporal risk of predation in male root voles (*Microtus oeconomus*). Journal of Mammalogy 86(5): 953–960. <https://doi.org/10.2307/4094441>
- Bian JH, Wu Y, Liu JK (2005b) Effect of predator-induced maternal stress during gestation on growth in root voles *Microtus oeconomus*. Acta Theriologica 50(4): 473–482. <https://doi.org/10.1007/BF03192640>
- Butler RK, Finn DP (2009) Stress-induced analgesia. Progress in Neurobiology 88: 184–202. <https://doi.org/10.1016/j.pneurobio.2009.04.003>
- Caetano JVO, Maia MR, Manica LT, Macedo R H (2014) Immune-related effects from predation risk in Neotropical blue-black grassquits (*Volatinia jacarina*). Behavioural Processes 109: 58–63. <https://doi.org/10.1016/j.beproc.2014.07.003>
- Caio M (2011) Modulation of nociceptive-like behavior in zebrafish (*Danio rerio*) by environmental stressors. Psychology and Neuroscience 4(1): 149–155. <https://doi.org/10.3922/j.psns.2011.1.017>
- Cao YF, Nie XH, He H, Du SY, Duszynski DW, Bian JH (2014) Gastrointestinal parasites of root voles, *Microtus oeconomus* (Rodentia: Muridae), from Haibei area, Qinghai Province, China. Comparative Parasitology 81(2): 185–190. <https://doi.org/10.1654/4675.1>
- Carreon N, Faulkes Z (2014) Position of larval tapeworms, *Polypocephalus* sp., in the ganglia of shrimp, *Litopenaeus setiferus*. Integrative and Comparative Biology 54(2): 143–148. <https://doi.org/10.1093/icb/ucu043>
- Chambers LK, Singleton GR, Hinds LA (1999) Fertility control of wild mouse populations: The effects of hormonal competence and an imposed level of sterility. Wildlife Research 26(5): 579–591. <https://doi.org/10.1071/WR98093>
- Colwell DD, Kavaliers M (1993) Evidence for involvement of endogenous opioid peptides in altered nociceptive responses of mice infected with *Eimeria vermiformis*. Journal of Parasitology 79(5): 751–756. <https://doi.org/10.2307/3283615>

- Cooch EG, White GC (2006) Program MARK: a gentle introduction. Available from <http://www.phidot.org/software/mark/docs/book>
- Duffy MA, Housley JM, Penczykowski RM, Caceres CE, Hall SR (2011) Unhealthy herds: indirect effects of predators enhance two drivers of disease spread. *Functional Ecology* 25(5): 945–953. <https://doi.org/10.1111/j.1365-2435.2011.01872.x>
- Furuya-da-Cunha EM, Souza RR, Canto-de-Souza A (2016) Rat exposure in mice with neuropathic pain induces fear and antinociception that is not reversed by 5-HT_{2C} receptor activation in the dorsal periaqueductal gray. *Behavioural Brain Research* 307, 250–257. <https://doi.org/10.1016/j.bbr.2016.04.007>
- Gooding EL, Kendrick MR, Brunson JF, Kingsley-Smith PR, Fowler AE, Frischer ME, Byers JE (2020) Black gill increases the susceptibility of white shrimp, *Penaeus setiferus* (Linnaeus, 1767), to common estuarine predators. *Journal of Experimental Marine Biology and Ecology* 524: 1–6. <https://doi.org/10.1016/j.jembe.2019.151284>
- Haye PA, Ojeda FP (1998) Metabolic and behavioral alterations in the crab *Hemigrapsus crenulatus* (Milne-Edwards 1837) induced by its acanthocephalan parasite *Profilicollis antarcticus* (Zdzitowiecki 1985). *Journal of Experimental Marine Biology and Ecology* 228(1): 73–82. [https://doi.org/10.1016/s0022-0981\(98\)00007-0](https://doi.org/10.1016/s0022-0981(98)00007-0)
- Hoey AS, McCormick MI (2004) Selective predation for low body condition at the larval-juvenile transition of a coral reef fish. *Oecologia* 139(1): 23–29. <https://doi.org/10.2307/40006407>
- Ives AR, Dobson AP (1987) Antipredatory behaviour and the population dynamics of simple predator-prey systems. *American Naturalist* 130(3): 431–447. <https://doi.org/10.1086/284719>
- Jiang Y, Wei S, Wang Z, Zhen Y, Cui R (1991) Productivity investigation of the root vole (*Microtus oeconomus*) population in the Haibei alpine bushland (*Potentilla fruticosa*) I. Population dynamics. *Acta Theriologica Sinica* 11(4): 270–278. <https://doi.org/10.16829/j.slx.1991.04.006>
- Joly DO, Messier F (2004) The distribution of *Echinococcus granulosus* in moose: evidence for parasite-induced vulnerability to predation by wolves? *Oecologia* 142(3): 586–590. <https://doi.org/10.1007/s00442-004-1802-1>
- Kavaliers M (1988) Evolutionary and comparative aspects of nociception. *Brain Research Bulletin* 21(6): 923–931. [https://doi.org/10.1016/0361-9230\(88\)90030-5](https://doi.org/10.1016/0361-9230(88)90030-5)
- Kavaliers M (1990) Responsiveness of deer mice to a predator, the short-tailed weasel: population differences and neuromodulatory mechanisms. *Physiological Zoology* 63(2): 388–407. <https://doi.org/10.1086/physzool.63.2.30158503>
- Kavaliers M, Colwell DD (1991) Sex differences in opioid and non-opioid predator-induced analgesia in mice. *Brain Research* 568(1–2): 173–177. [https://doi.org/10.1016/0006-8993\(91\)91394-G](https://doi.org/10.1016/0006-8993(91)91394-G)
- Kavaliers M, Colwell DD (1994) Parasite infection attenuates nonopioid mediated predator-induced analgesia in mice. *Physiology & Behavior* 55(3): 505–510. [https://doi.org/10.1016/0031-9384\(94\)90108-2](https://doi.org/10.1016/0031-9384(94)90108-2)
- Kavaliers M, Colwell DD (1995) Decreased predator avoidance in parasitized mice: neuromodulatory correlates. *Parasitology* 111(3): 257–263. <https://doi.org/10.1017/s0031182000081816>
- Kavaliers M, Colwell DD, Choleris E (2000) Parasites and behaviour: an ethopharmacological perspective. *Parasitology Today* 16(11): 464–468. [https://doi.org/10.1016/S0169-4758\(00\)01786-5](https://doi.org/10.1016/S0169-4758(00)01786-5)
- Kavaliers M, Colwell DD, Perrot-Sinal TS (1997) Opioid and non-opioid NMDA-mediated predator-induced analgesia in mice and the effects of parasitic infection. *Brain Research* 766(1–2), 11–18. [https://doi.org/10.1016/s0006-8993\(97\)00521-0](https://doi.org/10.1016/s0006-8993(97)00521-0)
- Klemola T, Korpimäki E, Koivula M (2002) Rate of population change in voles from different phases of the population cycle. *Oikos* 96(2): 291–298. <https://doi.org/10.1034/j.1600-0706.2002.960211.x>
- Krams I, Kivleniece I, Kuusik A, Krama T, Freeberg TM, Mänd R, Vrublevska J, Rantala MJ, Mänd M (2013) Predation selects for low resting metabolic rate and consistent individual differences in anti-predator behavior in a beetle. *Acta Ethologica* 16(3): 163–172. <https://doi.org/10.1007/s10211-013-0147-3>
- Krkošek M, Connors BM, Ford H, Peacock S, Mages P, Ford JS, Morton A, Volpe J, Hilborn R, Dill LM (2011) Fish farms, parasites, and predators: implications for salmon population dynamics. *Ecological Applications* 21(3): 897–914. <https://doi.org/10.1890/09-1861.1>
- Lamana MS, Miranda J, Tobaldini G, Fischer L, Tambeli CH (2018) Pain chronification and chronic pain impair a defensive behavior, but not the ability of acute pain to facilitate it, through the activation of an endogenous analgesia circuit. *Behavioral Neuroscience* 132(6): 614–623. <https://doi.org/10.1037/bne0000255>
- Lebreton JD, Burnham KP, Clobert J, Anderson DR (1992) Modeling survival and testing biological hypotheses using marked animals: a unified approach with case studies. *Ecological Monographs* 62(1): 67–118. <https://doi.org/10.2307/2937171>
- Liang JR, Zeng JX, Wang ZW, Han YC (1982) Studies on growth and development in the root vole (*Microtus oeconomus*). *Acta Biology Plateau Sinica* 1: 195–208.
- Lichtman AH, Fanselow MS (1990) Cats produce analgesia in rats in the tail-flick test: naltrexone sensitivity is determined by the nociceptive test stimulus. *Brain Research* 533(1): 91–94. [https://doi.org/10.1016/0006-8993\(90\)91800-v](https://doi.org/10.1016/0006-8993(90)91800-v)
- Lima SL, Dill LM (1990) Behavioral decisions made under the risk of predation: a review and prospectus. *Canadian Journal of Zoology* 68(4): 619–640. <https://doi.org/10.1139/z90-092>
- Li YN, Zhao XQ, Cao GM, Zhao L, Wang QX (2004) Analyses on climates and vegetation productivity background at

- Haibei Alpine Meadow Ecosystem Research Station. Plateau Meteorology 23(4): 558–566.
- Marino JA, Werner EE (2013) Synergistic effects of predators and trematode parasites on larval green frog (*Rana clamitans*) survival. Ecology 94(12): 2697–2708. <https://doi.org/10.1890/13-0396.1>
- Merkens M, Harestad AS, Sullivan TP (1991) Cover and efficacy of predator-based repellents for Townsend's vole. Journal of Chemical Ecology 17(2): 401–412. <https://doi.org/10.1007/BF00994341>
- Møller AP, Nielsen JT (2007) Malaria and risk of predation: a comparative study of birds. Ecology 88 (4): 871–881. <https://doi.org/10.1890/06-0747>
- Nie XH, Cao YF, Du SY, He H, Bian JH (2014) Intestinal parasite prevalence in root voles (*Microtus oeconomus*) from field enclosures. Acta Theriologica Sinica 34(2): 172–180. <https://doi.org/10.16829/j.slx.2014.02.009>.
- Ornstein K, Shimon A (1981) Pinch-induced catalepsy in mice. Journal of Comparative and Physiological Psychology 95(5): 827–835. <https://doi.org/10.1037/h0077827>.
- Pedersen AB, Greives TJ (2008) The interaction of parasites and resources cause crashes in a wild mouse population. Journal of Animal Ecology 77(2): 370–377. <https://doi.org/10.1111/j.1365-2656.2007.01321.x>
- Ramirez RA, Snyder WE (2009) Scared sick? Predator–pathogen facilitation enhances exploitation of a shared resource. Ecology 90(10): 2832–2839. <https://doi.org/10.2307/25592817>
- Rebecca MC (2003) Sex differences in drug- and non-drug-induced analgesia. Life Sciences 72(24): 2675–2688. [https://doi.org/10.1016/s0024-3205\(03\)00178-4](https://doi.org/10.1016/s0024-3205(03)00178-4)
- Rödel HG, Bora A, Kaetzke P, Khaschei M, Hutzelmeyer H, von Holst D (2004) Over-winter survival in subadult European rabbits: weather effects, density dependence, and the impact of individual characteristics. Oecologia 140(4): 566–576. <https://doi.org/10.1007/s00442-004-1616-1>
- Rodgers RJ (1995) Neuropharmacological aspects of adaptive pain inhibition in murine “victims” of aggression. Aggressive Behavior 21(1): 29–39. [https://doi.org/10.1002/1098-2337\(1995\)21:1<29::AID-AB2480210106>3.0.CO;2-A](https://doi.org/10.1002/1098-2337(1995)21:1<29::AID-AB2480210106>3.0.CO;2-A)
- Ryberg MP, Skov PV, Vendramin N, Buchmann K, Nielsen A, Behrens JW (2020) Physiological condition of Eastern Baltic cod, *Gadus morhua*, infected with the parasitic nematode *Contracaecum osculatum*. Conservation Physiology 8(1): coaa093. <https://doi.org/10.1093/conphys/coaa093>
- Shang GZ, Zhu YH, Wu Y, Cao YF, Bian JH (2019) Synergistic effects of predation and parasites on the overwinter survival of root voles. Oecologia 191: 83–96. <https://doi.org/10.1007/s00442-019-04455-4>
- Shang GZ, Yang YG, Zhu YH, Wu XQ, Cao YF, Wu Y, Bian JH (2020) A complex regulating pattern induced by the effects of predation and parasites on root vole (*Microtus oeconomus*) populations during the breeding season. Journal of Mammalogy 101(5): 1345–1355. <https://doi.org/10.1093/jmammal/gyaa084>
- Sheriff MJ, Peacor SC, Hawlena D, Thaker M (2020) Non-consumptive predator effects on prey population size: A dearth of evidence. Journal of Animal Ecology 89(6): 1302–1316. <https://doi.org/10.1111/1365-2656.13213>
- Steen H, Taitt M, Krebs CJ (2002) Risk of parasite-induced predation: an experimental field study on Townsend's voles (*Microtus townsendii*). Canadian Journal of Zoology 80(7): 1286–1292. <https://doi.org/10.1139/Z02-115>
- Sun RY, Zheng SW, Cui RX (1982) Home range of the root vole *Microtus oeconomus*. Acta Theriologica Sinica 2: 219–231.
- Sun P, Zhao XQ, Xu SX, Zhao TB, Zhao W (2002) Changes after snow of the population characteristics of root vole (*Microtus oeconomus*). Acta Theriologica Sinica 22(2): 318–320. <https://doi.org/10.16829/j.slx.1982.02.014>
- Sundell J (2006) Experimental tests of the role of predation in the population dynamics of voles and lemmings. Mammal Review 36(2): 107–141. <https://doi.org/10.1111/j.1365-2907.2006.00083.x>
- Tambeli CH, Fischer L, Monaliza SL, Menescal-de-Oliveira L, Parada CA (2012) The functional role of ascending nociceptive control in defensive behavior. Brain Research 1464: 24–29. <https://doi.org/10.1016/j.brainres.2012.05.010>
- Taraborelli PA, Moreno P, Srur A, Sandobal AJ, Martínez MG, Giannoni SM (2008) Different antipredator responses by *Microcavia australis* (Rodentia, Hystricognate, Caviidae) under predation risk. Behaviour. 145(6): 829–842. <https://doi.org/10.1163/156853908783929115>
- Thomson JS, Deakin AG, Cossins AR, Spencer JW, Young IS, Sneddon LU (2020) Acute and chronic stress prevents responses to pain in zebrafish: evidence for stress-induced analgesia. Journal of Experimental Biology 223(14): 1–11. <https://doi.org/10.1242/jeb.224527>
- Tompkins DM, Dunn AM, Smith MJ, Telfer S (2011) Wildlife diseases: from individuals to ecosystems. Journal of Animal Ecology 80(1): 19–38. <https://doi.org/10.1111/j.1365-2656.2010.01742.x>
- Wang ZL, Liu JK (2002a) Effect of silver fox odor on breeding and foraging of root voles. Acta Theriologica Sinica 22(1): 22–29. <https://doi.org/10.16829/j.slx.2002.01.005>
- Wang ZL, Liu JK (2002b) The effects of silver fox odor on behaviors of root voles. Acta Ecologica Sinica 22: 554–558.
- White GC, Burnham KP (1999) Program MARK: Survival estimation from populations of marked animals. Bird Study 46(Suppl.): S120–S139. <https://doi.org/10.1080/00063659909477239>
- Wirsing AJ, Steury TD, Murray DL (2002) Relationship between body condition and vulnerability to predation in red squirrels and snowshoe hares. Journal of Mammalogy 83(3): 707–715. <https://doi.org/10.2307/1383534>
- Yang YB, Shang GZ, Du SY, Zhang X, Wu Y, Bian JH (2018) Maternal density stress and coccidian parasitism: Synergistic effects on overwinter survival in root voles. Functional Ecology 32: 2181–2193. <https://doi.org/10.1111/1365-2435.13129>

Submitted: April 25, 2021

Accepted: June 24, 2021

Available online: July 7, 2021

Editorial responsibility: Carolina Arruda Freire

Author Contributions: BJH and WY designed the experiment;

YYG, SGZ, WXQ, CYF and CHQ conducted the experiments; YYG and SGZ both contributed to analyzing data and writing the paper. YYG and SGZ contributed equally to this paper.

Competing Interests: The authors have declared that no competing interests exist.

© 2021 Sociedade Brasileira de Zoologia. Published by Pensoft Publishers at <https://zoologia.pensoft.net>