



New records of *Anastrepha* Schiner, 1868 (Diptera, Tephritidae) in an urban forest fragment in Manaus, Amazonas, Brazil

Francisco Clóvis Costa-Silva¹, Agno Nonato Serrão Acioli¹, Neliton Marques da Silva¹, Keiko Uramoto², Marcoandre Savaris², Roberto Antonio Zucchi²

1 Laboratório de Entomologia e Acarologia Agrícola, Faculdade de Ciências Agrárias, Universidade Federal do Amazonas, Av. Rodrigo Otávio 6200, Setor Sul, Coroado, 69080-900, Manaus, AM, Brazil. **2** Laboratório de Taxonomia dos Insetos de Importância Agrícola, Escola Superior de Agricultura “Luiz de Queiroz”, Universidade de São Paulo, Av. Pádua Dias 11, 13418-900, Piracicaba, SP, Brazil.

Corresponding author: Neliton Marques da Silva, nmerinato@gmail.com.

Abstract

Four new records of *Anastrepha* species, *A. elongata* Fernández, 1953, *A. fraterculus* (Wiedemann, 1830) (sensu lato), *A. pseudanomala* Norrbom, 2002, and *A. sodalis* Stone, 1942, are discussed. *Anastrepha elongata* is recorded for the first time in Brazil. All species were collected in McPhail traps in an urban forest fragment on the campus of the Federal University of Amazonas, city of Manaus, state of Amazonas.

Keywords

Brazilian Amazon, diversity, fruit flies, McPhail trap, new records.

Academic editor: Tiago Kütter Krolow | Received 18 March 2020 | Accepted 9 June 2020 | Published 3 July 2020

Citation: Costa-Silva FC, Acioli ANS, Silva NM, Uramoto K, Savaris M, Zucchi RA (2020) New records of *Anastrepha* Schiner, 1868 (Diptera, Tephritidae) in an urban forest fragment in Manaus, Amazonas, Brazil. Check List 16 (4): 853–857. <https://doi.org/10.15560/16.4.853>

Introduction

The Amazon Rainforest poses a major challenge for studies on fruit flies, due to its enormous extent and the logistical difficulties in collecting these insects, resulting in inadequate knowledge of fruit flies in the Brazilian Amazon. Many records from this region have resulted from occasional collections (Zucchi et al. 1996). In the beginning of this century, 25 species of *Anastrepha* Schiner, 1868 were reported in the state of Amazonas (Silva and Ronchi-Teles 2000).

Because some *Anastrepha* species cause severe damage to marketable fruits and are highly important as quarantine pests, this genus has been studied mainly in agricultural areas (e.g. Malavasi and Zucchi 2000). However, studies in areas with undisturbed native vegetation are crucial for understanding biological and behavioral

aspects of fruit flies (Aluja et al. 2003). Likewise, studies in human-impacted areas may provide important information about the population dynamics of fruit fly species (Uramoto et al. 2003, 2004, 2005). For example, in studies carried out in three different areas (urban, research station, and agricultural), the highest diversity and similarity among several species of *Anastrepha* were observed in the urban and research-station areas (Oliveira et al. 2017).

The geographic diversity of *Anastrepha* species is related directly and positively to the size and continuity of forest fragments in urbanized areas (Garcia et al. 2017). Araujo et al. (2019) compared two sampling methods (trapping and fruit sampling) on the campus of the Luiz de Queiroz College of Agriculture (ESALQ;

Piracicaba, São Paulo, Brazil), and discussed that although fruit sampling has resulted in a greater diversity and abundance of species, it did not adequately represent the composition of the *Anastrepha* species in the area. This note reports four new records of *Anastrepha* species from an urbanized area in the city of Manaus, state of Amazonas.

Methods

This study was conducted in an urban forest fragment of the campus of the Federal University of Amazonas, Manaus, Amazonas, Brazil (Fig. 1) from September 2010 to September 2011 (Costa-Silva 2012). The predominant vegetation in the northern sector of the campus is a well-preserved Dense Ombrophilous Forest. Anthropogenic intervention in this sector comprises a cluster of buildings and some trails. The vegetation in the southern sector consists of a highly impacted Open Ombrophilous Forest as well as secondary forest, agroforestry crops, orchards, medicinal and olericultural plants, deforested areas, and buildings.

Fruit flies were collected in McPhail traps baited with 10% sugarcane molasses + borax, mounted 1.80 m high on the east side of trees and approximately 50 m apart. We used 80 traps, 40 in each sector. In the northern sector, traps were hung between buildings and along forest trails. In the southern sector, traps were hung between buildings and in the original ombrophilous forest,

secondary forest, and fruit orchards.

Morphological characterizations and illustrations for these four species are available online, as is the key used for identification (Norrbon et al. 2012). Voucher specimens are deposited at the Luiz de Queiroz Museum of Entomology (MELQ), Department of Entomology and Acarology, ESALQ, Piracicaba. The collection was carried out under permit from the National System of Biodiversity Information (SISBIO; authorization number 24539-1).

The wing pattern of each species was photographed in dorsal view under a Leica M205C stereomicroscope equipped with a Leica DFC 450 camera. Digital photographs were enhanced using Photoshop CS6 to correct the color and make minor corrections (e.g., remove debris). The distribution map was made using Quantum GIS 2.8.

Results

Four new records are recorded for the first time in the state of Amazonas, namely *Anastrepha elongata* Fernández, 1953 (new from Brazil), *A. fraterculus* (Wiedemann, 1830) (sensu lato), *A. pseudanomala* Norrbom, 2002, and *A. sodalis* Stone, 1942.

Anastrepha elongata Fernández, 1953

Figure 2A

New record. BRAZIL • 2♀; Amazonas, Manaus,




Figure 1. Collecting site of new records of species of *Anastrepha* in the campus of the Federal University of Amazonas, Manaus (*; white symbols), and their previous occurrences (black symbols) in Brazil.




Figure 2. Wing patterns of new records of species of *Anastrepha* from Amazonas. **A.** *A. elongata* Fernández, 1953. **B.** *A. fraterculus* (Wiedemann, 1830) (sensu lato). **C.** *A. pseudanomala* Norrbom, 2002. **D.** *A. sodalis* Stone, 1942. Scale bars = 2 mm.

campus of the Federal University of Amazonas; 03°05'37"S, 059°58'01.1"W; alt. 92 m; 29 Jul. 2011; McPhail trap; F.C. Costa-Silva leg.; ESALQENT00002, ESALQENT00003.

Previous records in Brazil. *Anastrepha elongata* is recorded here for the first time in Brazil.

Identification. *Anastrepha elongata* is a large species (mesonotum: 4 to 5 mm; aculeus: 9.25–12.5 mm), with all wing bands separate (Fig. 2A). Steyskal (1977) considered this taxon to be a junior synonym of *A. barnesi* Aldrich, 1925. However, Caraballo (1981) revalidated *A. elongata*, and discussed the morphological characters that distinguish it from *A. barnesi*, such as the highly distinct mesonotum bands, position of spiracles in the oviscap, teeth rows on the eversible membrane, and shorter aculeus tip.

***Anastrepha fraterculus* (Wiedemann, 1830) (sensu lato)**

Figure 2B

New record. BRAZIL • 2♀; Amazonas, Manaus, campus of the Federal University of Amazonas; 03°05'37.0"S, 59°58'01.1"W; alt. 92 m; 15 Jul. 2011; McPhail trap; F.C. Costa-Silva leg.; ESALQENT00001.

Previous records in Brazil. The *Anastrepha fraterculus* complex is widely distributed in Brazil. The cryptic species of this complex were recorded in 23 of the 26 Brazilian states (Zucchi and Moraes 2008), but the distributions of the individual cryptic species in the country are not known.

Identification. The *A. fraterculus* complex comprises several cryptic species (Selivon and Perondini 2007; Hernández-Ortiz et al. 2012). Therefore, the nominal *A. fraterculus* is used here in sensu lato. Recognition of the *A. fraterculus* complex is based on females, mainly on

the shape of the aculeus tip (0.2–0.3 mm long, with a distinct constriction before the serrate part). The wing pattern is variable, such as: (1) C-band and S-band connected, V-band with distal and proximal arms connected, but S-band separate (Fig. 2B); (2) C-band and S-band separate, V-band incomplete (distal and proximal arms separate), and S-band separate; and (3) V-band complete (distal and proximal arms connected) and connected to S-band. Hernández-Ortiz et al. (2012) discussed the wing patterns of seven *A. fraterculus* morphotypes from several countries in the Americas.

***Anastrepha pseudanomala* Norrbom, 2002**

Figure 2C

New record. BRAZIL • 4♀; Amazonas, Manaus, campus of the Federal University of Amazonas; 03°05'37.0"S, 059°58'01.1"W; alt. 92 m; 23 Sep. 2011; McPhail trap; F.C. Costa-Silva leg.; ESALQENT00004, ESALQENT00005, ESALQENT00006, ESALQENT00007.

Previous records in Brazil. Amapá (Jesus et al. 2010).

Identification. *Anastrepha pseudanomala* could be confused with *A. anomala* Stone, 1942 and *A. normalis* Norrbom, 2002, differing in having the longest aculeus tip (Norrbom 2002). It is a member of the *A. serpentina* group, in which the hyaline area in the br cell does not extend to the vein R_{4+5} (Fig. 2C).

***Anastrepha sodalis* Stone, 1942**

Figure 2D

New record. BRAZIL • 1♀; Amazonas, Manaus, campus of the Federal University of Amazonas; 03°05'37.0"S, 059°58'01.1"W; alt. 92 m; 7 Oct. 2011; McPhail trap; F.C. Costa-Silva leg.; ESALQENT00008.

Previous records in Brazil. Pará (Stone 1942; Menezes et al. 2000), Amapá (Baia et al. unpub. data – XXVI

Brazilian Congress of Entomology 2016), Piauí (Menezes et al. 2000).

Identification. *Anastrepha sodalis* is a large species (mesonotum: 3–4 mm; aculeus: 3–4 mm). All the wing bands are joined (Fig. 2D) and the aculeus tip has minute serrations (visible only with a compound microscope).

Discussion

Our results demonstrate how little is known about the diversity of fruit flies in the state, and reinforce the need to intensify collections of fruit flies, in traps and from fruits, throughout the state of Amazonas.

Nineteen species of *Anastrepha*, including the four newly recorded here, were collected on the campus of the Federal University of Amazonas (Costa-Silva 2012). These represent about 45% of the species known in the state of Amazonas. In that survey, carried out in a small urbanized area over a short period of time (one year), almost 20% of the species were new records for the state.

Anastrepha elongata was formerly known only from the type locality (Rancho Grande, Aragua State, Venezuela). Therefore, the collection of *A. elongata* in Manaus is the first record of this species in Brazil and the second in South America. No host plant is known for *A. elongata*. It belongs to the *A. leptozona* group.

The record of *A. fraterculus* (sensu lato) in the state of Amazonas is important, considering its status as a quarantine pest. Surveys of fruit flies should be intensified because studies on isolated populations of the *A. fraterculus* complex in northern Brazil should help to better understand the complex. Research is being carried out in several countries to clarify the identities of the species of the *A. fraterculus* complex. Several approaches have been used to clarify this complex, including isozymes, biological crosses, pheromones, molecular and morphometric analyses, and egg morphology (see Hendrichs et al. 2015; Schutze et al. 2017).

Additionally, the host plants exploited by members of the *A. fraterculus* complex throughout the Americas reinforce the hypothesis that the complex consists of several species, because although most of the hosts occur throughout the entire distribution area, not all hosts are common to all *A. fraterculus* populations (Hernández-Ortiz et al. 2020). In Brazil, larvae of *A. fraterculus* (sensu lato) breed in fruits of 116 host plants (Zucchi and Moraes 2008). The *A. fraterculus* complex occurs in several Brazilian states, especially in the southeastern and southern regions, and is uncommon in the northern states. *Anastrepha fraterculus* (sensu lato) is one of the most important and widespread pests of fruits in Brazil.

Anastrepha pseudanomala was described from females collected in Panama and Costa Rica. The first record in Brazil was in the state of Amapá, where it was breeding in *Couma utilis* (Mart.) Müll. Arg. (Apocynaceae) (Jesus et al. 2010). Therefore, the collection of *A. pseudanomala* in the state of Amazonas establishes the second record of this species in Brazil.

The original description of *A. sodalis* was based on a single female collected in Santarém, Pará state, Brazil, that is deposited in the Natural History Museum, London (Stone 1942). It was recorded in the states of Amapá (Baia et al. unpub. data – XXVI Brazilian Congress of Entomology 2016) and Piauí (Menezes et al. 2000). We report here the first occurrence of *A. sodalis* in the state of Amazonas. So far, this species has not been included in any of the groups of the genus *Anastrepha* (Norrbon et al. 1999), and has no host plant record.

The new records of *Anastrepha* species discussed here increase to 42 the number of *Anastrepha* species known in the state of Amazonas. This number corresponds to 52% of the species recorded in the Brazilian Amazon biome. The host is not known for about half of the species recorded in Amazonas (NMS pers. inf.). Collections in host fruits and traps should be intensified in order to adequately evaluate the diversity of fruit flies in the Amazon Rainforest ecosystem.

Acknowledgements

We thank the Postgraduate Program in Entomology of the Luiz de Queiroz College of Agriculture, University of São Paulo, and the Postgraduate Program in Tropical Agronomy of the Federal University of Amazonas. RAZ is a fellow of the National Council for Scientific and Technological Development. We also thank the anonymous reviewers and the academic editor for improving the manuscript.

Authors' Contributions

FCC, ANSA and NMS collected the specimens in the field. KU identified the specimens. MS constructed the map and Figure 2. NMS and RAZ conceived the study and analyzed the data. All authors wrote the final version of manuscript.

References

- Aluja M, Rull J, Sivinski J, Norrbom AL, Wharton RA, Macías-Ordóñez R, Díaz-Fleischer F, López M (2003) Fruit flies of the genus *Anastrepha* (Diptera: Tephritidae) and associated native parasitoids (Hymenoptera) in the tropical rain forest biosphere reserve of Montes Azules, Chiapas, Mexico. *Environmental Entomology* 32: 1377–1385. <https://doi.org/10.1603/0046-225X-32.6.1377>
- Araujo MR, Uramoto K, Ferreira ENL, Mesquita Filho W, Walder JMM, Savaris M, Zucchi RA (2019) Fruit fly (Diptera: Tephritidae) diversity and host relationship in diverse environments estimated with two sampling methods. *Environmental Entomology* 48: 227–233. <https://doi.org/10.1093/ee/nvy177>
- Caraballo JV (1981) Las moscas de frutas del genero *Anastrepha* Schiner, 1868 (Diptera: Tephritidae) de Venezuela. PhD thesis, Faculty of Agronomy, Central University of Venezuela, Caracas, 210 pp.
- Costa-Silva FC (2012) Biodiversidade de moscas-das-frutas (Diptera: Tephritidae) no campus da Universidade Federal do Amazonas, Manaus-AM. Master's thesis, Federal University of Amazonas, Manaus, 97 pp.

- Garcia AG, Araujo MR, Uramoto K, Walder JMM, Zucchi RA (2017) Geostatistics and geographic information system to analyze the spatial distribution of the diversity of *Anastrepha* species (Diptera: Tephritidae): the effect of forest fragment in an urban area. *Environmental Entomology* 46: 1189–1194. <https://doi.org/10.1093/ee/nvx145>
- Hendrichs J, Vera MT, de Meyer M, Clarke AR (2015) Resolving cryptic species complexes of major tephritid pests. *ZooKeys* 540: 5–29. <http://doi:10.3897/zookeys.540.9656>
- Hernández-Ortiz V, Barradas-Juanz N, Díaz-Castelazo C (2020) A review of the natural host plants of the *Anastrepha fraterculus* complex in the Americas. In: Pérez-Staples D, Díaz-Fleischer F, Montoya P, Vera MT (Eds) *Area-wide management of fruit fly pests*. CRC Press, Boca Raton, 90–122.
- Hernández-Ortiz V, Bartolucci A, Morales-Valles P, Frías D, Selivon D (2012) Cryptic species of the *Anastrepha fraterculus* complex (Diptera, Tephritidae): a multivariate approach for the recognition of South American morphotypes. *Annals of the Entomological Society of America* 105: 305–318. <https://doi.org/10.1603/AN11123>
- Jesus CR, Silva RA, Souza Filho MF, Deus EG, Zucchi RA (2010) First record of *Anastrepha pseudanomala* Norrbom (Diptera: Tephritidae) and its host in Brazil. *Neotropical Entomology* 39: 1059–1060. <https://doi.org/10.1590/S1519-566X2010000600036>
- Malavasi A, Zucchi RA (2000) *Moscas-das-frutas de importância econômica no Brasil: conhecimento básico e aplicado*. Editora Holos, Ribeirão Preto, 327 pp.
- Menezes RVS, Nunes EM, Branco RSC, Zucchi RA (2000) Piauí. In: Malavasi A, Zucchi RA (Eds) *Moscas-das-frutas de importância econômica no Brasil: conhecimento básico e aplicado*. Editora Holos, Ribeirão Preto, 213–215.
- Norrbom AL (2002) A revision of the *Anastrepha serpentina* species group (Diptera: Tephritidae). *Proceedings of the Entomological Society of Washington* 104: 390–436.
- Norrbom AL, Zucchi RA, Hernández-Ortiz V (1999) Phylogeny of the genera *Anastrepha* and *Toxotrypana* (Trypetinae: Toxotrypanini) based on morphology. In: Aluja M, Norrbom AL (Eds) *Fruit flies (Tephritidae): phylogeny and evolution of behavior*. CRC Press, Boca Raton, 299–342.
- Norrbom AL, Korytkowski CA, Zucchi RA, Uramoto K, Venable GL, McCormick J, Dallwitz MJ (2012) onwards. *Anastrepha* and *Toxotrypana*: descriptions, illustrations and interactive keys [version 2019-4-9]. <https://www.delta-intkey.com/anatox/index.htm>. Accessed on: 2019-12-4.
- Oliveira MBR, Ferreira ENL, Reigada C, Lopes GN, Lemos LJU, Uramoto K, Souza-Filho MF, Zucchi RA (2017) Composition of *Anastrepha* species (Diptera: Tephritidae) in habitats with different levels of anthropogenic activity. *Biotemas* 30: 61–69. <https://doi.org/10.5007/2175-7925.2017v30n3p61>
- Schutz MK, Virgilio M, Norrbom AL, Clarke AR (2017) Tephritid integrative taxonomy: where we are now, with a focus on the resolution of three tropical fruit fly species complexes. *Annual Review of Entomology* 62: 147–164. <https://doi.org/10.1146/annurev-ento-031616-035518>
- Selivon D, Perondini ALP (2007) Especies crípticas del complejo *Anastrepha fraterculus* en Brasil. In: Hernández-Ortiz V (Ed.) *Moscas de la Fruta en Latinoamérica (Diptera: Tephritidae): Diversidad, Biología y Manejo*. SyG Editores, Distrito Federal, 101–118.
- Silva NM, Ronchi-Teles B (2000) *Moscas-das-frutas nos estados brasileiros: Amapá, Amazonas, Pará, Rondônia e Roraima*. In: Malavasi A, Zucchi RA (Eds) *Moscas-das-frutas de importância econômica no Brasil: conhecimento básico e aplicado*. Editora Holos, Ribeirão Preto, 203–209.
- Steyskal GC (1977) Pictorial key to species of the genus *Anastrepha* (Diptera: Tephritidae). *Entomological Society of Washington, Washington, DC*, 35 pp.
- Stone A (1942) *The fruit flies of the genus Anastrepha*. United States Department of Agriculture, Miscellaneous Publication 439: 1–112.
- Uramoto K, Walder JMM, Zucchi RA (2003) Flutuação populacional de moscas-das-frutas do gênero *Anastrepha* Schiner, 1868 (Diptera: Tephritidae) no campus “Luiz de Queiroz”, Piracicaba, São Paulo. *Arquivos do Instituto Biológico* 70: 459–465.
- Uramoto K, Walder JMM, Zucchi RA (2004) Biodiversidade de moscas-das-frutas do gênero *Anastrepha* (Diptera, Tephritidae) no campus da ESALQ-USP, Piracicaba, São Paulo. *Revista Brasileira de Entomologia* 48: 409–414. <https://doi.org/10.1590/S0085-56262004000300018>
- Uramoto K, Walder JMM, Zucchi RA (2005) Quantitative analysis and distribution of the population of species in the genus *Anastrepha* (Diptera: Tephritidae) on “Luiz de Queiroz” campus, Piracicaba, SP, Brazil. *Neotropical Entomology* 34: 33–39. <https://doi.org/10.1590/S1519-566X2005000100005>
- Zucchi RA, Moraes RCB (2008) Fruit flies (Diptera: Tephritidae) in Brazil: *Anastrepha* species their host plants and parasitoids [updated 2019-11-28]. <http://www.lea.esalq.usp.br/anastrepha/>. Accessed on: 2020-3-17.
- Zucchi RA, Silva NM, Silveira Neto S (1996) *Anastrepha* species (Diptera; Tephritidae) from the Brazilian Amazon: distribution, hosts and lectotype designation. In: Steck GJ, MacPherson BA (Eds) *Fruit fly pests: a world assessment of their biology and management*. St. Lucie Press, Delray Beach, 259–264.