



Contribution to the knowledge of the beetle fauna (Insecta, Coleoptera) of Malta: new records of seven species with supporting DNA barcodes

Clare Marie Mifsud¹, Noel Vella¹, Adriana Vella^{1*}

¹ Conservation Biology Research Group, Biology Department, University of Malta, Msida, MSD 2080, Malta • CMM: cmifsud2@gmail.com

 <https://orcid.org/0000-0003-4605-7400> • NV: noel.vella@um.edu.mt  <https://orcid.org/0000-0003-0080-5709> • AV: adriana.vella@um.edu.mt

 <https://orcid.org/0000-0003-2246-5354>

* Corresponding author

Abstract

We report the presence of seven species of Coleoptera newly found on the Maltese Islands in the central Mediterranean. The first species records from the Maltese Islands include: *Ancylopus melanocephalus* (Olivier, 1808) (Endomychidae), *Aplidia transversa* (Fabricius, 1801) (Scarabaeidae), *Cercyon quisquilius* (Linnaeus, 1761) (Hydrophilidae), *Hyperaspis duvergeri* Fürsch, 1985 (Coccinellidae), *Lebia cruxminor* (Linnaeus, 1758) (Carabidae), *Smicronyx pauperculus* Wollaston, 1864 (Curculionidae), and *Oxytelus sculptus* Gravenhorst, 1806 (Staphylinidae). The morphological identification of each newly reported species was also confirmed through DNA barcoding.

Keywords

Carabidae, Coccinellidae, Curculionidae, Endomychidae, Hydrophilidae, Scarabaeidae, Staphylinidae

Academic editor: Pedro da Silva | Received 30 June 2021 | Accepted 27 September 2021 | Published 29 October 2021

Citation: Mifsud CM, Vella N, Vella A (2021) Contribution to the knowledge of the beetle fauna (Insecta, Coleoptera) of Malta: new records of seven species with supporting DNA barcodes. Check List 17 (5): 1443–1449. <https://doi.org/10.15560/17.5.1443>

Introduction

Coleoptera is the most diverse animal order on Earth. More than 386,000 extant beetle species have been described and represent around 25% of all described animal species (Bouchard et al. 2017). Due to this immense diversity, listing the actual number of described species from all world regions remains challenging, and estimating the number of undescribed species is even more so.

Cameron and Caruana Gatto (1907) comprehensively listed around 600 Coleoptera species from the Maltese Islands, and 100 years later the total number of species increased to 800 (Mifsud 2000a). An estimate of 1800 species of Coleoptera has been indicated for the Maltese Islands (Mifsud 2000a), and although an updated catalogue is still unavailable, progress has been made with updated species lists for several coleopteran families

(reviewed by Schembri 2003). The groups to which we add species newly recorded from the Maltese Islands have been reviewed and updated in previous works and include the families Endomychidae, Scarabaeidae, Hydrophilidae, Coccinellidae, Carabidae, Curculionidae, and Staphylinidae (Bordoni 1973; Schembri and Schembri 1982; Schembri 1993; Magrini and Schembri 1997; Mifsud and Vigna Taglianti 1999; Shockley et al. 2009; Mifsud and Colonnelli 2010; 2018; Pivotti et al. 2011; Przewoźny 2018).

Complete species lists from restricted areas such as islands are useful by providing information on biodiversity at a finer scale to better guide conservation efforts. Therefore, this research aims to extend our knowledge on the coleopteran fauna of the Maltese Islands to support biodiversity conservation efforts.

Methods

A beetle collection was compiled by the Conservation Biology Research Group, University of Malta (CBRG-UM) between 2015 and 2019. All beetles were either hand collected during the day or captured at night in UV light traps set in the field from May to October of each sampling year. Beetle specimens were collected from various habitat types including urban and rural areas across the islands of Malta and Gozo (Fig. 1). Beetle specimens were individually stored in labelled sampling bottles and placed at -20°C on the same day of collection until further processing. Each specimen was examined and photographed. Morphologically identified species were checked for their occurrence in Malta using the Fauna Europaea database portal (de Jong et al. 2014), published catalogues, and literature (Mifsud and Scupola 1998; Mifsud 2000b, 2002; Mifsud and Knizek 2009; Mifsud and Colonnelli 2010; Pivotti et al. 2011; Lillig et al. 2012; Mifsud and Jelínek 2012; Nardi and Mifsud 2015; Sabella and Mifsud 2016; Sogoh and Yoshitomi 2017; Haran 2018; Mifsud and Nardi 2020). Only beetles not previously recorded from the Maltese Islands are included in this work.

For DNA barcoding of voucher specimens, genomic DNA was extracted using the whole body of small beetle specimens (<3 mm body length). Three legs and one leg were used for DNA extraction from medium-sized beetle specimens (3–10 mm body length) and large specimens (>10 mm body length), respectively. Genomic DNA extraction was done using the GF-1 Tissue DNA Extraction Kit (Vivantis, Malaysia) following the manufacturer's manual. The 658 bp DNA barcode

region of the mitochondrial cytochrome *c* oxidase subunit I gene (COI) was amplified using a cocktail mix of the Folmer primers LCO1490/HCO2198 (Folmer et al. 1994) and LepF1/LepR1 (Hebert et al. 2004) appended with the universal M13 oligonucleotide tails. Amplification reactions were done in 25 μL reaction volumes, containing 1 μL genomic DNA (approximately 10 ng), 0.25 μL of each primer pair (0.5 μM , each), 1 μL of 25 mM MgCl_2 , 5 μL 1X FIREPOL Master Mix containing 2.5 mM Mg^{2+} , 200 μM of each dNTP and 1 U FIREPOL DNA polymerase (Solis BioDyne, Estonia) and 17.5 μL ddH₂O. PCR reactions were carried out on a Nexus Gradient Mastercycler (Effendorf, Hamburg, Germany) using the following temperature profile: 95 $^{\circ}\text{C}$ for 5 min; followed by 6 cycles of 95 $^{\circ}\text{C}$ for 45 s, 45 $^{\circ}\text{C}$ for 30 s, 72 $^{\circ}\text{C}$ for 1 min and 36 cycles of 95 $^{\circ}\text{C}$ for 45 s, 50 $^{\circ}\text{C}$ for 30 s, 72 $^{\circ}\text{C}$ for 1 min and final extension at 72 $^{\circ}\text{C}$ for 15 min. 2 μL of each PCR product and 100 bp DNA ladder (Solis BioDyne, Estonia) were visualised on a 1.5% agarose gel stained with ethidium bromide to confirm amplification and concentration. Good quality PCR products were subsequently purified and sequenced with the respective PCR forward and reverse primers using an ABI3730XL sequencer.

Quality check, editing, and alignments of the resulting DNA sequences were done in Geneious v. 11.1.2 (<https://www.geneious.com>; Kearse et al. 2012). DNA barcode sequences were aligned using MUSCLE (Edgar 2004), primer nucleotide sequences were removed, and chromatograms were checked for the presence of double peaks, stop codons or frameshifts which could indicate the amplification of nuclear mitochondrial (NUMT) pseudogenes. Molecular species identification was




Figure 1. Location of the Maltese Islands within the Mediterranean Sea and a distribution map of the newly recorded seven beetle species across the islands of Malta and Gozo.

carried out by comparing the newly generated DNA barcodes to those available in public databases. The public databases used include the Nucleotide collection (nr/nt) of the NCBI GenBank database (<https://www.ncbi.nlm.nih.gov/genbank>; Benson et al. 2012) and the species level barcode records available at the Barcode of Life Data System (BOLD, <http://www.boldsystems.org>; Ratnasingham and Hebert 2007). Comparison of the query DNA sequences to the databases was done using BLASTn v. 2.9.0 (Zhang et al. 2000; Morgulis et al. 2008) in GenBank and using the identification portal system in BOLD. A species delimitation genetic distance threshold value of 2% was used.

Results

Family Endomychidae

Ancylopus melanocephalus (Olivier, 1808)

Figure 2A

New records. MALTA – Malta • Żebbiegħ; 35°55'14.17" N, 014°22'58.38"E; 33 m a.s.l.; 1.VIII.2018, C. M. Mifsud leg.; UV light trap; Mediterranean stream dominated by *Arundo donax*; 2 ♂, CBRG-UM T10T11 • same locality; 8.IX.2018; 6 ♂, CBRG-UM T60; 4 ♀, CBRG-UM T59; GenBank: MT908984; 1 ♀, CBRG-UM BTS26.

Identification. Body elongate-oval, 5.0–5.5 mm long; elytra with three black maculae laterally and a longitudinal band along the sutural line; elytral basal edge thickened and raised. Female pronotum with lateral sulci connected medially by an arcuate ridge. Male mesofemur with a fringe of long hairs along inner edge. *Ancylopus melanocephalus* (Olivier, 1808) can be distinguished from *Ancylopus ceylonicus* Strohecker, 1971, the most similar species, by the mesotibia of males, which is slightly widened and without a tooth in *A. melanocephalus* (El Torkey and Al Dhafer 2015). The elytra are also a distinguishing feature between the two similar species. The elytra have three black maculae laterally and a longitudinal band along the sutural line in *A. melanocephalus* while in *A. ceylonicus* the elytral suture, apex, and side margin are black with a large mediolateral elytral spot in males (El Torkey and Al Dhafer 2015). Molecular species identification analyses resulted in the highest pairwise identity of 96.8% with *Ancylopus pictus asiaticus* Strohecker, 1972 (KU188382) from GenBank, while no species level barcode records were available from BOLD. The molecular species identification results confirm the genus of our specimen and represent the first DNA barcode for the species *A. melanocephalus*.

Distribution. Saudi Arabia (El Torkey and Al Dhafer 2015); Bulgaria, Hungary, Spain, Italy (Shockley et al. 2009), and Malta (newly recorded).

Family Scarabaeidae

Aplidia transversa (Fabricius, 1801)

Figure 2E

New record. MALTA – Malta • Attard; 35°53'27.34"N, 014°26'24.07"E; 90 m a.s.l.; 9.VI.2017; N. Vella leg.; urban habitat; GenBank: MH510725; 1 ♀, CBRG-UM Col77.

Identification. *Aplidia transversa* (Fabricius, 1801) can be distinguished from the other five species within the genus following the work of Baraud (1992). Body length 15.5 mm, dark red-brown coloration, base of the pronotum without long erect hairs, pronotum punctured in a fine, regular, and simple pattern, hair on elytra very short, shorter than the intervals between the punctures. Molecular species identification analyses resulted in the highest pairwise identity of 99.6% with *Aplidia transversa* (KM449893) from GenBank and also matched the BIN (Barcode Index Number) for *Aplidia transversa* [BOLD:ACD1022] from BOLD. The molecular species identification further confirms the morphological identification of the examined specimen.

Distribution. Albania, Austria, Bosnia and Herzegovina, Bulgaria, Croatia, Germany, Greece, France, Italy, Macedonia, Montenegro, Romania, Serbia, Slovenia, Switzerland, Turkey (Baraud 1992), and Malta (newly recorded).

Family Hydrophilidae

Cercyon quisquilius (Linnaeus, 1761)

New record. MALTA – Malta • Hal Far; 35°49'07.75"N, 014°30'42.26"E; 46 m a.s.l.; 20.VII.2018; C. M. Mifsud leg.; UV light trap; Mediterranean cliff and open habitat; GenBank: MT908996; 1 ♂, CBRG-UM BTS101.

Identification. Body length 2.5 mm; dorsal surface of head black; elytra brownish yellow without dark sutural spot. Mesoventral plate narrower, 6× as long as wide. Metaventrite without femoral lines and with raised pentagonal area markedly wide at midlength. *Cercyon quisquilius* (Linnaeus, 1761) can be confused with *Cercyon nigriceps* (Marsham, 1802) but can be distinguished from it in the coloration of the pronotum, which is blackish with diffuse yellowish areas on lateral margins in *C. quisquilius*, while in *C. nigriceps* it is reddish-brown, similar to the elytral coloration. *Cercyon quisquilius* is also larger than *C. nigriceps* (2.4–3.2 mm vs 1.0–2.1 mm; Arriaga-Varela et al. 2017). Our molecular species identification analyses using the DNA barcode sequence resulted in a pairwise identity of 99.8% with *C. quisquilius* (KU911068) from records on GenBank and also matched the BIN for *C. quisquilius* [BOLD:AAH0271] from BOLD, further confirming our morphological species identification.

Distribution. Native to the Palaearctic region, including Malta (newly recorded). Introduced to Australia, Mexico, Argentina, Hawaii, the Caribbean (Cuba), and China (Arriaga-Varela et al. 2017).




Figure 2. Dorsal habitus. **A.** *Ancylopus melanocephalus* (Olivier, 1808). **B.** *Smicronyx pauperculus* Wollaston, 1864. **C.** *Oxytelus sculptus* Gravenhorst, 1806. **D.** *Lebia cruxminor* (Linnaeus, 1758). **E.** *Aplidia transversa* (Fabricius, 1801).

Family Coccinellidae

Hyperaspis duvergeri Fürsch, 1985

New record. MALTA – Gozo • Fontana; 36°02'11.15"N, 014°14'06.17"E; 70 m a.s.l.; 21.V.2019; N. Vella leg.; Mediterranean stream dominated by *Arundo donax*; GenBank: MT909025; 1 sex indet., CBRG-UM P5_39.

Identification. Body uniformly oval, 3.2 mm long; pronotum black in the middle with orange lateral margins; elytra black with a single pair of orange spots in the posterior half of the elytra. These morphological characteristics separate *H. duvergeri* from other species within the genus *Hyperaspis* using the key by Biranvand et al. (2017). Our molecular species identification analyses resulted in the highest pairwise identity of 91.0% with *Hyperaspis postica* LeConte, 1880 (MF633070), from GenBank, while no species level barcode records were matched on BOLD. Our DNA barcode represents the first DNA barcode record for *H. duvergeri*.

Distribution. Iran (Moddarras-Awal 1997); Algeria, Morocco, Portugal, Spain, France, Italy, Croatia, Hungary (Kovář 2007), and Malta (newly recorded).

Family Carabidae

Lebia cruxminor (Linnaeus, 1758)

Figure 2D

New record. MALTA – Malta • Fawwara; 35°50'19.75"N, 014°24'40.06"E; 185 m a.s.l.; 12.VII.2018; C. M. Mifsud leg.; UV light trap; Mediterranean garrigue habitat; GenBank: MT909027; 1 sex indet., CBRG-UM BTS144.

Identification. Body length 6.0 mm; elytra with dark, transverse band and presence of a scutellar spot, apical margin of elytra dark; legs partially blackened. Our molecular species identification analyses resulted in the highest pairwise identity of 99.1% with *Lebia cruxminor* (Linnaeus, 1758) (KU911743) from GenBank. The molecular species identification on BOLD matched our sequence to the BIN [BOLD:ACB0195] assigned to *L. cruxminor*; further confirming our morphological species identification through DNA barcoding.

Distribution. Europe (de Jong et al. 2014), including Malta (newly recorded), Iran (Azadbakhsh and Nozari 2015), Russia (Egorov et al. 2020), and Japan (GBIF Secretariat 2021a).

Family Curculionidea

***Smicronyx pauperculus* Wollaston, 1864**

Figure 2B

New records. MALTA – Malta • Fawwara; 35°50'19.75" N, 014°24'40.06"E; 185 m a.s.l.; 31.VII.2018; C. M. Mifsud leg.; UV light trap; Mediterranean garrigue habitat; GenBank: MT909077; 1 sex indet., CBRG-UM BTS173 • same locality; GenBank: MH510770; 1 sex indet., CBRG-UM Col250.

Identification. Body length 2.0 mm. Body integument reddish except rostrum, antennae, prothorax, suture of elytra, and tarsi, which are black; vestiture of elytra consisting of elongate, brownish scales not concealing integument and white scales, thicker, forming transverse patches; rostrum moderately downcurved in lateral view (Haran 2018). Our molecular species identification analyses resulted in the highest pairwise identity of 98.6% with *Smicronyx pauperculus* Wollaston, 1864 (KU942313) from GenBank and matched the BIN for *S. pauperculus* [BOLD:ACS7419] from BOLD, thereby supporting the morphological species identification through DNA barcoding.

Distribution. Mediterranean basin and Africa (Haran et al. 2017; Haran 2018), including Malta (newly recorded).

Family Staphylinidae

***Oxytelus sculptus* Gravenhorst, 1806**

Figure 2C

New record. MALTA – Malta • Fawwara; 35°50'19.75" N, 014°24'40.06"E; 185 m a.s.l.; 12.VII.2018; C. M. Mifsud leg.; UV light trap; Mediterranean garrigue habitat; GenBank: MT909046; 1 sex indet., CBRG-UM, voucher ID BTS150.

Identification. Body length 3.0 mm; antennomere 4 notably baso-dished; antennomere 1 stout; supra-antennal ridges elevated (Lü and Zhou 2015). Our molecular species identification analyses resulted in the highest pairwise identity of 99.6% with *Oxytelus sculptus* Gravenhorst, 1806 (KM452399) from GenBank and matched the BIN for *O. sculpus* [BOLD:ABX3754] from BOLD, further supporting our morphological species identification.

Distribution. Palearctic, including Malta (newly recorded), and introduced into North and South America, New Zealand, Australia, and the Hawaiian Islands (Newton 2019).

Discussion

Seven species of coleopterans are newly recorded from Malta in this study. All species presented here are generally distributed on the European mainland or have a Mediterranean distribution (de Jong et al. 2014), and these species are expected to be present on the Maltese Islands. Our new data represent geographic range extensions for

four of these species. Our record of *Cercyon quisquilius* from Malta represents the first from the central Mediterranean, with the closest previously known occurrence being in northern Italy (GBIF Secretariat 2019). Our new data for *Lebia cruxminor* and *Oxytelus sculptus* represent the first records from a Mediterranean island, as both species have been previously recorded only from the Mediterranean mainland (GBIF Secretariat 2021a, 2021b). In addition, our discovery of *Smicronyx pauperculus* from Malta represents its first European records; the closest occurrences are in Israel (Haran et al. 2017).

Each of these newly recorded beetle species was only detected from a single locality on the Maltese Islands (Fig. 1), despite that our sampling efforts were throughout the islands. This observation is likely because these species are locally rare and have a restricted distribution in the islands. *Ancylopus melanocephalus* was collected in UV light traps. *Ancylopus melanocephalus* was recorded at a freshwater habitat, which is a rare habitat throughout Malta, and this habitat specificity possibly contributed to its delayed discovery. The closest geographical records of this species, where it has also been recorded in *Phragmites* marshes, are in southern mainland Italy and Sicily (Focarile 1964). *Lebia cruxminor* is distributed across the vast Palaearctic region (GBIF Secretariat 2021a); however, it is believed to be a rare beetle species throughout its native range (Alexander 2003; Dekoninck et al. 2019). We captured *L. cruxminor* in a UV light trap set up in a Mediterranean garrigue habitat. This species is associated with meadow habitats in Ireland (Alexander 2003) and grasslands in Belgium (Dekoninck et al. 2019). Although mainland European grasslands and Mediterranean garrigue habitats are composed of different plant communities, they share an open habitat type and are characteristically sparsely covered by shrubs and herbs.

In conclusion, our discoveries clearly demonstrate that the coleopteran fauna of the Maltese Islands merit further attention, as the lack of knowledge on island faunas is considered a crucial issue for biodiversity conservation (Gillespie and Will 2018). Our study provides new essential knowledge useful for conservation of biodiversity by providing accurate data on species that were previously overlooked. Additionally, our study contributes towards biodiversity conservation by providing DNA barcodes useful for the identification of coleopteran species and for various future biomonitoring applications across Malta, the central Mediterranean region, and beyond.

Acknowledgements

This work was financially supported by the ENDEAVOUR Scholarships Scheme, Malta [grant no. 168] awarded to CMM; and the BioCon_Innovate Research Excellence Grant from the University of Malta awarded to AV. We would like to thank the subject editor, the copy editor and the two anonymous reviewers for their constructive comments, suggestions, and improvements on earlier versions of this manuscript.

Authors' Contributions

Conceptualization: NV, AV. Formal analysis: CMM. Funding acquisition: CMM, NV, AV. Methodology: CMM. Resources: NV, AV. Supervision: AV. Writing – original draft: CMM, AV. Writing – review and editing: CMM, NV, AV.

References

- Alexander KNA (2003) The third Irish locality for *Lebia cruxminor* (L.) (Coleoptera: Carabidae) and its association with old fen meadow habitat. The Irish Naturalists' Journal 27 (8): 306–308.
- Arriaga-Varela E, Seidel M, Deler-Hernández A, Senderov V, Fikáček M (2017) A review of the *Cercyon* Leach (Coleoptera, Hydrophilidae, Sphaeriiniinae) of the Greater Antilles. ZooKeys 681: 39–93. <https://doi.org/10.3897/zookeys.681.12522>
- Azadbakhsh S, Nozari J (2015) Checklist of the Iranian ground beetles (Coleoptera; Carabidae). Zootaxa 4024 (1): 001–108. <https://doi.org/10.11646/zootaxa.4024.1.1>
- Baraud J (1992) Coléoptères Scarabaeoidea d'Europe. Fédération Française des Sociétés de Sciences Naturelles, Paris, Société Linnéenne, Lyon, France, 856 pp.
- Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW (2012) GenBank. Nucleic Acids Research 41 (D1): D36–D42. <https://doi.org/10.1093/nar/gks1195>
- Biranvand A, Tomaszewska W, Li W, Nicolas V, Shakarami J, Fekrat L, Hesami S (2017) Review of the tribe Chilocorini Mulsant from Iran (Coleoptera, Coccinellidae). ZooKeys 712: 43–68. <https://doi.org/10.3897/zookeys.712.20419>
- Bordoni A (1973) I Coleotteri Stafilinidi delle isole circumsiciliane. XXI contributo alla conoscenza degli Staphylinidae. Biogeographia – The Journal of Integrative Biogeography 3: 651–754. <https://doi.org/10.21426/B63110069>
- Bouchard P, Smith ABT, Douglas H, Gimmel ML, Brunke AJ, Kanda K (2017) Biodiversity of Coleoptera. In: Footitt RG, Adler PH (Eds.) Insect biodiversity: science and society. Second edition. Volume I. John Wiley & Sons, Oxford, UK, 337–417. <https://doi.org/10.1002/9781118945568.ch11>
- Cameron M, Caruana Gatto A (1907) A list of the Coleoptera of the Maltese Islands. Transactions of the Entomological Society of London 59 (3): 383–403.
- de Jong Y, Verbeek M, Michelsen V, Bjørn P, Los W, Steeman F, Bailly N, Basire C, Chylarecki P, Stloukal E, Hagedorn G, Wetzel F, Glöckler F, Kroupa A, Korb G, Hoffmann A, Häuser C, Kohlbecker A, Müller A, Güntsch A, Stoev P, Penev L (2014) Fauna Europaea – all European animal species on the web. Biodiversity Data Journal 2: e4034. <https://doi.org/10.3897/BDJ.2.e4034>
- DeKoninck W, Baugnée J, Bleich O, Bonamie G, Boosten G, Breaeckman A, Crèvecoeur L, Dahan L, Delwaide M, Drumont A, Felix R, Hendrickx R, Ignace D, Jacobs M, Lambrechts K, Lambrechts J, Limbourg P, Maquet D, Migon M, Minet G, Muls D, Potier J, Raemdonck H, Renner K, Renneson J, Scheers K, Stassen E, Struyve T, Thys N, Troukens W, Van Grimberge A, Van Kerckvoorde M, Van Malderen M and Hendrickx F (2019) Records of new and rare carabid beetles (Coleoptera: Carabidae) for Belgium collected or identified during the period 2008–2018. Bulletin de La Société Royale Belge d'Entomologie 155 (2): 19–63.
- Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Research 32 (5): 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Egorov LV, Ruchin AB, Semenov VB, Semionov OI, Semishin GB, Penev L (2020) Checklist of the Coleoptera of Mordovia State Nature Reserve, Russia. ZooKeys 962: 13–22. <https://doi.org/10.3897/zookeys.962.54477>
- El Torkey AM, Al Dhafer HM (2015) First record of a handsome fungus beetle (Coleoptera: Endomychidae) from the Arabian Peninsula. The Coleopterists Bulletin 69 (4): 824–827. <https://doi.org/10.1649/0010-065X-69.4.824>
- Focarile A (1964) La geonemia in Italia di *Ancylopus melanocephalus* (Oliv.) (Coleoptera Endomychidae). Bollettino della Società Entomologica Italiana 94: 132–138.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology and Biotechnology 3 (5): 294–299.
- Gillespie RG, Will K (2018) Biodiversity of Arthropods on Islands. In: Footitt RG, Adler PH (Eds.) Insect biodiversity: science and society. Volume II. John Wiley & Sons, Oxford, UK, 81–104. <https://doi.org/10.1002/9781118945582.ch4>
- GBIF (Global Biodiversity Information Facility) Secretariat (2019) *Cercyon quisquilius* (Linnaeus, 1760), Copenhagen, Denmark, GBIF Backbone Taxonomy. <https://www.gbif.org/species/1034424>. Accessed on: 2021-6-30.
- GBIF (Global Biodiversity Information Facility) Secretariat (2021a) *Lebia cruxminor* (Linnaeus, 1758), Copenhagen, Denmark, GBIF Backbone Taxonomy. <https://www.gbif.org/species/6097856>. Accessed on: 2021-6-9.
- GBIF (Global Biodiversity Information Facility) Secretariat (2021b) *Oxytelus sculptus* Gravenhorst, 1806. Copenhagen, Denmark, GBIF Backbone Taxonomy. <https://www.gbif.org/species/1039973>. Accessed on: 2021-6-10.
- Haran J (2018) A review of the genus *Smicronyx* Schoenherr (Coleoptera, Curculionidae, Curculioninae) in tropical Africa. Zootaxa 4508 (2): 267–287. <https://doi.org/10.11646/zootaxa.4508.2.9>
- Haran J, Schutte A, Friedman A (2017) A review of *Smicronyx* Schoenherr (Coleoptera, Curculionidae) of Israel, with description of two new species. Zootaxa 4237 (1): 17–40. <https://doi.org/10.11646/zootaxa.4237.1.2>
- Hebert PDN, Penton EH, Burns JM, Janzen DH, Hallwachs W (2004) Ten species in one: DNA barcoding reveals cryptic species in the Neotropical skipper butterfly *Astraptes fulgerator*. Proceedings of the National Academy of Sciences of the United States of America 101 (41): 14812–14817. <https://doi.org/10.1073/pnas.0406166101>
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics 28 (12): 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>
- Kovář I (2007) Coccinellidae. In: Löbl I, Smetana A (Eds.), Catalogue of Palaearctic Coleoptera. Volume 4. Elateroidea, Derodontoida, Bostrichoidea, Lymexyloidea, Cleroidea, Cucujoidea. Apollo Books, Stenstrup, Denmark, 71–74, 568–630.
- Lillig M, Barthel HB, Mifsud D (2012) An identification and informative guide to the Tenebrionidae of Malta (Coleoptera). Bulletin of the Entomological Society of Malta 5: 121–160.
- Lü L, Zhou HZ (2015) Revision of the genus *Oxytelus* Gravenhorst (Staphylinidae: Oxytelinae) in Southeast Asia. Zootaxa 3992 (1): 1–94. <https://doi.org/10.11646/zootaxa.3992.1.1>
- Magrini P, Schembri S (1997) A topographic catalogue of the Carabidae of the Maltese Islands (Coleoptera). Bollettino della Società Entomologica Italiana 129 (3): 213–234.
- Mifsud D, Vigna Taglianti A (1999) Notes on three carabids collected from the Maltese Islands (Coleoptera, Carabidae). The Central Mediterranean Naturalist 3 (1): 27–28.
- Mifsud D (2000a) Present knowledge of the entomofauna of the Maltese Islands. Entomologica Basiliensia 22: 75–86.
- Mifsud D (2000b) Ptinid beetles (Coleoptera, Ptinidae) from the Maltese Islands. The Central Mediterranean Naturalist 3 (2): 43–48.
- Mifsud D (2002) Longhorn beetles (Coleoptera, Cerambycidae) of the Maltese Islands (Central Mediterranean). The Central Mediterranean Naturalist 3 (4): 161–169.
- Mifsud D, Colonnelli E (2010) The Curculionidae of the Maltese Islands

- lands (Central Mediterranean) (Coleoptera). Bulletin of the Entomological Society of Malta 3: 55–143.
- Mifsud D, Colonnelli E (2018) Two new records of weevils from Malta, with a new synonymy (Coleoptera: Curculionoidea). Revista Gaditana de Entomologia 9 (1): 43–47.
- Mifsud D, Jelínek J (2012) The Mycetophagidae of the Maltese Islands (Coleoptera). Bulletin of the Entomological Society of Malta 5: 105–109.
- Mifsud D, Knizek M (2009) The bark beetles (Coleoptera: Scolytidae) of the Maltese Islands (Central Mediterranean). Bulletin of the Entomological Society of Malta 2 (1): 25–52.
- Mifsud D, Nardi G (2020) New records of adepagous water beetles (Coleoptera: Gyrinidae, Haliplidae, Dytiscidae) from the Maltese Islands. Bulletin of the Entomological Society of Malta 11: 119–126.
- Mifsud D, Scupola A (1998) The Tenebrionidae (Coleoptera) of the Maltese Islands (Central Mediterranean). Annali del Museo Civico di Storia Naturale Giacomo Doria 9: 191–229.
- Moddarras-Awal M (1997) List of agricultural pests and their natural enemies in Iran. Fourth Edition. Ferdowsi University Press, Mashhad, Iran, 429 pp.
- Morgulis A, Coulouris G, Raytselis Y, Madden TL, Agarwala R, Schäffer AA (2008) Database indexing for production Megablast searches. Bioinformatics 24 (16): 1757–1764. <https://doi.org/10.1093/bioinformatics/btn322>
- Nardi G, Mifsud D (2015) The Bostrichidae of the Maltese Islands (Coleoptera). ZooKeys 481: 69–108. <https://doi.org/10.3897/zookeys.481.8294>
- Newton AF (2019) StaphBase: Staphyliniformia world catalog database. Species 2000 ITIS Catalogue of Life, 2019 Annual Checklist. Leiden, the Netherlands. <http://www.catalogueoflife.org/annual-checklist/2019>. Accessed on: 2021-6-30.
- Pivotti I, Agoglitta R, Zunino M, Piattella E, Dellacasa M, Corallini C, Mifsud D (2011) The Scarabaeoidea of the Maltese Islands (Central Mediterranean) (Coleoptera). Bulletin of the Entomological Society of Malta 41: 85–124.
- Przewoźny M (2018) Catalogue of Palearctic Hydrophiloidea (Coleoptera). Internet version 2018-01-01, Adam Mickiewicz University, Poland. <http://www.waterbeetles.eu>. Accessed on: 2021-6-30.
- Ratnasingham S, Hebert PDN (2007) BOLD: the Barcode of Life Data System. Molecular Ecology Notes 7: 355–364. <https://doi.org/10.1111/j.1471-8286.2007.01678.x>
- Sabella G, Mifsud D (2016) The Pselaphinae (Coleoptera, Staphylinidae) of the Maltese Archipelago. Bulletin of the Entomological Society of Malta 8: 7–34.
- Schembri PJ (2003) Current state of knowledge of the Maltese non-marine fauna. In: Malta Environment and Planning Authority Annual Report and Accounts 2003. Malta Environment and Planning Authority, Floriana, Malta, 33–65.
- Schembri S (1993) The coccinellid fauna of the Maltese Islands (Central Mediterranean): an annotated list (Coleoptera Coccinellidae). Bollettino Della Societa Entomologica Italiana 125 (1): 45–52.
- Schembri SP, Schembri JL (1982) Recent records of Coleoptera Staphylinidae from the Maltese Islands. Lavori Della Societa' Veneziana Di Scienze Naturali 7: 45–53.
- Shockley FW, Tomaszewska KW, McHugh JV (2009) An annotated checklist of the handsome fungus beetles of the world (Coleoptera: Cucujoidea: Endomychidae). Zootaxa 1999 (1): 1–113. <https://doi.org/10.11646/zootaxa.1999.1.1>
- Sogoh K, Yoshitomi H (2017) A revision of the genus *Ancylopus* (Coleoptera, Endomychidae) of Japan. Elytra (New Series) 7 (2): 421–438.
- Zhang Z, Schwartz S, Wagner L, Miller W (2000) A greedy algorithm for aligning DNA sequences. Journal of Computational Biology 7 (1–2): 203–214. <https://doi.org/10.1089/10665270050081478>