



Elevational range extension of Drab Water Tyrant, *Ochthornis littoralis* (Pelzeln, 1868) (Passeriformes, Tyrannidae)

Jefferson García-Loor^{1,2}

1 Konrad Lorenz Research Center, Core Facility for Behavior and Cognition, University of Vienna, Grünau im Almtal, Austria

2 Department of Behavioral and Cognitive Biology, University of Vienna, Vienna, Austria • garcialoorjefferson@gmail.com  <https://orcid.org/0000-0002-7427-9652>

Abstract

The Drab Water Tyrant, *Ochthornis littoralis* (Pelzeln, 1868), is a small flycatcher occurring across lower Amazonia. Its usual habitat is the edges of rivers, and it is easily observed flying over sandbanks and among the trunks of fallen trees and branches. Here, I present a new locality in the Ecuadorian Andean foothills which represents a significant increase in this species' elevational range from the Amazon lowlands. The new record is also from a previously unknown habitat for the species.

Keywords

Amazonian lowlands, Andes, new location

Academic editor: Guilherme Sementili-Cardoso | Received 23 October 2021 | Accepted 18 April 2022 | Published 4 May 2022

Citation: García-Loor J (2022) Elevational range extension of Drab Water Tyrant, *Ochthornis littoralis* (Pelzeln, 1868) (Passeriformes, Tyrannidae). Check List 18 (3): 451–453. <https://doi.org/10.15560/18.3.451>

Introduction

The Drab Water Tyrant, *Ochthornis littoralis* (Pelzeln, 1868) (Passeriformes, Tyrannidae), is a small flycatcher occurring across northern South America in Colombia, Venezuela, Guyana, French Guyana, Brazil, Ecuador, Peru, and Bolivia. It is associated with sandbanks along the edges of rivers (Farnsworth and Langham 2019). Its usual altitudinal range is 400–900 m a.s.l. (Hilty 2003; Restall et al. 2006; Schulenberg et al. 2007; Ber Van Perlo 2009; Herzog et al. 2016; Freile and Restall 2018), and the highest record is from Bolivia. In Ecuador, *O. littoralis* occurs throughout the Amazonian lowlands below 400 m on the eastern slope of the Andes (Ridgely and Greenfield 2001; McMullan and Navarrete 2017; Freile and Restall 2018) (Fig. 1). In this region, this flycatcher species is considered to be a resident, and it

usually observed flying low between tree trunks and branches (Ridgely and Greenfield 2001). On 26 January 2018, five individuals were observed as part of routine transect observations for a bird-monitoring project, in a location that is here reported as a new elevation record for this species.

Methods

Field methods consisted of fixed-band line transects with a length of 1000 m. Two transects were placed across the facilities of the Topo River hydroelectric power plant, Tungurahua, Ecuador. The transects were conducted at 06:00–10:00 in the morning and at 15:00–18:00 in the afternoon. Although there was a mist-net catching




Figure 1. Distribution map of *Ochthornis littoralis* in Ecuador; the red star denotes the new record. Gray areas represent the species' distribution.

station in place, *Ochthornis littoralis* was not caught. The mist nets were opened during the same times mentioned above.

Results

Ochthornis littoralis (Pelzeln, 1868)

Figure 2

New record. ECUADOR – **Tungurahua** • Río Negro; Topo River Hydroelectric Power Plant; 01°21'43"S, 078°12'40"W; 1560 m alt.; 26.I.2018; Jefferson García Loor obs.; Lower Montane Evergreen Forest found on the eastern Andes; 5 adults individuals observed, no sexual differentiation.

The birds were observed around the concrete wall in the water-catchment area of the hydroelectric power plant. Two individuals of *O. littoralis* were observed pecking and foraging in moss on the ground (Fig. 2A). The other three individuals were observed at the same time on the road to the plant; they had a similar foraging behavior, pecking the ground for small insects (Fig. 2B). On the second day, I saw two individuals which were pecking for invertebrates on the ground.

The vegetation in the area is characterized by the presence of ferns, herbs, and human-maintained lawn. The river's edge showed signs of heavy erosion. The surroundings are a human-modified secondary open forest.

The vegetation height was ~5 m and the vegetation consisted mostly of succulents and a few trees. The south side of the water catchment area had less disturbance than the north side and had a slope of ~75° and ~8 m high vegetation of trees and shrubs.

Identification. The identification was based on the plumage pattern (Fig. 2), which is consistent with the descriptions in the literature: sandy brown body, browner crown, darker tail and wings, and facial pattern with pale supercilium. The position of the body in a more inclined-horizontal posture and foraging behavior were determinants. A similar species in Ecuador is River Tyrannulet (*Serpophaga hypoleuca* Sclater & Salvin, 1866), but it has a clearer neck, chest, and ventral area, and also lacks the facial patterns and dark shades on the wings; it is also a less frequently encountered species in the lower Amazon region than *O. littoralis*. Little Ground Tyrant (*Syrtidicola fluviatilis* (Sclater & Salvin, 1866)) is also similar, but its slimmer bill, longer tail, and absence of a pale supercilium differs from *O. littoralis* (Rigdelly and Greenfield 2001; McMullan and Navarrete, 2017; Freile and Restall 2018).

Discussion

This report expands the elevational range of *Ochthornis littoralis* by 1160 m in Ecuador (Freile and Restall 2018), 660 m higher than the highest known occurrence



Figure 2. *Ochthornis littoralis* recorded at 1560 m a.s.l. near the Topo River hydroelectric power plant, Tungurahua, Ecuador. **A.** Frontal view. **B.** Side view.

anywhere in its range. The general vegetation in the water-capture area of the hydroelectric plant was typical of the Low Montane Evergreen Forest found on the eastern Andes, which differs considerably from the Evergreen Forest of the Amazonian lowlands and other riparian ecosystems where the bird is usually observed (MAE 2013). Referring to the system of bioclimatic classification used for the Ecuadorian fauna, this species is now known to occupy not only the Oriental Tropical Area but also the Oriental Subtropical Area (Albuja et al. 2012). The habitat at the new locality presents human disturbance and lacks the typical sandbanks of the usual riparian habitats where *O. littoralis* usually occurs. The vegetation is also lower, and the current in the river is much greater.

Although the habitat differs at the site of the new record from what is typical for *O. littoralis*, the strong relationship between this species and water was observed at this site. Precipitation and temperature in the species' usual range is, on average, 200 mm and 25 °C, respectively, whereas in the new site reported here, temperatures average 20°C, although precipitation is the same (Albuja et al. 2012; MAE 2013). The species may be seasonally moving, following the rains and precipitation patterns. Young individuals may also be exploring new territories looking for food before to start the breeding season. The closest localities where *O. littoralis* has been recorded are approximately 70 km north or south from the new record, but considering that the Topo River is a tributary of the larger Pastaza River, where this species is known to occur, it is likely that the newly recorded individuals originated from the southern population.

Acknowledgements

I am grateful to Juan Freile, Manuel Sánchez, and Galo Buitrón for the confirmation of the bird identification of

the observed individuals. I also thank Pablo Medrano for his help with the map, and Sonia Kleindorfer, Elisa Bonaccorso, and Héctor Cadena for their comments and advice. KAYMANTA Consulting Group funded the fieldwork on which the manuscript was based. I thank the reviewers and academic editor for their comments and advice.

References

- Albuja, L, Almendáriz A, Barriga R, Montalvo LD, Cáceres F, Román JL (2012) Fauna de vertebrados del Ecuador. Instituto de Ciencias Biológicas. Escuela Politécnica Nacional, Quito, Ecuador, 490 pp.
- Farnsworth A, Langham G (2019) Drab Water-tyrant (*Ochthornis littoralis*). In: del Hoyo J, Elliott A, Sargatal J, Christie D A, de Juana E (Eds.) Handbook of the birds of the world alive. Lynx Edicions, Barcelona, Spain. <https://www.hbw.com/node/57409>. Accessed on: 2022-4-28.
- Freile J, Restall R (2018). Birds of Ecuador. Helm Field Guides, London, UK, 656 pp.
- Herzog S, Terril R, Jhan A, Remsen J, Maillard O, García-Soliz V, Macleod R, Maccormick A, Vidoz J (2016) Birds of Bolivia. Asociación Armonía, Santa Cruz de la Sierra, Bolivia, 324 pp.
- MAE (2013) Sistema de clasificación de ecosistemas del Ecuador continental. Subsecretaría de Patrimonio Natural, Quito, Ecuador, 232 pp.
- McMullan M, Navarrete L (2017) Fieldbook of the birds of Ecuador, including the Galapagos Islands and common mammals. 2nd edition. Ratty Ediciones, Quito, Ecuador, 228 pp.
- Ridgely R, Greenfield P (2001) The birds of Ecuador: status, distribution, and taxonomy. Vol. 1. Cornell University Press, Ithaca, USA, 848 pp.
- Schulenberg T, Stotz D, Lane D, O'Neill J, Parker T (2010) Birds of Peru. Revised edition. Princeton University Press, Princeton, USA, 656 pp.
- Van Perlo B (2009) A field guide to the birds of Brazil. Oxford University Press, Oxford, UK, 465 pp.