



# First record of *Uperodon globulosus* (Günther, 1864) (Anura, Microhylidae) in Rajshahi Division, Bangladesh

Md. Mokhlesur Rahman, Md. Mahabub Alam, Md. Fazle Rabbe

University of Dhaka, Department of Zoology, Dhaka 1000, Bangladesh.

**Corresponding author:** Md. Mokhlesur Rahman, [mmrahman48@du.ac.bd](mailto:mmrahman48@du.ac.bd)

## Abstract

*Uperodon globulosus* (Günther, 1864) has so far been recorded from 6 localities in Bangladesh over the past 3 decades. In June 2017, we found an adult female in a courtyard of a house in Jonail village (Natore district, Rajshahi division, Bangladesh), which is approximately 102 km southwest of the previous nearest previously known occurrence of this species.

## Key words

Balloon Frog; species distribution; Natore.

**Academic editor:** Perry L. Wood, Jr | Received 24 August 2017 | Accepted 4 January 2018 | Published 23 February 2018

**Citation:** Rahman MM, Alam, MM, Rabbe MF (2018) First record of *Uperodon globulosus* Günther, 1864 (Anura, Microhylidae) in Rajshahi Division, Bangladesh. Check List 14 (1): 277–280. <https://doi.org/10.15560/14.1.277>

## Introduction

*Uperodon globulosus* (Günther, 1864), a species belonging to the family Microhylidae and commonly known as the Balloon Frog, is characterized by its enormously distensible lungs that swell like a globular puffball, giving the appearance of an inflated balloon (Asmat 2009). This species is found in southern Nepal, northeastern India to Gujarat, southwestward to Karnataka and Kerala, India, and Bangladesh (Ahsan 2015). *Uperodon globulosus* is a nocturnal fossorial species that is found in forests and agricultural lands and breeds in shallow ponds and marshy wetlands following heavy rainfall (Khan 1987, Dutta et al. 2004, Asmat 2009, Ahsan 2015). These frogs move mainly by walking and/or hopping, and have the ability to float on water (Daniel 2002, Asmat 2009).

## Methods

We documented the species using a Canon PowerShot SX50 HS digital camera and deposited photographic

vouchers in the Professor Md. Kazi Zaker Husain Museum, Department of Zoology, University of Dhaka, Bangladesh. The specimen was not collected because it is protected by law. This species is categorized as Vulnerable under criteria B1ab(iii), according to IUCN Bangladesh (2015) and is a protected animal under Schedule I of the *Bangladesh Wildlife (Conservation and Security) Act*, 2012.

## Results

**New record.** Bangladesh, Rajshahi division, Natore district: Boraigram: Jonail village (24°16'39" N, 089°13'25" E; WGS84), Md. Mokhlesur Rahman, 18 June 2017, in courtyard of a house (photographic voucher number W14001a-b, 1 adult female).

**Identification.** Preliminary identification was done in the field by the authors, using the field guide by Hasan et al. (2014). This identification was later confirmed by Professor Mohammad Firoj Jaman. Prior to photographing the



**Figure 1.** Adult female *Uperodon globulosus* from Natore, Rajshahi division. Photograph by Md. Mokhlesur Rahman.

individual, it was observed inflating its body. This specimen agrees with the description in Asmat (2009), Hasan et al. (2014), and Khan (2015) by having a variable dorsal color within the range of variation known for the species (reddish brown to greenish grey or olive), a dirty whitish ventrum, a granular anal region, fingers not webbed (with the first finger shorter than the second), fingertips rounded, hind limbs short, and toes with rudimentary webs (Fig. 1).

## Discussion

Since its discovery in 1980 from Madhupur forest, Mymensingh (Khan 1982, 2004), *U. globulosus* has been recorded from 6 localities in Bangladesh and with few sightings over the last 3 decades (Table 1). Mahony et al. (2009) recorded 2 juveniles from the same forest of Madhupur track in Rasulpur, Tangail district, and they also recorded 2 males from Savar, Dhaka. Another individual


was found from a paddy field of Berakuthi village, Barua, Nilphamari (Sarkar et al. 2012). Most recently, Reza and Perry (2015) documented this species from Milonchori, Bandarban (Table 1).

The habitat we report is dissimilar to previously recorded habitats (Table 1). Our new record expands the known geographic distribution of this species; this record is approximately 176 km south of Nilphamari, 115 km northwest of Savar, 102 km southwest of Tangail, and 390 km northwest of Bandarban (Fig. 2). Moreover, this new record might help in promoting more studies on the ecology, behavior, and other aspects of *U. globulosus*. Considering that this species occurs in a variety of different types of habitats, we conclude that gaps between known occurrences of *U. globulosus* are due to limited sampling rather than environmental restrictions. With further fieldwork, we expect to find additional records between the currently known occur-

**Table 1.** Known locality records for *U. globulosus* in Bangladesh.

Place	Latitude	Longitude	Habitat	Reference
Madhupur forest, Mymensingh	24°42' N *	090°09' E *	Barren patch	Khan 1982
Savar, Dhaka	23°52' N	090°16' E	Semi-urban	Mahony et al. 2009
Rasulpur, Madhupur, Tangail	24°41' N	090°08' E	Forest edge	Mahony et al. 2009
Berakuthi, Barua, Nilphamari	25°49'33" N	088°49'45" E	Paddy field	Sarker et al. 2012
Milonchori Hillside resort, Bandarban	22°10'04" N	092°13'24" E	Tropical forest	Reza and Perry 2015
Jonail, Boraigram, Natore	24°16'39" N	089°13'25" E	Courtyard	Present study

\* Geographic coordinates approximate (subsequently derived here).



**Figure 2.** Map showing the localities where *Uperodon globulosus* has been recorded. Colored areas represent divisions within Bangladesh.

rences and to be better able to describe the distribution of *U. globulosus* in Bangladesh.

### Acknowledgements

We thank Professor Mohammad Firoj Jaman for confirmation of the identification of the species, Mr Md. Salauddin for assisting to prepare the distribution map, and the 2 anonymous reviewers for helpful comments and suggestions on the earlier version of the manuscript.

### Authors' Contributions

MMR and MMA collected the data, MMR and MFR wrote the text, and MMA made the analysis.

### References

Ahsan MF (2015) *Uperodon globulosus*. In: IUCN Bangladesh. Red List of Bangladesh Volume 4: Reptiles and Amphibians. IUCN Bangladesh Country Office, Dhaka, 228.  
 Asmat GSM (2009) *Uperodon globulosus*. In: Kabir SMH, Ahmad M, Ahmed ATA, Rahman AKA, Ahmed ZU, Begum ZNT, Hassan

- MA, Khondker M (Eds) Encyclopedia of Flora and Fauna of Bangladesh. Volume 25: Amphibians and Reptiles. Asiatic Society of Bangladesh, Dhaka, 27–28.
- Daniel JC (2002) The Book of Indian Reptiles and Amphibians. Bombay Natural History Society, Oxford University Press, Mumbai, 238 pp.
- Dutta S, Padhye A, Sengupta S, Sarker SU (2004) *Uperodon globulosus*. The IUCN Red List of Threatened Species. <http://doi.org/10.2305/IUCN.UK.2004.RLTS.T58022A11717889.en>. Accessed on: 2017-8-23.
- Günther ACLG (1864) The Reptiles of British India. Ray Society, London, 452 pp.
- Hasan MK, Khan MMH, Feeroz MM (2014) Amphibians and Reptiles of Bangladesh: A Field Guide. Arannayk Foundation, Dhaka, Bangladesh, 191 pp.
- IUCN Bangladesh (2015) Red List of Bangladesh Volume 4: Reptiles and Amphibians. IUCN Bangladesh Country Office, Dhaka, 320 pp.
- Khan MAR (1982) Wildlife of Bangladesh. The University of Dhaka, Dhaka, 174 pp.
- Khan MAR (1987) Wildlife of Bangladesh Volume 1: Amphibians and Reptiles. Bangla Academy, Dhaka, 169 pp.
- Khan MAR (2004) Checklist of the herpetofauna of Bangladesh. Cobra 57: 1–31.
- Khan MAR (2015) Wildlife of Bangladesh: Checklist and Guide. Chayabithi publication, Dhaka, 567 pp.
- Mahony S, Hasan MK, Kabir MM, Ahmed M, Hossain MK (2009) A catalogue of amphibians and reptiles in the collection of Jahangirnagar University, Dhaka, Bangladesh. Hamadryad 34 (1): 80–94.
- Reza AHMA, Perry G (2015) Herpetofaunal species richness in the tropical forests of Bangladesh. Asian Journal of Conservation Biology 4 (2): 100–108.
- Sarkar MAR, Howlader MSA, Kabir MA (2012) Geographic Distribution. *Uperodon globulosus*. Herpetological Review 43 (2): 301–302.