



First record of *Phragmatopoma caudata* Krøyer in Mörch, 1863 (Annelida, Sabellariidae) in the state of Piauí, northern Brazilian coast: westernmost record in Brazil

José Gerardo Ferreira Gomes Filho,¹ Marcus Vinícius Quixabeira Bezerra dos Santos,² Rosana Aquino de Souza³

1 Universidade Federal do Piauí, Curso de Engenharia de Pesca, Campus Ministro Reis Veloso, Av. São Sebastião, CEP: 64.202-020. Parnaíba, PI, Brazil. **2** Universidade Federal do Piauí, Departamento de Biologia, Centro de Ciências da Natureza, Campus Ministro Petrônio Portela, Bairro Ininga, CEP: 64049-550. Teresina, PI, Brasil. **3** Universidade Federal do Piauí, Curso de Ciências Biológicas, Campus Ministro Reis Veloso, Av. São Sebastião, CEP: 64.202-020. Parnaíba, PI, Brazil.

Corresponding author: José Gerardo Ferreira Gomes Filho, gerardogomes@ufpi.edu.br

Abstract

We report the first occurrence of *Phragmatopoma caudata* Krøyer in Mörch, 1863 in Piauí state, Brazil. This is the first record of this species in the APA Delta do Parnaíba, a national environmental protection area that comprises areas from 3 Brazilian states, Maranhão, Piauí and Ceará. This is also the northernmost and westernmost record of this species in Brazil to date, further extending by 415 km its known distribution in Brazil, and helping to fill a gap on the knowledge of its distribution in South America.

Key words

Polychaeta; Sabellida; reef-building polychaete; new record; extension range; Delta do Parnaíba.

Academic editor: Sérgio N. Stampar | Received 9 March 2017 | Accepted 24 June 2017 | Published 18 August 2017

Citation: Gomes-Filho JGF, Santos, MVQB, Aquino-Souza RA (2017) First record of *Phragmatopoma caudata* Krøyer in Mörch, 1863 (Annelida, Sabellariidae) in the state of Piauí, northern Brazilian coast: westernmost record in Brazil. Check List 13 (4): 343–348. <https://doi.org/10.15560/13.4.343>

Introduction

The coastline of the state of Piauí is 66 km long, and despite being the shortest coastline among the Brazilian states (Fig. 1), it is characterized by a wide variety of natural environments such as mangroves, dunes, lagoons, granitic and sandstone rocky outcrops, estuaries and deltas. These different environments, and the interplay between them, probably contribute to a high local biodiversity, although few efforts have been made to survey Piauí's marine invertebrate fauna. As a result, the coastline of Piauí is an area where there is still a great uncertainty regarding the richness and composition of

marine species, although all its extension is included within a national environmental protection area, the Delta do Parnaíba Environmental Protection Area (APA Delta do Parnaíba) (Fig. 2).

Phragmatopoma caudata Krøyer in Mörch, 1863 belongs to the family Sabellariidae Johnston, 1865, and like many species from this family is a reef-building species. Reef-building polychaetes, like other bioconstructing species, are able to increase heterogeneity of the substrata, generating new habitats, and thus contributing to biodiversity (Jones et al. 1994, Goldberg 2013). Observations on increases in taxonomic diversity and on reef associated species in Brazil and elsewhere have




Figure 1. Records of *Phragmatopoma caudata* in Brazilian states and the nearest westward record. (+) Previous records. Venezuela: Ilhas Tortuga e Ilhas Margarita (Kirtley 1994). Brazil: Ceará-CE (Fausto-Filho and Furtado 1970, Kirtley 1994, Fournier et al. 2010; Nunes et al. 2016), Rio Grande do Norte-RN (Fournier et al. 2010), Paraíba-PB (Fournier et al. 2010, Santos et al. 2011), Pernambuco-PE (Nunes et al. 2016), Bahia-BA (Santa-Isabel et al. 2000), Espírito Santo-ES (Nunes et al. 2016), Rio de Janeiro-RJ (Amaral 1987, Kirtley 1994, Lana and Bremec 1994, Moreno and Rocha 2006, Masi and Zalmon 2008, Masi et al. 2009, Occhioni et al. 2009, Fournier et al. 2010), São Paulo-SP (Amaral 1987, Morgado and Amaral 1989, Lana and Bremec 1994, Rocha 1995, Micheletti-Flores and Negreiros-Fransozo 1999, Fransozo et al. 2000, Occhioni et al. 2009, Fournier et al. 2010, Nunes et al. 2016), Paraná-PR (Lana and Bremec 1994, Bosa and Masunari 2002a, 2002b, Occhioni et al. 2009, Fournier et al. 2010, Nunes et al. 2016), Santa Catarina-SC (Kirtley 1994, Lana and Bremec 1994, Fournier et al. 2010) and Rio Grande do Sul-RS (Amaral 1987, Lana and Bremec 1994). (●) New record. Letters correspond to Brazilian states abbreviations.

provided evidence of the role of sabellariid reefs in supporting high levels of diversity (Micheletti-Flores and Negreiros-Fransozo 1999, Bosa and Masunari 2002a, 2002b, Dubois et al. 2002, 2006, Dias and Paula 2001, Sepúlveda et al. 2003, Pholer 2004, Ataíde et al. 2014). It is likely that *P. caudata* reefs contribute to invertebrate diversity in rocky areas of the APA Delta do Parnaíba, and therefore it is important to register and monitor local occurrence and abundances of this species.

Regarding other polychaete species, 5 belonging to 2 families were previously recorded at the APA Delta do Parnaíba. *Laeonereis culveri* (Webster, 1879) (Pamplin et al. 2007), *Alitta succinea* (Leuckart, 1847) and *Perinereis anderssoni* Kinberg, 1866 (Santos and Steiner 2006) and *Nereis riisei* Grube, 1857 (Santos and Lana 2003) belonging to the family Nereididae Blainville, 1818; and *Chloëia*

viridis Schmarda, 1861 (Amaral and Nonato 1994) belonging to the family Amphinomidae Lamarck, 1818.

The world known distribution of *P. caudata* is restricted




Figure 2. Location of Pedra do Sal Beach, in Piauí (02°48'13" S, 041°43'48" W), and limits of the APA Delta do Parnaíba.



Figure 3. *Phragmatopoma caudata* at Pedra do Sal Beach, Parnaíba, Brazil.

to the Western Atlantic and extends from Brazil to Florida, including locations in Venezuela, the Caribbean, Mexico and Texas (Kirtley 1994). The type locality is the Caribbean, but the exact location is unknown (Kirtley 1994). *Phragmatopoma lapidosa* Kinberg, 1867 is the main synonym of *P. caudata* as suggested by Kirtley (1994) and confirmed by molecular data (Nunes et al. 2016). The later work also provide evidence for the existence of genetic connectivity among populations across the whole distribution range of the species, and emphasizes the importance of new records of this species along this range in order to fill the gaps on its distribution knowledge and to confirm or deny the existence of discontinuities within its geographic limits (Nunes et al. 2016).

The present study reports the first record of *Phragmatopoma caudata* off the coast of the Piauí and gives detailed information on the morphology and characters used in its identification; the new data extend the known geographic distribution of this species in Brazil and improves the knowledge of the fauna of the APA Delta do Parnaíba. The coastline distance that separates the closest known record of the species in the country, in Fortaleza-Ceará (Fausto-Filho and Furtado 1970, Kirtley 1994), from the new record reported here is of approximately 415 km.

Methods

Sand reefs of *Phragmatopoma caudata* were first noticed in December 2010 (Fig. 3) at Pedra do Sal beach (02°48' 13" S, 041°43'48" W), municipality of Parnaíba, Brazil (Fig. 2). In April 2011 those reefs were studied in more details as follows.

Colonies were found on granitic natural substrata, in the intertidal region of Pedra do Sal beach. With the aid of a chisel and a mallet, 5 cubes of about 0.1 m edge (0.001 m³) were taken from random colonies. Subsequently, each cube was broken into smaller pieces and the individuals found were collected and fixed in 4% formaldehyde. In laboratory, 4% formaldehyde was replaced by 70% ethanol. With the aid of needles, scalpels, and a

microscope, some individuals were examined and their morphology was observed. The opercular structure of some individuals was dissected to expose the opercular paleae (outer, middle and inner), due to the necessity of observing their shape and relative position to assure species identification, following Kirtley (1994). Thirty individuals were examined in total, including individuals that were left intact and the ones that were dissected.

After collection and identification, 16 individuals used for identifications to serve as voucher specimens were deposited in 2 scientific collections in Brazil, Coleção do Instituto de Biologia da Universidade Federal do Rio de Janeiro (UFRJ) and Museu Nacional do Rio de Janeiro. These specimens are settled under the numbers IBUFRJ 3580 and IBUFRJ 3588. Fifty individuals were fixed in 98% ethanol for future molecular analysis.

Results

Thirty specimens were examined, with average length of 19 mm (min. 16 mm; max. 23 mm) (Fig. 4). The operculum presents paddle-shaped outer paleae slightly larger than 1 mm long, with plume-shaped appendage on top (Fig. 5); these paleae surround the whole opercular crown circumference (Fig. 6). The middle paleae are arranged radially, forming the opercular cone completely hiding the row of inner paleae (Fig. 6). The middle paleae are formed by two teeth that give them an uncinata shape; the superior tooth is approximately 3 times longer than the inferior one, with a pronounced downward curvature distally (Fig. 7). The inner paleae (Fig. 8) are inferior to the middle paleae (Fig. 9) and very similar to them, except by the relatively less robust shape and by a distal bifurcation formed by its 2 teeth, which have the same length.


Discussion

The material of *Phragmatopoma caudata* examined here does not differ from descriptions of this species from other localities (Kinberg 1867, Amaral 1987, Kirtley 1994).

This work reports for the first time the occurrence of *Phragmatopoma caudata* on the coast of the state of Piauí. Previous records show that the species has a wide distribution in Brazil, occurring from the state of Rio Grande do Sul to the state of Ceará, and that the nearest record to the west is in Venezuela (Fig. 1). The species also occurs in several locations in the Caribbean and in the Gulf of Mexico up to Florida (Kirtley 1994).

This work, therefore, extends the known distribution of this species across the Brazilian coast, becoming the westernmost record of *P. caudata* in the country (Fig. 1). It is also only the second location where the species is recorded in Brazil's northern coastline, after Kirtley (1994), Fausto-Filho and Furtado (1970) and Fournier et al. (2010) had recorded it for Fortaleza, Ceará (Fig. 1).

Fournier et al. (2010) examined the inter- and intra-specific variability on chemical composition of biocement among sand-reef-building polychaetes, including Brazil-



Figures 4–9. Adult of *Phragmatopoma caudata* from Pedra do Sal, Parnaíba. **4.** Entire animal. **5.** Lateral views of the outer palea. **6.** Upper view of the opercular crown. **7.** Middle palea. **8.** Inner palea. **9.** Relative position of paleae in association to form one of the sections of the opercular crown circumference.

ian populations previously recorded as *P. caudata* or *P. lapidosa* Kinberg 1867 from the following states: Santa Catarina, Paraná, São Paulo, Rio de Janeiro, Paraíba, Rio Grande do Norte and Ceará. These authors concluded that there is some degree of variation on the cement's biochemistry among populations. Another work (Faroni-Perez and Zara 2014) acknowledged differences on physiological and morphometric reproductive traits between Brazilian and Florida populations of *P. caudata*, particularly on oogenesis and oocyte development. Although these works revealed the existence of phenotypic variations among populations of *P. caudata* along its distribution, which could indicate intraspecific variation or possibly speciation, they did not deal with taxonomic relationships among *Phragmatopoma* spp. or with genetic connectivity between *P. caudata* populations.

Recent published data (Nunes et al. 2016) from DNA analysis pointed that populations found along the Brazilian coast, as well as those found in the rest of South America, up to Florida, and including the Caribbean Sea, are representatives of a single species (*P. caudata*), as previously suggested by Kirtley (1994). Therefore, *P. lapidosa* is a synonym of *P. caudata* as inferred by both morphological (Kirtley 1994) and molecular data (Drake et al. 2007, Nunes et al. 2016). It was also shown that

populations across the Brazilian coastline and the Caribbean have a strong genetic connection and do not show population structure. This result indicates that this species is able to overcome the major geographic barriers along South and Central American Atlantic coasts, although this does not discard the existence of areas where the species is absent. The occurrence of *P. caudata* in the state of Piauí is in accordance with the species dispersive ability.

This study therefore contributes to fill the distribution gap of this species observed in the northern coast of Brazil, although the blank on distribution persists between this new record and the nearest records to the west, which are from Tortuga Island and Margarita Island in Venezuela (Kirtley 1994) (Fig. 1). In addition, this is the first record of *P. caudata* in the APA Delta do Parnaíba, increasing the knowledge on the local biota and the ability of environmental agencies to manage this important marine reserve in the Brazilian northeastern coast.

Acknowledgements

The authors thank those members from the Grupo de Ecologia de Invertebrados Bentônicos (GEIB/UFPI) who helped during fieldwork. We thank Dr Jérôme Fournier for insights and helpful information on the taxonomic

status of *P. caudata* populations in Brazil. We also thank Dra A. Cecília Z. Amaral, Dr Paulo C. Lana and two anonymous reviewers for suggestions and comments on the manuscript. Finally we acknowledge Dr Paula Paiva and Gisele Occhioni for having received and held specimens in the IBUFRJ collection.

Authors' Contributions

JGFGF collected the material, JGGF, MVQBS and RAS identified the material, JGGF made the drawings, JGGF, MVQBS and RAS wrote the text.

References

- Amaral ACZ, Nonato EF (1994) Anelídeos poliquetos da costa brasileira. 5. Pisionidae, Chrysopetalidae, Amphinomidae e Euphrosinidae. *Revista Brasileira de Zoologia* 11 (2): 361–390. <https://doi.org/10.1590/S0101-81751994000200022>
- Amaral ACZ (1987) Uma breve caracterização de *Phragmatopoma lapidosa* Kinberg 1867 (Polychaeta, Sabellariidae). *Revista Brasileira de Zoologia* 3 (8): 471–474. <https://doi.org/10.1590/S0101-81751986000400001>
- Ataide MB, Venekey V, Rosa-Filho JS, Santos PJP (2014) Sandy reefs of *Sabellaria wilsoni* (Polychaeta: Sabellariidae) as ecosystem engineers for meiofauna in the Amazon coastal region, Brazil. *Marine Biodiversity* 44 (3): 403–430. <https://doi.org/10.1007/s12526-014-0248-x>
- Bosa CR, Masunari S (2002a) Crustáceos decápodos associados aos bancos de *Phragmatopoma caudata* (Kroyer) (Polychaeta, Sabellariidae) na Praia de Caiobá, Martinhos, Paraná. *Revista Brasileira de Zoologia* 19 (1): 117–133. <https://doi.org/10.1590/S0101-81752002000500006>
- Bosa CR, Masunari S (2002b) Peracáridos associados aos bancos de *Phragmatopoma caudata* (Kroyer) (Polychaeta, Sabellariidae) na Praia de Caiobá, Martinhos, Paraná. *Revista Brasileira de Zoologia* 19 (1): 135–147. <https://doi.org/10.1590/S0101-81752002000500007>
- Dias A, Paula J (2001) Associated fauna of Sabellaria alveolata colonies on the central coast of Portugal. *Journal of the Marine Biological Association of the United Kingdom* 81 (1): 169–170. doi: <https://doi.org/10.1017/S0025315401003538>
- Drake CA, McCarthy DA, von Dohlen CD (2007) Molecular relationships and species divergence among *Phragmatopoma* spp. (Polychaeta: Sabellariidae) in the Americas. *Marine Biology* 150: 345–358. doi: <https://doi.org/10.1007/s00227-006-0373-6>
- Dubois S, Retière C, Olivier F (2002) Biodiversity associated with *Sabellaria alveolata* (Polychaeta: Sabellariidae) reefs: Effects of human disturbances. *Journal of the Marine Biological Association of the United Kingdom* 82 (5): 817–826. <https://doi.org/10.1017/S0025315402006185>
- Dubois S, Commito JA, Retière C, Olivier F (2006) Effects of epibionts on *Sabellaria alveolata* (L.) biogenic reefs and their associated fauna in the Bay of Mount Saint-Michel. *Estuarine, Coastal and Shelf Science* 68: 635–646. <https://doi.org/10.1016/j.ecss.2006.03.010>
- Faroni-Perez L, Zara FJ (2014) Oogenesis in *Phragmatopoma* (Polychaeta: Sabellariidae): Evidence for morphological distinction among geographically remote populations. *Memoirs of Museum Victoria* 71: 53–65. <https://doi.org/10.24199/j.mmv.2014.71.06>
- Fausto-Filho J, Furtado E. (1970) Nota preliminar sobre a fauna das colônias de Sabellariidae do litoral do estado do Ceara (Annelida, Sedentaria). *Revista Brasileira de Biologia* 30 (2): 285–289.
- Fournier J, Etienne S, Le Cam JB (2010) Inter and intraspecific variability in the chemical composition of the mineral phase of cements from several tube-building polychaetes. *Geobios* 43 (2): 191–200. <https://doi.org/10.1016/j.geobios.2009.10.004>
- Fransozo A, Bertini G, Correa MOD (2000) Population biology and habitat utilization of the stone crab *Menippe nodifrons* in the Ubatuba region, São Paulo, Brazil. In: Vaupel Klein JC, Scharm FR (Eds) *The Biodiversity Crisis and Crustacea*, Crustacean Issues 12. Balkema, Rotterdam, 275–281.
- Goldberg WM (2013) *The biology of reefs and reef organisms*. University of Chicago Press, Chicago, 424 pp.
- Jones CG, Lawton JH, Shachak M (1994) Organisms as ecosystem engineers. *Oikos* 69(3): 373–386. doi: https://doi.org/10.1007/978-1-4612-4018-1_14
- Kinberg JGH (1867) *Annulata nova*. *Oversigt af Kongliga Vetenskaps-Akademiens Forhandlingar* 23: 337–357.
- Kirtley DW (1994) A Review and Taxonomic Revision of the Family Sabellariidae Johnston, 1865 (Annelida; Polychaeta). *Sabecon Press-Science Series*, Vero Beach, FL, 223 pp.
- Lana PC, Bremec CS (1994) Sabellariidae (Annelida, Polychaeta) from South America. *Mémoires du Museum National d'Histoire Naturelle* 162: 211–222.
- Masi BP, Zalmon IR (2008) Zonação de comunidade benthica do entremarés em molhes sob diferente hidrodinamismo na costa norte do estado do Rio de Janeiro, Brasil. *Revista Brasileira de Zoologia* 25 (4): 668–673. doi: <https://doi.org/10.1590/S0101-81752008000400011>
- Masi BP, Macedo IM, Zalmon IR (2009) Benthic community zonation in a breakwater on the North coast of the State of Rio de Janeiro, Brazil. *Brazilian Archives of Biology and Technology* 52 (3): 637–646. <https://doi.org/10.1590/S1516-89132009000300016>
- Micheletti-Flores CV, Negreiros-Fransozo ML (1999) Porcellanid crabs (Crustacea, Decapoda) inhabiting sand reefs built by *Phragmatopoma lapidosa* (Polychaeta, Sabellariidae) at Paranapuã Beach, São Vicente, São Paulo, Brazil. *Revista Brasileira de Biologia* 59 (1): 63–73. <https://doi.org/10.1590/S0034-71081999000100009>
- Moreno TR, Rocha RM (2006) Associated fauna of *Eudistoma carolinense* (Tunicata—Ascidacea) compared with other biological substrates with different architectures. *Journal of Coastal Research* SI39: 1695–1699.
- Morgado EH, Amaral ACZ (1989) Anelidos poliquetos da região de Ubatuba (SP)—padrões de distribuição geográfica. *Revista Brasileira de Zoologia* 6 (3): 535–568. <https://doi.org/10.1590/S0101-81751989000300013>
- Nunes F, Van Wormhoudt A, Perez-Faroni L, Fournier J (2016) Phylogeography of the reef-building polychaetes of the genus *Phragmatopoma* in the Western Atlantic region. *Journal of Biogeography*. <https://doi.org/10.1111/jbi.12938>
- Occhioni GE, Brasil ACS, Araújo AFB (2009) Morphometric study of *Phragmatopoma caudata* (Kroyer) Morch, 1863 (Polychaeta—Sabellida—Sabellariidae). *Zoologia* 26 (4): 739–743. <https://doi.org/10.1590/S1984-46702009000400019>
- Pamplin PAZ, Almeida TCM, Silva-Filho JP (2007) New record of *Laeonereis acuta* (Treadwell, 1923) (Nereididae: Polychaeta) in Northeast coast of Brazil. *Biota Neotropica* 7 (3): 1–3. <https://doi.org/10.1590/S1676-06032007000300035>
- Pohler SML (2004) The sabellariid worm colonies of Suva Lagoon, Fiji. *South Pacific Journal of Natural and Applied Sciences* 22 (1): 36–42.
- Rocha RM (1995) Abundance and distribution of sessile invertebrates under intertidal boulders (São Paulo, Brasil). *Boletim do Instituto Oceanográfico* 43 (1): 71–88. <https://doi.org/10.1590/S0373-55241995000100006>
- Santa-Isabel LM, Leão ZMAN, Peso-Aguiar MC (2000) Polychaetes from the Guarajuba coral reefs, Bahia, Brazil. *Bulletin of Marine Science* 67 (1): 645–653.
- Santos CSG, Lana PC (2003) Nereididae (Polychaeta) da costa nordeste do Brasil III. Gêneros *Ceratonereis* e *Nereis*. *Iheringia, Série Zoologia* 93 (1): 5–22.
- Santos AS, Riul P, Brasil ACS, Christoffersen ML (2011) Encrusting Sabellariidae (Annelida: Polychaeta) in rhodolith beds, with

- description of a new species of *Sabellaria* from the Brazilian coast. *Journal of the Marine Biological Association of the United Kingdom* 91 (2): 425–438. <https://doi.org/10.1017/S0025315410000780>
- Santos CSG, Steiner TM (2006) Família Nereididae. In: Amaral ACZ, Rizzo AE, Arruda EP (Eds) *Manual de Identificação dos Invertebrados Marinhos da Região Sudeste-Sul do Brasil*. Editora da USP, São Paulo, 121–133.
- Sepúlveda RD, Moreno RA, Carrasco FD (2003) Macroinvertebrate diversity associated to reefs of *Phragmatopoma moerchi* Kinberg, 1867 (Polychaeta: Sabellariidae) in the intertidal rocky shore at Cocholgüe, Chile. *Gayana* 67 (1): 45–54. <https://doi.org/10.4067/S0717-65382003000100007>