



# First record of Double-lined Mackerel, *Grammatorcynus bilineatus* (Rüppell, 1836) (Scombriformes, Scombridae) from Peninsular Malaysia

Ying Giat Seah<sup>1,2</sup>, Mohd Sharol Ali<sup>1</sup>, Siti Asmal Mat Piah<sup>1</sup>, Tun Nurul Aimi Mat Jaafar<sup>1</sup>

<sup>1</sup> Faculty of Fisheries and Food Science, Universiti Malaysia Terengganu, Kuala Nerus, Terengganu, Malaysia

<sup>2</sup> South China Sea Repository and Reference Centre, Institute Oceanography and Environment, Universiti Malaysia Terengganu, Kuala Nerus, Terengganu, Malaysia.

Corresponding author: Tun Nurul Aimi Mat Jaafar (tun\_aimi@umt.edu.my)

**Abstract.** Three individuals of *Grammatorcynus bilineatus* (Rüppell, 1836), Double-lined Mackerel, were recorded in local fish market in Terengganu, Malaysia. Although in Malaysia *G. bilineatus* previously has been recorded from East Malaysia, the species has not been recorded from Peninsular Malaysia. Therefore, the specimens described herein represents the first record of *G. bilineatus* from Peninsular Malaysia. Detailed morphometric and meristic data are presented along with brief diagnostic characters.

**Key words.** Distribution, marine fish, new record, range extension, South China Sea

**Seah YG, Ali MS, Mat Piah SA, Mat Jaafar TNA** (2024) First record of Double-lined Mackerel, *Grammatorcynus bilineatus* (Rüppell, 1836) (Scombriformes, Scombridae) from Peninsular Malaysia. Check List 20 (1): 53–57. <https://doi.org/10.15560/20.1.53>

## INTRODUCTION

Double-lined Mackerel, *Grammatorcynus bilineatus* (Rüppell, 1836), is one of two species of *Grammatorcynus* Gill, 1862 (Scombriformes, family Scombridae). It has a wide distribution spanning from the Red Sea to Andaman Sea in Indian Ocean, and from the Ryukyu Islands to the northern coast of Australia, including Marshall Islands and Fiji in Western Pacific (Lheknim 2019). Therefore, its geographical distribution spans the tropical and sub-tropical areas of the Indian and Pacific oceans from 30–20°S and from 030–180°E. This subtropical reef-associated fish species is usually found in shallow waters at depths of 15–50 m, where it either forms groups with other scombrids or is solitary (Froese and Pauly 2023).

*Grammatorcynus bilineatus* in the Malay language among fishermen is locally known as “Aya Dua Garis” (Lim et al. 2018) or “Kembung Tenggiri”. In Malaysia, this species has been recorded in only East Malaysia (or Malaysian Borneo) (GBIF 2023). A single occurrence was reported in Sabah based on human observation near Pulau Pom Pom, while the remaining three specimens were collected from a fish landing site in Semporna, facing the Celebes Sea and deposited in the collection of the Universiti Sains Malaysia. However, *G. bilineatus* has never been recorded from Peninsular Malaysia (or West Malaysia). Here, we provide a record of *G. bilineatus* collected at Terengganu water, which represents the first record of this species from Peninsular Malaysia, South China Sea.

## METHODS

Specimens of *Grammatorcynus bilineatus* were collected from a local fish market in Kuala Nerus, Terengganu, Malaysia. These specimens were caught by hook-and-line off Pulau Karah, Terengganu by local fishermen. We collected and fixed the specimens following the procedure of Motomura and Ishikawa (2013). Measurements and terminology follow Collette and Gillis (1992). The specimens were measured to the nearest 0.01 mm using Vernier calipers. Morphometric measurements, including total length (**TL**), fork length (**FL**) and standard length (**SL**), are in millimeters, while other measurements are expressed as a percentage of either fork length or head length. The species was identified following Collette and Gillis (1992). The specimens were deposited at the South China Sea Repository and Reference Center, Universiti Malaysia Terengganu, Malaysia (**UMTF**).



Academic editor: Zeehan Jaafar

Received: 4 September 2023

Accepted: 6 January 2024

Published: 17 January 2024

Copyright © The authors. This is an open-access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0)

## RESULTS

Order Scombriformes  
 Family Scombridae Rafinesque, 1815  
 Genus *Grammatorcynus* Gill, 1862

### *Grammatorcynus bilineatus* (Rüppell, 1836)

Figure 1, Table 1

*Thynnus bilineatus* Rüppell 1836: 39, pl. 12 fig. 2 (type locality: Massawa, Eritrea, Red Sea)

**New record.** MALAYSIA – **Terengganu** · Kuala Nerus; 05°26'24.57"N, 103°03'26.65"E; 20.VII.2022; Mat Jaafar TNA leg.; hook and line; 3 spec., ♂, 213.0–264.0 mm SL; UMTF 10391 (Figure 1b), UMTF 10392 (Figure 1a), UMTF 10393.

**Identification.** *Grammatorcynus bilineatus* can be distinguished from *G. bicarinatus* Collette in Carpenter & Niem, 2000, (Shark Mackerel) by combination of the following characters: more gill rakers (18–24; 12–15 in *G. bicarinatus*), larger eye size (4.0–6.0% FL, vs. 3.1–4.6% in *G. bicarinatus*), and smaller maximum size (600 mm FL vs. 1100 mm in *G. bicarinatus*) (Collette and Gillis 1992). Other useful morphological features for identification are the elongate and slightly compressed body, which is covered by moderately small, cycloid scales. The mouth is relatively small, and the jaws have slender, conical teeth on vomer, palatines, and upper surface of broad tongue. The upper jaw reaches to about the middle of the eye. Two dorsal fins are separated by a short space and are followed by finlets. The pectoral fins are rather short. The caudal fin is deeply forked and with three caudal keels; one of these keels is not very prominent and covered with a row of pored scales, and the other two keels at the base of the caudal fin are much smaller. Two lateral lines are present; the upper originates from the dorsal part of the operculum and runs along the upper half of the body to almost anterior to the median caudal keel, and the lower lateral line branches off from a point below the third dorsal spine then descends to lower half of the body; this lower lateral line re-joins the upper lateral line at the base of the caudal fin. The body color is metallic bluishgreen on the back and upper sides, silvery on the lower sides, and silvery white without black spots on the belly. Detailed morphometric measurements and meristic counts are provided in Table 1.

## DISCUSSION

Our specimen agrees with the original description of *Grammatorcynus bilineatus*. Before 1983, *G. bilineatus* was sometimes confused with *G. bicarinatus*. Collette (1983) and Collette and Nauen (1983) demonstrated that *G. bilineatus* can be morphologically distinguished from the only other *Grammatorcynus* species, *G. bicarinatus*. The latter species is restricted to the waters of northern Australia and southern New Guinea. Collette and Gillis (1992) later gave osteological evidence that confirm the separation of these *Grammatorcynus* species.

The presence of *G. bilineatus* in Peninsular Malaysia was anticipated, given its widespread distribution near coral reefs in the Indo-West Pacific (Collette and Gillis 1992). Furthermore, the coastal waters off Pulau Karah, Terengganu, are known for their abundant coral reefs and associated sandy bottoms. *Grammatorcynus bilineatus* is rare in Malaysian waters. Previously, it has been documented only in the Celebes Sea of East Malaysia. Our new record extends the known geographic range of *G. bilineatus* to Peninsular Malaysia

**Figure 1.** *Grammatorcynus bilineatus*, Kuala Nerus, Terengganu, Malaysia. **a.** Fresh specimen, UMTF 10392, 264 mm SL. **b.** Preserved specimen, UMTF 10391, 252 mm SL.




**Table 1.** Morphometric measurements and meristic counts of *G. bilineatus* collected from Kuala Nerus, Terengganu.

Character	Present study			Mean	SD	Collette and Gillis (1992)
	UMTF 10391	UMTF 10392	UMTF 10393			
<b>Total length (mm)</b>	319.0	338.0	260.0	305.7	40.7	—
Fork length (mm)	264.0	278.0	225.0	255.7	27.5	226–575
Standard length (mm)	252.0	264.0	213.0	243.0	26.7	—
<b>As % of fork length</b>						
Snout–anal fin	60.2	60.8	61.4	60.8	0.6	58.1–64.1
Snout–2 <sup>nd</sup> dorsal fin	56.2	55.6	54.4	55.4	0.9	52.8–61.9
Snout–1 <sup>st</sup> dorsal fin	31.0	31.1	30.5	30.9	0.3	27.6–32.2
Snout–pelvic fin	25.7	25.6	26.3	25.8	0.4	23.6–30.6
Snout–pectoral fin	22.7	22.7	23.7	23.0	0.6	19.9–24.5
Pectoral fin–pelvic fin	11.4	10.8	12.5	11.6	0.8	9.0–11.1
Head length	22.0	22.1	22.7	22.2	0.4	19.7–23.6
Max body depth	21.8	21.5	22.0	21.7	0.2	16.4–23.4
Max body width	12.7	12.7	12.6	12.7	0.1	9.1–13.6
Pectoral fin length	12.5	12.8	7.6	11.0	2.9	10.6–14.2
Pelvic fin length	7.7	7.3	7.2	7.4	0.3	7.0–8.7
Pelvic fin insertion–vent	30.7	32.8	33.7	32.4	1.5	26.2–35.4
Pelvic fin tip–vent	24.7	25.9	26.8	25.8	1.1	22.8–27.3
1 <sup>st</sup> dorsal fin base	22.3	20.4	22.2	21.6	1.1	21.1–26.1
2 <sup>nd</sup> dorsal fin height	9.4	10.1	8.9	9.5	0.6	8.8–11.6
2 <sup>nd</sup> dorsal fin base	10.8	10.3	10.7	10.6	0.3	7.9–11.8
Anal fin height	8.3	9.5	7.2	8.3	1.1	8.2–11.4
Anal fin base	8.8	9.2	8.9	8.9	0.2	7.3–10.1
Snout (fleshy)	7.5	7.1	7.2	7.2	0.2	5.8–9.0
Snout (bony)	8.0	7.7	7.7	7.8	0.2	6.0–8.0
Maxilla length	9.8	10.1	10.4	10.1	0.3	8.9–10.7
Postorbital	8.5	9.5	9.5	9.2	0.6	7.8–9.8
Orbit (fleshy)	5.0	4.5	5.4	5.0	0.5	4.0–6.0
Orbit (bony)	4.9	5.1	6.1	5.3	0.6	5.3–8.8
Interorbital	7.1	7.3	7.6	7.3	0.2	5.6–7.4
2 <sup>nd</sup> dorsal fin –caudal	42.5	40.7	41.3	41.5	0.9	42.7–49.6
<b>As % of head length</b>						
Snout (fleshy)	34.1	32.0	31.5	32.5	1.4	24.8–39.7
Snout (bony)	36.4	34.9	34.1	35.1	1.2	28.1–35.7
Maxilla length	44.8	45.8	45.6	45.4	0.5	42.0–48.0
Postorbital	38.6	43.1	41.9	41.2	2.3	35.0–45.0
Orbit (fleshy)	22.9	20.4	23.9	22.4	1.8	19.1–25.7
Orbit (bony)	22.4	23.2	26.4	24.0	2.1	25.2–38.1
Interorbit	32.4	33.3	33.3	33.0	0.5	25.3–32.7
<b>Counts</b>				<b>Modes</b>		
Dorsal fin spines	XII	XII	XII	XII	0.0	XI–XIII (XII)
Dorsal fin rays	10	10	11	10	0.6	10–12 (11)
Dorsal finlets	6	6	6	6	0.0	6–7 (6)
Anal fin soft rays	11	11	10	11	0.6	11–13 (12)
Anal finlets	6	7	6	6	0.6	5–7 (6)
Pectoral fin rays	24	24	24	24	0.0	22–26 (24–25)
Pelvic fin rays	5	5	5	5	0.0	—
<b>Gill rakers</b>						
Upper	5	6	6	6	0.6	—
Lower	15	15	16	15	0.6	—
Total	20	21	22	21	1.0	18–24 (21)

and the southern South China Sea (Figure 2). The emergence and expansion of *G. bilineatus* to Peninsular Malaysia may have been caused by natural displacement as well as by changes in oceanographic conditions.

Current knowledge of fish species inhabiting the continental shelf of Malaysia remains limited (Chong et al. 2010). In the past five years, scientific papers and books have revealed interesting discoveries, including 31 species newly reported from the east coast of Peninsular Malaysia (Mat Jaafar et al. 2019; Okamoto et al. 2019; Motomura et al. 2021; Seah et al. 2020, 2021; Yusuf et al. 2021; Mohd Nur et al. 2022; Mat Piah et al. 2023). Further exploration of Malaysia's marine fishes will contribute valuable data to enhance the understanding of Malaysia's biodiversity.



**Figure 2.** Map of records of *Grammatorcynus bilineatus* in the Indo-West Pacific Ocean: square = previous records; star = new record from the east coast of Peninsular Malaysia.

## ACKNOWLEDGEMENTS

We sincerely acknowledge Mrs. Suzana Sukri for providing us with information on the locality of the specimens. We also express our heartfelt thanks to the Universiti Malaysia Terengganu for funding, Mr. Yuzwan Mohamad for his efforts in preparing the maps, as well as the entire staff of the Faculty of Fisheries and Food Science and the South China Sea Repository and Reference Centre for their valuable research assistance. We also wish to express our special thanks to the reviewers and academic editor.

## ADDITIONAL INFORMATION

### Conflict of interest

The authors declare that no competing interests exist.

### Ethical statement

No ethical statement is reported.

### Funding

This research received support from Universiti Malaysia through the Research Intensified Grant Scheme with vote number 55192/2.

### Author contributions

Conceptualization: YGS, TNAMJ. Formal analysis: YGS. Funding acquisition: YGS. Investigation: YGS, TNAMJ. Methodology: MSA, SAMP. Validation: YGS, MSA. Writing – original draft: TNAMJ. Visualization: MSA. Writing – review and editing: YGS, MSA, SAMP, TNAMJ.

### Author ORCID iDs

Ying Giat Seah  <https://orcid.org/0000-0002-2976-4448>

Siti Asmal Mat Piah  <https://orcid.org/0009-0000-9301-5292>

Tun Nurul Aimi Mat Jaafar  <https://orcid.org/0000-0002-3032-0545>

## REFERENCES

**Collette BB** (1983) Recognition of two species of double-lined mackerels (*Grammatorcynus*, Scombridae). Proceedings of the Biological Society Washington 96: 715–718.

- Collette BB, Nauen CE** (1983) FAO species catalogue. Vol. 2. Scombrids of the world. Annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date. FAO Fisheries Synopsis Number 125, FAO, Rome, Italy, 137 pp. <https://www.fao.org/3/ac478e/ac478e00.htm>. Accessed on: 2023-1-8.
- Collette BB, Gillis GB** (1992) Morphology, systematics, and biology of the double-lined mackerels (*Grammatorcynus*, Scombridae). Fishery Bulletin 90: 13–53.
- Chong VC, Lee PKY, Lau CM** (2010) Diversity, extinction risk and conservation of Malaysian fishes. Journal of Fish Biology 76: 2009–2066. <https://doi.org/10.1111/j.1095-8649.2010.02685.x>
- Froese R, Pauly D** (2023) FishBase, version (02/2023). <https://www.fishbase.org>. Accessed on: 2023-6-13.
- GBIF** (2023) Global Biodiversity Information Facility occurrences, version 13 June 2023. <https://doi.org/10.15468/dl.zn953r>. Accessed on: 2023-6-13
- Lheknim V** (2019) First record of double-lined mackerel *Grammatorcynus bilineatus* (Rüppell, 1836) from the Gulf of Thailand. Phuket Marine Biological Center Research Bulletin 76: 19–23.
- Lim APK, Ahmad A, Nor Azman Z, Mohd Saki N** (2018) Field guide to fishes and crustaceans of the Southeast Asian region. SEAFDEC/MFRDMD, Kuala Terengganu, Malaysia, 246pp.
- Mat Jaafar TNA, Lee JN, Mazlan AG, Sheikh Abdul Kadir ST, Seah YG** (2019) *Opistognathus nigromarginatus* Rüppell, 1830 (Perciformes, Opistognathidae), Bridled Jawfish: a first record from Malaysia. Check List 15 (5): 883–886. <https://doi.org/10.15560/15.5.883>
- Mat Piah SA, Lee HH, Abd Latiff NA, Wong LL, Okamoto J, Motomura H, Kimura S, Seah YG** (2023) First record and distribution extension of two emperor fishes, *Lethrinus semicinctus* and *Lethrinus olivaceus* to the east coast of Peninsular Malaysia. Sains Malaysiana 52 (4): 1037–1046. <http://doi.org/10.17576/jsm-2023-5204-01>
- Mohd Nur NF, Abdullah S, Seah YG, Mohd Nor SA, Tan MP, Habib A, Mat Piah R, Mat Jaafar TNA** (2022) DNA barcoding of commercially important trevallies, *Carangoides* spp. (Carangiformes: Carangidae): a baseline report of species from Malaysia. Thalassas 38: 227–253. <https://doi.org/10.1007/s41208-022-00405-9>
- Motomura H, Kimura S, Seah YG, Sheikh Abdul Kadir ST, Ghaffar MA** (2021) Reef and shore fishes of Bidong Island, off east coast of Malay Peninsula. The Kagoshima University Museum, Kagoshima, Japan, 80 pp.
- Motomura H, Ishikawa S** (2013) Fish collection building and procedures manual. English edition. The Kagoshima University Museum, Kagoshima and the Research Institute for Humanity and Nature, Kyoto, Japan, 70 pp.
- Okamoto J, Sheikh Abdul Kadir ST, Motomura H, Mat Jaafar TNA, Seah YG** (2019) First records of the sole, *Aseragodes kobensis* (Steindachner, 1896) (Pleuronectiformes, Soleidae), from Malaysia. Check List 15 (6): 991–995. <https://doi.org/10.15560/15.6.991>
- Seah YG, Mat Jaafar TNA, Ali MS** (2020) Field guide to trawl fishes near Bidong Island. Penerbit Universiti Malaysia Terengganu, Kuala Nerus, Malaysia, 144 pp.
- Yusuf Y, Seah YG, Izarenah MR, Lee JN** (2021) First record of Spotted Knifejaw, *Oplegnathus punctatus* (Temminck & Schlegel, 1844) (Oplegnathidae) in the southern South China Sea. Check List 17 (4): 1195–1198. <https://doi.org/10.15560/17.4.1195>