



First record of *Microspingus cinereus* (Bonaparte, 1850), Cinereous Warbling Finch (Passeriformes, Thraupidae), from the state of Rio de Janeiro, Brazil

GABRIEL MAGALHÃES TAVARES^{1*}, JOÃO RAFAEL GOMES DE ALMEIDA E MARINS²,
MANOEL TULER FILHO³, CARLOS NEI ORTÚZAR-FERREIRA⁴

1 Programa de Pós-graduação em Zoologia, Universidade Federal de Mato Grosso, Cuiabá, MT, Brasil • GMT: gabriel_magalhaestavares@hotmail.com <https://orcid.org/0000-0001-9948-7012>

2 Secretaria Municipal de Meio Ambiente e Desenvolvimento Sustentável, Prefeitura de Barra Mansa, RJ, Brazil • JRGAM: joaorgam.smmadsbm@gmail.com <https://orcid.org/0000-0001-6784-1005>

3 Independent researcher, Barra Mansa, RJ, Brasil • MTF: manoel.tuler@hotmail.com

4 Programa de Pós-graduação em Biologia Animal, Universidade Federal Rural do Rio de Janeiro, Seropédica, RJ, Brazil • CNOF: carlosneiortuzarferreira@gmail.com <https://orcid.org/0000-0002-0177-5591>

* Corresponding author

Abstract. We document from the state of Rio de Janeiro, Brazil, the first record of *Microspingus cinereus* (Bonaparte, 1850), which was observed in the municipality of Barra Mansa in October 2020. The species was originally documented in central Brazil, potentially linked to Cerrado habitats. However, recent observations from eBird and WikiAves databases indicate that *M. cinereus* has been spreading towards the southeast for several years. It has been considered Vulnerable for almost two decades due to the loss of grassland habitats. Our discovery expands the known geographic range of *M. cinereus* to eastern Brazil and may have importance for future biogeographic studies and the conservation of this species.

Keywords. Conservation, geographic distribution, ornithology

Academic editor: Rodolpho Credo Rodrigues

Received 29 June 2023, accepted 24 October 2023, published 6 November 2023

Tavares GM, Almeida e Marins JRG, Filho MT, Ortúzar-Ferreira CN (2023) First record of *Microspingus cinereus* (Bonaparte, 1850), Cinereous Warbling Finch (Passeriformes, Thraupidae), from the state of Rio de Janeiro, Brazil. Check List 19 (6): 827–832. <https://doi.org/10.15560/19.6.827>

Introduction

Located in the eastern Brazil, the first region to be colonized and exploited, the Atlantic Forest has accumulated a broad and relatively early knowledge of its avifauna as compared to other Brazilian biomes; the state of Rio de Janeiro one of the most studied states of Brazil with this biome (Pacheco et al. 2008). Currently, 806 bird species are known to occur in the state of Rio de Janeiro; this number represents 41% of the total bird species occurring in Brazil and approximately 79% of the total bird species in the Atlantic Forest (Bergallo et al. 2000; Moreira-Lima 2013; Pacheco et al. 2021; Gagliardi and Serpa 2022). Despite extensive knowledge of the avifauna of Rio de Janeiro state, new bird

records have recently been made from the state (Vecchi and Alves 2008; Guido et al. 2016; Alves et al. 2017; Del-fino and Carlos 2022).

Microspingus cinereus (Bonaparte, 1850), Cinereous Warbling Finch, is a Neotropical passeriform that occur sparsely in non forest habitats of central Brazil (Costa and Rodrigues 2013; Marques-Santos et al. 2014). Classified as endemic to the Cerrado biome (Silva and Bates 2002), *M. cinereus* was considered to be globally Vulnerable between 2000 and 2017 (IUCN 2022). During this time, the population was suspected to have declined due to a reduction of suitable habitat (Collar et al. 1992). However, more recent studies suggest that this species can tolerate and even benefit from degraded areas, such as abandoned pastures (Simon et al. 1999;

Ribon 2002; Vasconcelos 2007), degraded Cerrado vegetation, fields, and orchards (Lopes et al. 2010), and its global conservation status was reassessed in 2018 to Least Concern (IUCN 2022). Here, we document the first record of *M. cinereus* from the state of Rio de Janeiro, Brazil.

Methods

Our observation was made during a birdwatching near the locality known as Robertão waterfall (22°21'37.66"S, 044°06'56.77"W) in the municipality of Barra Mansa, Rio de Janeiro state, Brazil. The region of the record is dominated by pasture, with fragments of native forests of differing sizes, shapes, and conservation levels; these fragmented forests are sparsely distributed over the landscape (Fig. 1). We used a Canon PowerShot SX540 HS camera for photographing the birds, and the resulting photograph (Fig. 2) was deposited in the WikiAves citizen-science database (Filho 2020). We produced the map (Fig. 3) in QGIS v. 3.18.1 (QGIS Development Team 2021) and used the BirdLife International (2022) map for *Microspingus cinereus* geographical distribution to compose this species' previously known distribution (Jaramillo and Juana 2021).

Results

Microspingus cinereus (Bonaparte, 1850)

Figure 2

New record. BRAZIL – Rio de Janeiro • Barra Mansa; 22°21'37.66"S, 044°06'56.77"W; 491 m elev.; 18.X.2020; Manoel Tuler Filho obs.; ♂ and ♀; WA4026228.

Identification. This species is a small, grayish finch and is characterized by having pale, plumbeous upperparts juxtaposed with a slightly darker loreal area. Its wings and tail are tinged with a dusky hue, subtly accented by a gray border, while the tail features a distinctive white tip (at the extremities of its rectrices). The throat and underparts are conspicuously white, offering a stark contrast to its blackish bill and reddish iris. Immature individuals of this species exhibit a prevalent brown wash on their heads.

A similar species is *Neothraupis fasciata* (Lichtenstein, 1823), White-banded Tanager, can be distinguished by the presence of a discernible black mask, while its juveniles exhibit a notable prevalence of brown pigmentation. This finch produces a series of high-pitched and vigorously spirited warbling sounds (Jaramillo and Juana 2021).

Discussion

Knowledge of the biology and natural history of *Microspingus cinereus* is sparse (Costa and Rodrigues 2013). Several authors have claimed that *M. cinereus* is naturally rare, has low density (Ridgely and Tudor 1989; Stotz et al. 1996; Lopes et al. 2009), or occurs irregularly (Melo-Júnior 1998). This pattern of occurrence is possibly due to a semi-nomadic behavior in response to fire dynamics, which occurs naturally in the Cerrado biome (Aguiar and Camargo 2004).

Microspingus cinereus is a recent colonizer in eastern Brazil and has been observed in non-native pastures in areas formerly covered by Atlantic Forest (Simon et



Figure 1. Site where *Microspingus cinereus* was observed in Barra Mansa, Rio de Janeiro state, Brazil. Photograph: Manoel Tuler Filho.



Figure 2. *Microspingus cinereus* photographed in field on 18 October 2020 in the municipality of Barra Mansa, Rio de Janeiro state, Brazil. Photograph: Manoel Tuler Filho.

al. 1999; Vasconcelos et al. 1999; Vasconcelos and D'angelo-Neto 2007). These pastures are generally renewed by frequent burning, a common practice in Brazil (our observation), which creates a similar environment for *M. cinereus* as in the Cerrado, and thus may have contributed to the expansion of this species to eastern Brazil. However, Lopes et al. (2010) reported a specimen of *M. cinereus* collected in 1936 in the municipality of Viçosa (Minas Gerais) and deposited in the National Museum of Rio de Janeiro (uncatalogued), which suggests that the species has been present in eastern Brazil for much longer.

Records of this species in the literature are sparse, limited to a few individuals per site, and temporally isolated (Ridgely and Tudor 1989; Stotz et al. 1996; Simon et al. 1999; Vasconcelos et al. 1999; Ribon 2002; Vasconcelos 2007; Vasconcelos and D'angelo-Neto 2007; Lopes et al. 2010; Rodrigues et al. 2011; Costa and Rodrigues 2012; Wischhoff et al. 2012; Costa and Rodrigues 2013). *Microspingus cinereus* may be extirpated in Mato Grosso (no records since 1904) and in Mato Grosso do Sul (one record in 1937); it had been thought to be extirpated in São Paulo, since there were no records since 1901, but in August 2012 *M. cinereus* was recorded in the municipality of Divinolândia (Coeti 2012). This sighting shows the importance of citizen science in gaining knowledge of the geographic distribution of species, especially of rare species.

Citizen science once again has been pivotal in advancing our understanding of the geographic distribution of *M. cinereus*, with the first sighting, reported here, of this species from the state of Rio de Janeiro. However, further research is necessary to determine whether the observed pair of individuals are permanent, breeding residents or transient and nomadic. Such an investigation holds significance due to this species' seminomadic behavior, particularly in response to fire dynamics, as expounded upon by Mello-Júnior (1998).

While fire is a natural phenomenon within the Cerrado biome (Aguiar and Camargo 2004), in the Atlantic Forest, it is primarily associated with the renewal of pastures for livestock. There are many records of this species in rocky fields in the Serra do Espinhaço and surrounding mountain ranges, such as Serra do Cipó, Serra do Caraça, and Serra do Curral, among others (Vasconcelos et al. 2008). However, recent data provided in the WikiAves and eBird platforms show that there are many records along the Serra da Mantiqueira, including in Itatiaia National Park, between the states of Minas Gerais and Rio de Janeiro. Hence, it warrants consideration whether anthropogenic practices have facilitated this species' expansion into the Atlantic Forest biome and concurrently influenced its seminomadic tendencies within this specific ecological context. These inquiries require a thorough investigation to uncover their complexities and implications, particularly




Figure 3. Geographic distribution of *Microspingus cinereus* in Brazil according to BirdLife (2022) with new record shown.

concerning the influence of human activities on the expansion or contraction of species distributions.

Acknowledgements

We thank the anonymous reviewers and academic editor for their helpful comments on the manuscript.

Author Contributions

Conceptualization: GMT, JRGAM. Data curation: GMT, JRGAM, CNOF. Investigation: MTF. Visualization: MTF, GMT. Formatting: CNOF. Writing – original draft: GMT. Writing – review and editing: GMT, JRGAM, CNOF

References

- Aguiar LMS, Camargo, AJA (2004) Cerrado: ecologia e caracterização. Embrapa, Brasília, Brazil, 249 pp.
- Alves MAS, Vecchi MB, Vallejos LM, Ribeiro EA, Martins-Silva J, Clair RSS (2017) New records of bird species from Ilha Grande, state of Rio de Janeiro, southeastern Brazil. Check List 12 (6): 2017. <https://doi.org/10.15560/12.6.2017>
- Bergallo HG, Rocha CFD, Van Sluys M, Alves MAS (2000) O status atual da fauna do estado do Rio de Janeiro: considerações finais. In: Bergallo HG, Rocha CFD, Alves MAS, Van Sluys M (Orgs). A fauna ameaçada de extinção do estado do Rio de Janeiro. Vol. 1. 1 ed. Rio de Janeiro, EDUERJ, 145–150 pp.

- BirdLife International** (2022) Species factsheet: *Microspingus cinereus*. <http://datazone.birdlife.org/species/factsheet/cinereous-warbling-finch-microspingus-cinereus>. Accessed on: 2022-12-15.
- Coeti JB** (2012) [WA719392, *Microspingus cinereus* Bonaparte, 1850] WikiAves—a enciclopédia das aves do Brasil. <https://www.wikiaves.com.br/719392>. Accessed on: 2023-02-01.
- Collar NJ, Gonzaga LP, Krabbe N, Madroño Nieto A, Naranjo LG, Parker TA, Wege DC** (1992) Threatened birds of the Americas: the ICBP/IUCN Red Data book. International Council for Bird Preservation, Cambridge, UK, 1150 pp.
- Costa LM, Rodrigues M** (2012) Bird community structure and dynamics in the campos rupestres of southern Espinhaço Range, Brazil: diversity, phenology and conservation. *Revista Brasileira de Ornitologia* 20 (2): 132–147.
- Costa LM, Rodrigues M** (2013) Notes on residency, home range, and natural history of the ‘Vulnerable’ Cinereous Warbling-Finch, *Poospiza cinerea*. *The Wilson Journal of Ornithology* 125 (2): 433–438. <https://doi.org/10.1676/12-115.1>
- Delfino HC, Carlos, CJ** (2022) On the wrong side of the Atlantic: first record of wild Greater Flamingo *Phoenicopterus roseus* in Brazil and in the Americas? *Bulletin of the British Ornithologists’ Club* 142 (2): 231–238. <https://doi.org/10.25226/bboc.v142i2.2022.a7>
- Filho MT** (2020) [WA4026228, *Microspingus cinereus* Bonaparte, 1850] WikiAves—a enciclopédia das aves do Brasil. <https://www.wikiaves.com.br/4026228>. Accessed on: 2023-02-01.
- Gagliardi RL, Serpa GA** (2022) Avifauna completa do estado do Rio de Janeiro. *Táxeus—listas de espécies*. <https://www.taxeus.com.br/lista/82>. Accessed on: 2023-01-26
- Guido RM, Carnevali RG, Vecchi MB, Alves MAS** (2016) First records of the Gull-billed Tern, *Gelochelidon nilotica* (Gmelin, 1789) (Aves: Sternidae), from Rio de Janeiro state, Brazil. *Check List* 12 (2): 1878. <https://doi.org/10.15560/12.2.1878>
- IUCN** (International Union for the Conservation of Nature) (2022) The IUCN Red List of threatened species. Version 2021-3. <https://www.iucnredlist.org>. Accessed on: 2022-12-15.
- Jaramillo A, de Juana E** (2021) Cinereous Warbling Finch (*Microspingus cinereus*), version 1.1. In: del Hoyo J, Elliott A, Sargatal J, Christie DA, de Juana E (Eds.) *Birds of the world*. Cornell Lab of Ornithology, Ithaca, USA. <https://doi.org/10.2173/bow.ciwfin1.01.1>
- Lopes LE, Malacco GB, Alteff EF, Vasconcelos MF, Hoffmann D, Silveira LF** (2010) Range extensions and conservation of some threatened or little known Brazilian grassland birds. *Bird Conservation International* 20 (1): 84–94. <https://doi.org/10.1017/S0959270909990190>
- Lopes LE, Pinho JB, Bernardon B, Oliveira FF, Bernardon G, Ferreira LP, Vasconcelos MF, Maldonado-Coelho M, Nóbrega PFA, Rubio TC** (2009) Aves da Chapada dos Guimarães, Mato Grosso, Brasil: uma síntese histórica do conhecimento. *Papéis Avulsos de Zoologia* 49: 9–47.
- Marques-Santos F, Wischhoff U, Rodrigues M** (2014) New insights on the rarity of the vulnerable Cinereous Warbling-finch (Aves, Emberizidae) based on density, home range, and habitat selection. *Brazilian Journal of Biology* 74 (4): 795–802. <https://doi.org/10.1590/1519-6984.02813>
- Melo-Júnior TA** (1998) *Poospiza cinerea* Bonaparte, 1850. In: Machado, ABM (Ed.) *Livro Vermelho das espécies ameaçadas de extinção da fauna de Minas Gerais*. Belo Horizonte, Fundação Biodiversitas, 387–389.
- Moreira-Lima L** (2013) Aves da Mata Atlântica: riqueza, composição, status, endemismo e conservação. Master’s thesis, Universidade de São Paulo (USP), São Paulo, Brazil, 526 pp.
- Pacheco JE, Parrini R, Lopes LE, Vasconcelos MF** (2008) A avifauna do Parque Estadual do Ibitipoca e áreas adjacentes, Minas Gerais, Brasil, com uma revisão crítica dos registros prévios e comentários sobre biogeografia e conservação. *Cotinga* 30: 16–32.
- Pacheco JE, Silveira LF, Aleixo A, Agne CE, Bencke GA, Bravo GA, Brito GRR, Cohn-Haft M, Maurício GN, Naka LN, Olmos F, Posso SR, Lees AC, Figueiredo LFA, Carrano E, Guedes RC, Cesari E, Franz I, Schunck F, Piacentini VQ** (2021) Annotated checklist of the birds of Brazil by the Brazilian Ornithological Records Committee, second edition. *Ornithology Research* 29: 94–105. <https://doi.org/10.1007/s43388-021-00058-x>
- QGIS Development Team** (2021) QGIS Geographic Information System. Open Source Geospatial Foundation Project. <http://qgis.osgeo.org>.
- Ribon R** (2002) Colonization of eastern Brazil by the Cinereous Warbling-Finch, with some comments on its natural history. In: Abstracts for the Third North American Ornithological Conference. American Ornithological Union, New Orleans, USA, 585–585.
- Ridgely RS, Tudor G** (1989) *Birds of South America: the oscine passerines*. University of Texas Press, Austin, USA, 516 pp.
- Rodrigues M, Freitas GHS, Costa LM, Dias DF, Varela MLM, Rodrigues LC** (2011) Avifauna, Alto do Palácio, Serra do Cipó National Park, state of Minas Gerais, southeastern Brazil. *Check List* 7 (2): 151–161. <https://doi.org/10.15560/7.2.151>
- Silva JMC, Bates JM** (2002) Biogeographic patterns and conservation in the South American Cerrado: a tropical savanna hotspot. *BioScience* 52 (3): 225–233. [https://doi.org/10.1641/0006-3568\(2002\)052\[0225:bpacit\]2.0.co;2](https://doi.org/10.1641/0006-3568(2002)052[0225:bpacit]2.0.co;2)
- Simon JE, Ribon R, Mattos GD, Abreu CRM** (1999) A avifauna do Parque Estadual da Serra do Brigadeiro, Minas Gerais. *Revista Árvore* 23 (1): 33–48.
- Stotz DF, Fitzpatrick JW, Parker III T, Moskovits DK** (1996) *Neotropical birds: ecology and conservation*. University of Chicago Press, Chicago, USA, 478 pp.
- Vasconcelos MF** (2007) Aves observadas no Parque Paredão da Serra do Curral, Belo Horizonte, Minas Gerais, Brasil. *Atualidades Ornitológicas* 136: 6–11.
- Vasconcelos MF, D’angelo-Neto S** (2007) Padrões de distribuição e conservação da avifauna na região central da Cadeia do Espinhaço e áreas adjacentes, Minas Gerais, Brasil. *Cotinga* 28: 27–44.
- Vasconcelos MF, Lopes L, Machado C, Rodrigues M** (2008) As aves dos campos rupestres da Cadeia do Espinhaço:

- diversidade, endemismo e conservação. *Megadiversidade* 4: 221–241.
- Vasconcelos MF, Maldonado-Coelho M, Durães R** (1999) Notas sobre algumas espécies de aves ameaçadas e pouco conhecidas da porção meridional da Cadeia do Espinhaço, Minas Gerais. *Melopsittacus* 2: 44–50.
- Vecchi MB, Alves MAS** (2008) New records of the restinga antwren *Formicivora littoralis* Gonzaga and Pacheco (Aves, Thamnophilidae) in the state of Rio de Janeiro, Brazil: inland extended range and threats. *Brazilian Journal of Biology* 68: 391–395. <https://doi.org/10.1590/S1519-69842008000200022>
- Wischoff U, Marques-Santos F, Rodrigues M** (2012) Nesting of the Cinereous Warbling Finch (*Poospiza cinerea*) in southeastern Brazil. *Wilson Journal of Ornithology* 124: 166–169.