



New record of *Sphaenorhynchus prasinus* Bokermann, 1973 (Anura, Hylidae) from northeastern Brazil

Washington Luiz Silva Vieira¹, Kleber Silva Vieira¹, Fagner Ribeiro Delfim², Frederico Gustavo Rodrigues França³, Rômulo Romeu Nóbrega Alves⁴, Paulo Fernando Guedes Pereira Montenegro¹, Gentil Alves Pereira Filho⁴

¹ Laboratório de Ecofisiologia Animal, Departamento de Sistemática e Ecologia, Centro de Ciências Exatas e da Natureza, Universidade Federal da Paraíba, João Pessoa, Paraíba, Brazil

² Coleção Herpetológica, Departamento de Sistemática e Ecologia, Centro de Ciências Exatas e da Natureza, Universidade Federal da Paraíba, João Pessoa, Paraíba, Brazil

³ Departamento de Engenharia e Meio Ambiente, Centro de Ciências Aplicadas e Educação, Universidade Federal da Paraíba, Rio Tinto, Paraíba, Brazil

⁴ Departamento de Biologia, Universidade Estadual da Paraíba, Av. Baraúnas 351, Bodocongó, Campina Grande, PB, 58430335, Brazil

Corresponding author: Washington LS Vieira (wlsvieira@yahoo.com.br)

Abstract. The distribution of *Sphaenorhynchus prasinus* Bokermann, 1973 in Brazil extends from eastern Bahia to eastern and central Minas Gerais, northern Espírito Santo, with isolated records in Alagoas and Pernambuco. Here, we present a new record of the occurrence of *S. prasinus* and the potential distribution of the species throughout the Brazilian Atlantic Forest. Our record increases the distribution of the species by 944 km from the type locality in the municipality of Ilhéus, state of Bahia and by 80 km from the northernmost record in Igarassu, Pernambuco.

Key words. Amphibia, Bokermann's Lime Treefrog, Brazilian Atlantic Forest, distribution extension, Pernambuco Endemism Center

Vieira WL, Vieira KS, Delfim FR, França FGR, Alves RRN, Montenegro PFG, Pereira Filho GA

(2024) New record of *Sphaenorhynchus prasinus* Bokermann, 1973 (Anura, Hylidae) from northeastern Brazil. *Check List* 20 (6): 1320–1325. <https://doi.org/10.15560/20.6.1320>

INTRODUCTION

The genus *Sphaenorhynchus* Tschudi, 1838 constitutes a monophyletic clade (Pyron and Wiens 2011; Duellman et al. 2016; Faivovich et al. 2018; Araujo-Vieira et al. 2019; Pereira et al. 2022) comprising 15 Neotropical species of small to medium-sized green tree frogs distributed in the Amazon and Orinoco basins of South America, Trinidad and Tobago, Guianas, and eastern Brazil (Duellman et al. 2016; Araujo-Vieira et al. 2019; 2020; Frost 2024). Three species—*S. carneus* (Cope, 1868), *S. dorisae* (Goin, 1957), and *S. lacteus* (Daudin, 1800)—are distributed west of South America in Peru, Ecuador, Colombia, Guyana, and the central Amazon in Brazil (Pereira et al. 2022; Frost 2024). *Sphaenorhynchus lacteus* also occurs in northeastern Brazil with records in the states of Maranhão and Piauí (Caramaschi et al. 2009; Benício et al. 2011), as well as in Trinidad and Tobago (named as *Hyla orophila* by Kenny 1969). *Sphaenorhynchus platycephalus* (Werner, 1894) is known from Serra do Mar and Serra da Mantiqueira in the states of Rio de Janeiro, São Paulo, and Minas Gerais in southeastern Brazil, at elevations above 1,000 m (Araujo-Vieira et al. 2018; Frost 2024). The other species, *S. botocudo* Caramaschi, Almeida & Gasparini, 2009; *S. bromelicola* Bokermann, 1966; *S. cammaeus* Roberto, Araujo-Vieira, Carvalho-e-Silva, & Ávila, 2017; *S. canga* Araujo-Vieira, Lacerda, Pezzuti, Leite, Assis & Cruz, 2015; *S. caramaschii* Toledo, Garcia, Lingnau & Haddad, 2007; *S. mirim* Caramaschi, Almeida & Gasparini, 2009; *S. palustris* Bokermann, 1966; *S. pauloalvini* Bokermann, 1973; *S. planicola* (Lutz & Lutz, 1938); *S. prasinus* Bokermann, 1973, and *S. surdus* (Cochran, 1953), are distributed throughout the Atlantic Forest domain in Brazil and have been recorded from the northern portion of the state of Rio Grande do Sul to the state of Pernambuco (Frost 2024).

Sphaenorhynchus prasinus Bokermann, 1973 is characterized by its medium size (snout–vent length (SVL) 28–31 mm), truncated snout in dorsal and lateral views, dark canthal line which extends from the tip of the snout to the anterior corner of the eye, indistinct tympanum, absence of a dorsolateral white line from the posterior corner of the eye to the sacral region, well-developed and smooth dermal fold on the ventrolateral surface of the forearms and tarsi, dermal fold on the elbow and round calcar appendage, and presence of a white dermal fold with rounded lateral margins in the subcloacal region (Bokermann 1973). The distribution of *S. prasinus* in Brazil extends from eastern Bahia to eastern and central Minas Gerais,



Academic editor: Natan Medeiros Maciel

Received: 2 April 2024

Accepted: 15 November 2024

Published: 26 November 2024

Copyright © The authors. This is an open-access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0)

and northern Espírito Santo, and there are isolated records in Alagoas and Pernambuco (Araujo-Vieira et al. 2016; Pereira et al., 2022; Frost 2024). In this context, we present a new record of the occurrence of *S. prasinus* and the potential distribution of the species throughout the Brazilian Atlantic Forest, especially in the Pernambuco Endemism Center, the most threatened sector of the Brazilian Atlantic Forest.

METHODS

Field research was carried out as part of the herpetofaunistic survey research project in a remnant of Atlantic Forest (Açude de Reis Forest) with 1495 ha belonging to Companhia Usina São João. This remnant forest is located in the rural area of the municipality of Santa Rita, Paraíba state, northeastern Brazil. Specimens of *S. prasinus* were collected through active time-constrained searches on 21.VIII.2023 in an area of a Semideciduous Seasonal Forest at 16–17 m in altitude. This area has lentic water bodies, including an artificial reservoir and temporary pools, and lotic water bodies consisting of small streams, and the remnant forest is surrounded by sugarcane monoculture. The collected specimens were anesthetized and euthanized with a subcutaneous injection of 2% lidocaine following to the guidelines on the Practice of Euthanasia of the Conselho Nacional de Controle de Experimentação Animal – Concea and Conselho Nacional de Biologia – CFBio Ordinance no. 148/2012. Afterwards, specimens were fixed in 10% formaldehyde and stored in 70% ethanol, according to Heyer et al. (1994). but before fixation, we collected muscle and liver tissue samples for storage in 99.6% ethanol. The collected specimens are housed in the Herpetological Collection of the Universidade Federal da Paraíba (CHUFPB). Collection permissions were granted by the Instituto Chico Mendes de Conservação e Biodiversidade (SISBio no. 78910-1) and also by the Animal Ethics Committee of the Universidade Federal da Paraíba (CEUA no. 7355310322).

A review of the geographic distribution of *S. prasinus* was carried out based on information available at the Global Biodiversity Information Facility (GBIF 2023) and in the literature (Feio et al. 1999; Feio and Caramaschi 2002; Juncá 2006; Toledo et al. 2007; Caramaschi et al. 2009; Santos and Moura 2009; Silva et al. 2013; Araujo-Vieira et al. 2015; 2016; Dubeux et al. 2020; Pereira et al. 2022). Based on this information, showing the occurrence of *S. prasinus* (containing 28 records) was produced using QGIS v. 3.32.0. Furthermore, the potential geographic distribution of the species was then carried out using the maximum-entropy method (MaxEnt v. 3.4.4), using only presence records in the model production (Phillip et al. 2006). A set of 19 bioclimatic variables (annual averages 1970–2000) served as input data: Bioclimatic Variations used to Model Production: BIO1 = Annual Mean Temperature; BIO2 = Mean Diurnal Range (Mean of monthly (max. temp–min. temp)); BIO3 = Isothermality (BIO2/BIO7) ($\times 100$); BIO4 = Temperature Seasonality (standard deviation $\times 100$); BIO5 = Max. Temperature of Warmest Month; BIO6 = Min. Temperature of Coldest Month; BIO7 = Temperature Annual Range (BIO5–BIO6); BIO8 = Mean Temperature of Wettest Quarter; BIO9 = Mean Temperature of Driest Quarter; BIO10 = Mean Temperature of Warmest Quarter; BIO11 = Mean Temperature of Coldest Quarter; BIO12 = Annual Precipitation; BIO13 = Precipitation of Wettest Month; BIO14 = Precipitation of Driest Month; BIO15 = Precipitation Seasonality (Coefficient of Variation); BIO16 = Precipitation of Wettest Quarter; BIO17 = Precipitation of Driest Quarter; BIO18 = Precipitation of Warmest Quarter; BIO19 = Precipitation of Coldest Quarter. These variables were obtained from the WorldClim portal (Fick and Hijmans 2023), and they derive from monthly precipitation and temperature values, whose purpose is to produce biologically significant information (Fick and Hijmans 2017).

The set of bioclimatic variables was reduced to two main multivariate geographic axes. The underlying objective was to find linear combinations that produced uncorrelated indices in order of importance, capable of describing the variation of environmental data, thus, reducing the multicollinearity of the variables. The generated rasters then served as the main environmental layer to be associated with the observation data to produce a predictive feature by maximum entropy. We estimated the accuracy of our model by probability through a standard accuracy indicator (ROC/AUC). Probable autocorrelative situations, internal to the observation data, were verified and selected locally (LISA) and on a larger scale (MORAN I); we use “spdep” packages for spatial dependence analysis and the “spatialEco” for additional functions in R v. 4.3.1. The average test AUC for the replicate runs was 0.982, and the standard deviation of 0.012. None of the records were dispersed beyond the likely known distribution limits of the species nor did they exhibit discrepant locality predictions.

All climate layers are in 2.5 arc-minute resolution ($4.625 \times 4.625 = 21.39 \text{ km}^2/\text{pixel}$ at the equator; GFSC/SIIL) at a resolution considered adequate for the data used.


RESULTS

Sphaenorhynchus prasinus Bokermann, 1973

Figures 1, 2

New records. BRAZIL – PARAÍBA · Municipality of Santa Rita, Remnant of Atlantic Forest Açude dos Reis, Companhia Usina São João; 07°08'25.05"S, 035°01'07.43"W; 17 m alt; 21.VIII.2023; G.A.P. Filho, F.G.R.


Figure 1. Specimens of *Sphaenorhynchus prasinus* recorded in the Açude de Reis Atlantic Forest remnant in the municipality of Santa Rita, Paraíba, Brazil. **A.** Young individual (CHUFPB29190). **B.** Adult individual (CHUFPB29191) Photos: Gentil A.P. Filho and Frederico G.R. França.



França leg.; specimens on herbaceous vegetation (grasses) at a height of 60–100 cm above ground, near lentic water body; sex indet; CHUFPB 29190. • same locality; 21.VIII.2023; same collectors; same microhabitat; sex indet; CHUFPB 29191.

Identification. The individuals were identified according to the morphological traits described by Bokermann (1973) and Araujo-Vieira et al. (2016). The specimens presented SVL ranging from 15.58 mm (CHUFPB29190) to 25.97 mm (CHUFPB29191) (Figure 1), truncated snout in dorsal and lateral views, presence of a dark canthal line that extends from the tip of the snout to the anterior corner of the eye, indistinct tympanum, absence of a dorsolateral white line from the posterior corner of the eye to the sacral region, well-developed and smooth dermal fold on the ventrolateral surface of the forearms and tarsi, dermal fold on the elbow, rounded calcar appendage and presence of a white dermal fold with rounded lateral margins on the subcloacal region. The coloration of the dorsum and limbs in life were light green, with round, irregular dark-brown spots and a light-green venter. When preserved in alcohol, the specimens presented a yellowish, pale color on the dorsum and limbs, and a whitish venter. The iris was golden, with many brown reticulations. The dermal ornamentation on the ventrolateral margin of the forearms, tarsus, and subcloacal region was white. The chromatic characterization of both living and preserved *Sphaenorhynchus prasinus* specimens was performed based on the standardization suggested by the colors catalogue for field biologists (Köhler 2012).

Figure 2. A. Geographic distribution map showing records of *Sphaenorhynchus prasinus* in Brazil. Black circles indicate the previously known locations and red star indicate the new record from municipality of Santa Rita, Paraíba, Northeastern Brazil (this study). **B.** Potential geographic distribution of *S. prasinus* in the Brazilian Atlantic Forest. Warmer colors mean higher environmental suitability value. 1 = Santa Rita; 2 = Paudalho; 3 = Igarassu; 4 e 5 = Recife; 6 = São Lourenço da Mata; 7 = Quebrangulo; 8 e 9 = Maceió; 10 = Coruripe; 11 = Mata de São João (Reserva Sapiranga); 12 = Mata de São João (Reserva Legal Fazenda Camurujipe); 13 = Cariranha; 14 = Itagibá; 15 = Ilhéus; 16 = Uruçuca; 17 = Teixeira de Freitas; 18 = Pedro Canário; 19 = Teófilo Otoni; 20 = Nanuque; 21 = Ecoporanga; 22 = Conceição da Barra; 23 = São Mateus; 24 = Almenara; 25 = Linhares; 26 = Aimorés; 27 = São Roque do Canaã; 28 = Marilândia.



DISCUSSION

The northeastern region of Brazil harbors eight species belonging to the genus *Sphaenorhynchus*, with two of them (*S. cammaeus* and *S. prasinus*) occurring in the Pernambuco Endemism Center (Roberto et al. 2017), the most fragmented and threatened region of the Atlantic Forest (Almeida and Souza 2023). *Sphaenorhynchus cammaeus* is known only from the type locality, Pedra Talhada Biological Reserve, municipality of Quebrangulo, state of Alagoas, while *S. prasinus* is widely distributed in the Atlantic Forest, occurring in the states of Minas Gerais, Espírito Santo, Bahia, Alagoas, and Pernambuco (Pereira et al. 2022; Frost 2024).

Although the study of the amphibian and reptile fauna in the Pernambuco Endemism Center has progressively increased in recent years, there is little information on the community ecology and distribution patterns of most species (Vasconcelos et al. 2014; Vieira et al. 2023; Pereira-Filho et al. 2023). The available data on the natural history of *S. prasinus*, as well as other species of the genus, indicate that these tree frogs inhabit degraded open areas, floodplain forests, forest edges, puddles, and temporary lagoons, where males vocalize while on floating vegetation or partially submerged in water (Araujo-Vieira et al. 2016).

Our record increases the distribution of the species by 944 km from the type locality in the municipality of Ilhéus, state of Bahia and by 80 km from the northernmost record in Igarassu, Pernambuco. The potential geographic distribution suggests that the species is widely distributed in the northern portion of the Atlantic Forest from the Doce river basin to the north of the São Francisco River in the Pernambuco Endemism Center. The new occurrence of *S. prasinus* in the state of Paraíba represents the northernmost limit of the distribution of the species and highlights the need for more support for research in the region and the intensification of future studies, which could reveal new records, as well as the existence of new species.

ACKNOWLEDGEMENTS

We thank the Companhia Usina São João for its support and dedication in preserving the Brazilian Atlantic Forest. We are in debt to Mr. Severino Dores Alves da Silva for his support and dedication in protecting the Açude de Reis Forest. We are also thankful to the anonymous reviewers and the academic editor, and to Fapesq and the National Council for Scientific and Technological Development (CNPQ) for funding this study (see below).

ADDITIONAL INFORMATION

Conflict of interest

The authors declare that no competing interests exist

Ethical statement

No ethical statement is reported.

Funding

This study was financially supported by Fapesq and the National Council for Scientific and Technological Development (CNPQ) for researching fellowship number 303972/20238. RRNA thanks the CNPQ for productivity research grant number 307011/2022. WLSV thanks the Fapesq (Edital no. 09/2021 Demanda Universal).

Author contributions

Conceptualization: WLSV, GAPF, FGRF, FRD. Data curation: FRD, WLSV, GAPF, RRNA, KSV, FGRF. Formal analysis: WLSV, KSV, FRD, RRNA, PFGPM, GAPF, FGRF. Funding acquisition: WLSV. Investigation: WLSV, GAPF, FRD, FGRF. Methodology: WLSV, KSV, FRD, RRNA, PFGPM, GAPF, FGRF. Supervision: GAPF, FGRF, WLSV, FRD, KSV. Project administration: GAPF, FGRF, WLSV, FRD. Writing – original draft: WLSV, KSV, GAPF. Writing – review and editing: WLSV, KSV, FRD, RRNA, PFGPM, GAPF, FGRF.

Author ORCID iDs

Washington Luiz Silva Vieira  <https://orcid.org/0000-0002-0673-262X>

Kleber Silva Vieira  <https://orcid.org/0000-0001-7664-6236>

Fagner Ribeiro Delfim  <https://orcid.org/0000-0002-5841-9571>

Frederico Gustavo Rodrigues França  <https://orcid.org/0000-0001-6989-2455>

Rômulo Romeu Nobrega Alves  <https://orcid.org/0000-0001-6824-0797>

Paulo Fernando Guedes Pereira Montenegro  <https://orcid.org/0000-0002-3256-8691>

Gentil Alves Pereira Filho  <https://orcid.org/0000-0002-0164-3898>

Data availability

All data that support the findings of this study are available in the main text.

REFERENCES

- Almeida AM, Souza AF** (2023) Northern Atlantic Forest: Conservation Status and Perspectives. In: Pereira-Filho GA, França FGR, Alves RRN, Vasconcellos A (Eds.) Animal biodiversity and conservation in Brazil's northern Atlantic Forest. Springer, Cham, Switzerland, 7–22. https://doi.org/10.1007/978-3-031-21287-1_2
- Araujo-Vieira K, Lacerda JVA, Pezuti TL, Leite FS, Assis CL, Cruz CAG** (2015) A new species of hatchet-faced treefrog *Sphaenorhynchus* Tschudi (Anura: Hylidae) from Quadrilátero Ferrífero, Minas Gerais, southeastern Brazil. *Zootaxa* 4059: 96–114. <https://doi.org/10.11646/zootaxa.4059.1.5>
- Araujo-Vieira K, Nascimnto FA, Roberto IJ, Tiburcio ICS, Lisboa BS, Ávila RW** (2016) A review of the geographical distribution of *Sphaenorhynchus prasinus* Bokermann, 1973 (Anura: Hylidae) in the state of Alagoas, northeastern Brazil. *Boletim do Museu de Biologia Mello Leitão* 38: 265–274.
- Araujo-Vieira K, Caramaschi U, Grillitsch H, Grant T, Faivovich J** (2018) On the identity of *Sphaenorhynchus platycephalus* (Werner, 1894) (Anura: Hylidae). *South American Journal of Herpetology* 13: 73–84. <https://doi.org/10.2994/sajh-d-17-00053.1>
- Araujo-Vieira K, Blotto BL, Caramaschi U, Haddad CFB, Faivovich J, Grant T** (2019) A total evidence analysis of the phylogeny of hatchet-faced treefrogs (Anura: Hylidae: *Sphaenorhynchus*). *Cladistics* 35: 469–486. <https://doi.org/10.1111/cla.12367>
- Araujo-Vieira K, Luna MC, Caramaschi U, Haddad CFB** (2020) A new genus of lime treefrogs (Anura: Hylidae: Sphaenorhynchini). *Zoologischer Anzeiger* 286 81–89. <https://doi.org/10.1016/j.jcz.2020.04.002>
- Benício R, Silva G, Fonseca M** (2011) Amphibia, Anura, Hylidae, *Sphaenorhynchus lacteus* (Daudin, 1800): first record of the genus and species for the state of Piauí, Brazil. *Check List* 7: 196–197. <https://doi.org/10.15560/7.2.196>
- Bokermann WCA** (1973) Duas novas espécies de *Sphaenorhynchus* da Bahia (Anura, Hylidae). *Revista Brasileira de Biologia* 33: 589–594.
- Caramaschi U, Almeida AP, Gasparini JL** (2009) Description of two new species of *Sphaenorhynchus* (Anura, Hylidae) from the state of Espírito Santo, southeastern Brazil. *Zootaxa* 2115: 34–46. <https://doi.org/10.11646/zootaxa.2115.1.3>
- Dubeux MJM, Gonçalves U, Ramos EB, Melo AH, Silva A, Mott T** (2020) New records of anuran amphibians from the Pedra Talhada biological reserve, a seasonal Semideciduous Atlantic Forest fragment in northeastern Brazil. *Herpetology Notes* 13: 997–1002.
- Duellman WE, Marion AB, Hedges SB** (2016) Phylogenetics, classification, and biogeography of the treefrogs (Amphibia: Anura: Arboranae). *Zootaxa* 4104: 1–109. <https://doi.org/10.11646/zootaxa.4104.1.1>
- Faivovich J, Pereyra MO, Luna MC, Hertz A, Blotto BL, Vásquez-Almazán CR, McCranie JR, Sánchez DA, Baêta D, Araujo-Vieira K, Köhler G, Kubicki B, Campbell JA, Frost DR, Wheeler WC, Haddad CFB** (2018) On the monophyly and relationships of several genera of Hylini (Anura: Hylidae: Hylinae), with comments on recent taxonomic changes in hylids. *South American Journal of Herpetology* 13: 1–32. <http://doi.org/10.2994/sajh-d-17-00115.1>
- Feio RN, Caramaschi U** (2002) Contribuição ao conhecimento da herpetofauna do nordeste do estado de Minas Gerais, Brasil. *Phyllomedusa* 1: 105–111.
- Feio RN, Santos PS, Caramaschi U** (1999) New records of amphibians from Parque Estadual do Rio Doce, state of Minas Gerais, Brazil. *Herpetological Review* 30: 56–57
- Fick SE, Hijmans RJ** (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* 37: 4302–4315. <https://doi.org/10.1002/joc.5086>
- Fick S, Hijmans RJ** (2023). WorldClim version 2.1 climate data for 1970–2000. <https://worldclim.org/data/worldclim21.html>. Accessed on: 2023-09-17.
- Frost DR** (2024) Amphibian Species of the World, version 6.2. <https://amphibiansoftheworld.amnh.org/index.php>. American Museum of Natural History, New York, USA. Accessed on: 2024-10-30.
- GBIF** (Global Biodiversity Information Facility) (2023) <https://www.gbif.org>. Accessed on: 2023-09-06.
- Heyer WR, Donnelly MA, McDiarmid RW, Hayek L, Foster MS** (1994) Measuring and monitoring biological diversity: standard methods for amphibians. Smithsonian Institution Press. Washington DC, USA & London, UK. 364p.
- Juncá FA** (2006) Diversidade e uso de hábitat por anfíbios anuros em duas localidades de Mata Atlântica, no norte do Estado da Bahia. *Biota Neotropica* 6 (2). <https://doi.org/10.1590/s1676-06032006000200018>
- Kenny JS** (1969) The Amphibia of Trinidad. *Studies on the Fauna of Curaçao and other Caribbean Islands* 29 (1): 1–78.
- Köhler G** (2012) Color catalogue for field biologists. Herpeton, Offenbach, Germany, 49 pp.
- Pereira EA, Ceron K, Silva HR, Santana DJ** (2022) The dispersal between Amazonia and Atlantic Forest during the Early Neogene revealed by the biogeography of the treefrog tribe Sphaenorhynchini (Anura, Hylidae). *Ecology and Evolution* 12: e8754. <https://doi.org/10.1002/ece3.8754>
- Pereira-Filho GA, Guedes TB, França RC, Freitas MA, Lourenço-de-Moraes R, Mesquita DO, Nunes PMS, Delfim FR, Dubeux MJM, França FGR** (2023) Composition, species richness, and conservation of the reptiles of the highly threatened northern Brazilian Atlantic Forest. In: Pereira-Filho GA, França FGR, Alves RRN, Vasconcellos A (Eds.) Animal biodiversity and conservation in Brazil's northern Atlantic Forest. Springer, Cham, Switzerland, 169–184. https://doi.org/10.1007/978-3-031-21287-1_11
- Phillip SJ, Anderson RP, Schapire RE** (2006) Maximum entropy modeling of species geographic distributions. *Ecological Modelling* 190: 231–259. <https://doi.org/10.1016/j.ecolmodel.2005.03.026>
- Pyron RA, Wiens JJ** (2011) A large-scale phylogeny of Amphibia including over 2,800 species, and a revised classification of extant frogs, salamanders, and caecilians. *Molecular Phylogenetics and Evolution* 61: 543–583. <https://doi.org/10.1016/j.ympev.2011.06.012>

- Santos EMD, Moura GJBD** (2009) Geographic distribution: *Sphaenorhynchus prasinus*. *Herpetological Review* 40: 362–363.
- Silva ASFL, Siqueira S, Zina J** (2013) Checklist of amphibians in a transitional area between the Caatinga and the Atlantic Forest, centraisouthern Bahia, Brazil. *Check List* 9: 725–732. <https://doi.org/10.15560/9.4.725>
- Toledo LF, Garcia PCA, Iingnau R, Haddad CFB** (2007) A new species of *Sphaenorhynchus* (Anura; Hylidae) from Brazil. *Zootaxa* 1658: 57–68. <https://doi.org/10.11646/zootaxa.1658.1.4>
- Vasconcelos TS, Prado VHM, da Silva FR, Haddad CFB** (2014) Biogeographic distribution patterns and their correlates in the diverse frog fauna of the Atlantic Forest hotspot. *PLoS ONE* 9: e104130. <https://doi.org/10.1371/journal.pone.0104130>
- Vieira WLSV, Moura GJB, Macedo FV, Santana GG, Mott T, Dubeux MJM, Nascimento FAG, Vieira KS** (2023) Species richness, distribution pattern, and conservation of amphibians in the northern portion of the Brazilian Atlantic Forest. In: Pereira-Filho GA, França FGR, Alves RRN, Vasconcellos A (Eds.) *Animal biodiversity and conservation in Brazil's northern Atlantic Forest*. Springer, Cham, Switzerland, 147–167. https://doi.org/10.1007/978-3-031-21287-1_10