

## Corrigendum to Carrasco-Rueda et al. (2021): “Noteworthy records of bats (Mammalia, Chiroptera) from southeastern Peru”

Farah Carrasco-Rueda<sup>1</sup>, Yolanda Alcarraz<sup>2</sup>, Diego J. Zavala<sup>3</sup>, Luiggi Carrasco-Escudero<sup>4,5</sup>, Hugo T. Zamora<sup>3</sup>

<sup>1</sup> Keller Science Action Center, Field Museum of Natural History, Chicago, Illinois, 60613, USA

<sup>2</sup> Centro de Ornitología y Biodiversidad (CORBIDI), Cl Santa Rita de Casia Nro 117 Huertos De San Antonio, Santiago de Surco, 15038, Lima, Peru

<sup>3</sup> Museo de Historia Natural Universidad Nacional de San Agustín, Av Daniel Alcides Carrión S/N, Arequipa, Peru

<sup>4</sup> Programa de Conservación de Murciélagos del Perú (PCMP), Piura, Peru

<sup>5</sup> Facultad de Ciencias, Universidad Nacional de Piura, Urb. Miraflores S/N, Castilla, 20002, Piura, Peru

Corresponding author: Farah Carrasco-Rueda (farahcarrasco@gmail.com)

**Carrasco-Rueda F, Zavala DJ, Alcarraz Y, Carrasco-Escudero L, Zamora HT** (2024) Corrigendum to Noteworthy records of bats (Mammalia, Chiroptera) from southeastern Peru”. *Check List* 20 (3): 586–587. <https://doi.org/10.15560/20.3.586>

Following publication of Carrasco-Rueda et al. (2021), we were made aware of a possible mistake in Table 2 of that publication. The measurements of greatest length of the skull (GLS) were bigger than the greatest length of the skull to inner incisors (GLSI) reported for two of the specimens (CEBIOMAS 0515 and CEBIOMAS 0513) of *Molossus alvarezi*. As indicated in the methods section, the measurement of GLS for the species of the Molossidae family were made following Freeman (1981), however in the case of *Molossus alvarezi* we followed González-Ruiz et al. (2011). We mistakenly did not indicate which authors were followed to measure the GLS in Table 2. Therefore, Table 2 is republished here including GLS measurements following both authors for all Molossidae species, and we include Figure A showing the different measures of the greatest length of the skull.

We would like to acknowledge José L. Poma-Urey from the Museo Noel Kempff Mercado for pointing out to us the error in Table 2.

### REFERENCE

**Carrasco-Rueda F, Zavala DJ, Alcarraz Y, Carrasco-Escudero L, Zamora HT** (2021) Noteworthy records of bats (Mammalia, Chiroptera) from southeastern Peru”. *Check List* 17 (1): 171–180. <https://doi.org/10.15560/17.1.171>



The corrected Table 2 and Figure A appears on the next page.

Editor: Felipe Moreira

Received: 16 April 2024

Accepted: 16 April 2024

Published: 8 May 2024

Copyright © The authors. This is an open-access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0)

**Table 2.** Measurements (mm) of the molossid specimens collected.

Code	<i>Molossus coibensis</i>		<i>Molossus alvarezi</i>		<i>Eumops maurus</i>	<i>Molossops temminckii</i>
	CEBIOMAS 0519	CEBIOMAS 0521	CEBIOMAS 0515	CEBIOMAS 0513	CEBIOMAS 0517	CEBIOMAS 0518
	male	male	female	female	female	male
FA	36.00	35.70	49.00	49.30	54.3	31.30
GLS following Freeman (1981)	15.99	16.06	19.14	19.71	20.30	14.87
GLS following González-Ruiz et al. (2011)	17.42	17.74	20.65	20.85	21.70	15.57
GLSI	16.78	17.24	20.23	20.65	21.51	15.65
CIL	15.78	15.91	18.73	19.48	20.24	14.45
CBL	15.20	15.26	18.17	18.86	19.22	14.04
BB	9.36	9.25	9.68	9.91	9.84	7.97
ROL	6.42	6.37	7.58	7.63	8.72	6.16
ZB	11.23	11.27	11.91	12.45	12.80	9.93
PB	4.10	3.96	3.72	3.92	4.40	4.34
PL	5.95	5.93	6.95	7.37	8.98	7.20
MTRL	6.10	5.99	7.13	7.45	8.22	5.78
M3–M3	4.76	4.79	5.16	5.31	5.71	3.98
LMA	11.74	11.89	13.66	14.48	15.74	10.85
MANDL	6.97	6.88	8.20	8.41	8.93	6.17
C-C	4.71	4.68	5.27	5.57	5.32	4.18

**Figure A.** Skull of *Molossus alvarezi*, specimen CEBIOMAS 0513, showing how to measure the greatest length of the skull (GLS), following different authors, and the general measure of the greatest length of the skull to inner incisive (GLSI).

