

NOTES ON GEOGRAPHIC DISTRIBUTION

Reptilia, Elapidae, *Bungarus niger*: Distribution extension and first record for the state of Uttarakhand, India, with notes on snakebites in the Gori River valley

Emmanuel Theophilus¹

Ashok Captain²

Frank Tillack³

Ulrich Kuch⁴

¹ Village Sarmoli, Post Munsiri 262554, District Pithoragarh, Uttarakhand, India.

² 3/1 Boat Club Road, Pune 411 001, Maharashtra, India.

³ Draesekestraße 8, D-12487 Berlin, Germany.

⁴ Biodiversity and Climate Research Centre.

Senckenberganlage 25, D-60325 Frankfurt am Main, Germany. E-mail: ulrich.kuch@senckenberg.de

Elapid snakes of the genus *Bungarus*, widely known as kraits, are distributed throughout the Oriental faunal region with the exception of the Philippines. Wherever they occur, these nocturnal active predators are among the most dangerous and medically important venomous snakes (Warrell 1999). Thirteen species of krait are currently recognized (Slowinski 1994; Kuch et al. 2005; Kuch and Mebs 2007). While some of these are common and wide-ranging, most krait species are actually rare in collections and have remained largely unknown both ecologically and toxicologically.

The Greater Black Krait, *Bungarus niger*, was described as a new species based on specimens from northeastern India (Dibrugarh and Sadiya in Assam, and Tindharia/Pashok near Darjeeling; Wall 1908), and is distinguished from the sympatric Lesser Black Krait, *Bungarus lividus*, by the state of its vertebral dorsal scales, which are distinctly enlarged in *B. niger* and all other species of *Bungarus*, but not or only feebly enlarged in *B. lividus*, and by a higher number of ventral and subcaudal scales (216-231 and 47-57 in *B. niger* vs. 209-221 and 35-42 in *B. lividus*; Wall 1908; 1909; 1910a; 1910b; 1924). *Bungarus niger* has since been found to be widely distributed in northeastern India (Smith 1943; Tillack and Grossmann 2001; Whitaker and Captain 2004), where it was recorded from several additional localities in West Bengal (Shaw et al. 2000), Sikkim (Wall 1924), Assam (including Cachar and Jalpaiguri districts; Wall 1908; 1910a;

1910b; Inglis et al. 1920; Grosselet et al. 2004), Meghalaya (West Garo Hills; Mathew 1992; 1995), and Arunachal Pradesh (Athreya 2005; Borang et al. 2005; Sanyal and Gayen 2006). The species is also listed for the northeastern Indian state of Nagaland (Dasgupta and Raha 2006; no voucher specimen cited). In addition, *Bungarus niger* was collected in Bhutan (Bauer and Günther 1992), western Nepal (Tillack and Grossmann 2001), and Bangladesh (Khan 2004). Vertical distribution records of *B. niger* have placed this species at elevations from 200-400 m above sea level in Bhutan (Bauer and Günther 1992) and in the district of Cachar, Assam (Grosselet et al. 2004), to 1,370-1,450 m in West Bengal (Wall 1908) and western Nepal (Naudanda, Zone Gandaki; Tillack and Grossmann 2001).

Here we report the first record of *B. niger* from the Indian state of Uttarakhand based on a specimen killed on 19 September 2005 at night by a shopkeeper who encountered the krait at the doorstep of his home at Bungapani (30°57.460' N, 80°18.248' E; 1,007 m), a very small rural marketplace in Dharchula Tehsil (district of Pithoragarh). The preserved specimen (Figures 1 and 2) was deposited in the collection of the Bombay Natural History Society (BNHS 3372). The subadult male has a snout-vent length of 560 mm and a tail length of 120 mm; smooth dorsal scales arranged in 15/15/15 rows with a vertebral row of distinctly enlarged scales; 227 ventrals; an undivided anal scale; approximately 57 undivided subcaudals (54 plus space for three scales in a

NOTES ON GEOGRAPHIC DISTRIBUTION

damaged part); 7/7 supralabials, 3+4/3+4 entering orbit; 1/1 preocular; 2/2 postoculars; and 1+2/1+2 temporals. This record extends the known range of *Bungarus niger* approximately 370 km west from its previous westernmost collecting locality in Nepal (Naudanda, Zone Gandaki; Tillack and Grossmann 2001).

Founded on 9 November 2000, the state of Uttarakhand (until recently also known as state of Uttaranchal) comprises 51,125 km² (28°43' to 31°27' N, 77°34' to 81°02' E) of the northern regions of India that had previously been part of Uttar Pradesh state, bordering Nepal in the east, Tibet in the northeast, and the Indian state of Himachal Pradesh in the northwest (Figure 3). Its territory is largely mountainous and includes some of India's highest peaks as well as the headwaters of two of the subcontinent's great rivers, Ganga and Yamuna. Situated close to the western border of Nepal, Bungapani is located on a dry river-bed about 100 meters from the Gori River. The Gori River basin is an exceptional and biologically highly diverse landscape whose number of listed angiosperm species alone exceeds 2,190 (Mathur et al. 2003; 2005; Theophilus 2003).

This great diversity is due to extreme altitude gradients within a small surface area (over 7,400 m above sea level at Nanda Devi East to 560 m above sea level at the confluence of the Gori and Kali rivers at Jauljibi, within a basin of 120 km length and an average width of 25 km), and due to its location at the transition between the Western and the Central Himalayas and Tibet. Between the settlement where the krait was found and the river lie open fields that are among the widest in the whole Gori River valley and are used for grazing livestock. The location is riparian and sub-tropical, and the dominant forest type of the location is a transition between *Shorea robusta* and *Pinus roxburghii*.

While there are very few trees near the site where the snake was found, tree species in its vicinity comprise *Ficus religiosa* near the settlement, a few scattered *Sapium insigne* along the river-bed, and *Toona ciliata* with a host of epiphytic orchids next to the river. Trees towards the mountain slope include *Pinus roxburghii*, *Macaranga pustulata*, *Engelhardtia spicata*, *Bauhinea*

purpurea, *Ficus auriculata*, *Ficus roxburghii*, *Ougenia dalbergioides*, *Boehmeria rugulosa*, and *Bombax ceiba*. The shrubs and tall herbs include *Boehmeria platyphylla*, *Pyracantha crenulata*, and *Asclepias curassavica*. The synanthropes in the vicinity include *Cassia tora*, *Urtica dioica*, and *Eupatorium* sp. (Osmaston 1927; Polunin and Stainton 1984). The other snake species caught and taxonomically confirmed within a 2 km radius of Bungapani so far are the King Cobra (*Ophiophagus hannah*), the Green Pitviper (*Cryptelytrops septentrionalis*), and two non-venomous species, *Ptyas mucosa* and *Amphiesma stolata*.

Death by snakebite is not uncommon in the area and all along the sub-tropical belt of the Gori River. Black snakes are generally believed to be venomous and feared more than others among local communities, with some recent anecdotal cases reporting people dying from snakebites within two hours. Apart from *Ophiophagus hannah* and *Bungarus niger* that are now known to inhabit this valley, the Common Krait (*Bungarus caeruleus*) has been found in the neighbouring valley west of the Gori River at a similar altitude, but its presence has not been confirmed for the Gori Basin yet. Most snakebites in the Bungapani area in the sub-tropical belt along the Gori River are caused by *Cryptelytrops septentrionalis*, so far apparently without cases of fatal envenoming. The majority of victims of *C. septentrionalis* bites are women cutting grass for livestock fodder during the post-monsoon period.

The wide geographical distribution of *Bungarus niger* along the foothills of the Himalayas and adjacent plains from the easternmost states of India to Uttarakhand in its far west, in combination with its quite extensive vertical distribution, suggests that *B. niger* may be an overlooked cause of snakebite mortality in this region of India, Nepal, and Bhutan. As it is a well-known fact that antivenom raised against the venom of a particular krait species may not be effective against that of another (Warrell et al. 1983; Chanhom et al. 1999), it remains to be demonstrated whether the polyvalent antivenoms that are produced in India (against the venoms of *Bungarus caeruleus*, *Daboia russelii*, *Echis*

NOTES ON GEOGRAPHIC DISTRIBUTION

carinatus, and *Naja naja*; see also Simpson and Norris 2007) are capable of neutralizing *Bungarus niger* venom. Studies on the venom of different

populations of this wide-ranging species and its medical importance in the Himalayan region of India, Nepal, and Bhutan are clearly indicated.



Figure 1. *Bungarus niger* from Bungapani, district of Pithoragarh, state of Uttarakhand, India (BNHS 3372); dorsal view. Photo by A. Captain.



Figure 2. *Bungarus niger* from Bungapani, district of Pithoragarh, state of Uttarakhand, India (BNHS 3372); ventral view. Photo by A. Captain.

NOTES ON GEOGRAPHIC DISTRIBUTION



Figure 3. Map indicating the geographical position of *Bungarus niger* records in the western Himalayas. 1 (red star), new record from Bungapani, Dharchula Tehsil, district of Pithoragarh, state of Uttarakhand, India. 2 (red square), Naudanda, district of Kaski, Zone Gandaki, Nepal, previously the westernmost record of this species.

Acknowledgements

We are grateful to Manohar Maher (Bungapani, India) for collecting the dead *Bungarus niger* and bringing it from Bungapani to Munsiri for identification, and to C. J. McCarthy and E. N. Arnold (The Natural History Museum, London, United Kingdom), A. Hängi (*Naturhistorisches Museum Basel*, Basel, Switzerland), F. Tiedemann and R. Gemel (*Naturhistorisches Museum Wien*, Vienna, Austria), V. B. Giri (Bombay Natural History Society, Mumbai, India), and R. Günther (*Museum für Naturkunde der Humboldt-Universität zu Berlin*, Berlin, Germany), for the loan of specimens under their care and assistance during research visits. M. Barts (Berlin, Germany) kindly assisted in the preparation of the map. Research on *B. niger* by UK was funded by Wellcome Trust Research Grant GR079027MA to J. B. Harris (Principal Investigator), M. A. Faiz, U. Kuch, and D. A. Warrell, and by the LOEWE Programme of the State Government of Hessen.

Literature cited

- Athreya, R. 2005. Herpetofauna of W. Arunachal Pradesh - checklist. Version 13 March 2005. Electronic database accessible at <http://www.clsp.jhu.edu/people/zak/ramana/wapListHerp.htm>. Captured on 22 March 2005.
- Bauer, A. and R. Günther. 1992. A preliminary report on the reptile fauna of the Kingdom of Bhutan with the description of a new species of scincid lizard (Reptilia: Scincidae). *Asiatic Herpetological Research* 4: 23-36.
- Borang, A., B. B. Bhatt, S. B. Chaudhury, A. Borkotoki, and P. T. Bhutia. 2005. Checklist of the snakes of Arunachal Pradesh, Northeast India. *Journal of the Bombay Natural History Society* 102 (1): 19-26.
- Chanhome, L., N. Wongtongkam, O. Khaw, N. Pakmanee, T. Omori-Satoh, and V. Sitprija. 1999. Genus specific neutralization of *Bungarus* snake venom by Thai Red Cross banded krait antivenom. *Journal of Natural Toxins* 8: 135-140.

NOTES ON GEOGRAPHIC DISTRIBUTION

- Dasgupta, G. and S. Raha. 2006. Reptilia; p. 433-460 In J. R. B. Alfred (ed.). Fauna of Nagaland. Calcutta: Zoological Survey of India State Fauna Series.
- Grosselet, O., M. Vauché, A. Gupta, and A. Susmita. 2004. *Bungarus niger* Wall, 1908 (Reptilia: Serpentes: Elapidae): extension of range to Cachar District, Assam, India. Russian Journal of Herpetology 11(1): 10-11.
- Inglis, C. M., W. L. Travers, H. V. O' Donel, and E. O. Shebbeare. 1920. A tentative list of the vertebrates of the Jalpaiguri District, Bengal. Part III. Journal of the Bombay Natural History Society 27(1): 151-162.
- Khan, M. A. R. 2004. Checklist of the herpetofauna of Bangladesh. Cobra 57: 1-31.
- Kuch, U. and D. Mebs. 2007. The identity of the Javan Krait, *Bungarus javanicus* Kopstein, 1932: evidence from mitochondrial and nuclear DNA sequence analyses and morphology. Zootaxa 1426: 1-26.
- Kuch, U., D. Kizirian, Q. T. Nguyen, R. Lawson, M. A. Donnelly, and D. Mebs. 2005. A new species of krait (Squamata: Elapidae) from the Red River system of northern Vietnam. Copeia 2005(4): 817-832.
- Leviton, A. E., R. H. Gibbs Jr., E. Heal, and C. E. Dawson. 1985. Standards in herpetology and ichthyology: Part I. Standard symbolic codes for institutional resource collections in herpetology and ichthyology. Copeia 1985(4): 802-832.
- Mathew, R. 1992. Additional records of snakes from northeast India, with four new records. Records of the Zoological Survey of India 91(3-4): 287-292.
- Mathew, R. 1995. Reptilia; p. 379-454 In A. K. Gosh (ed.). Fauna of Meghalaya. Part 1. (Vertebrates). Calcutta. Zoological Survey of India State Fauna Series.
- Mathur, V.B., R. Raza, P. Lal, M. K. Agarwal, and S. Singait. 2003. Biodiversity characterization in Middle Goriganga Valley, Askot Wildlife Sanctuary, Uttarakhand State. Technical report submitted to Indian Institute of Remote Sensing, Dehradun. Electronic database accessible at <http://www.wii.gov.in/publications/researchreports/2005/diversity&rarityinfloralwh.pdf>. Captured on 7 December 2006.
- Mathur, V. B., K. J. Gaston, and R. H. Raza. 2005. Diversity and rarity in floral and avifaunal assemblages in the Western Himalaya: a study of patterns and mechanisms to devise viable biodiversity conservation strategies. Electronic database accessible at <http://www.wii.gov.in/publications/researchreports/2200/diversity&rarityinfloralwh.pdf>. Captured on 7 December 2006.
- Osmaston, A. E. 1927. A Forest Flora for Kumaon. Allahabad: Government Press, United Provinces. xxxiv + 605 p. [reprint 1978 by Bishen Singh Mahendra Pal Singh, Dehra Dun].
- Polunin, O. and A. Stainton. 1984. Flowers of the Himalaya. Delhi: Oxford University Press. 580 p.
- Sanyal, D. P. and N. C. Gayen. 2006. Reptilia; p. 247-284 In J. R. B. Alfred (ed.). Fauna of Arunachal Pradesh. Part 1. Calcutta: Zoological Survey of India State Fauna Series.
- Shaw, G. E., E. O. Shebbeare, and P. E. Barker. 2000. The Snakes of Sikkim and Bengal. Delhi: Asiatic Publ. House. vi+125 p.
- Simpson, I. D. and R. L. Norris. 2007. Snakes of medical importance in India: Is the concept of the "Big 4" still relevant and useful? Wilderness and Environmental Medicine 18: 2-9.
- Slowinski, J. B. 1994. A phylogenetic analysis of *Bungarus* (Elapidae) based on morphological characters. Journal of Herpetology 28(4): 440-446.
- Smith, M. A. 1943. The Fauna of British India, Ceylon, and Burma, Including the Whole of the Indo-Chinese Sub-region. Reptilia and Amphibia. Vol. III-Serpentes. London: Taylor & Francis. xii + 583 p.
- Theophilus, E. 2003. Angiosperms; p. 91-113 In Foundation for Ecological Security (ed.). A Biodiversity Log and Strategy Input Document for the Gori River Basin, Western Himalayan Ecoregion, Uttarakhand. A Sub State Process under the National Biodiversity Strategy and Action Plan of India. Munsiri, India.
- Tillack, F. and W. Grossmann. 2001. Ein neuer Nachweis zur Schlangenfauna Nepals: *Bungarus niger* Wall, 1908 (Reptilia: Serpentes: Elapidae). Sauria 23(1): 3-9.
- Wall, F. 1908. A popular treatise on the common Indian snakes. Part VIII. Journal of the Bombay Natural History Society 18(4): 711-735.
- Wall, F. 1909. Notes on snakes from the neighbourhood of Darjeeling. Journal of the Bombay Natural History Society 19(2): 337-357.
- Wall, F. 1910a. Notes on snakes collected in upper Assam. Part II. Journal of the Bombay Natural History Society 19(4): 825-845.
- Wall, F. 1910b. Notes on snakes collected in the Jalpaiguri District. Journal of the Bombay Natural History Society 19(4): 897-900.
- Wall, F. 1924. A hand-list of the snakes of the Indian Empire. Part IV. Journal of the Bombay Natural History Society 30(1): 12-24.
- Warrell, D. A. 1999. WHO/SEARO guidelines for the clinical management of snake bites in the Southeast Asian region. Southeast Asian Journal of Tropical Medicine and Public Health 30(Suppl. 1): 1-85.
- Warrell, D. A., S. Looareesuwan, N. J. White, R. D. G. Theakston, M. J. Warrell, W. Kosakarn, and H. A. Reid. 1983. Severe neurotoxic envenoming by the Malayan krait *Bungarus candidus* (Linnaeus): response to antivenom and anticholinesterase. British Medical Journal 286: 678-680.
- Whitaker, R. and A. Captain. 2004. Snakes of India. The Field Guide. Chennai. Draco Books. xiv+479 p.

NOTES ON GEOGRAPHIC DISTRIBUTION

Received April 2008
Accepted September 2008
Published online October 2008

Appendix 1

Voucher list [museum acronyms follow Leviton et al. (1985)]: BHUTAN: no locality (NMBA 22737); Phuntsholing (NMBA 22736). INDIA: West Bengal: Darjeeling (NMW 27738.2); district of Darjeeling (BMNH 1937.4.3.3); Uttarakhand: district of Pithoragarh: Bungapani (BNHS 3372). NEPAL: Zone Gandaki: district of Kaski: Naudanda (ZMB 69998).