

NOTES ON GEOGRAPHIC DISTRIBUTION

Cnidaria, Hydrozoa, Anthoathecata, Pandeidae, *Stomotoca atra*: Distribution Extension

Sérgio N. Stampar¹
Guilherme Kodja²

¹Departamento de Zoologia, IB-USP, Universidade de São Paulo, Rua do Matão, travessa 14, nº 101, Cidade Universitária, CEP 05508-900, São Paulo, SP, Brazil. E-mail: stampar@usp.br

²Instituto Laje Viva, Rua Joaquim Floriano 466, sala 2207, Itaim, CEP 04534-002, São Paulo, SP, Brazil.

The genus *Stomotoca* L. Agassiz, 1862 is characterized by medusae with two opposite marginal tentacles and a stomach on a broad peduncle, extending beyond bell margin (Kramp 1961). The genus has only four species: *Stomotoca atra* L. Agassiz 1862, *S. geogrinae* Hadzi 1913, *S. mira* Fewkes, 1883 and *S. pietschi* Martin, 1975 (Bouillon and Boero 2000; van der Land et al. 2001). Some of these species have their sessile polyps known and their life cycles described (Larson 1982; Boero and Bouillon 1989; Boero et al. 1991). A number of works was performed about the bodily processes responses of these species (e.g. Lenhoff 1964; Mackie 1975; Mackie and Singla 1975; Schwab 1977).

The species *Stomatoca atra* was normally reported to Pacific waters (Kramp 1961; Wrobel and Mills 1998). However, there was recorded to Papua New Guinea (Boero and Bouillon 1989) and to Indian Ocean (Navas 1971; Vannucci and Navas 1973; Navas-Pereira and Vannucci 1991).

In this note we present records of two specimens of *Stomotoca atra* found in the water column close to the Laje de Santos (24°15'48"S, 46°12'00"W), a parcel ca. 40 km off Santos (State of São Paulo, Brazil) (Figure 1) on January 10th, 2006 (Figure 2) and December 23th, 2006 (Figure 3). Laje de Santos is a State Marine Conservation Park under the supervision of the NGO Instituto Laje Viva. The photographs (Figure 2 and 3) were made underwater using digital resources.




Figure 1. Map showing the location of Laje de Santos, State of São Paulo, Brazil.

The jellyfish (Figure 2 and 3) photographed in Brazilian waters greatly resemble the description of *Stomotoca atra* by Agassiz (1862): umbrella bell-shaped, mouth not crenulated, gonads simple folds, numerous rudimentary tentacles. The arrangement of tentacles and long manubrium with conspicuous gonads in regular transverse folds, besides the high cone bell shape, clearly characterize this species.




Figure 2. Alive specimen of *Stomotoca atra* L. Agassiz, 1862, photographed underwater in the Laje de Santos. Scale: 4 cm.

NOTES ON GEOGRAPHIC DISTRIBUTION



Figure 3. *Stomotoca atra* L. Agassiz, 1862 swimming in the Laje de Santos. Scale: 3 cm.

A new record of a given species is always an interesting issue. However, when this record occurs in another ocean and different environments (NE and tropical Pacific waters according to Boero & Bouillon 1989; Wrobel and Mills 1998) it becomes more important, because distributional patterns and biology of the species can be emphasized. The distribution of the species *Stomotoca atra*, with this new record, is very hard to understand, because the ocean currents are not coincident with this allocation (see more in Valiela 1995). That can be hypothesized as the first evidence of a cryptogenic or a report of an extremely rare species in Brazilian waters. Bouillon et al. 1991 discuss about this distribution (NE Pacific, Papua New Guinea and Indian Ocean) and the taxonomy of *S. atra*. They conclude that is really one species and it is necessary more studies about life cycle and other aspects to understand something about the zoogeography of this species.

This work again emphasizes the importance of photographic records in ocean areas. The Laje de Santos is a sanctuary in which sampling is very much restricted. In this way, high quality photography is very useful to record species and the identification is performed without sampling in a protected area.

Acknowledgments

We thank Dr. A. C. Morandini (NUPEM/UFRJ) for criticism in the text and Mr. Armando de Luca Junior for providing a photo. We are deeply indebted to Dr. F. L. da Silveira (IB-USP) for providing laboratory facilities and for valuable comments on the manuscript. We thanks anonymous referee for important remarks on the text. We are grateful to Instituto Laje Viva and Departamento de Zoologia (IB-USP) by supporting this work.

Literature cited

- Agassiz, L. 1862. Contributions to the natural history of the United States of America. IV. Second monograph, In five parts, Acalephs in general, Ctenophorae, Discophorae, Hydroidae, homologues of the Radiata. Boston: Little, Brown & Co. viii, 380, pls XX-XXXV p.
- Boero, F. and J. Bouillon. 1989. The life cycles of *Octotiarra russelli* and *Stomotoca atra* (Cnidaria, Anthomedusae, Pandeidae). *Zoologica Scripta* 18(1):1-7.
- Boero, F., J. Bouillon, and C. Gravili. 1991. The life cycle of *Hydrichthys mirus* (Cnidaria:Hydrozoa: Anthomedusae: Pandeidae). *Zoological Journal of the Linnean Society* 101: 189-199.
- Bouillon, J., F. Boero, and G. Seghers. 1991. Notes additionnelles sur les méduses de papouasie Nouvelle-Guinée (Hydrozoa, Cnidaria) IV. *Cahiers de Biologie Marine* 32(3): 387-411.
- Kramp, P. L. 1961. Synopsis of the medusae of the world. *Journal of the Marine Biological Association of the United Kingdom* 40:7-469.
- Larson, R. J. 1982. Life history of the hydromedusa *Stomotoca pterophylla* Haeckel and its ichthyoparasitic hydroid. *In* The Atlantic Bar-

NOTES ON GEOGRAPHIC DISTRIBUTION

- rier Reef Ecosystem at Currie Bow Cay, Belize, I: Structure and Communities (K. Rützler and I. G. Macintyre, ed.). Smithsonian Contributions to Marine Science 12: 433–439.
- Lenhoff, H. M. 1964. Reversible inhibition of swimming in *Stomatoca atra* by mesogleal extracts of some other medusae. Biological Bulletin 126(1): 115-120.
- Mackie, G. O. 1975. Neurobiology of *Stomatoca*. II. Pacemakers and conduction pathways. Journal of Neurobiology 6 (4): 357-378.
- Mackie, G. O. and C. L. Singla. 1975. Neurobiology of *Stomatoca*. I. Action Systems Journal of Neurobiology 6 (4): 339-356.
- Navas, D. 1971. New records of Hydromedusae from the Indian Ocean. Contribuições do Instituto Oceanográfico de São Paulo, ser. Oceanografia Biológica 22: 1-33.
- Navas-Pereira, D. and M. Vannucci. 1991. The hydromedusae and water masses of the Indian Ocean. Boletim do Instituto Oceanográfico 39(1): 25-60.
- Schwab, W. E. 1977. Electrical responses of *Stomatoca atra* induced by an extract of *Aequorea aequorea*. Nature 267: 701-702.
- Valiela, I. 1995. Marine Ecological Processes. 2nd ed. New York: Springer-Verlag. 686 p.
- van der Land, J., W. Vervoort, S. D. Cairns, and P. Schuchert. 2001. Hydrozoa. In European register of marine species: a check-list of the marine species in Europe and a bibliography of guides to their identification (M. J. Costello, C. S. Emblow and R. White, ed.). Collection Patrimoines Naturels 50: 112-120.
- Vannucci, M. and D. Navas. 1973. Distribution of hydromedusae in the Indian Ocean. In: Zeitschel, B. (ed.), The Biology of the Indian Ocean. pp. 273-282. Springer-Verlag, Berlin, Heidelberg, New York.
- Wrobel, D. and C. E. Mills. 1998. Pacific coast pelagic invertebrates. A guide to the common gelatinous animals. Monterey: Sea Challengers and Monterey Bay Aquarium. i-iv, 1-108 p.

Received January 2007

Accepted March 2007

Published online March 2007