

## LISTS OF SPECIES

### Fish, Ribeirão do Feijão Basin, São Carlos, São Paulo, Brazil.

Oscar Akio Shibatta  
Ângela Teresa Silva-Souza

*Departamento de Biologia Animal e Vegetal. Centro de Ciências Biológicas, Universidade Estadual de Londrina. CEP 86051-990, Londrina, PR, Brazil. E-mail: shibatta@uel.br.*

**Abstract:** The Ribeirão do Feijão Basin is the main water supplier for the municipality of São Carlos, in the state of São Paulo, Brazil. Field work was carried out from February to November 2005, using sieves, casting nets, and drag nets. A total of 7,286 specimens of fish were collected, belonging to 30 species, 13 families and six orders. The richest order was Characiformes with 14 species, followed by Siluriformes with seven.

#### Introduction

The municipality of São Carlos is located in the central region of the state of São Paulo, Brazil, at the coordinates 21°30' to 22°30' S, and 47°30' to 48°30' W. With an area of 1,132 km<sup>2</sup>, São Carlos has a rich hydrological network and is situated among the headways of many streams (São Paulo 2008).

The Ribeirão do Feijão, an affluent of Jacaré-Guaçu River that drains to Tietê River is included in the upper Paraná River Basin. It constitutes the main source of water supply for the city of São Carlos. Therefore, its entire basin requires constant monitoring with respect to the quality of its waters and biotic diversity.

Fishes constitute one of the most representative groups for the study of environmental quality, because they provide an overall perspective of the aquatic environment. They not only occupy various levels of the trophic chain but are also relatively easy to identify in critical situations (Karr 1981).

Since the ichthyofauna of the Ribeirão do Feijão Basin still unknown, a survey of its species was carried out, as support for subsequent studies that are being conducted. A list of the species resulting from this survey is presented in this paper.

#### Material and methods

For the survey, besides the main water course of Ribeirão do Feijão, five tributaries (Córrego Quebra Canela, Córrego dos Macacos, Ribeirão Laranja Azeda, Córrego São José, and Córrego do Monjolinho) were sampled, totaling 12 stations of

collections. A detailed description and geographic localization of each of these stations were presented by Silva-Souza et al. (2005). The fish samplings were performed seasonally, in the period of February to November 2005, using drag nets with a mesh size of 2.0 mm between adjacent knots, casting nets with a mesh size of 2.0 cm between opposing knots and sieves with a mesh size of 2.0 mm between adjacent knots. For each stretch, a portion of 50 m was delineated, and each end was blocked by a screen with mesh size of 2.0 mm between adjacent knots, to prevent the escape of specimens. The Jaccard index (Hauer and Lamberti 1996) was used to compare the community found in this study with that described in other works.

Voucher specimens were deposited in the fish collection of *Museu de Zoologia da Universidade Estadual de Londrina* (MZUEL), state of Paraná, Brazil. Collects were authorized by IBAMA (DIREN 083/2003).

#### Results and discussion

A total of 7,286 specimens were collected, 398 in Córrego Quebra Canela, 4,508 in Ribeirão do Feijão, 523 in Córrego dos Macacos, 160 in Ribeirão Laranja Azeda, 1,057 in Córrego São José, and 50 in Córrego do Monjolinho. The order with largest number of species was Characiformes (14 species, five families), followed by Siluriformes (seven species, four families), Perciformes (five species, one family), Cyprinodontiformes (two species, one family), and Gymnotiformes and Synbranchiformes (only one species each), totaling 30 species, 13 families in six orders (Table 1).

## LISTS OF SPECIES

**Table 1.** Fish captured in the Ribeirão do Feijão Basin, municipality of São Carlos, February to November 2005.

---

### Order Characiformes

#### Family Parodontidae

*Apareiodon piracicabae* (Eigenmann, 1907)

*Parodon nasus* Kner, 1859

#### Family Characidae

*Astyanax altiparanae* Garutti & Britski, 2000

*Astyanax* sp. aff. *A. fasciatus* (Cuvier, 1819)

*Astyanax paranae* Eigenmann, 1914

*Bryconamericus iheringii* (Boulenger, 1887)

*Hyphessobrycon anisitsi* (Eigenmann, 1907)

*Hyphessobrycon eques* (Steindachner, 1882)

*Oligosarcus pintoii* Campos, 1945

*Serrapinnus notomelas* (Eigenmann, 1915)

#### Family Curimatidae

*Cyphocharax modestus* (Fernández-Yepez, 1948)

#### Family Crenuchidae

*Characidium gomesi* Travassos, 1956

*Characidium* cf. *zebra* Eigenmann, 1909

#### Family Erythrinidae

*Hoplias malabaricus* (Bloch, 1794)

---

### Order Siluriformes

#### Family Callichthyidae

*Callichthys callichthys* (Linnaeus, 1758)

*Corydoras aeneus* (Gill, 1858)

#### Family Heptapteridae

*Cetopsorhamdia iheringi* Schubart & Gomes, 1959

*Rhamdia quelen* (Quoy & Gaimard, 1824)

#### Family Loricariidae

*Corumbataia cuestae* Britski, 1997

*Hypostomus ancistroides* (Ihering, 1911)

#### Family Trichomycteridae

*Trichomycterus* sp.

---

### Order Gymnotiformes

#### Family Gymnotidae

*Gymnotus* spp. 1-4

---

### Order Perciformes

#### Family Cichlidae

*Australoheros facetum* (Jenyns, 1842)

*Cichlasoma paranaense* Kullander, 1983

*Geophagus brasiliensis* (Quoy & Gaimard, 1824)

*Oreochromis niloticus* (Linnaeus, 1758)

*Tilapia rendalli* (Boulenger, 1897)

---

### Order Cyprinodontiformes

#### Family Poeciliidae

*Phalloceros* sp.

*Poecilia reticulata* Peters, 1859

---

### Order Synbranchiformes

#### Family Synbranchidae

*Synbranchus marmoratus* Bloch, 1785

---

## LISTS OF SPECIES

This is a common tendency for rivers of the Neotropical region, as observed by Lowe-McConnell (1999) and Perin et al. (2007). A comparison based on the Jaccard similarity index of the results of the present study with those found by Gomiero and Braga (2006) for Corumbataí River and Jacaré-Pepira River, showed a similarity of only 30 %. These authors noted that the similarity between the rivers basins studied was low, due to the variability of the habitats and to environmental conditions. The same can be used to explain the differences between the present study and that from Gomiero and Braga (2006), since no endemism was observed.

Probably due to the fact that the environments sampled in the Ribeirão do Feijão Basin were typically streams, only small-sized species were collected, contrary to what Gomiero and Braga (2006) collected, which includes anostomids and characids (Salmininae) with a standard length over 15.0 cm.

In the Ribeirão do Feijão Basin, species were not identified for two genera: *Trichomycterus*, which could be a new species, and *Gymnotus*, which showed a color polymorphism that could signify that these latter fishes are different syntopic species. Color pattern was utilized by Castro et al.

(2003; 2004) to identify the species of *Gymnotus* from the streams of Paranapanema and Rio Grande basins, in which four and three species (*G. cf. carapo*, *G. cf. inaequilabiatus*, *G. cf. sylvius*, and *Gymnotus* sp.) were identified, respectively. A recent study (Graça and Pavanelli 2007), however, demonstrated even the presence of *G. pantanal* and *G. paraguensis* in the upper Paraná River Basin. Considering the four color pattern of the specimens of *Gymnotus* collected, maybe they represent four different cryptic species. Therefore, more detailed studies such as a hole, using anatomy, cytogenetic, and molecular biology approaches are needed to help resolve this question.

The introduced species *Tilapia rendalli*, *Oreochromis niloticus*, and *Poecilia reticulata* were also found in the Ribeirão do Feijão Basin. It is possible that the first two originated from fish farming stations and the last due to its discard by aquarium enthusiasts.

In general, the Ribeirão do Feijão Basin appears to be poor in the number of species, when compared to the richness recorded for streams of the Paranapanema, with 52 species (Castro et al. 2003), and Rio Grande, with 64 species (Castro et al. 2004).

---

### Acknowledgements

We thank the *Instituto Internacional de Ecologia de São Carlos* for logistic support and fish collections; Edson Santana and Wanner Galves for their help with the fish samplings; and FAPESP (process#98/10924-3) for financial support. We are also grateful to Dr. A. Leyva for language revising and Leonardo F. S. Ingenito, Francisco Langeani Neto, and Vinícius Abilhoa for valuable comments that improved the text.

### Literature cited

- Castro, R. M. C., L. Casatti, H. F. Santos, K. M. Ferreira, A. C. Ribeiro, R. C. Benine, G. Z. P. Dardis, A. L. A. Melo, R. Stopiglia, T. X. Abreu, F. A. Bockmann, M. Carvalho, F. Z. Gilbran, and F. C. T. Lima. 2003. Estrutura e composição da ictiofauna de riachos do rio Paranapanema, sudeste e sul do Brasil. *Biota Neotropica* 3(1): 1-31.
- Castro, R. M. C., L. Casatti, H. F. Santos, A. L. A. Melo, L. S. F. Martins, K. M. Ferreira, F. Z. Gilbran, R. C. Benine, M. Carvalho, A. C. Ribeiro, T. X. Abreu, F. A. Bockmann, G. Z. Pelição, R. Stopiglia, and F. Langeani. 2004. Estrutura e composição da ictiofauna de riachos da bacia do rio Grande no estado de São Paulo, sudeste do Brasil. *Biota Neotropica* 4(1): 1-39.
- Gomiero, L. M. and F. M. S. Braga. 2006. Fish, Corumbataí and Jacaré-Pepira river basins, São Paulo State, Brazil. *Check List* 2(1): 57-66.
- Graça, W. J. and C. S. Pavanelli. 2007. Peixes da planície de inundação do alto rio Paraná e áreas adjacentes. Maringá, EDUEM. 241 p.
- Hauer, F. R. and G. A. Lambert. 1996. *Methods in Stream Ecology*. San Diego, Academic Press. 674 p.

## LISTS OF SPECIES

Karr, J. R. 1981. Assessment of biotic integrity using fish communities. *Fisheries* 6(6): 21-27.

Lowe-McConnell, R. H. 1999. Estudos ecológicos de comunidades de peixes tropicais. São Paulo, EDUSP. 534 p.

Perin, L., O. A. Shibatta, and P. S. Bernarde. 2007. Fish, Machado River Basin, Cacoal urban area, state of Rondônia, Brazil. *Check List* 3(2): 94-97.

São Paulo, Governo do Estado. 2008. Fundação Sistema Estadual de Análise de Dados – SEADE. Database accessible at <http://www.seade.gov.br/>. Captured on February 2008.

Silva-Souza, Â. T., D. F. Rosim, O. A. Shibatta, T. Matsumura-Tundisi, and J. G. Tundisi. 2005. Análise inicial da qualidade ambiental da bacia do córrego do Feijão utilizando peixes como indicadores (Políticas Públicas FAPESP processo 98/10924 – 2ª fase), p. 29-36, *In Rede Mercocidades* (org.). São Carlos, Anais da II Mostra de Ciências e Tecnologia em Políticas Públicas Municipais.

Received November 2007

Accepted January 2008

Published online March 2008