

## NOTES ON GEOGRAPHIC DISTRIBUTION

### Reptilia, Squamata, Colubridae, *Chironius exoletus*: distribution extension, new state record.

Fernando J. M. Rojas-Runjaic<sup>1</sup>  
Edwin E. Infante Rivero<sup>2</sup>

<sup>1</sup>Museo de Historia Natural La Salle, Sección de Herpetología. Apartado Postal 1930, Caracas 1010-A, Venezuela. E-mail: rojas\_runjaic@yahoo.com


<sup>2</sup>Museo de Biología Universidad del Zulia. Apartado Postal 526, Maracaibo 4011, Venezuela.

The genus *Chironius* Fitzinger currently comprises 13 species (Dixon et al. 1993) of diurnal terrestrial-arboreal colubrids, of medium to large size (over 200 cm), cylindrical or compressed body, with 10 or 12 dorsal scales rows to the midbody, dorsal scales largest, smooth or keeled, with apical pits, head differentiated from neck, eyes prominent, pupil round, and moderate to very long tail. This genus is distributed from Nicaragua, Central America, San Vincent island in the Caribbean sea, to southern Brazil and Argentina in South America (Peters and Orejas-Miranda 1970; Dixon et al. 1993). In Venezuela the genus is widely distributed from

sea level to 2500 m (Lancini 1982), and six species currently are known: *C. carinatus* (Linnaeus, 1758), *C. exoletus* (Linnaeus, 1758), *C. fuscus* (Linnaeus, 1758), *C. monticola* Roze, 1952, *C. multiventris* Schmidt & Walker, 1943, and *C. scurrulus* (Wagler, 1824) (Lancini and Kornacker 1986; Dixon et al. 1993; La Marca 1997).

*Chironius exoletus* has a very wide distribution through South America (northeastern Argentina, eastern and Amazonian Brazil, Bolivia, eastern Peru and Ecuador, Colombia, Venezuela, Guyana, Surinam and French Guiana), to Panama and Costa Rica in Central America. In Venezuela it has been only reported from the states of Amazonas, Bolivar and Delta Amacuro (McDiarmid and Paolillo 1988; Dixon et al. 1993; Rivas and La Marca 2001), all south of Orinoco River.

During a field herpetological survey carried out in “El Progreso” farm, upper basin of Socuy river, western slope Sierra de Perijá, Jesús Enrique Lossada municipality, Zulia state, northwestern Venezuela (c. 10°47' N, 72°26' W, c. 800 m a.s.l.) (Figure 1), between 23 to 26 February 2006, we found a specimen of *C. exoletus* (Figure 2). The snake was active at 11:00 h dwelling on herbaceous vegetation at the edge of primary forest.



**Figure 1.** White dot shown “El Progreso” farm, first locality record form Zulia state (Venezuela), northward in the country and South America from *Chironius exoletus*.

## NOTES ON GEOGRAPHIC DISTRIBUTION



**Figure 2.** Juvenile male of *Chironius exoletus* (MHNLS 17931). Photo by F. Rojas-Runjaic.

Our specimen is a juvenile male (hemipenes partially everted) snout-vent length 293 mm; tail length 185 mm; nasals divided; internasals, prefrontals, and parietals paired; frontal single; loreal present; preoculars 1/1, postoculars 2/2, temporals 1+2, supralabials 9(5+6), infralabials 11(6+7), two pairs of chin shields; infralabials one through six in contact with anterior chin shields, six and seven in contact with posterior chin shields; 12-12-8 scale row formula, dorsal scales with apical pits and only paravertebral rows keeled; ventral scales 150, subcaudals 160 divided, anal plate divided, and 28 maxillary teeth. These characters allow us to identify this specimen as *C. exoletus*, and to discriminate this from *C. carinatus* (a sympatric species and very common in the Sierra de Perijá), and the all others species of the genus.

This is the first record for the species from Zulia state in Venezuela, northward in the country and South America. The nearest locality reported is Arboledas, Norte de Santander, Colombia, c. 380

km NE (Dixon et al. 1993), and c. 1200 km NW from Monte Duida, Amazonas state, Venezuela (McDiarmid and Paolillo 1988). The voucher is deposited in the herpetological collection of the Museo de Historia Natural La Salle (Caracas, Venezuela) with number MHNLS 17931.

### Acknowledgments

We thank Cesar Barrio-Amoros, Marco Natera and Gilson Rivas for suggestions and revision of an early draft of the manuscript. The necessary collecting permit (N° 01-03-03-1146) was granted by Ministerio del Ambiente y de los Recursos Naturales (MARN) to Fernando Rojas-Runjaic.

### Literature cited

- Dixon, J. R., J. A. Wiest and J. M. Cei. 1993. Revision of the Neotropical snake genus *Chironius* Fitzinger (Serpentes, Colubridae). *Monografie dei Museo Regionale di Scienze Naturali di Torino* 13: 279 p.
- La Marca, E. 1997. Lista actualizada de los reptiles de Venezuela. Pp. 123-142. In E. La Marca (ed.), *Vertebrados actuales y fósiles de Venezuela. Serie Catálogo Zoológico de Venezuela. Vol. I. Mérida. Museo de Ciencia y Tecnología de Mérida.*
- Lancini, A. R. 1982. Serpientes de Venezuela. Distribución geográfica y altitudinal de géneros de serpientes en Venezuela. *Memórias do Instituto Butantan* 46: 95-103.
- Lancini A. R and P. M. Kornacker. 1986. *Die Schlangen von Venezuela.* Caracas. Editorial Armitano. 381 p.
- McDiarmid, R. W. and A. Paolillo. 1988. Herpetological collections: Cerro de la Neblina. Pp. 667-670. In C. Brewer-Carías (ed.), *Cerro de la Neblina. Resultados de la expedición 1983-1987.* Caracas. Fudeci. 922 p.
- Peters, J. A. and B. Orejas-Miranda. 1970. *Catalogue of the Neotropical Squamata. Part I. Snakes.* United States National Museum Bulletin 297: viii + 347 p.
- Rivas F., G. and E. La Marca. 2001. Geographic Distribution. *Chironius scurrulus.* *Herpetological Review* 32: 58-59.

Received September 2006

Accepted November 2006

Published online November 2006