

NOTES ON GEOGRAPHIC DISTRIBUTION

**Serpentes, Dipsadidae, Echinantherini, *Taeniophallus poecilopogon* (Cope, 1863):  
Rediscovery in Argentina**

Santiago Javier Nenda<sup>1</sup>  
Diego Omar Di Pietro<sup>2</sup>

<sup>1</sup> *División Herpetología, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”.*  
Ángel Gallardo 470. CP 1405. Ciudad Autónoma de Buenos Aires, Argentina. E-mail: santiagojnenda@yahoo.com.ar

<sup>2</sup> *Museo de La Plata, División Zoología Vertebrados, Sección Herpetología.*  
Paseo del Bosque s/n. CP 1900. La Plata, Buenos Aires, Argentina.

The colubrid snake genus *Taeniophallus* Cope, 1895 has a complex taxonomic history (Myers 1974; Di-Bernardo and Lema 1987; Myers and Cadle 1994; Cacivio et al. 1999; Schargel et al. 2005; Santos-Jr et al. 2008; Zaher et al. 2009) and currently two species are considered to occur in Argentina: *Taeniophallus occipitalis* and *Taeniophallus poecilopogon* (Koslowsky 1898; Serié 1936; Peters and Orejas-Miranda 1970; Myers 1974; Williams and Francini 1991; Di-Bernardo 1992; Cei 1993; Scrocchi et al. 2000; Cabrera 2001; Giraudo 2001; Giraudo and Scrocchi 2002).


*Taeniophallus poecilopogon* has the southernmost distribution of the genus, inhabiting open savannahs in Pampean and Espinal phytogeographic provinces (Di-Bernardo and Lema 1987; Cei 1993; Cabrera 2001; Giraudo 2001; Giraudo and Scrocchi 2002), and is known from southern Brazil, Paraguay, Argentina, and Uruguay (Koslowsky 1898; Devincenzi 1925; Peters and Orejas-Miranda 1970; Myers 1974; Di-Bernardo and Lema 1987; Di-Bernardo 1992; Cei 1993; Gonzáles and Hamilton 1998; Achaval-Elena 2001; Giraudo 2001; Giraudo and Scrocchi 2002; Achaval and Olmos 2003; Carreira et al. 2005).

In Argentina *Taeniophallus poecilopogon* has been reported in the literature for Buenos Aires, Corrientes, Entre Ríos, and Misiones provinces (Koslowsky 1898; Serié 1936; Freiberg 1939; Abalos and Mischis 1975; Williams and Francini 1991; Cei 1993; Scrocchi et al. 2000; Giraudo 2001; Giraudo and Scrocchi 2002).

Despite these multiple reports, specimens of *Taeniophallus poecilopogon* from Argentina are poorly represented in herpetological collections, and only two specimens are known until now. One of them was housed in the Herpetological Collection of the *Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”*, MACN without numeration, and with “Argentina” as the only locality data. Unfortunately ongoing curatorial efforts on this collection failed to find this specimen and it should be considered lost for the time being. Probably the records from Corrientes and Misiones are based on this specimen (Giraudo 2001; Giraudo and Scrocchi 2002), and the record for Entre Ríos (Freiberg 1939) is not supported by specimens in museum collections. The other voucher specimen is deposited in the *Museo di Torino*, MZUT R-819, from Las Flores, Buenos Aires province (Figure 1A), illustrated by Jan and Sordelli (1866, Volume I, Livraison 16, Plate 1, Figure 3) and by Cei (1993, Figure 99: C, D; p. 669), and represents a historical record of *Taeniophallus poecilopogon* in Argentina.

In the present contribution we report and describe the second specimen of *Taeniophallus poecilopogon* from Argentina with precise locality and complete data: MACN 39227, an adult roadkill, ca. 273 mm body length and ca. 86 mm tail length, the tail represents 24 percent of total length, found in the *Ruta Provincial* # 11, 10 km south from Magdalena city, Partido de Magdalena, Buenos Aires province, Argentina (35°08'32" S, 57°24'43" W) (Figure 1B); collected on August 28<sup>th</sup> 2008.

## NOTES ON GEOGRAPHIC DISTRIBUTION



**Figure 1.** Distribution map of *Taeniophallus poecilopogon* in Argentina. A. (\*) locality of MZUT R-819. B. Detailed map of the locality of MACN 39227, arrow points to collection place in Ruta Provincial N° 11, Partido de Magdalena, Buenos Aires province, Argentina.

The specimen MACN 39227 has 17 smooth dorsal scales without reduction and without apical pits, 165 ventrals (the highest number reported for the species), cloacal plate divided, 57 subcaudals, 7(3+4)/7(3+4) supralabials, 8(4)/8(4) infralabials, one loreal in contact with second supralabial only, 1+2/1+2 oculars, and 1+2/1+2 temporal scales.

The color pattern of the specimen MACN 39227 is longitudinally striped; a conspicuous white line over the canthus rostralis reaches the temporal region, where it is interrupted by dark coloration on the top of the head; dark head cap extends around and below the corner of the mouth; heavily dappled labials and chins; edges of the venter and sides of the body and tail are dark to about the middle on the ca. 4<sup>th</sup> dorsal scales; a bright red coloration

appears on the ca. 20<sup>th</sup> ventral scale and extends into the tail.

The locality is situated 110 km (airline) west from Montevideo Department, Uruguay (the southernmost locality in Uruguay, separated from Magdalena by the Río de la Plata); and 182 km northeast from Las Flores, Buenos Aires province, Argentina. The place where MACN 39227 has been found belongs to the Talares (*Celtis tala*) district in the Espinal phytogeographic province (Cabrera 1968), and is a rural zone with cattle raising and woodlands of *Eucalyptus* spp., located in the margins of the biosphere reserve “Parque Costero del Sur” (Figure 2). In this reserve, where remnants of particular woodlands of *Celtis tala* and the endemic *Phytolacca tetramera* are mixed with natural grasslands and wetlands, the species is very likely to be present.

## NOTES ON GEOGRAPHIC DISTRIBUTION



**Figure 2.** Habitat of *T. poecilopogon* near “Parque Costero del Sur” biosphere reserve, Partido de Magdalena, Buenos Aires province, Argentina.

---

### Acknowledgments

We are grateful to G. Koppers for English editing, to J. Williams, C. Borteiro, F. Kolenc, for share with us their valuable bibliography, F. Agnolin, and S. Lucero for helping at the field, and to D. Barrasso, R. Cajade, J. Faivovich, B. Blotto, A. Sheinkman for permanent support during this study.

---

### Literature cited

- Abalos, J. W. and C. C. Mischis. 1975. Elenco sistemático de los ofidios argentinos. *Boletín Academia Nacional de Ciencias de Córdoba* 51(1-2): 55-76.
- Achaval-Elena, F. 2001. Actualización sistemática y mapas de distribución de los reptiles del Uruguay. *Smithsonian Herpetological Information Service* 129: 1-37.
- Achaval, F. and A. Olmos. 2003. *Anfibios y Reptiles del Uruguay*. 2da Edición corregida y ampliada. Montevideo: Graphis Impresora. 136 p.
- Cabrera, A. L. 1968. Vegetación de la Provincia de Buenos Aires; p. 101-122 *In* A. L. Cabrera (ed.). *Flora de la Provincia de Buenos Aires*. Buenos Aires: INTA.
- Cabrera, M. R. 2001. Composición comparativa de la fauna de serpientes (Reptilia, Squamata) de cuatro provincias biogeográficas del Dominio Chaqueño. *Revista del Museo Argentino de Ciencias Naturales n.s.* 3(2): 109-118.
- Cacivio, P. M., G. J. Scrocchi, and A. Giraudo. 1999. *Echianthera occipitalis* (Serpentes: Colubridae) nuevos datos de escamación, hemipenes y distribución. *Cuadernos de Herpetología* 13(1-2): 81-87.
- Carreira, S., M. Meneghel, and F. Achaval. 2005. *Reptiles de Uruguay*. Montevideo: D.I.R.A.C. Facultad de Ciencias, Universidad de la República. 639 p.
- Cei, J. M. 1993. Reptiles del noroeste, nordeste y este de la Argentina. *Herpetofauna de las selvas subtropicales, puna y pampas*. Monografía XIV. Torino: Museo Regionale di Scienze Naturali di Torino. 929 p.
- Devincenzi, G. J. 1925. *Fauna Erpetológica del Uruguay*. *Anales del Museo de Historia Natural de Montevideo, Serie II* 2: 1-65.
- Di-Bernardo, M. 1992. Revalidation of the genus *Echianthera* Cope, 1894, and its conceptual amplification (Serpentes, Colubridae). *Comunicações do Museu de Ciências da PUCRS, Série Zoologia* 5(13): 225-256.

## NOTES ON GEOGRAPHIC DISTRIBUTION

- Di-Bernardo, M. and T. de Lema. 1987. O género *Rhadinea* Cope, 1863, no Brasil meridional. I. *Rhadinea poecilopogon* Cope, 1863 (Serpentes, Colubridae). Acta Biologica Leopoldensia 9(2): 203-224.
- Freiberg, M. A. 1939. Enumeración sistemática de los reptiles de Entre Ríos y lista de los ejemplares que los representan en el Museo de Entre Ríos. Memorias del Museo de Entre Ríos, Zoología 11: 1-28.
- Giraud, A. R. 2001. Serpientes de la selva paranaense y del chaco húmedo. Taxonomía, biogeografía y conservación. Buenos Aires: L.O.L.A. 285 p.
- Giraud, A. R. and G. J. Scrocchi. 2002. Argentinian snakes: an annotated checklist. Smithsonian Herpetological Information Service 132: 1-53.
- González, E. M. and S. Hamilton. 1998. Un nuevo registro de *Echinanthera poecilopogon* (Cope, 1863) (Reptilia: Squamata) en Uruguay. Cuadernos de herpetología 12(2): 37.
- Jan, G. and F. Sordelli. 1866. Iconographie générale des ophiidiens. Atlas. Volume I, Livraison 16. Paris: J. B. Bailliére et Fills.
- Koslowsky, J. 1898. Enumeración sistemática y distribución geográfica de los reptiles argentinos. Revista del Museo de La Plata 8: 161-200.
- Myers, C. W. 1974. The systematics of *Rhadinaea* (Colubridae), a genus of new world snakes. Bulletin of the American Museum of Natural History 153: 1-262.
- Myers, C. W. and J. E. Cadle. 1994. A New Genus for South American Snakes Related to *Rhadinaea obtusa* Cope (Colubridae) and Resurrection of *Taeniophallus* Cope for the "*Rhadinaea*" brevirostris Group. American Museum Novitates (3102): 1-33.
- Peters, J. A. and B. R. Orejas-Miranda. 1970. Catalogue of Neotropical Squamata. Part I. Snakes. Washington D.C.: Bulletin of U.S. Natural Museum. 347 p.
- Santos-Jr, A. P., M. Di-Bernardo, and T. de Lema. 2008. New species of the *Taeniophallus occipitalis* group (Serpentes, Colubridae) from eastern Amazonia, Brasil. Journal of Herpetology 42(3): 419-426.
- Schargel, W. E., G. Rivas-Fuenmayor, and C. W. Myers. 2005. An enigmatic new Snake from Cloud Forest of the Peínsula de Paria, Venezuela (Colubridae: Genus *Taeniophallus*?). American Museum Novitates (3484): 1-22.
- Scrocchi, G. J., I. Aguer, V. Arzamendia, P. Cacivio, H. Carcacha, M. Chiaraviglio, A. Giraud, S. Kretzschmar, G. Leynaud, M. S. López, L. Rey, T. Waller, and J. Williams. 2000. Categorización de las serpientes de Argentina; p. 75-93 In E. O. Lavilla, G. J. Scrocchi, and E. Richard (ed.). Categorización de los Anfibios y Reptiles de Argentina. San Miguel de Tucumán: Asociación Herpetológica Argentina.
- Serié, P. 1936. Nueva enumeración sistemática de los ofidios argentinos. Obra Cincuentenario, Revista del Museo de La Plata 2: 33-61.
- Williams, J. D. and F. Francini. 1991. A checklist of the Argentine snakes. Torino, Bollettino del Museo Regionale di Scienze Naturali di Torino 9(1): 55-90.
- Zaher, H., F. Gobbi-Grazziotin, J. E. Cadle, R. W. Murphy, J. C. de Moura-Leite, and S. L. Bonatto. 2009. Molecular phylogeny of advanced snakes (Serpentes, Caenophidia) with an emphasis on South American Xenodontines a revised classification and descriptions of new taxa. Papéis Avulsos de Zoologia 49(11): 115-153.

Received June 2009

Accepted July 2009

Published online August 2009